Sermorelin 5mg

QUICK GUIDE

The original approved GHRH analog ♥



WHAT IS SERMORELIN?

Sermorelin is the first 29 amino acids of growth-hormone-releasing hormone, the active part. It was FDA-approved as Geref for children with GH deficiency, then discontinued in 2008 for business reasons, not safety. It now comes only from compounders.

Strongest evidence: A · Approved use

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 5 mg vial.

This yields 2 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog ladder 200→500 mcg nightly (grade E)

WEEK	DOSE (Once nightly)	UNITS (U-100 SYRINGE)
Weeks 1–4	200 mcg	10 units
Weeks 5–8	300 mcg	15 units
Weeks 9–12	400 mcg	20 units
Weeks 13–16	500 mcg	25 units

Cycle: catalog: 12–16 weeks on, 6–8 weeks off.

POTENTIAL BENEFITS

- **History:** Approved 1997 (Geref); withdrawn 2008, not for safety. A · Approved use
- **Natural GH pulses:** Raises the body's own GH with feedback intact. A · Approved use
- **Adult use:** No adult outcome trials. N/E · Not enough evidence

POTENTIAL SIDE EFFECTS

- Injection-site reactions
- Flushing, headache (GHRH class)
- Fluid retention possible at higher GH/IGF-1

Unit math: every unit value the catalog prints matches (5 mg vial 2 mg/mL; 10 mg vial 4 mg/mL).

Active cancer (raises GH/IGF-1)

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade A: Approved use

For educational use only.

PEPPERSDVO