Survodutide 10mg

QUICK GUIDE

GLP-1 / glucagon dual agonist for weight and liver ♥



WHAT IS SURVODUTIDE?

Survodutide is a once-weekly dual GLP-1 and glucagon receptor agonist from Boehringer Ingelheim and Zealand. It reached about 16.6% weight loss in phase 3, and it's being developed for fatty liver disease (MASH) as well as obesity.

Strongest evidence: **B · Human trials**

RECONSTITUTION

Add 1.5 mL bacteriostatic water to the 10 mg vial.

This yields 6.7 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 1.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog ladder 1.6→4.8 mg, 0.8 mg steps (grade E)

WEEK	DOSE (Once weekly, same day each week)	UNITS (U-100 SYRINGE)
Weeks 1–4	1.6 mg	24 units
Weeks 5–8	2.4 mg	36 units
Weeks 9–12	3.2 mg	48 units
Weeks 13–16	4 mg	60 units
Week 17 onward	4.8 mg	72 units

Phase 3 maintenance doses are 3.6 and 6.0 mg; edit the last step to match.

POTENTIAL BENEFITS

- **Weight loss:** Up to −16.6% at 76 weeks (SYNCHRONIZE-1). B · Human trials
- **Liver fat:** Up to −63.1% liver fat in a substudy. B · Human trials
- **Type 2 diabetes:** HbA1c down up to 1.21% (SYNCHRONIZE-2). B · Human trials
- **Tolerability:** 25% stopped for adverse events in phase 2. B · Human trials

POTENTIAL SIDE EFFECTS

- Nausea, vomiting, diarrhea, constipation (dose-related)
- Phase 2: 25% stopped for adverse events (mostly GI)
- Heart-rate rise possible (glucagon class)

Catalog maximum 6.4 mg is above the highest phase 3 dose (6.0 mg).

The catalog's '19% at 4.8 mg' is the actual-treatment analysis. The planned-treatment result was −14.9%.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO