Tesamorelin / Ipamorelin 10/3 10mg
QUICK GUIDE

Catalog blend: untested as a combination ♥



WHAT IS IT?

Tesamorelin / Ipamorelin 10/3 is a pre-made catalog blend of Tesamorelin, Ipamorelin. Each ingredient keeps its own evidence grade (Tesamorelin A, Ipamorelin B). No study of this combination was found.

Strongest evidence: E · Community / vendor

RECONSTITUTION

Add 3 mL acetic acid + BAC water to the 10 mg vial.

This yields 3.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 3 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Per catalog protocol)	UNITS (U-100 SYRINGE)
Weeks 1–6	770 mcg	23.1 units
Weeks 7–12	1.53 mg	45.9 units

Cycle: Catalog cycle.

POTENTIAL BENEFITS

- **Tesamorelin:** 10 mg per vial · grade A — A · Approved use
- **Ipamorelin:** 3 mg per vial · grade B — B · Human trials

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

Units only. Computed: 23 units = 0.77 mg tesamorelin + 0.23 mg ipamorelin; 46 units = 1.53 mg + 0.46 mg.

The label tesamorelin dose is 1.28–1.4 mg daily.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.


PEPPERSDVO