Tesamorelin 10mg

QUICK GUIDE

The approved GHRH analog for visceral fat ♥



WHAT IS TESAMORELIN?

Tesamorelin is a stabilized form of growth-hormone-releasing hormone. It makes the pituitary release more of its own GH in natural pulses. It's FDA-approved (Egrifta) to cut abdominal visceral fat in people with HIV-associated lipodystrophy.

Strongest evidence: A · Approved use

RECONSTITUTION

Add 3 mL acetic acid + BAC water to the 10 mg vial.

This yields 3.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 3 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Label: 1.28 mg daily (Egrifta WR, grade A)

WEEK	DOSE (Once daily (label); 5 days a week)	UNITS (U-100 SYRINGE)
Weeks 1–16	1.28 mg	38.4 units

Cycle: catalog: 12–16 weeks on, 4–8 weeks off (84–112 / 28–56 days). The label has no cycling.

POTENTIAL BENEFITS

- **Visceral fat:** −18% vs +2% placebo at 26 weeks (label, study 1). A · Approved use
- **Natural GH pulses:** Raises the body's own GH, keeping feedback intact. A · Approved use
- **Glucose:** HbA1c ≥6.5% in 5% vs 1% on placebo. A · Approved use
- **Fluid retention:** Edema, joint pain, carpal tunnel. A · Approved use

POTENTIAL SIDE EFFECTS

- Injection-site reactions
- Arthralgia, peripheral edema, carpal tunnel (fluid retention)
- Raised glucose / HbA1c
- Raised IGF-1

Catalog doses 1–2 mg, 5 days a week. The label is 1.28 mg (WR) or 1.4 mg (SV) daily; the trials used 2 mg daily.

Unit math: every unit value the catalog prints matches. Diluent totals: 10 mg in 3 mL, 20 mg in 5 mL.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade A: Approved use

For educational use only.


PEPPERSDVO