BPC-157 / TB-500 ("Wolverine") 20mg
QUICK GUIDE

Catalog blend: untested as a combination ♥



WHAT IS IT?

BPC-157 / TB-500 ("Wolverine") is a pre-made catalog blend of BPC-157, TB-500. Each ingredient keeps its own evidence grade (BPC-157 C, TB-500 D). No study of this combination was found.

Strongest evidence: E · Community / vendor

RECONSTITUTION

Add 2 mL bacteriostatic water to the 20 mg vial.

This yields 10 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Per catalog protocol)	UNITS (U-100 SYRINGE)
Weeks 1–2	600 mcg	6 units
Weeks 3–8	800 mcg	8 units

Cycle: Catalog cycle.

POTENTIAL BENEFITS

- **BPC-157:** 10 mg? per vial · grade C — C · Human observational
- **TB-500:** 10 mg? per vial · grade D — D · Animal / lab

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

Ambiguous strength: "10MG" with 6 units = 600 mcg only works if the vial holds 10 mg of EACH (20 mg total). If it's 10 mg total, every stated mcg dose is double the real one. The per-mL ingredient table didn't survive capture.

This sheet assumes 20 mg total (10 + 10) because that's what makes the catalog's numbers consistent.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.


PEPPERSDVO