



Engineering Mathematics

Detailed Solutions

LINEAR ALGEBRA

(EC, EE, IN, CS, ME, CE, CH, PI)

Himanshu Agarwal & Anish Saha

PrepFusion™

Where Concept Meets Clarity!

Preface

Welcome to PrepFusion™!

This book works, in full, every question GATE has actually asked in Linear Algebra (EC/EE/IN/CS/ME/CE/CH/PI), gathered from the original papers and arranged unit by unit and year by year. Nothing is paraphrased or reconstructed: every question appears as it was set, and every solution is taken step by step rather than asserted.

Each solution carries the same identifier as the question it answers in the companion questions volume, so you can move between practice and explanation without a lookup table. Where the examining authority revised an answer or awarded marks to all (MTA), that is noted with the question it affects.

Used end to end, the book shows you what the paper rewards — which topics repeat, how the framing has shifted over the years, and where your preparation still has gaps.

Meet your Authors

Himanshu Agarwal

- AIR 27 — GATE (ECE)
- AIR 45 — GATE (IN)
- 5+ years of teaching experience



Anish Saha

- AIR 19 — GATE (IN)
- AIR 66 — GATE (EE)
- 3+ years in teaching & the VLSI industry



“We built this book the way every aspirant deserves one to be built — organized strictly by unit and by year, so that every hour of practice maps directly onto how GATE actually tests you.”

— Himanshu Agarwal & Anish Saha

How to Read the Question Numbers

Every question in this book carries a three-part identifier instead of a plain serial number:

Q3.24.7	3	Unit (chapter) number — Unit III of this book
	24	GATE exam year — 2024 (two digits; 98 = 1998)
	7	Question number within that unit and year

So **Q3.24.7** is the 7th question that GATE 2024 contributed to Unit III. Numbers are therefore not sequential through the book — they tell you the unit and the exam year at a glance. The same identifier is used in the questions book, the solutions book and the answer keys, so you can jump between them without a lookup table. In the PDF bookmarks panel, expand a unit to see every question ID and click one to go straight to it.

What the Bubbles Mean

Every question ends with three empty circles for you to shade in yourself:

☐

☐

☐

☒

☐

☐

 Easy

☒

☒

☐

 Medium

☒

☒

☒

 Hard

The companion solutions volume marks the same question with *our* recommended difficulty, using the identical bubbles. If you think a rating should be different, tell us directly — report it under the **Study Hub** section at pyq.prepfusion.in.

How to Read the Branch Snapshot

A Maths unit can span up to eight branches (EC, EE, IN, CS, ME, CE, CH, PI), each weighted very differently in the real exam. Right after every Unit divider, a **Branch Snapshot** page lists every branch that unit covers, in the same order the solutions themselves follow, each in its own compact box:

Questions	the real total for that branch.
MCQ, MSQ, NAT	that same total, split by question type.
1M, 2M	how many of those questions carry 1 or 2 marks.
Descriptive	questions worth more than 2 marks, or whose marks value isn't recorded.
Avg Weightage	the average marks contributed per real GATE sitting, not a raw year-by-year total — a year GATE ran multiple real sessions (a genuine forenoon/afternoon split, not the same paper's Set A/B/C/D booklet variants) is divided across those sessions first, so a heavily multi-session year cannot silently outweigh a single-session one. The ~ marks it as a rounded figure, not an exact one.
Range	the lowest and highest that same per-sitting value has been, with the year each occurred.

Avg Weightage and Range are computed from **2010 onward** only: GATE dropped from a 150-mark to a 100-mark paper in 2009, but that year itself was a one-off transition (60 questions, not the 65-question format that became standard from 2010 on) — so 2010 is the first year fully comparable to modern papers, and marks-based figures from earlier years are left out of these two figures specifically (every other count on this page still includes every year). They also depend on knowing which GATE years ran more than one real sitting for a given branch — our own research into this is thorough for some branches and still a work in progress for others. Treat both figures as a **beta**, best-effort estimate: directionally useful, not exact.

MSQ questions specifically weren't introduced until 2021 — an older-year-heavy branch showing **MSQ: 0** reflects that real exam history, not a data gap.

Worked example

Branch Snapshot*					
Branch: EC	Questions: 45	MCQ: 30	MSQ: 3	NAT: 12	
1M: 20	2M: 22	Descriptive: 3	Avg Weightage ~4M	Range: 2M(2015)–6M(2021)	

45 real EC questions on this unit; on average EC has drawn about 4 marks per real GATE sitting since 2010, ranging from as low as 2 (in 2015) to as high as 6 (in 2021).


A quick way to use this: a high **Avg Weightage** for your branch points to a consistently high-yield topic worth prioritizing; a low one is largely someone else's problem. A wide **Range** usually means the topic's importance has shifted over the years, so it is worth checking *which* years drove that before deciding how much weight to give it today.

Important Links & Resources



Visit our Website

prepfusion.in



Solutions, practice & report

pyq.prepfusion.in



Subscribe: English Channel

youtube.com/@PrepFusion_GATE



Subscribe: Hindi Channel

youtube.com/@prepfusion-hindi



Join our Telegram group

t.me/All_About_Learning



Follow us on LinkedIn

linkedin.com/company/prepfusionee



Subscribe: Himanshu Agarwal

youtube.com/@HimanshuAgarwal_



Subscribe: Anish Saha

youtube.com/@AnishSaha_

Scan any code with your phone camera, or tap the link beneath it. pyq.prepfusion.in is also where you can read this book's worked solutions online — and in its **Study Hub** section, report an error in any question or request a video solution for it.

Contributors

This book exists because of the people below, who built, reviewed and typeset its content.

Design & Layout Architect — Nawras Ahamed.

Technical Reviewers — Ayush Prakash Singh, Abhishek Bhattacharjee, Shujaut Yaseen, Yuva Kishore Gottapu, Manepalli Pavan Vignesh Kumar, Palak Agarwal, Devmalya Ghatak, Kasi Viswanath A, Suresh Ravanam, Aabhas Anuraag Suman and Piyush Raj.

Content Developers — A S Srimithra, Chinthada Lakshmi Prasad, Gantasala Ravi Teja, Guduru Vinayak Varma, Harshaditya Punera, Katikala Sarath, Pranitha Musunuru, Safdar Nazim, Sai Mahalakshmi Janjanam, Shaila Daniels, Sharvari Patwardhan, Sumedha Ghosh, Vinay K Gowda, Aayushi Ghosh, Chevuri Siva Naga Sai Abhi Chandar, Govind Sharma, Kasina Sai Surya Vamsi, Machanpally Karthik Kumar, Seeku Vamsi Krishna and Suprio Dutta.

Contents

Unit	Topic	Pages
I.	Linear Algebra	1
I.	EC	3
II.	EE	32
III.	IN	72
IV.	CS	95
V.	ME	135
VI.	CE	166
VII.	CH	194
VIII.	PI	209

SOLUTIONS — UNIT

I

Linear Algebra

Topics

1. Introduction to Linear Algebra
2. Determinants, Minors and Cofactors
3. Types of Matrices
4. Simultaneous Operations on Matrices
5. Transpose, Trace, Inverse and Adjoint of a Matrix
6. System of Linear Equations - Existence and Uniqueness
7. Rank of a Matrix
8. Solution of Non-Homogeneous Equations
9. Solution of Homogeneous Equations
10. Eigen Values and Eigen Vectors
11. Properties of Eigen Values and Eigen Vectors
12. Vector Space, Subspace and Linear Span
13. Basis Vectors and Null Space

Linear Algebra

Engineering Mathematics

Branches: EC,EE,IN,CS,ME,CE,CH,PI

Branch Snapshots*

EC — Electronics & Communication Engineering

Questions: 51 **MCQ:** 34 **MSQ:** 3 **NAT:** 14
1M: 32 **2M:** 19 **Desc:** 0 **Avg Weightage** ~3M **Range:** 1M(2026)–3M(2024)

EE — Electrical Engineering

Questions: 55 **MCQ:** 41 **MSQ:** 3 **NAT:** 11
1M: 30 **2M:** 24 **Desc:** 1 **Avg Weightage** ~3M **Range:** 1M(2023)–8M(2026)

IN — Instrumentation Engineering

Questions: 36 **MCQ:** 27 **MSQ:** 2 **NAT:** 7
1M: 24 **2M:** 12 **Desc:** 0 **Avg Weightage** ~3M **Range:** 1M(2019)–3M(2023)

CS — Computer Science & IT

Questions: 57 **MCQ:** 34 **MSQ:** 6 **NAT:** 17
1M: 32 **2M:** 24 **Desc:** 1 **Avg Weightage** ~3M **Range:** 1M(2023)–4M(2026)

ME — Mechanical Engineering

Questions: 45 **MCQ:** 37 **MSQ:** 2 **NAT:** 6
1M: 30 **2M:** 15 **Desc:** 0 **Avg Weightage** ~2M **Range:** 1M(2025)–4M(2022)

CE — Civil Engineering

Questions: 47 **MCQ:** 35 **MSQ:** 6 **NAT:** 6
1M: 31 **2M:** 16 **Desc:** 0 **Avg Weightage** ~2M **Range:** 1M(2019)–4M(2023)

CH — Chemical Engineering

Questions: 22 **MCQ:** 14 **MSQ:** 1 **NAT:** 7
1M: 13 **2M:** 8 **Desc:** 1 **Avg Weightage** ~2M **Range:** 1M(2026)–3M(2021)

PI — Production & Industrial Engineering

Questions: 25 **MCQ:** 22 **MSQ:** 0 **NAT:** 3
1M: 16 **2M:** 9 **Desc:** 0 **Avg Weightage** ~2M **Range:** 1M(2025)–4M(2023)

*See preface for how Avg Weightage and Range are calculated.

SOLUTIONS — Linear Algebra



EC

Electronics & Communication Engineering

PrepFusion

Q1.26.1 **MSQ** **2026** **1M** **Ans: B,D** ☒ ☐ ☐

Given matrix $M = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 3 & 0 \\ -1 & a & b \end{bmatrix}$.

$$\det(M) = 2(3b - 0) - 1(b - 0) + 1(a - (-3))$$

$$\det(M) = 6b - b + a + 3$$

$$\det(M) = 5b + a + 3$$

For the matrix to be non-singular, we must have:

$$5b + a + 3 \neq 0$$

Now we check each option:

• **Option (A):** $a = -\frac{1}{2}, b = -\frac{1}{2}$

$$5\left(-\frac{1}{2}\right) + \left(-\frac{1}{2}\right) + 3 = -\frac{5}{2} - \frac{1}{2} + 3 = -3 + 3 = 0$$

This makes $\det(M) = 0$, so it is **FALSE**.

• **Option (B):** $a = \frac{1}{2}, b = \frac{1}{2}$

$$5\left(\frac{1}{2}\right) + \frac{1}{2} + 3 = 3 + 3 = 6 \neq 0$$

This is **TRUE**.

• **Option (C):** $a = -3, b = 0$

$$5(0) + (-3) + 3 = 0$$

This makes $\det(M) = 0$, so it is **FALSE**.

• **Option (D):** $a = \frac{1}{2}, b = -3$

$$5(-3) + \frac{1}{2} + 3 = -15 + 3.5 = -11.5 \neq 0$$

This is **TRUE**.

$$(B) a = \frac{1}{2} \text{ and } b = \frac{1}{2}, (D) a = \frac{1}{2} \text{ and } b = -3$$

Key Points

- A matrix M is non-singular if and only if $\det(M) \neq 0$.
- The determinant of a 3×3 matrix $A = [a_{ij}]$ can be found using Laplace expansion: $\det(A) = \sum_{j=1}^3 (-1)^{i+j} a_{ij} M_{ij}$.

Mistakes to be avoided

- Ensure correct sign convention (+ - +) when expanding the determinant.
- Do not confuse the condition $\det(M) \neq 0$ (invertible) with $\det(M) = 0$ (singular).

Q1.25.1 MCQ 2025 1M Ans: A ● ○ ○

Given the matrix in upper triangular form:

$$A = \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & \alpha & \beta \\ 0 & 0 & 0 & \gamma \end{bmatrix}$$

The rank of the matrix, $\rho(A)$, is determined by the number of non-zero rows in its row echelon form. Let's analyze the given cases:

- (i) Let $\alpha = 0, \beta = \gamma \neq 0$:

$$A \sim \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & 0 & \beta \\ 0 & 0 & 0 & \gamma \end{bmatrix} \sim \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & 0 & \beta \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\Rightarrow \rho(A) = 3$. This satisfies the condition of rank being at least three.

- (ii) Let $\alpha = \beta = \gamma = 0$:

$$A \sim \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\Rightarrow \rho(A) = 2$. This does not satisfy the condition.

- (iii) Let $\alpha \neq 0, \beta = \gamma = 0$:

$$A \sim \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\Rightarrow \rho(A) = 3$. This satisfies the condition.

- (iv) Let $\alpha = \beta = \gamma \neq 0$:

$$A \sim \begin{bmatrix} 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 \\ 0 & 0 & \alpha & \alpha \\ 0 & 0 & 0 & \alpha \end{bmatrix}$$

$\Rightarrow \rho(A) = 4$. This satisfies the condition.

Since cases (i), (iii), and (iv) result in a rank ≥ 3 , the correct combination matches Option (a).

(A) Only (i), (iii), and (iv)

Q1.24.1 NAT 2024 1M Ans: 5 ● ○ ○

Let $X = \begin{bmatrix} 2 & -3 & \alpha \end{bmatrix}$, $Y = \begin{bmatrix} 3 & -1 & 3 \end{bmatrix}$ and $Z = \begin{bmatrix} 1 & -5 & 7 \end{bmatrix}$. Then $A = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} 2 & -3 & \alpha \\ 3 & -1 & 3 \\ 1 & -5 & 7 \end{bmatrix}$. Given that X, Y and Z vectors do not form a basis for $\mathbb{R}^3$. $\Rightarrow X, Y$ and Z are not linearly independent vectors. $\Rightarrow |A| = 0$

$$\begin{vmatrix} 2 & -3 & \alpha \\ 3 & -1 & 3 \\ 1 & -5 & 7 \end{vmatrix} = 0$$

$$2(-7 + 15) + 3(21 - 3) + \alpha(-15 + 1) = 0$$

$$2(8) + 3(18) + \alpha(-14) = 0$$

$$16 + 54 - 14\alpha = 0$$

$$70 - 14\alpha = 0$$

$$14\alpha = 70$$

$$\alpha = \boxed{5}$$

Key Points

- Vectors are linearly dependent if the determinant of the matrix formed by them is zero ($|A| = 0$).

Q1.24.2 **MSQ** **2024** **2M** **Ans: A,B** ☒ ☒ ☐

Given matrix $M = \begin{bmatrix} 1 & k \\ 2 & 1 \end{bmatrix}$ where $k > 0$.

Method 1

We first determine the eigenvalues λ by solving the characteristic equation $\det(M - \lambda I) = 0$:

$$\begin{vmatrix} 1 - \lambda & k \\ 2 & 1 - \lambda \end{vmatrix} = 0$$

$$(1 - \lambda)^2 - 2k = 0$$

$$1 - \lambda = \pm\sqrt{2k} \Rightarrow \lambda = 1 \pm \sqrt{2k}$$

The eigenvector $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ satisfies $(M - \lambda I)X = 0$:

$$\begin{bmatrix} \mp\sqrt{2k} & k \\ 2 & \mp\sqrt{2k} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

From the first row: $\mp\sqrt{2k}x_1 + kx_2 = 0 \Rightarrow \frac{x_2}{x_1} = \pm\frac{\sqrt{2k}}{k} = \pm\sqrt{\frac{2}{k}}$.

- For $\lambda = 1 - \sqrt{2k}$, the ratio is $-\sqrt{2/k}$, which matches Option (A).
- For $\lambda = 1 + \sqrt{2k}$, the ratio is $\sqrt{2/k}$, which matches Option (B).

$$\boxed{(A) \begin{bmatrix} 1 \\ -\sqrt{2/k} \end{bmatrix}, (B) \begin{bmatrix} 1 \\ \sqrt{2/k} \end{bmatrix}}$$

Method 2

We can verify the eigenvectors directly by checking if $MX = \lambda X$.

Check Option (A): $X_1 = \begin{bmatrix} 1 \\ -\sqrt{2/k} \end{bmatrix}$:

$$MX_1 = \begin{bmatrix} 1 & k \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -\sqrt{2/k} \end{bmatrix} = \begin{bmatrix} 1 - \sqrt{2k} \\ 2 - \sqrt{2/k} \end{bmatrix} = (1 - \sqrt{2k}) \begin{bmatrix} 1 \\ \frac{2 - \sqrt{2/k}}{1 - \sqrt{2k}} \end{bmatrix}$$

Simplifying the second component: $\frac{\sqrt{2/k}(\sqrt{2k}-1)}{1-\sqrt{2k}} = -\sqrt{2/k}$. Since $MX_1 = (1-\sqrt{2k})X_1$, Option (A) is an eigenvector.

Check Option (B): $X_2 = \begin{bmatrix} 1 \\ \sqrt{2/k} \end{bmatrix}$:

$$MX_2 = \begin{bmatrix} 1 & k \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ \sqrt{2/k} \end{bmatrix} = \begin{bmatrix} 1 + \sqrt{2k} \\ 2 + \sqrt{2/k} \end{bmatrix} = (1 + \sqrt{2k}) \begin{bmatrix} 1 \\ \sqrt{2/k} \end{bmatrix}$$

Since $MX_2 = (1 + \sqrt{2k})X_2$, Option (B) is an eigenvector.

$$\boxed{(A) \begin{bmatrix} 1 \\ -\sqrt{2/k} \end{bmatrix}, (B) \begin{bmatrix} 1 \\ \sqrt{2/k} \end{bmatrix}}$$

Key Points

- Eigenvectors X satisfy the linear transformation property $MX = \lambda X$.
- Any scalar multiple of an eigenvector is also an eigenvector.

Mistakes to be avoided

- Be careful with algebraic simplification involving square roots of variables like k .

Q1.23.1 MCQ 2023 1M Ans: C ● ● ○

Given that v_k is an eigenvector of matrix B corresponding to the eigenvalue λ_k :

$$Bv_k = \lambda_k v_k$$

$$B = P^{-1}AP$$

$$PB = AP$$

$$PBv_k = APv_k$$

Substituting $PB = AP$ on the left side and using the property that λ_k is a scalar:

$$P\lambda_k v_k = A(Pv_k)$$

$$\lambda_k(Pv_k) = A(Pv_k)$$

$$\Rightarrow A(Pv_k) = \lambda_k(Pv_k)$$

By the definition of eigenvalues and eigenvectors ($Ax = \lambda x$), the eigenvalue and eigenvector of A are λ_k and Pv_k respectively.

$$\boxed{(C) \{\lambda_k \mid 1 \leq k \leq n\} \text{ and } \{Pv_k \mid 1 \leq k \leq n\}}$$

Q1.23.2 MCQ 2023 2M Ans: A ● ○ ○

Let x be a 3×1 real column vector:

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

The length l of the vector is defined as $l = \sqrt{x^T x}$. Squaring both sides, we get:

$$l^2 = x^T x = \begin{pmatrix} x_1 & x_2 & x_3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$l^2 = x_1^2 + x_2^2 + x_3^2$$

Now, consider the matrix $P = xx^T$:

$$P = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \begin{pmatrix} x_1 & x_2 & x_3 \end{pmatrix} = \begin{pmatrix} x_1^2 & x_1x_2 & x_1x_3 \\ x_2x_1 & x_2^2 & x_2x_3 \\ x_3x_1 & x_3x_2 & x_3^2 \end{pmatrix}$$

$$\text{Trace}(P) = x_1^2 + x_2^2 + x_3^2$$

$$\text{Trace}(P) = l^2$$

$$(C) l^2$$

Q1.22.1 MCQ 2022 1M Ans: C ☒ ☐ ☐

The augmented matrix $[A | B]$ for the given system of linear equations is:

$$[A | B] = \begin{bmatrix} 1 & -\sqrt{2} & 3 & 1 \\ -1 & \sqrt{2} & -3 & 3 \end{bmatrix}$$

Applying the row operation $R_2 \rightarrow R_2 + R_1$:

$$= \begin{bmatrix} 1 & -\sqrt{2} & 3 & 1 \\ 0 & 0 & 0 & 4 \end{bmatrix}$$

$$\rho(A) = 1$$

$$\rho(A | B) = 2$$

Since $\rho(A) \neq \rho(A | B)$, the system of equations is inconsistent. Therefore, there is no solution for x .

$$(C) \text{ no solutions for } x$$

Key Points

- **Rouché–Capelli Theorem:** A system of linear equations $Ax = b$ is consistent if and only if the rank of the coefficient matrix A is equal to the rank of the augmented matrix $[A | B]$.
- If $\rho(A) = \rho(A | B) = n$ (number of variables), the system has a unique solution.
- If $\rho(A) = \rho(A | B) < n$, the system has infinitely many solutions.
- If $\rho(A) < \rho(A | B)$, the system has no solution.

Q1.22.2 MCQ 2022 2M Ans: A ☒ ☒ ☐

Given:

$$A = \alpha V_1 V_1^T + \beta V_2 V_2^T$$

Multiplying by V_1 on both sides:

$$AV_1 = \alpha V_1 V_1^T V_1 + \beta V_2 V_2^T V_1$$

Since $V_1^T V_1 = 1$ and $V_2^T V_1 = 0$ (orthogonal vectors):

$$AV_1 = \alpha V_1 (1) + \beta V_2 (0)$$

$$AV_1 = \alpha V_1$$

Comparing with the standard eigenvalue equation $A\mathbf{x} = \lambda\mathbf{x}$, we find: One of the eigenvalue $= \alpha$ Similarly, we can verify the other eigenvalues by multiplying by V_2 and a third vector V_3 orthogonal to both V_1 and V_2 :

$$AV_2 = \alpha V_1 V_1^T V_2 + \beta V_2 V_2^T V_2 = \alpha V_1(0) + \beta V_2(1) = \beta V_2$$

$$AV_3 = \alpha V_1(0) + \beta V_2(0) = 0$$

The eigenvalues are $\alpha, \beta, 0$.

(A) $0, \alpha, \beta$

Q1.21.1 NAT 2021 1M Ans: 8 ☒ ☐ ☐

Given that the three vectors $(1.0, -1.0, 2.0)$, $(7.0, 3.0, x)$, and $(2.0, 3.0, 1.0)$ in $\mathbb{R}^3$ are linearly dependent.

$$\begin{vmatrix} 1 & -1 & 2 \\ 7 & 3 & x \\ 2 & 3 & 1 \end{vmatrix} = 0$$

Expanding the determinant along the first row:

$$1(3 - 3x) - (-1)(7 - 2x) + 2(21 - 6) = 0$$

$$3 - 3x + 7 - 2x + 2(15) = 0$$

$$10 - 5x + 30 = 0$$

$$40 - 5x = 0$$

$$5x = 40$$

$$x = \frac{40}{5} = 8$$

Key Points

- Vectors are linearly dependent if at least one vector can be expressed as a linear combination of the others.
- For n vectors in $\mathbb{R}^n$, they are linearly dependent if and only if the determinant of the matrix they form is zero.

Q1.21.2 NAT 2021 2M Ans: 10 ☒ ☒ ☐

Given matrix:

$$A = \begin{bmatrix} x & -3 \\ 3 & 4 \end{bmatrix}$$

The characteristic equation of a given 2×2 matrix A is defined as $|A - \lambda I| = 0$:

$$\lambda^2 - \text{Trace}(A)\lambda + |A| = 0$$

$$\lambda^2 - (x + 4)\lambda + (4x + 9) = 0$$

The roots (eigenvalues) are given by:

$$\lambda = \frac{(x + 4) \pm \sqrt{(x + 4)^2 - 4(1)(4x + 9)}}{2(1)}$$

It is given that the characteristic equation has repeated roots. The condition for repeated roots in a quadratic equation $a\lambda^2 + b\lambda + c = 0$ is the discriminant $D = b^2 - 4ac = 0$:

$$(x + 4)^2 - 4(1)(4x + 9) = 0$$

$$x^2 + 16 + 8x - 16x - 36 = 0$$

$$x^2 - 8x - 20 = 0$$

Solving for x :

$$x = \frac{8 \pm \sqrt{64 + 80}}{2} = \frac{8 \pm \sqrt{144}}{2} = \frac{8 \pm 12}{2}$$

$$x = 10 \text{ or } -2$$

Since x is stated to be a real positive number:

10

Key Points

- For a 2×2 matrix, it is $\lambda^2 - S_1\lambda + S_2 = 0$, where S_1 is the trace and S_2 is the determinant.
- Repeated Eigenvalues occur when the discriminant of the characteristic equation is zero ($D = 0$).
- For $ax^2 + bx + c = 0$, Discriminant $D = b^2 - 4ac$.

Mistakes to be avoided

- **Ignoring Constraints:** Failing to select the positive value $x = 10$ over $x = -2$ as specified in the problem.

Q1.20.1 MCQ 2020 1M Ans: C ☒ ☐ ☐

Given six vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_6$ in $\mathbb{R}^4$.

- **Statement (A):** It is not necessary that these vectors span $\mathbb{R}^4$. This is **TRUE**. While the minimum number of vectors needed to span $\mathbb{R}^4$ is 4, having 6 vectors does not guarantee a span if they are linearly dependent (e.g., all 6 could lie on a single line).
- **Statement (B):** These vectors are not linearly independent. This is **TRUE**. In a vector space of dimension n , any set containing more than n vectors must be linearly dependent. Since $6 > 4$, the set is definitely linearly dependent.
- **Statement (C):** Any four of these vectors form a basis for $\mathbb{R}^4$. This is **FALSE**. For four vectors to form a basis for $\mathbb{R}^4$, they must be linearly independent. There is no guarantee that any subset of four vectors from the given six will be linearly independent (e.g., some could be zero vectors or multiples of others).
- **Statement (D):** If $\{\mathbf{v}_1, \mathbf{v}_3, \mathbf{v}_5, \mathbf{v}_6\}$ spans $\mathbb{R}^4$, then it forms a basis for $\mathbb{R}^4$. This is **TRUE**. In an n -dimensional space, a set of n vectors that spans the space is automatically a basis.

The false statement is (C).

(C) Any four of these vectors form a basis for $\mathbb{R}^4$.

Key Points

- **Basis Dimension:** A basis for $\mathbb{R}^n$ must contain exactly n linearly independent vectors.
- **Linear Dependence:** Any set of m vectors in $\mathbb{R}^n$ where $m > n$ is linearly dependent.
- **Spanning Set:** A set of n vectors in an n -dimensional space forms a basis if and only if it spans the space.

Mistakes to be avoided

- Do not assume that having "enough" vectors (4 or more) automatically guarantees they span $\mathbb{R}^4$ or contain a basis.
- Remember that linear independence is a requirement for a basis, not just the count of vectors.

Q1.20.2 MCQ 2020 2M Ans: A ● ● ○

Consider the augmented matrix $(A | B)$ for the given system of linear equations:

$$(A | B) = \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 2 & 4 & b_2 \\ 3 & 7 & b_3 \\ 3 & 9 & b_4 \end{array} \right]$$

Applying row operations $R_2 \rightarrow R_2 - 2R_1$, $R_3 \rightarrow R_3 - 3R_1$, and $R_4 \rightarrow R_4 - 3R_1$:

$$\Rightarrow (A | B) \sim \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & 0 & b_2 - 2b_1 \\ 0 & 1 & b_3 - 3b_1 \\ 0 & 3 & b_4 - 3b_1 \end{array} \right]$$

Interchanging rows $R_2 \leftrightarrow R_3$:

$$\Rightarrow (A | B) \sim \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & 1 & b_3 - 3b_1 \\ 0 & 0 & b_2 - 2b_1 \\ 0 & 3 & b_4 - 3b_1 \end{array} \right]$$

Applying row operation $R_4 \rightarrow R_4 - 3R_2$:

$$\Rightarrow (A | B) \sim \left[\begin{array}{cc|c} 1 & 2 & b_1 \\ 0 & 1 & b_3 - 3b_1 \\ 0 & 0 & b_2 - 2b_1 \\ 0 & 0 & 6b_1 - 3b_3 + b_4 \end{array} \right]$$

Here, $\rho(A) = 2$. To have a solution, $\rho(A) = \rho(A | B)$.

$$b_2 - 2b_1 = 0 \implies b_2 = 2b_1$$

$$6b_1 - 3b_3 + b_4 = 0$$

Thus, the system has a solution if $b_2 = 2b_1$ and $6b_1 - 3b_3 + b_4 = 0$.

$$(A) \quad b_2 = 2b_1 \text{ and } 6b_1 - 3b_3 + b_4 = 0$$

Q1.19.1 NAT 2019 1M Ans: 3 ● ○ ○

$$A = \begin{bmatrix} 2 & 2 & 3 & 3 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

The matrix A is an upper triangular matrix because all entries below the main diagonal are zero. Hence, the eigenvalues are the elements of principle diagonal.

$$\lambda_1 = 2, \quad \lambda_2 = 1, \quad \lambda_3 = 3, \quad \lambda_4 = 2$$

$$\text{Distinct eigenvalues} = \{1, 2, 3\}$$

The total number of distinct eigenvalues is 3.

$$\boxed{3}$$

Key Points

- For triangular, or diagonal matrices, the eigenvalues are the diagonal elements.

Q1.18.1 MCQ 2018 1M Ans: C ● ○ ○

For a square matrix, the following properties regarding eigen values and eigen vectors hold true:

- If a matrix $M_{4 \times 4}$ has four distinct eigen values, then the matrix $M_{4 \times 4}$ will have four linearly independent eigen vectors.
- If a matrix $M_{4 \times 4}$ has repeated eigen values, then the eigen vectors of the matrix may or may not be linearly independent.
- If a matrix $M_{4 \times 4}$ is non-singular, then the eigen values of the matrix may or may not be repeated.

From the theoretical properties of linear algebra, statement $S_2 \Rightarrow S_1$ is correct.

(C) S_2 implies S_1

Q1.18.2 NAT 2018 1M Ans: 2 ● ○ ○

The condition for a homogeneous system $Ax = O$ to have infinitely many solutions is that the matrix A must be singular, i.e., $|A| = 0$, or the rank $\rho(A) < n$, where n is the number of variables.

Consider $|A| = 0$:

$$\begin{aligned} \begin{vmatrix} k & 2k \\ k^2 - k & k^2 \end{vmatrix} &= 0 \\ \Rightarrow k^3 - 2k(k^2 - k) &= 0 \\ \Rightarrow k^3 - 2k^3 + 2k^2 &= 0 \\ \Rightarrow 2k^2 - k^3 &= 0 \\ \Rightarrow k^2(2 - k) &= 0 \\ \Rightarrow k &= 0, 2 \end{aligned}$$

Therefore, the system $Ax = O$ will have infinitely many solutions for two distinct real values of k .

2

Q1.17.1 NAT 2017 1M Ans: 4 ● ● ○

Given the matrix A :

$$A = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Applying elementary row transformations to reduce the matrix to Echelon form: Step 1: $R_4 \rightarrow R_4 + R_1$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Step 2: $R_2 \leftrightarrow R_3$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Step 3: $R_4 \rightarrow R_4 + R_2$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Step 4: $R_4 \rightarrow R_4 + R_3$

$$\Rightarrow A \sim \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix}$$

Step 5: $R_5 \rightarrow R_5 + R_4$

$$\Rightarrow A = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

The above matrix is in Row Echelon Form. The rank of matrix A is the number of non-zero rows in its Echelon form. Number of non-zero rows = 4.

4

Key Points

- The rank of a matrix is defined as the number of non-zero rows in its Row Echelon Form.
- Elementary row operations (row addition, row swapping) do not change the rank of a matrix.

Q1.17.2 MCQ 2017 1M Ans: C ● ● ○

Given the matrix A :

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 5 & 1 & 2 & 3 & 4 \\ 4 & 5 & 1 & 2 & 3 \\ 3 & 4 & 5 & 1 & 2 \\ 2 & 3 & 4 & 5 & 1 \end{bmatrix}$$

The characteristic equation is $|A - \lambda I| = 0$:

$$\begin{vmatrix} 1-\lambda & 2 & 3 & 4 & 5 \\ 5 & 1-\lambda & 2 & 3 & 4 \\ 4 & 5 & 1-\lambda & 2 & 3 \\ 3 & 4 & 5 & 1-\lambda & 2 \\ 2 & 3 & 4 & 5 & 1-\lambda \end{vmatrix} = 0$$

Applying the column transformation $C_1 \rightarrow C_1 + C_2 + C_3 + C_4 + C_5$:

$$\Rightarrow \begin{vmatrix} 15-\lambda & 2 & 3 & 4 & 5 \\ 15-\lambda & 1-\lambda & 2 & 3 & 4 \\ 15-\lambda & 5 & 1-\lambda & 2 & 3 \\ 15-\lambda & 4 & 5 & 1-\lambda & 2 \\ 15-\lambda & 3 & 4 & 5 & 1-\lambda \end{vmatrix} = 0$$

Taking $(15 - \lambda)$ common from C_1 :

$$\Rightarrow (15 - \lambda) \begin{vmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 1-\lambda & 2 & 3 & 4 \\ 1 & 5 & 1-\lambda & 2 & 3 \\ 1 & 4 & 5 & 1-\lambda & 2 \\ 1 & 3 & 4 & 5 & 1-\lambda \end{vmatrix} = 0$$

$$15 - \lambda = 0 \Rightarrow \lambda = 15$$

(C) 15

Q1.17.3 MCQ 2017 1M Ans: C ● ○ ○

Given the matrix M :

$$M = \begin{bmatrix} 5 & 10 & 10 \\ 1 & 0 & 2 \\ 3 & 6 & 6 \end{bmatrix}$$

Step 1: $R_2 \rightarrow 5R_2 - R_1$ and $R_3 \rightarrow 5R_3 - 3R_1$

$$\Rightarrow M \sim \begin{bmatrix} 5 & 10 & 10 \\ 0 & -10 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The resulting matrix is in Row Echelon Form. The rank of a matrix is defined as the number of non-zero rows in its Echelon form. Number of non-zero rows = 2. $\therefore$ Rank of $M = 2$.

(C) 2

Q1.16.1 MCQ 2016 1M Ans: C ● ○ ○

Given that:

$$M^4 = I$$

By multiplying both sides by M^4 , we get:

$$M^8 = M^4 \Rightarrow M^8 = I$$

$$M^8 \cdot M^{-1} = I \cdot M^{-1} \Rightarrow M^7 = M^{-1}$$

Similarly,

$$M^{12} = M^8 = I \Rightarrow M^{11} = M^{-1}$$

$$M^{16} = M^{12} = I \Rightarrow M^{15} = M^{-1}$$

In general, we observe that the power of M that equals M^{-1} follows the pattern:

$$M^{-1} = M^{4k+3}, \text{ where } k \text{ is any natural number } (0, 1, 2, \dots)$$

(C) M^{4k+3}

Key Points

- For an invertible matrix M , $M \cdot M^{-1} = I$.
- If $M^n = I$, then $M^{-1} = M^{n-1}$.

Q1.16.2 NAT 2016 1M Ans: 1 ☒ ☐ ☐

Given that one of the eigen values of the matrix $A_{3 \times 3}$ is 0. This implies that the determinant of the matrix must be zero:

$$\det(A) = 0$$

$$3\{(-63 + 7x) + 52\} - 2\{(-81 + 9x) + 78\} + 4\{-36 + 42\} = 0$$

$$3\{7x - 11\} - 2\{9x - 3\} + 4\{6\} = 0$$

$$21x - 33 - 18x + 6 + 24 = 0$$

$$3x - 3 = 0$$

$$\therefore x = \boxed{1}$$

Key Points

- A matrix is singular (determinant is zero) if and only if at least one of its eigen values is zero.
- The product of all eigen values of a matrix is equal to its determinant.

Q1.16.3 NAT 2016 2M Ans: 3 ☒ ☐ ☐

Given the matrix A :

$$A = \begin{bmatrix} a & 0 & 3 & 7 \\ 2 & 5 & 1 & 3 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & b \end{bmatrix}$$

Given $\text{trace}(A) = 14$:

$$a + 5 + 2 + b = 14$$

$$\Rightarrow a + b = 7 \quad \dots (1)$$

The determinant of a block triangular matrix is the product of the determinants of its diagonal blocks.

$$\det(A) = \begin{vmatrix} a & 0 \\ 2 & 5 \end{vmatrix} \times \begin{vmatrix} 2 & 4 \\ 0 & b \end{vmatrix}$$

Given $\det(A) = 100$:

$$(5a - 0) \times (2b - 0) = 100$$

$$\Rightarrow 10ab = 100$$

$$\Rightarrow ab = 10 \quad \dots (2)$$

Using the algebraic identity $(a - b)^2 = (a + b)^2 - 4ab$:

$$(a - b)^2 = (7)^2 - 4(10)$$

$$(a - b)^2 = 49 - 40 = 9$$

$$|a - b| = \sqrt{9} = 3$$

$$\boxed{3}$$

Key Points

- The trace of a matrix is the sum of the principal diagonal elements.
- For a 2×2 matrix, $\det \begin{bmatrix} p & q \\ r & s \end{bmatrix} = ps - qr$.

Q1.16.4 MCQ 2016 1M Ans: D ● ○ ○

We know that the determinant of a matrix A is equal to the product of its eigen values.

$$\det(A) = \text{product of eigen values}$$

Based on the given matrix properties:

$$\sigma^2 - \omega x = \sigma^2 + \omega^2$$

$$\Rightarrow \omega^2 = -\omega x$$

$$\Rightarrow \omega^2 + \omega x = 0$$

$$\therefore x = -\omega$$

$$(D) -\omega$$

Key Points

- The determinant of any square matrix is always equal to the product of all its eigen values.
- Eigen values are the roots of the characteristic equation $\det(A - \lambda I) = 0$.

Q1.16.5 MCQ 2016 2M Ans: D ● ● ●

Method 1

Let the vector u be expressed as a linear combination of the basis vectors e_1 , e_2 , and e_3 :

$$u = x_1 e_1 + x_2 e_2 + x_3 e_3$$

Given the vector $(4, 3, -3)$ and the basis vectors, we have:

$$(4, 3, -3) = x_1(1, 0, 2) + x_2(0, 1, 0) + x_3(-2, 0, 1)$$

Equating the components, we get the following system of linear equations:

$$x_1 - 2x_3 = 4 \quad \dots (1)$$

$$x_2 = 3 \quad \dots (2)$$

$$2x_1 + x_3 = -3 \quad \dots (3)$$

On solving these equations: From (2), we directly have $x_2 = 3$. To find x_1 and x_3 , we use equations (1) and (3). Multiplying (3) by 2 and adding to (1):

$$(x_1 - 2x_3) + 2(2x_1 + x_3) = 4 + 2(-3)$$

$$5x_1 = -2 \Rightarrow x_1 = -\frac{2}{5}$$

Substituting x_1 into (3):

$$2\left(-\frac{2}{5}\right) + x_3 = -3 \Rightarrow x_3 = -3 + \frac{4}{5} = -\frac{11}{5}$$

Thus, the vector u is:

$$\therefore u = -\frac{2e_1}{5} + 3e_2 - \frac{11e_3}{5}$$

$$(D) u = -\frac{2}{5}e_1 + 3e_2 - \frac{11}{5}e_3$$

Method 2

The problem can be solved using a matrix approach $AX = B$, where A is the matrix formed by the basis vectors as columns.

$$\begin{bmatrix} 1 & 0 & -2 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ -3 \end{bmatrix}$$

Using Cramer's Rule or matrix inversion: The determinant $\Delta = 1(1 - 0) - 0 + (-2)(0 - 2) = 1 + 4 = 5$.

To find x_1 :

$$\Delta_1 = \begin{vmatrix} 4 & 0 & -2 \\ 3 & 1 & 0 \\ -3 & 0 & 1 \end{vmatrix} = 4(1) - 0 + (-2)(0 - (-3)) = 4 - 6 = -2$$

$$x_1 = \frac{\Delta_1}{\Delta} = -\frac{2}{5}$$

To find x_3 :

$$\Delta_3 = \begin{vmatrix} 1 & 0 & 4 \\ 0 & 1 & 3 \\ 2 & 0 & -3 \end{vmatrix} = 1(-3) - 0 + 4(0 - 2) = -3 - 8 = -11$$

$$x_3 = \frac{\Delta_3}{\Delta} = -\frac{11}{5}$$

From the second row of the matrix, $x_2 = 3$.

$$\therefore u = -\frac{2e_1}{5} + 3e_2 - \frac{11e_3}{5}$$

$$(D) u = -\frac{2}{5}e_1 + 3e_2 - \frac{11}{5}e_3$$

Key Points

- Matrix inversion or Cramer's Rule are efficient for solving 3×3 systems in competitive exams like GATE.

Mistakes to be avoided

- Ensure the basis vectors are placed as columns in the matrix A , not rows.
- Be careful with signs when calculating the determinant.

Q1.15.1

NAT

2015

1M

Ans: 2

☒ ☐ ☐

$$[A|B] = \begin{bmatrix} 1 & -2 & 3 & -1 \\ 1 & -3 & 4 & 1 \\ -2 & 4 & -6 & k \end{bmatrix}$$

Step 1: $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 + 2R_1$:

$$\Rightarrow [A|B] \sim \begin{bmatrix} 1 & -2 & 3 & -1 \\ 0 & -1 & 1 & 2 \\ 0 & 0 & 0 & k-2 \end{bmatrix}$$

The system of equations is consistent if and only if the rank of the coefficient matrix A is equal to the rank of the augmented matrix $[A|B]$. From the Echelon form, $\text{rank}(A) = 2$ because there are two non-zero rows in the coefficient part. For consistency, we must have:

$$k - 2 = 0 \Rightarrow k = \boxed{2}$$

Key Points

- A system $AX = B$ is consistent if $\text{rank}(A) = \text{rank}([A|B])$.
- If $\text{rank}(A) = \text{rank}([A|B]) < \text{number of variables}$, the system has infinitely many solutions.
- If $\text{rank}(A) \neq \text{rank}([A|B])$, the system is inconsistent (no solution).

Q1.15.2 NAT 2015 1M Ans: 16.5 to 17.5 ☒ ☐ ☐

To find the value of P , we consider the eigen value equation $AX = \lambda X$:

$$\begin{pmatrix} 4 & 1 & 2 \\ P & 2 & 1 \\ 14 & -4 & 10 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \lambda \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

From the first row:

$$\begin{aligned} 4(1) + 1(2) + 2(3) &= \lambda(1) \\ 4 + 2 + 6 &= \lambda \Rightarrow \lambda = 12 \end{aligned}$$

From the second row:

$$\begin{aligned} P(1) + 2(2) + 1(3) &= \lambda(2) \\ P + 4 + 3 &= 2\lambda \\ P + 7 &= 2(12) \\ P + 7 &= 24 \\ \therefore P &= \boxed{17} \end{aligned}$$

Q1.15.3 MCQ 2015 1M Ans: B ☒ ☒ ☐

A complex square matrix $A_{n \times n}$ is called a Hermitian matrix if $A^\theta = A$, where $A^\theta = (\overline{A})^T$ denotes the conjugate transpose. Given the matrix:

$$A = \begin{bmatrix} 10 & 5+j & 4 \\ x & 20 & 2 \\ 4 & 2 & -10 \end{bmatrix}$$

It is given that the eigen values of the matrix are real. Since the matrix contains complex entries (specifically $a_{12} = 5 + j$), it must be a Hermitian matrix for its eigen values to be real.

For a matrix to be Hermitian, the condition $a_{ij} = \overline{a_{ji}}$ must be satisfied for all i, j :

- $a_{13} = a_{31} = 4$
- $a_{23} = a_{32} = 2$
- $a_{21} = \overline{a_{12}}$

Given $a_{12} = 5 + j$, then a_{21} must be the complex conjugate:

$$x = \overline{5 + j} = 5 - j$$

$$\boxed{\text{(B) } 5 - j}$$

Key Points

- A Hermitian matrix is defined by $A = A^{\theta}$.
- The eigen values of any Hermitian matrix (and real symmetric matrix) are always real.

Mistakes to be avoided

- Do not confuse Hermitian matrices with skew-Hermitian matrices, where $A^{\theta} = -A$.

Q1.15.4 MCQ 2015 1M Ans: C ☒ ☐ ☐

$$|A^T A^{-1}| = |A^T| |A^{-1}|$$

Using the properties $|A^T| = |A|$ and $|A^{-1}| = \frac{1}{|A|}$:

$$\Rightarrow |A^T A^{-1}| = |A| \cdot \frac{1}{|A|}$$

$$\therefore |A^T A^{-1}| = 1$$

Given the specific case for matrix A , the determinant is calculated as:

$$|A| = 1 + \tan^2 x = \sec^2 x$$

Since $|A| = \sec^2 x \neq 0$ (for defined values of $\tan x$), the inverse exists and the product of the determinants remains unity.

$$|A^T A^{-1}| = 1$$

(C) 1

Key Points

- The determinant of a product of matrices is the product of their individual determinants: $|AB| = |A||B|$.
- The determinant of a matrix is equal to the determinant of its transpose: $|A| = |A^T|$.
- The determinant of the inverse of a matrix is the reciprocal of the determinant of the original matrix: $|A^{-1}| = \frac{1}{|A|}$.

Mistakes to be avoided

- Ensure $|A| \neq 0$ before assuming the existence of A^{-1} .

Q1.14.1 NAT 2014 1M Ans: 1 ☒ ☐ ☐

Let λ be an eigenvalue of matrix A . According to the property of eigenvalues, if λ is an eigenvalue of A , then λ^n is an eigenvalue of A^n .

Given that:

$$A^2 = I$$

$$\lambda^2 = 1$$

$$\lambda = \pm 1$$

$$\lambda = \boxed{1}$$

Key Points

- If λ is an eigenvalue of A , then λ^n is an eigenvalue of A^n .

Q1.14.2 MCQ 2014 1M Ans: D ● ○ ○

The fundamental property of matrix algebra states that matrix multiplication is generally not commutative. This means for any two square matrices M and N of the same order, the product MN is not necessarily equal to the product NM .

$$MN \neq NM$$

Therefore, the statement aligns with option (D).

$$(D) MN = NM$$

Q1.14.3 NAT 2014 2M Ans: 1 ● ● ○

Given the matrix expression for P :

$$P = I_6 + \alpha J_6$$

$$\Rightarrow P = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \alpha \\ 0 & 1 & 0 & 0 & \alpha & 0 \\ 0 & 0 & 1 & \alpha & 0 & 0 \\ 0 & 0 & \alpha & 1 & 0 & 0 \\ 0 & \alpha & 0 & 0 & 1 & 0 \\ \alpha & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

To find the value of α that makes the matrix singular ($|P| = 0$), we observe the relationship between the rows.

If we set $\alpha = 1$:

$$R_1 = [1, 0, 0, 0, 0, 1]$$

$$R_6 = [1, 0, 0, 0, 0, 1]$$

Since $R_1 = R_6$, the rows are linearly dependent. According to the properties of determinants, if any two rows (or columns) of a matrix are identical, the determinant of that matrix is zero.

$$\Rightarrow |P| = 0$$

Thus, the required value of α is 1.

$$1$$

Key Points

- Determinant Property:** If a matrix has two identical rows or two identical columns, its determinant is 0.

Q1.14.4 NAT 2014 1M Ans: 200 ● ○ ○

$$\text{Det}(A \cdot B) = \text{Det}(A) \cdot \text{Det}(B)$$

$$\text{Det}(A \cdot B) = 5 \times 40$$

$$\text{Det}(A \cdot B) = 200$$

Q1.14.5 MCQ 2014 2M Ans: B ● ● ○

$$(A|B) = \begin{pmatrix} 2 & 1 & 3 & 5 \\ 3 & 0 & 1 & -4 \\ 1 & 2 & 5 & 14 \end{pmatrix}$$

Step 1: $R_1 \leftrightarrow R_3$

$$(A|B) \sim \begin{pmatrix} 1 & 2 & 5 & 14 \\ 3 & 0 & 1 & -4 \\ 2 & 1 & 3 & 5 \end{pmatrix}$$

Step 2: $R_2 \rightarrow R_2 - 3R_1$ and $R_3 \rightarrow R_3 - 2R_1$

$$(A|B) \sim \begin{pmatrix} 1 & 2 & 5 & 14 \\ 0 & -6 & -14 & -46 \\ 0 & -3 & -7 & -23 \end{pmatrix}$$

Step 3: $R_3 \rightarrow 2R_3 - R_2$

$$(A|B) \sim \begin{pmatrix} 1 & 2 & 5 & 14 \\ 0 & -6 & -14 & -46 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Here, $\rho(A) = \rho(AB) = 2 < 3 = \text{Number of variables}$.

The system is consistent and has infinitely many solutions.

(B) infinitely many solutions

Q1.14.6 NAT 2014 2M Ans: 49 ● ● ○

Let the matrix A be defined as:

$$A = \begin{pmatrix} x & y \\ y & 14 - x \end{pmatrix}$$

$$\text{Det } A = x(14 - x) - y^2$$

To maximize the value of the determinant, $y = 0$.

$$A = \begin{pmatrix} x & 0 \\ 0 & 14 - x \end{pmatrix}$$

$$|A| = f(x) = x(14 - x) = 14x - x^2$$

$$f'(x) = 14 - 2x$$

$$14 - 2x = 0 \implies x = 7$$

$$f''(x) = -2$$

Since $f''(x) < 0$, the function has a maximum at $x = 7$.

Substituting $x = 7$ back into the determinant function:

$$f(7) = 14(7) - (7)^2 = 98 - 49 = 49$$

Thus, the maximum value of the determinant is **49**.

Q1.14.7 MCQ 2014 2M Ans: B ● ○ ○

For an upper triangular, lower triangular or diagonal matrix, the eigen values are the diagonal elements of the matrix. Hence, Option A is TRUE.

Consider the matrix:

$$A = \begin{pmatrix} -1 & 0 \\ 0 & -3 \end{pmatrix}$$

A is a real symmetric matrix (since $A = A^T$). $\Rightarrow$ The eigenvalues of matrix A are -1 and -3 . Hence Option B is FALSE.

Options C and D are standard theorems (referring to properties such as the eigenvalues of a Hermitian matrix being real, or the product of eigenvalues equaling the determinant).

(B) If A is real symmetric, the eigenvalues of A are always real and positive

Key Points

- Triangular Matrices: The eigenvalues of any triangular (upper or lower) or diagonal matrix are the entries on its main diagonal.
- Real Symmetric Matrices: The eigenvalues of a real symmetric matrix are always *real*, but they can be positive, negative, or zero.
- Characteristic Equation: The eigenvalues are the roots of the polynomial equation $\det(A - \lambda I) = 0$.

Q1.13.1 MCQ 2013 1M Ans: A ● ○ ○

Given the matrix A:

$$A = \begin{pmatrix} 3 & 5 & 2 \\ 5 & 12 & 7 \\ 2 & 7 & 5 \end{pmatrix}$$

To find the minimum eigenvalue, we can check the properties of the matrix rows or calculate the determinant.

Observing the rows of the matrix: Row 1 (R_1): $[3, 5, 2]$ Row 2 (R_2): $[5, 12, 7]$ Row 3 (R_3): $[2, 7, 5]$

Notice that $R_1 + R_3 = [3 + 2, 5 + 7, 2 + 5] = [5, 12, 7]$. Since $R_1 + R_3 = R_2$, the rows are linearly dependent. When the rows (or columns) of a matrix are linearly dependent, the matrix is singular, which implies its determinant is zero:

$$|A| = 0$$

$$\lambda_1 \lambda_2 \lambda_3 = |A|$$

$$\lambda_1 \lambda_2 \lambda_3 = 0$$

This means at least one of the eigenvalues must be zero. Looking at the options provided, the smallest possible eigenvalue is 0.

(A) 0

Key Points

- The product of the eigenvalues of a matrix is equal to the determinant of the matrix.
- Linear Dependency: If any row can be expressed as a linear combination of other rows, the determinant is zero.
- Singularity: A matrix with a zero eigenvalue is a singular matrix.

Mistakes to be avoided

- Do not immediately start calculating the characteristic equation $|A - \lambda I| = 0$ for a 3×3 matrix; always check for row/column dependencies first to save time.

Q1.13.2 **MCQ** **2013** **2M** **Ans: B** ● ● ○
Method 1

$$\text{Let } A = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}_{4 \times 1} \text{ and } B = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}_{1 \times 4}.$$

The product AB results in a 4×4 matrix:

$$AB = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

$$\text{Then, } I_4 + AB = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{pmatrix}.$$

Applying the identity with $m = 4$ and $n = 1$:

$$|I_4 + AB| = |I_1 + BA|$$

Calculate BA :

$$BA = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = [1 + 1 + 1 + 1] = [4]$$

$$\text{Thus, } |I_1 + BA| = |[1] + [4]| = |[5]| = 5.$$

(B) 5

Method 2

Alternatively, we can evaluate the determinant using row and column operations.

$$|A| = \begin{vmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{vmatrix}$$

Step 1: $C_1 \rightarrow C_1 + C_2 + C_3 + C_4$:

$$|A| = \begin{vmatrix} 5 & 1 & 1 & 1 \\ 5 & 2 & 1 & 1 \\ 5 & 1 & 2 & 1 \\ 5 & 1 & 1 & 2 \end{vmatrix}$$

Step 2: $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$, and $R_4 \rightarrow R_4 - R_1$:

$$|A| = \begin{vmatrix} 5 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix}$$

Since this is an upper triangular matrix, the determinant is the product of the diagonal elements:

$$|A| = 5 \times 1 \times 1 \times 1 = 5$$

(B) 5

Key Points

- **Sylvester's Determinant Identity:** $\det(I_m + AB) = \det(I_n + BA)$. This is extremely useful when one dimension (n) is much smaller than the other (m).
- **Determinant Properties:** Adding a multiple of one column/row to another does not change the determinant.
- **Triangular Matrices:** The determinant of an upper or lower triangular matrix is the product of its diagonal entries.

Q1.12.1 MCQ 2012 2M Ans: B ● ● ●

Given the matrix A :

$$A = \begin{pmatrix} -5 & -3 \\ 2 & 0 \end{pmatrix}$$

The characteristic equation of matrix A using $|A - \lambda I| = 0$:

$$\begin{vmatrix} -5 - \lambda & -3 \\ 2 & 0 - \lambda \end{vmatrix} = 0$$

$$(-5 - \lambda)(-\lambda) - (-3)(2) = 0$$

$$5\lambda + \lambda^2 + 6 = 0$$

$$\lambda^2 + 5\lambda + 6 = 0$$

According to the Cayley-Hamilton theorem, every square matrix satisfies its own characteristic equation.

Therefore:

$$A^2 + 5A + 6I = 0$$

$$A^2 = -5A - 6I$$

$$A^3 = -5A^2 - 6A$$

$$A^3 = -5(-5A - 6I) - 6A$$

$$A^3 = 25A + 30I - 6A$$

(B) $19A + 30I$

Key Points

- **Characteristic Equation:** For a 2×2 matrix, it can be simplified as $\lambda^2 - \text{Tr}(A)\lambda + |A| = 0$.
- **Cayley-Hamilton Theorem:** This theorem allows us to express higher powers of a matrix (A^n) as a linear combination of lower powers ($I, A, A^2, \dots, A^{n-1}$).
- **Matrix Inverse:** This theorem can also be used to find A^{-1} by multiplying the characteristic equation by A^{-1} .

Q1.11.1 MCQ 2011 2M Ans: B ● ● ○

Given $AX = B$.

$$[A|B] = \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 4 & 6 & 20 \\ 1 & 4 & \lambda & \mu \end{array} \right)$$

Step 1: $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$[A|B] \sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 3 & 5 & 14 \\ 0 & 3 & \lambda - 1 & \mu - 6 \end{array} \right)$$

Step 2: $R_3 \rightarrow R_3 - R_2$:

$$[A|B] \sim \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 3 & 5 & 14 \\ 0 & 0 & \lambda - 6 & \mu - 20 \end{array} \right)$$

The system has no solution if the rank of the coefficient matrix A is not equal to the rank of the augmented matrix $[A|B]$ ($\rho(A) \neq \rho[A|B]$).

This condition occurs when:

- $\lambda - 6 = 0 \implies \lambda = 6$ (This makes $\rho(A) = 2$)
- $\mu - 20 \neq 0 \implies \mu \neq 20$ (This makes $\rho[A|B] = 3$)

Under these conditions, $\rho(A) = 2 \neq 3 = \rho[A|B]$. Therefore, the system has no solution if $\lambda = 6$ and $\mu \neq 20$.

(B) $\lambda = 6, \mu \neq 20$

Key Points

• Consistency Criteria:

- Unique Solution: $\rho(A) = \rho[A|B] = n$ (number of variables).
- Infinite Solutions: $\rho(A) = \rho[A|B] < n$.
- No Solution: $\rho(A) \neq \rho[A|B]$.

Q1.10.1 MCQ 2010 1M Ans: C ☒ ☐ ☐

The eigenvalues of a real skew-symmetric matrix are always either zero or purely imaginary. This occurs because the characteristic equation of a skew-symmetric matrix involves purely imaginary roots or a zero root when the matrix is of odd order.

Therefore, option (c) is correct.

(C) either zero or pure imaginary

Q1.09.1 MCQ 2009 2M Ans: D ☒ ☒ ☐

The characteristic equation of a square matrix A is obtained by solving the determinant equation:

$$|A - \lambda I| = 0$$

$$(3 - \lambda)(\lambda^2 + 2\lambda + 10) = 0$$

i. $3 - \lambda = 0 \implies \lambda_1 = 3$

ii. $\lambda^2 + 2\lambda + 10 = 0$

Solving the quadratic equation:

$$\lambda = \frac{-2 \pm \sqrt{2^2 - 4(1)(10)}}{2(1)}$$

$$\lambda = \frac{-2 \pm \sqrt{4 - 40}}{2}$$

$$\lambda = \frac{-2 \pm \sqrt{-36}}{2}$$

$$\lambda = \frac{-2 \pm 6j}{2} = -1 \pm 3j$$

Thus, the eigenvalues of the matrix are 3, $-1 + 3j$, and $-1 - 3j$.

$$(D) \ 3, -1 + 3j, -1 - 3j$$

Q1.08.1 MCQ 2008 1M Ans: C ☒ ☐ ☐

Given the 2×2 matrix P :

$$P = \begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{bmatrix}, \text{ where } P_{ij} \neq 0 \quad \forall i, j$$

It is stated that one of the eigenvalues of matrix P is zero. Let the eigenvalues be λ_1 and λ_2 . If $\lambda_1 = 0$, then:

$$\lambda_1 \cdot \lambda_2 = \text{Det}(P)$$

$$0 \cdot \lambda_2 = \text{Det}(P) \implies \text{Det}(P) = 0$$

A matrix with a zero determinant is a singular matrix. For a 2×2 matrix, the determinant is calculated as:

$$|P| = P_{11}P_{22} - P_{12}P_{21}$$

$$P_{11}P_{22} - P_{12}P_{21} = 0$$

Thus, option (C) is correct.

$$(C) \ p_{11}p_{22} - p_{12}p_{21} = 0$$

Q1.08.2 MCQ 2008 1M Ans: B ☒ ☐ ☐

Given the system of linear equations:

$$4x + 2y = 7$$

$$2x + y = 6$$

$$2x + y = 7/2$$

$$2x + y = 6$$

Consider the augmented matrix $(A|B)$:

$$(A|B) = \left[\begin{array}{cc|c} 2 & 1 & 7/2 \\ 2 & 1 & 6 \end{array} \right]$$

Step 1: $R_2 \rightarrow R_2 - R_1$

$$(A|B) \sim \left[\begin{array}{cc|c} 2 & 1 & 7/2 \\ 0 & 0 & 5/2 \end{array} \right]$$

- $\rho(A) = 1$
- $\rho(A|B) = 2$

Since $\rho(A) \neq \rho(A|B)$, the system has no solution.

$$(B) \text{ no solution}$$

Q1.06.1 MCQ 2006 1M Ans: C ● ○ ○

Given the 3×3 matrix A :

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$$

Applying $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$A \sim \begin{bmatrix} 1 & 1 & 1 \\ 0 & -2 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\therefore \rho(A) = 2$$

$$\boxed{(C) \ 2}$$

Q1.06.2 MCQ 2006 2M Ans: A ● ● ○

Method 1

We know that for any square matrix, the sum of its eigenvalues is equal to the trace of the matrix.

By checking the given options, the matrix in option (a) is $\begin{bmatrix} 6 & 2 \\ 2 & 6 \end{bmatrix}$. Trace = $6 + 6 = 12$. Sum of given eigenvalues = $\lambda_1 + \lambda_2 = 8 + 4 = 12$.

Since the sum of eigenvalues matches the trace, option (a) is correct.

$$\boxed{(A) \begin{bmatrix} 6 & 2 \\ 2 & 6 \end{bmatrix}}$$

Method 2

Given eigenvalues $\lambda_1 = 8, \lambda_2 = 4$ and corresponding eigenvectors:

$$V_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, V_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

A matrix A can be reconstructed using the diagonalization formula $A = PDP^{-1}$, where P is the modal matrix consisting of eigenvectors and D is the diagonal matrix of eigenvalues.

$$P = [V_1 \ V_2] = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad D = \begin{bmatrix} 8 & 0 \\ 0 & 4 \end{bmatrix}$$

First, find P^{-1} :

$$P^{-1} = \frac{1}{|P|} \text{adj}(P) = \frac{1}{-1-1} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} = \frac{1}{-2} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

Now, compute A :

$$\begin{aligned} A &= \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 8 & 0 \\ 0 & 4 \end{bmatrix} \left(\frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right) \\ A &= \begin{bmatrix} 8 & 4 \\ 8 & -4 \end{bmatrix} \left(\frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \right) = \frac{1}{2} \begin{bmatrix} 8+4 & 8-4 \\ 8-4 & 8+4 \end{bmatrix} \\ A &= \frac{1}{2} \begin{bmatrix} 12 & 4 \\ 4 & 12 \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 2 & 6 \end{bmatrix} \end{aligned}$$

$$\boxed{(A) \begin{bmatrix} 6 & 2 \\ 2 & 6 \end{bmatrix}}$$

Key Points

- Trace Property: $\sum \lambda_i = \text{Tr}(A)$.
- Determinant Property: $\prod \lambda_i = |A|$.
- Diagonalization: $A = PDP^{-1}$ is valid if A has n linearly independent eigenvectors.

Mistakes to be avoided

- Order of Matrices: Ensure the order in PDP^{-1} is strictly followed; matrix multiplication is not commutative.

Q1.06.3 MCQ 2006 2M Ans: C ● ○ ○

Given that $X = \begin{bmatrix} 101 \\ 101 \end{bmatrix}$ is an eigenvector of the matrix $A = \begin{bmatrix} 4 & 2 \\ 2 & 4 \end{bmatrix}$ corresponding to an eigenvalue λ .

By the fundamental definition:

$$AX = \lambda X$$

$$\begin{bmatrix} 4 & 2 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 101 \\ 101 \end{bmatrix} = \lambda \begin{bmatrix} 101 \\ 101 \end{bmatrix}$$

$$\begin{bmatrix} (4 \times 101) + (2 \times 101) \\ (2 \times 101) + (4 \times 101) \end{bmatrix} = \lambda \begin{bmatrix} 101 \\ 101 \end{bmatrix}$$

$$\begin{bmatrix} 404 + 202 \\ 202 + 404 \end{bmatrix} = \lambda \begin{bmatrix} 101 \\ 101 \end{bmatrix}$$

$$\begin{bmatrix} 606 \\ 606 \end{bmatrix} = \lambda \begin{bmatrix} 101 \\ 101 \end{bmatrix}$$

$$6 \begin{bmatrix} 101 \\ 101 \end{bmatrix} = \lambda \begin{bmatrix} 101 \\ 101 \end{bmatrix}$$

$$\lambda = 6$$

$$(C) \ 6$$

Q1.05.1 MCQ 2005 2M Ans: C ● ○ ○

Given that A is an orthogonal matrix.

By the fundamental definition of an orthogonal matrix, a square matrix A is orthogonal if its product with its transpose A^T results in the identity matrix I .

$$AA^T = I = A^T A$$

$$(AA^T)^{-1} = (I)^{-1}$$

$$\therefore (AA^T)^{-1} = I$$

Thus, option (c) is correct.

$$(C) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Key Points

- Orthogonal Matrix Property: For an orthogonal matrix, the inverse is equal to its transpose ($A^{-1} = A^T$).
- Identity Matrix Inverse: The identity matrix is its own inverse ($I^{-1} = I$).
- Determinant: The determinant of an orthogonal matrix is always ± 1 .

Q1.05.2 MSQ 2005 2M Ans: C ● ● ○

Given matrix A:

$$A = \begin{bmatrix} -4 & 2 \\ 4 & 3 \end{bmatrix}$$

The eigenvalues are found by solving $|A - \lambda I| = 0$:

$$(-4 - \lambda)(3 - \lambda) - 8 = 0$$

$$\lambda^2 + \lambda - 12 - 8 = 0 \implies \lambda^2 + \lambda - 20 = 0$$

$$(\lambda + 5)(\lambda - 4) = 0 \implies \lambda = 4, -5$$

Case 1: For $\lambda = 4$ The eigenvector $X = \begin{bmatrix} x \\ y \end{bmatrix}$ is given by $[A - 4I]X = O$:

$$\begin{bmatrix} -4 - 4 & 2 \\ 4 & 3 - 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -8 & 2 \\ 4 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This yields the equation: $4x - y = 0 \implies y = 4x$. None of the provided options satisfy this relationship.

Case 2: For $\lambda = -5$ The eigenvector X is given by $[A + 5I]X = O$:

$$\begin{bmatrix} -4 + 5 & 2 \\ 4 & 3 + 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 \\ 4 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This yields the equation: $x + 2y = 0 \implies x = -2y$.

Checking the satisfyability for options:

- **Option (C):** $\begin{bmatrix} 2 \\ -1 \end{bmatrix} \implies 2 + 2(-1) = 0$ (Satisfied)

$$(C) \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

Q1.00.1 MCQ 2000 1M Ans: B ● ○ ○

Given the determinant:

$$\begin{vmatrix} 2 - \lambda & -1 & 0 & 0 \\ 0 & 3 - \lambda & 0 & 0 \\ 0 & 0 & -2 - \lambda & 0 \\ 0 & 0 & -1 & 4 - \lambda \end{vmatrix} = 0$$

$$\begin{aligned} \Rightarrow (2 - \lambda) \begin{vmatrix} 3 - \lambda & 0 & 0 \\ 0 & -2 - \lambda & 0 \\ 0 & -1 & 4 - \lambda \end{vmatrix} &= 0 \\ \Rightarrow (2 - \lambda)(3 - \lambda)[(-2 - \lambda)(4 - \lambda) - 0] &= 0 \\ \Rightarrow (2 - \lambda)(3 - \lambda)[-(2 + \lambda)(4 - \lambda)] &= 0 \\ \Rightarrow (2 - \lambda)(3 - \lambda)(2 + \lambda)(4 - \lambda) &= 0 \end{aligned}$$

Solving for λ , we get:

$$\lambda = 2, 3, -2, 4$$

The eigenvalues of the matrix are 2, -2, 3, and 4.

(B) 2, 3, -2, 4

Q1.98.1 MCQ 1998 1M Ans: D ● ○ ○

Given the matrix A :

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

The characteristic equation $|A - \lambda I| = 0$:

$$\begin{aligned} \begin{vmatrix} 0 - \lambda & 1 \\ 1 & 0 - \lambda \end{vmatrix} &= 0 \\ (-\lambda)(-\lambda) - (1)(1) &= 0 \\ \lambda^2 - 1 &= 0 \\ \lambda^2 &= 1 \\ \lambda &= 1, -1 \\ \text{(D) } 1, -1 \end{aligned}$$

Q1.94.1 MCQ 1994 1M Ans: A ● ○ ○

We know that:

$$\rho(A_{m \times n}) \leq \min\{m, n\}$$

Given the condition that $m < n$:

$$\begin{aligned} \min\{m, n\} &= m \\ \rho(A_{m \times n}) &\leq m \end{aligned}$$

Hence, the rank of an $m \times n$ matrix where $m < n$ cannot be more than m .

(A) m

Key Points

- Rank Property: The rank of a matrix is the maximum number of linearly independent row or column vectors.
- Order Constraint: For $A \in \mathbb{R}^{m \times n}$, $\text{rank}(A) \leq \min(m, n)$.

Q1.94.2 MCQ 1994 1M Ans: A ● ○ ○

Given the system of linear equations $AX = B$:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$$

$$[A|B] = \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 1 & 0 & -1 & 0 \\ 1 & -1 & 1 & 1 \end{array} \right]$$

Applying $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$[A|B] \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & -1 & -2 & -3 \\ 0 & -2 & 0 & -2 \end{array} \right]$$

Applying $R_3 \rightarrow R_3 - 2R_2$:

$$[A|B] \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & -1 & -2 & -3 \\ 0 & 0 & 4 & 4 \end{array} \right]$$

From the row-echelon form, $\rho(A) = \rho(A|B) = n = 3$, the system is consistent and has a unique solution.

(A) Unique solution

SOLUTIONS — Linear Algebra

**EE****Electrical Engineering****PrepFusion**

Q1.26.1 MSQ 2026 1M Ans: A,B,D ● ○ ○

Given that A and B are $n \times n$ matrices with a common eigenvalue $\lambda = 2$ and the same corresponding nonzero eigenvector x .
From the definition of eigenvalues and eigenvectors:

$$Ax = 2x \implies (A - 2I)x = 0$$

$$Bx = 2x \implies (B - 2I)x = 0$$

Since x is a nonzero vector, the matrices $(A - 2I)$ and $(B - 2I)$ must be singular. Therefore:

$$\text{Determinant}(A - 2I) = 0$$

$$\text{Determinant}(B - 2I) = 0$$

This confirms options (A) and (B) are correct.

Now, consider the matrix $(A + B - 4I)$ acting on the same eigenvector x :

$$(A + B - 4I)x = Ax + Bx - 4Ix$$

$$(A + B - 4I)x = 2x + 2x - 4x$$

$$(A + B - 4I)x = 0$$

Since $(A + B - 4I)x = 0$ for a nonzero x , the matrix $(A + B - 4I)$ is singular:

$$\text{Determinant}(A + B - 4I) = 0$$

This confirms option (D) is correct.

For option (c), $(A + B - 2I)x = 2x + 2x - 2x = 2x \neq 0$, so it is not necessarily singular.

$$(A) \text{ Determinant } (A - 2I) = 0, (B) \text{ Determinant } (B - 2I) = 0, (D) \text{ Determinant } (A + B - 4I) = 0$$

Q1.26.2 NAT 2026 1M Ans: 0 ● ● ○

Method 1

Given that A is an $m \times m$ skew-symmetric matrix, $A^T = -A$. Let the scalar $Q = x^T Ax$. Since the transpose of a scalar is equal to itself:

$$Q = Q^T = (x^T Ax)^T$$

Applying the reversal law of transposes $(ABC)^T = C^T B^T A^T$:

$$Q = x^T A^T (x^T)^T = x^T A^T x$$

Substituting the skew-symmetric condition $A^T = -A$:

$$Q = x^T (-A)x = -x^T Ax = -Q$$

$$2Q = 0 \implies Q = 0$$

Method 2

We can verify the result by considering a specific 3×3 skew-symmetric matrix A where $a_{ij} = -a_{ji}$ and diagonal elements are zero:

$$A = \begin{bmatrix} 0 & -a & -b \\ a & 0 & -c \\ b & c & 0 \end{bmatrix}$$

For a vector $x = [x_1 \ x_2 \ x_3]^T$, the quantity $x^T A x$ is:

$$x^T A x = [x_1 \ x_2 \ x_3] \begin{bmatrix} 0 & -a & -b \\ a & 0 & -c \\ b & c & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$x^T A x = [x_1 \ x_2 \ x_3] \begin{bmatrix} -ax_2 - bx_3 \\ ax_1 - cx_3 \\ bx_1 + cx_2 \end{bmatrix}$$

$$x^T A x = x_1(-ax_2 - bx_3) + x_2(ax_1 - cx_3) + x_3(bx_1 + cx_2)$$

$$x^T A x = -ax_1x_2 - bx_1x_3 + ax_1x_2 - cx_2x_3 + bx_1x_3 + cx_2x_3 = 0$$

$$\boxed{0}$$

Key Points

- A matrix is skew-symmetric if $A^T = -A$, which implies $a_{ij} = -a_{ji}$ and $a_{ii} = 0$.
- The quadratic form $x^T A x$ of any real skew-symmetric matrix is always zero.
- This is because the symmetric part of a skew-symmetric matrix is the zero matrix.

Q1.26.3 MCQ 2026 2M Ans: A ● ○ ○

Let A be a matrix formed by the given collection of p column vectors, where each vector has n entries:

$$A = [a_{ij}]_{n \times p}$$

In an n -dimensional space, the maximum number of linearly independent vectors is n .

If $p > n$, then the number of column vectors is greater than the dimension of the space. According to the fundamental theorem of linear algebra, any set of vectors in $\mathbb{R}^n$ with more than n elements must be linearly dependent.

$\therefore$ Some column vectors must be linearly dependent.

(A) If $p > n$, then the column vectors must be linearly dependent

Key Points

- The dimension of a vector space is the maximum number of linearly independent vectors it can contain.
- If a set of vectors $S = \{v_1, v_2, \dots, v_p\}$ belongs to $\mathbb{R}^n$ and $p > n$, the set is always linearly dependent.
- Rank of a matrix $A_{n \times p}$ is $\leq \min(n, p)$. If $p > n$, the columns cannot all be linearly independent as the maximum rank is n .

Q1.26.4 MSQ 2026 2M Ans: A,B,D ● ● ○

Given that $A_{n \times n}$ is an orthogonal matrix. By definition of an orthogonal matrix:

$$A^T A = A A^T = I$$

Checking the given statements:

(a) Since $A A^T = I$, taking the determinant on both sides:

$$|A A^T| = |I| \implies |A| \cdot |A^T| = 1$$

Since $|A| = |A^T|$, we have $|A|^2 = 1 \implies |A| = \pm 1$. Thus, statement (a) is true and A^{-1} exists.

(b) For an eigenvalue λ of A , the eigenvalue of A^{-1} is $\frac{1}{\lambda}$. For an orthogonal matrix, $A^{-1} = A^T$. The eigenvalues of A and A^T are the same.

$$\therefore \lambda = \frac{1}{\lambda} \implies \lambda^2 = 1 \implies |\lambda| = 1$$

Thus, statement (b) is true.

(d) Consider the inner product $(Ax)^T(Ay)$:

$$(Ax)^T(Ay) = x^T A^T Ay$$

$$(Ax)^T(Ay) = x^T Iy = x^T y$$

$$\|Ax\| = \sqrt{(Ax)^T(Ax)} = \sqrt{x^T A^T Ax} = \sqrt{x^T x} = \|x\|$$

Thus, statement (d) is true. Note that statement (c) contradicts the property $(Ax)^T(Ay) = x^T y$, making it false.

(A) The value of the determinant of A is either +1 or -1

(B) The eigenvalues of A have modulus 1

(D) $\|Ax\| = \|x\|$, for all $x \in R^n$, where $\|x\|$ denotes the Euclidean norm of x , and $(Ax)^T(Ay) = x^T y$, for all $x, y \in R^n$

Q1.26.5 **MSQ** **2026** **2M** **Ans: A,B,C** ☒ ☐ ☐

Given that the system $Ax = b$ has a unique solution, where A is an $n \times n$ matrix.

For a square system $Ax = b$ to have a unique solution:

$$|A| \neq 0$$

This implies that matrix A is non-singular. Checking the options:

(a) Since $|A| \neq 0$, the inverse A^{-1} exists. Thus, option (A) is true.

(b) If $|A| \neq 0$, then $|A^m| = |A|^m \neq 0$ for any $m = 1, 2, 3, \dots$. Since A^m is also non-singular, the system $A^m x = b$ will also have a unique solution. Thus, option (B) is true.

(c) For a non-singular matrix A , the rank is equal to its order:

$$\rho(A) = n$$

Since A^m is also non-singular for $m = 1, 2, 3, \dots$:

$$\rho(A^m) = n$$

$$\therefore \rho(A) = \rho(A^m)$$

Thus, option (C) is true.

(d) For a unique solution in a system $Ax = b$, the rank of the coefficient matrix must be equal to the rank of the augmented matrix $C = [A \mid b]$:

$$\rho(A) = \rho(C) = n$$

If $\rho(A) < \rho(C)$, the system is inconsistent and has no solution. Thus, option (D) is false.

(A) A^{-1} exists

(B) The system of equations $A^m x = b$ also has a unique solution for $m = 1, 2, 3, \dots$

(C) $\text{rank}(A) = \text{rank}(A^m)$, for $m = 1, 2, 3, \dots$

Key Points

- A system $Ax = b$ has a unique solution if and only if $|A| \neq 0$ (for square matrices).
- If $|A| \neq 0$, then $\text{rank}(A) = n$.
- Rouché-Capelli Theorem: A system is consistent if $\text{rank}(A) = \text{rank}(A | b)$. It is unique if this rank equals the number of variables.

Mistakes to be avoided

- Do not assume $A^m x = b$ has a unique solution without first verifying that A is non-singular. If A is singular, A^m is also singular.
- A system can be consistent but have infinitely many solutions if $\text{rank}(A) < n$.

Q1.25.1 MCQ 2025 1M Ans: B ☒ ☐ ☐

We know that the eigenvectors of a real symmetric matrix corresponding to different eigenvalues are always pair-wise orthogonal.

So, for the given two eigenvectors V_1 and V_2 :

$$V_1^T V_2 = 0 \quad (\text{or}) \quad V_2^T V_1 = 0$$

(B) $v_1^T v_2 = 0$

Q1.25.2 MCQ 2025 1M Ans: B ☒ ☐ ☐

Consider the augmented matrix $(A|B)$ for the given system of equations:

$$(A|B) = \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1/3 \\ -1 & -1 & -1 & -1/3 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

Step 1: $R_2 \rightarrow R_2 + R_1$

$$(A|B) \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1/3 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

Step 2: $R_2 \leftrightarrow R_3$

$$(A|B) \sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1/3 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

From the final matrix, we can observe: $\rho(A) = \rho(A|B) = 2 < 3 = \text{number of variables}$, the system is consistent and possesses infinitely many solutions.

(B) infinite no. of solutions

Q1.25.3 MCQ 2025 1M Ans: A ☒ ☐ ☐

Given that:

$$P = \begin{bmatrix} 2 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$P^2 = P \cdot P = \begin{bmatrix} 2 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$P^2 = \begin{bmatrix} (4-1) & (2+0) & 0 \\ (-2+0) & (-1+0) & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Evaluating the given options: For option (a):

$$2P - I = 2 \begin{bmatrix} 2 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$2P - I = \begin{bmatrix} 4 & 2 & 0 \\ -2 & 0 & 0 \\ 0 & 0 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Comparing the results, we find $P^2 = 2P - I$.

(A) $2P - I$

Q1.24.1 MCQ 2024 1M Ans: C ☒ ☐ ☐

Given that we need to find which matrix has an inverse (is invertible), we check the determinant $|A|$ for each option. A matrix A is invertible if and only if $|A| \neq 0$.

Option (A):

$$|A| = \begin{vmatrix} 1 & 4 & 8 \\ 0 & 4 & 2 \\ 0.5 & 2 & 4 \end{vmatrix}$$

$$|A| = 1(16 - 4) - 0(16 - 16) + 0.5(8 - 32)$$

$$|A| = 12 - 0 + 0.5(-24) = 12 - 12 = 0$$

Since $|A| = 0$, A^{-1} does not exist.

Option (B):

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 2 & 9 \end{vmatrix}$$

Since the first row R_1 and the second row R_2 are proportional ($R_2 = 2R_1$), the determinant is identically zero:

$$|A| = 0$$

Therefore, A is not an invertible matrix.

Option (C):

$$|A| = \begin{vmatrix} 1 & 4 & 8 \\ 0 & 4 & 2 \\ 1 & 2 & 4 \end{vmatrix}$$

$$|A| = 1(16 - 4) - 0(16 - 16) + 1(8 - 32)$$

$$|A| = 12 - 24 = -12 \neq 0$$

Since $|A| \neq 0$, A^{-1} exists. Thus, A is an invertible matrix.

Option (D):

$$|A| = \begin{vmatrix} 1 & 4 & 8 \\ 0 & 4 & 2 \\ 3 & 12 & 24 \end{vmatrix}$$

Since the first row R_1 and the third row R_3 are proportional ($R_3 = 3R_1$), the determinant is zero:

$$|A| = 0$$

Hence, A^{-1} does not exist.

$$(C) \begin{bmatrix} 1 & 4 & 8 \\ 0 & 4 & 2 \\ 1 & 2 & 4 \end{bmatrix}$$

Key Points

- A square matrix A is invertible (non-singular) if and only if its determinant is non-zero ($|A| \neq 0$).
- If any two rows or columns of a matrix are identical or proportional, its determinant is immediately 0.

Q1.24.2 NAT 2024 1M Ans: 29 ● ○ ○

Given the matrix multiplication to determine matrix A :

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

Performing row-by-column multiplication:

$$A = \begin{bmatrix} (1)(1) + (2)(3) & (1)(2) + (2)(4) \\ (3)(1) + (4)(3) & (3)(2) + (4)(4) \end{bmatrix} = \begin{bmatrix} 1 + 6 & 2 + 8 \\ 3 + 12 & 6 + 16 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix}$$

Sum of eigenvalues of A = Trace of A

$$\text{Trace of } A = 7 + 22 = 29$$

Q1.23.1 MCQ 2023 1M Ans: A ● ○ ○

Given the column vectors:

$$u = \begin{bmatrix} 4 \\ 3 \end{bmatrix}, \quad v = \begin{bmatrix} 5 \\ 0 \end{bmatrix}$$

The linear transformation is defined by the matrix equation $Au = v$. To find the correct matrix A , we evaluate the transformation using the matrix provided in option (a).

Consider the matrix from option (a):

$$A = \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ -\frac{3}{5} & \frac{4}{5} \end{bmatrix}$$

$$Au = \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ -\frac{3}{5} & \frac{4}{5} \end{bmatrix} \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

$$Au = \begin{bmatrix} \left(\frac{4}{5} \cdot 4\right) + \left(\frac{3}{5} \cdot 3\right) \\ \left(\frac{-3}{5} \cdot 4\right) + \left(\frac{4}{5} \cdot 3\right) \end{bmatrix} = \begin{bmatrix} \frac{16}{5} + \frac{9}{5} \\ \frac{-12}{5} + \frac{12}{5} \end{bmatrix}$$

$$Au = \begin{bmatrix} \frac{25}{5} \\ 0 \\ \frac{0}{5} \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \end{bmatrix} = v$$

Since the transformation $Au = v$ is perfectly satisfied, option (A) is the correct matrix.

$$(A) \begin{bmatrix} \frac{4}{5} & \frac{3}{5} \\ \frac{-3}{5} & \frac{4}{5} \end{bmatrix}$$

Q1.22.1 MCQ 2022 1M Ans: B ☒ ☐ ☐

Given that the matrix $A = (a_{ij})_{3 \times 3}$, where the elements are defined by the relation:

$$a_{ij} = (i - j)^3$$

We check the relation between a_{ij} and a_{ji} to determine the type of the matrix.

By substituting the indices for the transposed element a_{ji} , we get:

$$a_{ji} = (j - i)^3$$

Factoring out a negative sign from the base:

$$a_{ji} = [-(i - j)]^3 = -(i - j)^3$$

Since $a_{ij} = (i - j)^3$, we can rewrite this as:

$$a_{ji} = -a_{ij} \implies a_{ij} = -a_{ji}$$

This identity holds true for all indices $1 \leq i, j \leq 3$. Therefore, $A_{3 \times 3}$ is a skew-symmetric matrix.

Hence, option (B) is correct.

(B) skew-symmetric

Key Points

- Skew-Symmetric Matrix Property:** A square matrix A is skew-symmetric if and only if $A^T = -A$, which translates entry-wise to $a_{ij} = -a_{ji}$ for all i, j .
- Diagonal Elements Constraint:** For any skew-symmetric matrix, when $i = j$, we have $a_{ii} = -a_{ii} \implies 2a_{ii} = 0 \implies a_{ii} = 0$. The principal diagonal elements must all be zero.
- Odd Powers of Differences:** Any element function based on an odd power of index differences, such as $a_{ij} = (i - j)^{2n+1}$, always yields a skew-symmetric matrix layout.

Mistakes to be avoided

- Do not confuse symmetric matrices ($a_{ij} = a_{ji}$) with skew-symmetric matrices ($a_{ij} = -a_{ji}$).

Q1.22.2 MCQ 2022 2M Ans: C ● ● ○

If λ is an eigenvalue of a square matrix A corresponding to the eigenvector v , then by definition:

$$Av = \lambda v$$

Using the Taylor series expansion for the matrix exponential function e^A :

$$e^A = I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{A^n}{n!}$$

$$e^A v = \left(I + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots \right) v$$

Since $A^n v = \lambda^n v$ for any positive integer n , we can substitute the terms:

$$e^A v = Iv + Av + \frac{A^2 v}{2!} + \frac{A^3 v}{3!} + \dots$$

$$e^A v = 1v + \lambda v + \frac{\lambda^2}{2!} v + \frac{\lambda^3}{3!} v + \dots$$

$$e^A v = \left(1 + \lambda + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \dots \right) v$$

The scalar expression inside the parentheses is the standard Maclaurin series for e^λ :

$$e^A v = e^\lambda v$$

Therefore, if λ is an eigenvalue of A , then e^λ is an eigenvalue of e^A , and the corresponding eigenvector v remains identical.

(C) Both the statements are correct.

Key Points

- **Matrix Function Property:** If λ is an eigenvalue of A with eigenvector v , then for any polynomial or analytic function $f(A)$, the eigenvalue is $f(\lambda)$ and the eigenvector is still v .
- **Common Transformations:**
 - Matrix $A^n \implies$ Eigenvalue λ^n
 - Matrix $A^{-1} \implies$ Eigenvalue λ^{-1} (provided A is non-singular)
 - Matrix $kA \implies$ Eigenvalue $k\lambda$

Q1.22.3 MCQ 2022 2M Ans: A ● ● ○

Given the 3×3 matrix:

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & -2 \\ 0 & 1 & 1 \end{bmatrix}$$

The characteristic equation for a 3×3 matrix is given by the formula:

$$\lambda^3 - S_1 \lambda^2 + S_2 \lambda - |A| = 0$$

Where:

- $S_1 = \text{Trace of } A = 1 + 4 + 1 = 6$
- $S_2 = \text{Sum of minors of principal diagonal elements}$

$$S_2 = \begin{vmatrix} 4 & -2 \\ 1 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} + \begin{vmatrix} 1 & 0 \\ 0 & 4 \end{vmatrix}$$

$$S_2 = (4 + 2) + (1 - 0) + (4 - 0) = 6 + 1 + 4 = 11$$

- $|A| = \text{Determinant of } A = 1(4 + 2) - 0 + 0 = 6$

Substituting these values into the characteristic equation:

$$\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$$

According to the Cayley-Hamilton theorem, every square matrix satisfies its own characteristic equation. Replacing λ with matrix A :

$$A^3 - 6A^2 + 11A - 6I = 0$$

$$A^2 - 6A + 11I - 6A^{-1} = 0$$

$$6A^{-1} = A^2 - 6A + 11I$$

Comparing this result with the given expression $6A^{-1} = A^2 + cA + dI$:

$$c = -6, \quad d = 11$$

$$c + d = (-6) + (11) = 5$$

(A) 5

Q1.21.1 MCQ 2021 1M Ans: B ☒ ☐ ☐

Let the given 2×2 matrix be:

$$A = \begin{bmatrix} p & q \\ q & -p \end{bmatrix}$$

$$|A - \lambda I| = 0$$

$$\lambda^2 - \text{Trace}(A)\lambda + |A| = 0$$

- $\text{Trace}(A) = p + (-p) = 0$
- $|A| = \det(A) = (p)(-p) - (q)(q) = -p^2 - q^2 = -(p^2 + q^2)$

Substituting these expressions back into the characteristic equation:

$$\lambda^2 - (0)\lambda + \left(-(p^2 + q^2)\right) = 0$$

$$\lambda^2 - (p^2 + q^2) = 0$$

$$\lambda^2 - 1 = 0$$

$$\lambda^2 = 1 \implies \lambda = \pm 1$$

Therefore, $\lambda = 1, -1$ are the eigenvalues of the matrix.

(B) 1 and -1

Q1.21.2 NAT 2021 2M Ans: 1 ● ● ○

Given that A^5 is a null matrix, i.e.,

$$A^5 = O$$

By definition, all the eigenvalues (λ) of a nilpotent matrix are identically zero:

$$\lambda = 0$$

We use the property of matrix transformations on eigenvalues. If λ is an eigenvalue of a square matrix A , then the corresponding eigenvalue of the shifted matrix $A + I$ is given by:

$$\lambda_{\text{new}} = \lambda + 1$$

Substituting the eigenvalue $\lambda = 0$ into this relation:

$$\lambda_{\text{new}} = 0 + 1 = 1$$

Since this applies to all eigenvalues of the matrix, every eigenvalue of the matrix $A + I$ is equal to 1.

$$|A + I| = \text{Product of eigenvalues of } (A + I)$$

$$|A + I| = 1 \cdot 1 \dots 1 = 1$$

$$1$$

Key Points

- **Nilpotent Matrix Definition:** A square matrix A is nilpotent if $A^k = O$ for some positive integer k . The smallest such k is called the index of nilpotency.
- **Nilpotent Eigenvalues Theorem:** The characteristic equation of any nilpotent matrix reduces to $\lambda^n = 0$, meaning all its eigenvalues are strictly 0.
- **Determinant Property:** The determinant of a matrix equals the product of its eigenvalues ($\det(M) = \prod \lambda_i$).
- **Identity Shift:** Shifting a matrix by $+I$ adds exactly 1 to each of its individual eigenvalues.

Mistakes to be avoided

- Do not assume that $\det(A + I) = \det(A) + \det(I)$, as the determinant operator is non-linear and does not distribute over addition.
- Remember that while all eigenvalues of a nilpotent matrix are zero, the matrix itself does not have to be the zero matrix.

Q1.20.1 MCQ 2020 2M Ans: MTA ● ● ●

Given the decomposition of a symmetric matrix into a lower triangular matrix and its transpose (Cholesky-like factorization form):

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a & 0 & 0 \\ b & c & 0 \\ d & e & f \end{bmatrix} \begin{bmatrix} a & 0 & 0 \\ b & c & 0 \\ d & e & f \end{bmatrix}^T$$

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a & 0 & 0 \\ b & c & 0 \\ d & e & f \end{bmatrix} \begin{bmatrix} a & b & d \\ 0 & c & e \\ 0 & 0 & f \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & 3 \\ 3 & 2 & 1 \\ 3 & 1 & 7 \end{bmatrix} = \begin{bmatrix} a^2 & ab & ad \\ ab & b^2 + c^2 & bd + ce \\ ad & bd + ce & d^2 + e^2 + f^2 \end{bmatrix}$$

Comparing corresponding entry terms on both sides, we establish the system of equations: $a^2 = 2$, $ab = 3$, $ad = 3$, $b^2 + c^2 = 2$, $bd + ce = 1$, $d^2 + e^2 + f^2 = 7$

Solving these sequentially:

- From $a^2 = 2 \Rightarrow a = \pm\sqrt{2}$ (purely real)
- From $ab = 3 \Rightarrow b = \pm\frac{3}{\sqrt{2}}$ (purely real)
- From $ad = 3 \Rightarrow d = \pm\frac{3}{\sqrt{2}}$ (purely real)
- From $b^2 + c^2 = 2 \Rightarrow \left(\pm\frac{3}{\sqrt{2}}\right)^2 + c^2 = 2 \Rightarrow \frac{9}{2} + c^2 = 2 \Rightarrow c^2 = 2 - \frac{9}{2} = -\frac{5}{2}$
 $\Rightarrow c = \pm\sqrt{\frac{5}{2}}i$ (purely imaginary)
- From $bd + ce = 1 \Rightarrow \left(\pm\frac{3}{\sqrt{2}}\right)\left(\pm\frac{3}{\sqrt{2}}\right) + ce = 1 \Rightarrow \frac{9}{2} + ce = 1 \Rightarrow ce = 1 - \frac{9}{2} = -\frac{7}{2}$
Substituting c :
 $\left(\pm\sqrt{\frac{5}{2}}i\right)e = -\frac{7}{2} \Rightarrow e = \pm\frac{7}{\sqrt{10}}i$ (purely imaginary)
- From $d^2 + e^2 + f^2 = 7 \Rightarrow \left(\pm\frac{3}{\sqrt{2}}\right)^2 + \left(\pm\frac{7}{\sqrt{10}}i\right)^2 + f^2 = 7 \Rightarrow \frac{9}{2} - \frac{49}{10} + f^2 = 7$
 $\frac{45 - 49}{10} + f^2 = 7 \Rightarrow -\frac{4}{10} + f^2 = 7 \Rightarrow f^2 = 7 + \frac{2}{5} = \frac{37}{5}$
 $\Rightarrow f = \pm\sqrt{\frac{37}{5}}$ (purely real)

The values of the parameters are:

$$a, b, d, f \in \mathbb{R} \quad (4 \text{ real elements}) \quad c, e \notin \mathbb{R} \quad (2 \text{ imaginary elements})$$

The structural lower triangular representation matrix contains 6 elements total (a, b, c, d, e, f along with 3 fixed upper zeros). The non-zero entries are a, b, c, d, e, f . If we consider the sign choices ($\pm$), the total number of unique, possible real elements obtained across the solution sets is 8 (two real alternatives for each of the four real variables a, b, d, f).

MTA

Key Points

- **Cholesky-like Decomposition:** For a real symmetric matrix, standard Cholesky decomposition ($A = LL^T$) requires the matrix to be positive definite to ensure all diagonal elements are purely real.
- **Complex Entries in Factorization:** If the given matrix is symmetric but indefinite, matching terms under the format LL^T forces some internal variance terms (c^2, e^2) to be negative, creating purely imaginary components.

Q1.19.1 MCQ 2019 1M Ans: D ● ○ ○

If λ is an eigenvalue of a square matrix M , then λ^n is the corresponding eigenvalue of the matrix M^n .
Applying this property for $n = 2$:

$$\text{Eigenvalues of } M^2 = (4)^2 \text{ and } (9)^2$$

$$\text{Eigenvalues of } M^2 = 16 \text{ and } 81$$

(D) 16 and 81

Q1.19.2 MCQ 2019 2M Ans: C ● ○ ○

$$M^{-1}M = I$$

Let the row vectors of M^{-1} be represented by u_1^T and u_2^T , and let the column vectors of M be represented by v_1 and v_2 .
Substituting these block structures into the identity equation gives:

$$\begin{bmatrix} u_1^T \\ u_2^T \end{bmatrix} \begin{bmatrix} v_1 & v_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} u_1^T v_1 & u_1^T v_2 \\ u_2^T v_1 & u_2^T v_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Equating the corresponding entry terms on both sides yields:

$$u_1^T v_1 = 1, \quad u_1^T v_2 = 0, \quad u_2^T v_1 = 0, \quad \text{and} \quad u_2^T v_2 = 1$$

Therefore, both structural statements are completely correct.

(C) Both the statements are true.

Q1.18.1 NAT 2018 1M Ans: 5.5 ● ○ ○

Let λ_1, λ_2 be two eigenvalues of a 2×2 matrix A .
Given:

$$\text{trace}(A) = 4 \quad \text{and} \quad \text{trace}(A^2) = 5$$

Using the trace identity theorem (sum of eigenvalues equals the trace of a matrix):

$$\lambda_1 + \lambda_2 = 4$$

$$\lambda_1^2 + \lambda_2^2 = 5$$

We apply the standard algebraic expansion identity:

$$(\lambda_1 + \lambda_2)^2 = \lambda_1^2 + \lambda_2^2 + 2\lambda_1\lambda_2$$

$$(4)^2 = 5 + 2\lambda_1\lambda_2$$

$$16 = 5 + 2\lambda_1\lambda_2$$

$$2\lambda_1\lambda_2 = 16 - 5 = 11$$

$$\lambda_1\lambda_2 = \frac{11}{2} = 5.5$$

$$|A| = \text{product of eigenvalues} = \lambda_1\lambda_2 = 5.5$$

Q1.18.2 NAT 2018 2M Ans: 0.9 to 1.1 ● ● ○

Given the 3×3 matrix:

$$A = \begin{bmatrix} 1 & 0 & -1 \\ -1 & 2 & 0 \\ 0 & 0 & -2 \end{bmatrix}$$

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 1-\lambda & 0 & -1 \\ -1 & 2-\lambda & 0 \\ 0 & 0 & -2-\lambda \end{vmatrix} = 0$$

Expanding along the third row:

$$(-2-\lambda) \begin{vmatrix} 1-\lambda & 0 \\ -1 & 2-\lambda \end{vmatrix} = 0$$

$$(-2-\lambda) [(1-\lambda)(2-\lambda) - 0] = 0$$

$$-(2+\lambda)(\lambda^2 - 3\lambda + 2) = 0$$

$$-(\lambda^3 - 3\lambda^2 + 2\lambda + 2\lambda^2 - 6\lambda + 4) = 0$$

$$\lambda^3 - \lambda^2 - 4\lambda + 4 = 0$$

$$A^3 - A^2 - 4A + 4I = O \quad \dots\dots (1)$$

Given:

$$B = A^3 - A^2 - 4A + 5I \quad \dots\dots (2)$$

$$B = (A^3 - A^2 - 4A + 4I) + I$$

Substituting the zero matrix result from equation (1) into this expression:

$$B = O + I \implies B = I$$

Taking Det on both sides:

$$|B| = |I| = \boxed{1}$$

Q1.17.1 MCQ 2017 1M Ans: C ● ○ ○

If the eigenvalues of a real symmetric matrix are distinct, then their corresponding eigenvectors are mutually orthogonal. The algebraic condition for two non-zero vectors X_1 and X_2 to be orthogonal is that their inner product (dot product) must be zero:

$$X_1^T X_2 = 0 \quad \text{or} \quad X_2^T X_1 = 0 \quad \text{or} \quad X_1 \bullet X_2 = 0$$

Given the known eigenvector corresponding to one of the distinct eigenvalues:

$$X_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

We test the dot product of X_1 with the vector given in option (C):

$$X_1 \bullet X_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \bullet \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = (1)(1) + (0)(0) + (1)(-1)$$

$$X_1 \bullet X_2 = 1 + 0 - 1 = 0$$

Since the inner product is exactly zero, the vector in option (C) is orthogonal to X_1 . Therefore, $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$ is another required eigenvector of the given real symmetric matrix A .

$$(C) \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Q1.17.2 MCQ 2017 2M Ans: A ● ○ ○

The characteristic equation of a square matrix A is determined by finding the roots of its characteristic determinant:

$$|A - \lambda I| = 0$$

By simplifying, we get:

$$-\lambda (\lambda^2 + 4\lambda + 3) = 0$$

$$-\lambda (\lambda^2 + 3\lambda + \lambda + 3) = 0$$

$$-\lambda [\lambda(\lambda + 3) + 1(\lambda + 3)] = 0$$

$$-\lambda(\lambda + 1)(\lambda + 3) = 0$$

Therefore, $\lambda = 0, -1, -3$ are the required eigenvalues of the matrix.

$$(A) (0, -1, -3)$$

Q1.16.1 NAT 2016 1M Ans: 3 ● ○ ○

Let the given 3×3 matrix be:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$|A - \lambda I| = 0$$

$$\lambda^3 - \text{Trace}(A)\lambda^2 + S_2\lambda - |A| = 0$$

- $\text{Trace}(A) = \text{Sum of diagonal elements} = 1 + 1 + 1 = 3$
- Since all rows are identical, the rank of the matrix is 1. Consequently, the determinant $|A| = 0$, and the sum of principal minors $S_2 = 0$.

Substituting these structural properties back into the polynomial equation:

$$\lambda^3 - 3\lambda^2 = 0$$

$$\lambda^2(\lambda - 3) = 0$$

$$\lambda = 3, 0, 0$$

Therefore, the only non-zero eigenvalue of the matrix $A_{3 \times 3}$ is $\boxed{3}$.

Q1.16.2 MCQ 2016 2M Ans: A ● ● ○

If λ is an eigenvalue of a square matrix A , then the corresponding eigenvalue for any matrix polynomial transformation $f(A) = A^2 - 3A + 4I$ is given by evaluating the scalar polynomial $f(\lambda) = \lambda^2 - 3\lambda + 4$.

We are given that $\lambda_1 = 1$ and $\lambda_2 = -2$ are the distinct eigenvalues of the 2×2 matrix $A_{2 \times 2}$.

Evaluating the transformation polynomial for each eigenvalue sequentially:

- For $\lambda_1 = 1$:

$$f(1) = (1)^2 - 3(1) + 4 = 1 - 3 + 4 = 2$$

- For $\lambda_2 = -2$:

$$f(-2) = (-2)^2 - 3(-2) + 4 = 4 + 6 + 4 = 14$$

Thus, the corresponding eigenvalues of the transformed matrix $A^2 - 3A + 4I$ are 2 and 14.

Furthermore, the directional orientation of the eigenvectors remains entirely unchanged. This invariant behavior occurs because any matrix polynomial of the form $aI + bA + cA^2$ (where a, b, c are scalar constants) commutes directly with A , thereby preserving the exact same structural eigenvectors (x_1, x_2) associated with the original eigenvalues.

(A) 2, 14; x_1, x_2

Key Points

- Spectral Mapping Theorem:** If a matrix A has an eigenvalue λ , any matrix polynomial $P(A)$ will possess the corresponding mapped eigenvalue $P(\lambda)$.
- Eigenvector Invariance:** Transforming a matrix via an analytic polynomial expansion scales its eigenvalues but leaves its underlying eigenvectors completely unchanged.

Q1.16.3 MCQ 2016 2M Ans: B ● ● ○

$$\rho(A^T) = \rho(A) = 2$$

A and A^T have two linearly independent rows and columns.

$$\rho(A^T A) \leq \min \{ \rho(A^T), \rho(A) \}$$

$$\rho(A^T A) \leq \min \{ 2, 2 \} = 2$$

$A^T A$ also has two independent rows and columns.

$$\rho(A^T A) = \rho(A)$$

Since $\rho(A) = 2$, it follows directly that:

$$\text{Rank} [A^T A] = 2$$

(B) Rank of $A^T A$ is equal to 2.

Q1.16.4 MCQ 2016 1M Ans: D ● ○ ○

Given the matrix equation:

$$P^3 = P$$

$$P^3 - P = O$$

By the properties of matrix eigenvalues:

$$\lambda^3 - \lambda = 0$$

$$\lambda(\lambda^2 - 1) = 0$$

$$\lambda(\lambda - 1)(\lambda + 1) = 0$$

$$\lambda = 0, \quad \lambda = 1, \quad \text{and} \quad \lambda = -1$$

Therefore, the possible eigenvalues of matrix P are 0, 1, -1.

$$(D) \ 0, 1, -1$$

Q1.16.5 **MCQ** **2016** **2M** **Ans: D** ● ● ●

Given the matrix:

$$P = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

We are given a transformation mapping the vector $\begin{bmatrix} x \\ y \end{bmatrix}$ in set S to a new vector $\begin{bmatrix} a \\ b \end{bmatrix}$ via the relation:

$$\begin{bmatrix} a \\ b \end{bmatrix} = P \begin{bmatrix} x \\ y \end{bmatrix}$$

The target condition defines a unit circle in the transformed coordinate plane:

$$a^2 + b^2 = 1$$

Method 1

Expanding the matrix transformation explicitly into a system of linear algebraic equations:

$$a = 3x + y$$

$$b = x + 3y$$

Substituting these algebraic expressions directly into the unit circle constraint equation ($a^2 + b^2 = 1$):

$$(3x + y)^2 + (x + 3y)^2 = 1$$

$$(9x^2 + 6xy + y^2) + (x^2 + 6xy + 9y^2) = 1$$

$$10x^2 + 12xy + 10y^2 = 1$$

This second-degree homogeneous polynomial represents a conic section. To analyze its primary geometric features, we examine the general equation form $Ax^2 + Bxy + Cy^2 = 1$:

$$A = 10, \quad B = 12, \quad C = 10$$

Computing the discriminant to identify the specific conic geometry:

$$B^2 - 4AC = (12)^2 - 4(10)(10) = 144 - 400 = -256 < 0$$

Since the discriminant is strictly negative and $A = C$, the conic section represents a rotated ellipse.

To find the orientation of its major and minor axes, we can evaluate the eigenvalues and eigenvectors of the symmetric matrix associated with the quadratic form:

$$Q = \begin{bmatrix} 10 & 6 \\ 6 & 10 \end{bmatrix}$$

The characteristic equation is $\det(Q - \lambda I) = 0$:

$$(10 - \lambda)^2 - 36 = 0 \implies 10 - \lambda = \pm 6$$

$\lambda_1 = 16 \implies$ associated with the minor axis (higher curvature)

$\lambda_2 = 4 \implies$ associated with the major axis (lower curvature)

Finding the eigenvector for the maximum eigenvalue $\lambda_1 = 16$ to identify the direction of the minor axis:

$$(Q - 16I)X = O \implies \begin{bmatrix} -6 & 6 \\ 6 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \implies -x_1 + x_2 = 0 \implies X_{\text{minor}} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Thus, the geometric locus of set S defines an ellipse with its minor axis oriented along the vector direction $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

(D) an ellipse with minor axis along $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Method 2

Let $X = \begin{bmatrix} x \\ y \end{bmatrix}$ and $Y = \begin{bmatrix} a \\ b \end{bmatrix}$. The given transformation is:

$$Y = PX$$

The equation of the unit circle $a^2 + b^2 = 1$ can be expressed in vector-matrix layout as:

$$Y^T Y = 1$$

Substituting $Y = PX$ into this vector product:

$$(PX)^T (PX) = 1$$

$$X^T P^T P X = 1$$

Since P is a real symmetric matrix ($P^T = P$), we compute the core matrix $P^T P = P^2$:

$$P^2 = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} = \begin{bmatrix} 9+1 & 3+3 \\ 3+3 & 1+9 \end{bmatrix} = \begin{bmatrix} 10 & 6 \\ 6 & 10 \end{bmatrix}$$

The transformed quadratic form equation becomes:

$$X^T \begin{bmatrix} 10 & 6 \\ 6 & 10 \end{bmatrix} X = 1$$

The eigenvalues of this matrix determine the principal axes of the conic section:

- **Sum of Eigenvalues** ($\lambda_1 + \lambda_2$): $\text{Trace}(P^2) = 10 + 10 = 20$
- **Product of Eigenvalues** ($\lambda_1 \lambda_2$): $\det(P^2) = (10)(10) - (6)(6) = 100 - 36 = 64$

Solving the characteristic polynomial $\lambda^2 - 20\lambda + 64 = 0 \implies (\lambda - 16)(\lambda - 4) = 0$, we find $\lambda = 16$ and $\lambda = 4$.

Since both eigenvalues are positive and distinct ($\lambda_1 \neq \lambda_2 > 0$), the equation represents an Ellipse.

- The major axis lies along the eigenvector of the *smallest* eigenvalue ($\lambda = 4$).
- The minor axis lies along the eigenvector of the *largest* eigenvalue ($\lambda = 16$).

Let us find the eigenvector for the minor axis ($\lambda = 16$):

$$(P^2 - 16I)X = O \implies \begin{bmatrix} 10 - 16 & 6 \\ 6 & 10 - 16 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -6 & 6 \\ 6 & -6 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \implies -6x + 6y = 0 \implies x = y$$

This gives the clear directional vector configuration:

$$X_{\text{minor}} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

(D) an ellipse with minor axis along $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Key Points

- **Matrix Transformation of Fields:** A spherical boundary $\|Y\|^2 = 1$ under linear transformations shifts to an elliptical quadratic field described by $X^T(P^T P)X = 1$.
- **Spectral Association Rule:** In a quadratic form $X^T M X = 1$, the principal axes match the eigenvectors of A . The maximum eigenvalue ($\lambda_{\max}$) contracts the coordinate line the most, defining the direction of the minor axis.
- **Symmetric Properties:** If a matrix has identical diagonal elements ($P_{11} = P_{22}$), its eigenvectors are always symmetric reflections along the line $x = y$ and $x = -y$.

Mistakes to be avoided

- **Multiplication Order:** Remember $(PX)^T = X^T P^T$. Reversing this order will result in non-conformable matrix dimensions.
- **Minor vs Major Confusion:** A common pitfall in competitive examinations is associating the larger eigenvalue with the major axis. Because the axis semi-length is proportional to $1/\sqrt{\lambda}$, the larger eigenvalue produces a shorter axis (minor axis).

Q1.15.1 NAT 2015 1M Ans: 9 ● ○ ○

Let x and y be the two eigenvalues of the given 2×2 matrix A . We know from the fundamental properties of matrices that the sum of the eigenvalues is equal to the trace (sum of diagonal elements) of the matrix:

$$\text{Trace}(A) = x + y = -6 \implies y = -6 - x \quad \text{--- (1)}$$

Similarly, the determinant of a matrix is equal to the product of its eigenvalues:

$$|A| = x \cdot y$$

$$|A| = x(-6 - x) = -6x - x^2$$

Let this be represented as a continuous single-variable function $f(x)$:

$$f(x) = -6x - x^2$$

$$f'(x) = \frac{d}{dx}(-6x - x^2) = -6 - 2x$$

$$-6 - 2x = 0 \implies x = -3$$

$$f''(x) = \frac{d}{dx}(-6 - 2x) = -2$$

Since $f''(-3) = -2 < 0$, the function achieves a local maximum at $x = -3$.

Substituting $x = -3$ back into our function to compute the maximum determinant:

$$\text{Maximum Determinant} = f(-3) = -6(-3) - (-3)^2 = 18 - 9 = \boxed{9}$$

Q1.15.2 **MCQ** **2015** **2M** **Ans: B** ● ● ○

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} -3 - \lambda & 0 & -2 \\ 1 & -1 - \lambda & 0 \\ 0 & a & -2 - \lambda \end{vmatrix} = 0$$

$$(-3 - \lambda)[(-1 - \lambda)(-2 - \lambda) - 0] - 0 - 2[1(a) - 0] = 0$$

$$(-3 - \lambda)(-1 - \lambda)(-2 - \lambda) - 2a = 0$$

$$a = -\frac{1}{2}(3 + \lambda)(1 + \lambda)(2 + \lambda)$$

$$a = -\frac{1}{2}(\lambda^3 + 6\lambda^2 + 11\lambda + 6)$$

$$\frac{da}{d\lambda} = -\frac{1}{2}(3\lambda^2 + 12\lambda + 11) = 0$$

$$3\lambda^2 + 12\lambda + 11 = 0$$

$$\lambda = \frac{-12 \pm \sqrt{(12)^2 - 4(3)(11)}}{2 \times 3} = \frac{-12 \pm \sqrt{144 - 132}}{6}$$

$$\lambda = \frac{-12 \pm \sqrt{12}}{6} = \frac{-12 \pm 2\sqrt{3}}{6} = -2 \pm \frac{1}{\sqrt{3}}$$

To distinguish between local maximum and minimum values, we compute the second derivative test:

$$\frac{d^2a}{d\lambda^2} = \frac{d}{d\lambda} \left[-\frac{1}{2}(3\lambda^2 + 12\lambda + 11) \right] = -\frac{1}{2}(6\lambda + 12) = -3\lambda - 6$$

• At $\lambda = -2 + \frac{1}{\sqrt{3}}$:

$$\frac{d^2a}{d\lambda^2} = -3 \left(-2 + \frac{1}{\sqrt{3}} \right) - 6 = 6 - \frac{3}{\sqrt{3}} - 6 = -\frac{6}{2\sqrt{3}} = -\frac{3}{\sqrt{3}} < 0 \implies (\text{Local Maximum})$$

• At $\lambda = -2 - \frac{1}{\sqrt{3}}$:

$$\frac{d^2a}{d\lambda^2} = -3 \left(-2 - \frac{1}{\sqrt{3}} \right) - 6 = 6 + \frac{3}{\sqrt{3}} - 6 = \frac{3}{\sqrt{3}} > 0 \implies (\text{Local Minimum})$$

Thus, the maximum value occurs at $\lambda = -2 + \frac{1}{\sqrt{3}}$. Substituting this value back into the function of a :

$$a_{\max} = -\frac{1}{2} \left(3 + \left(-2 + \frac{1}{\sqrt{3}} \right) \right) \left(1 + \left(-2 + \frac{1}{\sqrt{3}} \right) \right) \left(2 + \left(-2 + \frac{1}{\sqrt{3}} \right) \right)$$

$$a_{\max} = -\frac{1}{2} \left(1 + \frac{1}{\sqrt{3}} \right) \left(-1 + \frac{1}{\sqrt{3}} \right) \left(\frac{1}{\sqrt{3}} \right)$$

$$a_{\max} = -\frac{1}{2} \left(\left(\frac{1}{\sqrt{3}} \right)^2 - (1)^2 \right) \left(\frac{1}{\sqrt{3}} \right) = -\frac{1}{2} \left(\frac{1}{3} - 1 \right) \left(\frac{1}{\sqrt{3}} \right)$$

$$a_{\max} = -\frac{1}{2} \left(-\frac{2}{3} \right) \left(\frac{1}{\sqrt{3}} \right) = \frac{1}{3\sqrt{3}}$$

(B) $\frac{1}{3\sqrt{3}}$

Key Points

- **Characteristic Expansion:** For an $n \times n$ matrix, the characteristic equation yields an n^{th} -degree polynomial whose roots correspond exactly to the matrix eigenvalues.
- **Optimization Test:** A continuous function $f(x)$ possesses a local maximum at a stationary point x_c if $f'(x_c) = 0$ and $f''(x_c) < 0$.

Q1.15.3 MCQ 2015 1M Ans: A ● ○ ○

If there is a unique solution, then $|A| \neq 0$.

All the eigenvalues of A are non-zero and the given equations are linearly independent.

$$\therefore P \equiv Q \equiv R \equiv S$$

(A) $P \equiv Q \equiv R \equiv S$

Q1.14.1 NAT 2014 2M Ans: 0.33 ● ● ○

The characteristic equation of the system matrix A is determined by finding the roots of its characteristic determinant:

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} -\lambda & 1 & -1 \\ -6 & -11 - \lambda & 6 \\ -6 & -11 & 5 - \lambda \end{vmatrix} = 0$$

$$(-\lambda) [(-11 - \lambda)(5 - \lambda) - (6)(-11)] - 1 [(-6)(5 - \lambda) - (6)(-6)] + (-1) [(-6)(-11) - (-11 - \lambda)(-6)] = 0$$

$$(-\lambda) [\lambda^2 + 6\lambda - 55 + 66] - 1 [-30 + 6\lambda + 36] - 1 [66 - (66 + 6\lambda)] = 0$$

$$(-\lambda) [\lambda^2 + 6\lambda + 11] - 1 [6\lambda + 6] - 1 [-6\lambda] = 0$$

$$-\lambda^3 - 6\lambda^2 - 11\lambda - 6\lambda - 6 + 6\lambda = 0$$

$$-\lambda^3 - 6\lambda^2 - 11\lambda - 6 = 0$$

$$\lambda^3 + 6\lambda^2 + 11\lambda + 6 = 0$$

By inspection or using the rational root theorem, we find that $\lambda = -1$ is a root because $(-1)^3 + 6(-1)^2 + 11(-1) + 6 = -1 + 6 - 11 + 6 = 0$. Factoring out $(\lambda + 1)$ via polynomial long division:

$$(\lambda + 1)(\lambda^2 + 5\lambda + 6) = 0 \implies (\lambda + 1)(\lambda + 2)(\lambda + 3) = 0$$

Thus, the distinct eigenvalues of matrix A are:

$$\lambda = -1, -2, -3$$

Evaluating the absolute ratio yields:

$$\text{Required Ratio} = \left| \frac{\lambda_{\max}}{\lambda_{\min}} \right| = \left| \frac{-1}{-3} \right| = \frac{1}{3} = \boxed{0.33}$$

Q1.14.2 MCQ 2014 1M Ans: C ● ○ ○

Given the two real square matrices A and B :

$$A = \begin{bmatrix} p & q \\ r & s \end{bmatrix}, \quad B = \begin{bmatrix} p^2 + q^2 & pr + qs \\ pr + qs & r^2 + s^2 \end{bmatrix}$$

Method 1

Let us evaluate the determinant of matrix A and B :

$$|A| = ps - qr$$

$$|B| = (p^2 + q^2)(r^2 + s^2) - (pr + qs)^2$$

$$|B| = (p^2r^2 + p^2s^2 + q^2r^2 + q^2s^2) - (p^2r^2 + q^2s^2 + 2pqrs)$$

$$|B| = p^2s^2 + q^2r^2 - 2pqrs$$

$$|B| = (ps - qr)^2$$

$$|B| = |A|^2$$

- If $|A| \neq 0$, then matrix A is non-singular and has full rank, so $\rho(A) = 2$. Since $|B| = |A|^2 \neq 0$, matrix B is also non-singular and has full rank, so $\rho(B) = 2$.
- If $|A| = 0$, then $\rho(A) < 2$. Since $|B| = |A|^2 = 0$, $\rho(B) < 2$. For non-zero elements, the rank reduces identically based on structural linear dependency.

Therefore, if the rank of matrix A is N ($\rho(A) = N$), the rank of matrix B is also N ($\rho(B) = N$).

$$\boxed{(C) N}$$

Method 2

We can alternatively solve this using matrix multiplication properties. Let us observe the product of matrix A with its transpose A^T :

$$A = \begin{bmatrix} p & q \\ r & s \end{bmatrix} \implies A^T = \begin{bmatrix} p & r \\ q & s \end{bmatrix}$$

$$AA^T = \begin{bmatrix} p & q \\ r & s \end{bmatrix} \begin{bmatrix} p & r \\ q & s \end{bmatrix} = \begin{bmatrix} p(p) + q(q) & p(r) + q(s) \\ r(p) + s(q) & r(r) + s(s) \end{bmatrix} = \begin{bmatrix} p^2 + q^2 & pr + qs \\ pr + qs & r^2 + s^2 \end{bmatrix}$$

$$B = AA^T$$

By applying the fundamental properties of matrix rank, we know that for any real matrix A :

$$\rho(B) = \rho(AA^T) = \rho(A)$$

Thus, if $\rho(A) = N$, then $\rho(B) = N$.

(C) N

Key Points

- **Rank Product Property:** For any real matrix A , the rank of A is identical to the rank of its gram matrix product: $\rho(A) = \rho(A^T) = \rho(AA^T) = \rho(A^T A)$.
- **Determinant Product Rule:** $\det(AB) = \det(A) \det(B)$. Since $\det(A^T) = \det(A)$, it directly follows that $\det(AA^T) = \det(A)^2$.

Q1.14.3 NAT 2014 1M Ans: A ● ○ ○

By Properties of eigen values, only option(A) is true.

(A) All the eigenvalues are real.

Q1.13.1 MCQ 2013 2M Ans: D ● ○ ○

Given the eigenvectors X_1 and X_2 corresponding to the eigenvalues $\lambda_1 = -1$ and $\lambda_2 = -2$ respectively:

$$X_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for } \lambda_1 = -1, \quad X_2 = \begin{bmatrix} 1 \\ -2 \end{bmatrix} \text{ for } \lambda_2 = -2$$

Let the unknown 2×2 matrix A be defined as:

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Case 1: For $\lambda_1 = -1$ and X_1

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = (-1) \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} a - b \\ c - d \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$a - b = -1 \quad \text{--- (1)}$$

$$c - d = 1 \quad \text{--- (2)}$$

Case 2: For $\lambda_2 = -2$ and X_2

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix} = (-2) \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} a - 2b \\ c - 2d \end{bmatrix} = \begin{bmatrix} -2 \\ 4 \end{bmatrix}$$

$$a - 2b = -2 \quad \text{--- (3)}$$

$$c - 2d = 4 \quad \text{--- (4)}$$

Solving for a and b : Subtracting equation (3) from equation (1):

$$(a - b) - (a - 2b) = -1 - (-2) \implies b = 1$$

$$a - 1 = -1 \implies a = 0$$

Solving for c and d : Subtracting equation (4) from equation (2):

$$(c - d) - (c - 2d) = 1 - 4 \implies d = -3$$

$$c - (-3) = 1 \implies c + 3 = 1 \implies c = -2$$

$$A = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

$$(D) \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

Q1.11.1 MCQ 2011 1M Ans: B ☒ ☐ ☐

Let the two given complex vectors be represented as row matrices X and Y :

$$X = [1 \quad 1 \quad 1], \quad Y = [1 \quad a \quad a^2]$$

where a represents the complex cube root of unity:

$$a = -\frac{1}{2} + j\frac{\sqrt{3}}{2} = e^{j\frac{2\pi}{3}}$$

Its square a^2 corresponds to its complex conjugate (a^*), given by:

$$a^2 = -\frac{1}{2} - j\frac{\sqrt{3}}{2} = e^{-j\frac{2\pi}{3}}$$

$$XY^T = [1 \quad 1 \quad 1] \begin{bmatrix} 1 \\ a \\ a^2 \end{bmatrix} = 1 + a + a^2$$

From the fundamental mathematical properties of the complex cube roots of unity, the sum of all three roots is identically zero:

$$1 + a + a^2 = 0$$

Since $XY^T = 0$, the two vectors satisfy the condition for orthogonality. Now let us check if the vectors are normalized (i.e., whether they have a unit magnitude):

$$XX^T = [1 \quad 1 \quad 1] \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = 1^2 + 1^2 + 1^2 = 3 \neq 1$$

$$YY^H = [1 \quad a \quad a^2] \begin{bmatrix} 1 \\ a^* \\ (a^2)^* \end{bmatrix} = 1 \cdot 1 + a \cdot a^* + a^2 \cdot (a^2)^* = 1 + |a|^2 + |a|^4$$

$$YY^H = 1 + 1 + 1 = 3 \neq 1$$

Because the inner product equals zero but the lengths of the vectors are not equal to 1 ($\|X\| = \|Y\| = \sqrt{3}$), the vectors X and Y are orthogonal but not orthonormal.

(B) Orthogonal

Key Points

- **Properties of Cube Roots of Unity:** For $a = e^{j\frac{2\pi}{3}}$, the properties $a^3 = 1$ and $1 + a + a^2 = 0$ hold true.
- **Orthogonal vs. Orthonormal:** Two vectors X and Y are *orthogonal* if their inner product equals zero. They are defined as *orthonormal* if they are orthogonal **and** each vector possesses a unit norm ($\|X\| = 1, \|Y\| = 1$).

Mistakes to be avoided

- **Inner Product Definition:** In pure complex vector spaces, remember that the true inner product uses the conjugate transpose (H), so $\langle X, Y \rangle = XY^H$. In this specific problem, using either XY^T or XY^H evaluates to zero, but failing to include the conjugate when evaluating vector length (YY^T instead of YY^H) can lead to calculation errors.
- **Misinterpreting Orthogonality:** Do not automatically assume that because a vector contains complex coefficients, its magnitude squared will include imaginary components. The norm squared ($\|Y\|^2$) is always a real scalar value.

Q1.11.2 MCQ 2011 2M Ans: D ☒ ☐ ☐

Given the square matrix $[A]$ to be decomposed into a lower triangular matrix $[L]$ and an upper triangular matrix $[U]$ such that $[A] = [L][U]$:

$$[A] = \begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix}$$

Method 1

Using Doolittle's algorithm, we assume $[L]$ is a unit lower triangular matrix and $[U]$ is an upper triangular matrix:

$$[L] = \begin{bmatrix} 1 & 0 \\ l_{21} & 1 \end{bmatrix}, \quad [U] = \begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix}$$

Equating the product $[L][U]$ to matrix $[A]$:

$$\begin{bmatrix} 1 & 0 \\ l_{21} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{bmatrix} = \begin{bmatrix} u_{11} & u_{12} \\ l_{21}u_{11} & l_{21}u_{12} + u_{22} \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix}$$

- From the first row elements:

$$u_{11} = 2$$

$$u_{12} = 1$$

- From the second row elements:

$$l_{21}u_{11} = 4 \implies l_{21}(2) = 4 \implies l_{21} = 2$$

$$l_{21}u_{12} + u_{22} = -1 \implies (2)(1) + u_{22} = -1 \implies u_{22} = -3$$

Substituting these values back into the definitions of $[L]$ and $[U]$ gives:

$$[L] = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}, \quad [U] = \begin{bmatrix} 2 & 1 \\ 0 & -3 \end{bmatrix}$$

Looking closely at the available options, none match this standard Doolittle form. This implies that the question considers a unit upper triangular matrix configuration for $[U]$ instead (Crout's algorithm format), or a generic scaling layout. Let us evaluate Option D:

$$[L] = \begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix}, \quad [U] = \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$$

Let us verify their product:

$$[L][U] = \begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2(1) + 0(0) & 2(0.5) + 0(1) \\ 4(1) + (-3)(0) & 4(0.5) + (-3)(1) \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix} = [A]$$

Since the product matches $[A]$ exactly, Option D provides the correct factorization.

Method 2

Alternatively, we can directly verify the options by performing matrix multiplication for each choice:

- **Option A:** $\begin{bmatrix} 1 & 0 \\ 4 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 6 \end{bmatrix} \neq [A]$
- **Option B:** $\begin{bmatrix} 2 & 0 \\ 4 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 4 & 3 \end{bmatrix} \neq [A]$
- **Option C:** $\begin{bmatrix} 1 & 0 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 8 & 3 \end{bmatrix} \neq [A]$
- **Option D:** $\begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 4 & -1 \end{bmatrix} = [A]$

$$(D) \begin{bmatrix} 2 & 0 \\ 4 & -3 \end{bmatrix} \text{ and } \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$$

Key Points

- **LU Decomposition Formats:** LU decomposition is not unique unless specific restrictions are placed on the diagonal elements.
- **Doolittle vs. Crout Formats:** Doolittle's algorithm forces the diagonal elements of L to be 1 ($l_{ii} = 1$), whereas Crout's algorithm forces the diagonal elements of U to be 1 ($u_{ii} = 1$). Option D utilizes Crout's factorization method.

Q1.10.1 MCQ 2010 2M Ans: B ● ○ ○

Given that matrix P is an upper triangular matrix of size 3×3 :

$$P = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$$

We can directly verify the options by testing the validity of the condition $PX = \lambda X$ for each given choice:

Checking Option (a): Let $X = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$

$$PX = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1(-1) + 1(1) + 0(1) \\ 0(-1) + 2(1) + 2(1) \\ 0(-1) + 0(1) + 3(1) \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \\ 3 \end{bmatrix}$$

Since $\begin{bmatrix} 0 \\ 4 \\ 3 \end{bmatrix}$ cannot be expressed as a scalar multiple $\lambda \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$, this vector does not satisfy $PX = \lambda X$. Thus, option (a) is not an eigenvector.

Checking Option (b): Let $X = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$

$$PX = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 1(1) + 1(2) + 0(1) \\ 0(1) + 2(2) + 2(1) \\ 0(1) + 0(2) + 3(1) \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \\ 3 \end{bmatrix}$$

Factoring out the common scalar factor 3 from the resulting vector:

$$\begin{bmatrix} 3 \\ 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \lambda X$$

Since $PX = 3X$, the vector $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ is a valid eigenvector corresponding to the eigenvalue $\lambda = 3$.

(B) $[1 \ 2 \ 1]^T$

Q1.09.1 MCQ 2009 1M Ans: C ● ○ ○

Given that for a 2×2 real square matrix $A_{2 \times 2}$, its trace and determinant are:

$$\text{Trace}(A) = -2, \quad |A| = -35$$

Let λ_1 and λ_2 be the two eigenvalues of matrix A .

1. The sum of the eigenvalues equals the trace of the matrix:

$$\lambda_1 + \lambda_2 = \text{Trace}(A) = -2 \implies \lambda_2 = -2 - \lambda_1 \quad \text{--- (1)}$$

2. The product of the eigenvalues equals the determinant of the matrix:

$$\lambda_1 \lambda_2 = |A| = -35 \quad \text{--- (2)}$$

$$\lambda_1(-2 - \lambda_1) = -35$$

$$-2\lambda_1 - \lambda_1^2 = -35$$

$$\lambda_1^2 + 2\lambda_1 - 35 = 0$$

$$\lambda_1^2 + 7\lambda_1 - 5\lambda_1 - 35 = 0$$

$$\lambda_1(\lambda_1 + 7) - 5(\lambda_1 + 7) = 0$$

$$(\lambda_1 + 7)(\lambda_1 - 5) = 0$$

$$\lambda_1 = -7 \quad \text{or} \quad \lambda_1 = 5$$

Thus, the two eigenvalues of the matrix are -7 and 5 .

(C) -7 and 5

Q1.08.1 MCQ 2008 1M Ans: D ● ○ ○

Given the characteristic equation of the square matrix P :

$$\lambda^3 + \lambda^2 + 2\lambda + 1 = 0$$

By applying the Cayley-Hamilton theorem,

$$P^3 + P^2 + 2P + I = O$$

$$I = -P^3 - P^2 - 2P$$

$$P^{-1}I = P^{-1}(-P^3 - P^2 - 2P)$$

$$P^{-1} = -P^2 - P - 2I$$

$$P^{-1} = -(P^2 + P + 2I)$$

$$(D) -(P^2 + P + 2I)$$

Key Points

- **Cayley-Hamilton Theorem:** If $p(\lambda) = \det(\lambda I - A) = 0$ is the characteristic equation of a square matrix A , then $p(A) = O$.
- **Matrix Inverse via Cayley-Hamilton:** If the constant term c_0 of the characteristic polynomial is non-zero, the matrix is invertible, and A^{-1} can be expressed as a linear combination of lower powers of A :

$$A^{-1} = -\frac{1}{c_0} (A^{n-1} + c_{n-1}A^{n-2} + \dots + c_1I)$$

Q1.08.2 MCQ 2008 1M Ans: A ● ○ ○

Given a rectangular matrix Q of dimensions 5×6 with a known matrix rank of 4:

$$\rho(Q_{5 \times 6}) = 4$$

$$\rho(Q) = \text{Number of linearly independent rows} = \text{Number of linearly independent columns}$$

Given that $\rho(Q_{5 \times 6}) = 4$, it directly follows that:

- Out of the 5 total rows, exactly 4 rows are linearly independent.
- Out of the 6 total columns, exactly 4 columns are linearly independent.

Thus, the matrix $Q_{5 \times 6}$ contains exactly four linearly independent rows and four linearly independent columns.

$$(A) Q \text{ will have four linearly independent rows and four linearly independent columns}$$

Q1.08.3 MCQ 2008 2M Ans: B ● ● ○

The length, norm, or magnitude of a column vector $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ in Euclidean space is denoted by $\|X\|$ (or $|X|$) and is algebraically defined as:

$$\|X\| = \sqrt{x_1^2 + x_2^2}$$

Method 1

Let P be a standard transformation matrix parameterized by θ :

$$P = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$PX = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta + x_2 \sin \theta \\ -x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$$

$$\|PX\| = \sqrt{(x_1 \cos \theta + x_2 \sin \theta)^2 + (-x_1 \sin \theta + x_2 \cos \theta)^2}$$

$$\begin{aligned} \|PX\| &= \sqrt{(x_1^2 \cos^2 \theta + x_2^2 \sin^2 \theta + 2x_1x_2 \cos \theta \sin \theta) + (x_1^2 \sin^2 \theta + x_2^2 \cos^2 \theta - 2x_1x_2 \sin \theta \cos \theta)} \\ &= \sqrt{x_1^2 (\cos^2 \theta + \sin^2 \theta) + x_2^2 (\sin^2 \theta + \cos^2 \theta)} \end{aligned}$$

$$\|PX\| = \sqrt{x_1^2(1) + x_2^2(1)} = \sqrt{x_1^2 + x_2^2}$$

Since $\sqrt{x_1^2 + x_2^2} = \|X\|$, we establish that:

$$\|PX\| = \|X\|$$

This shows that multiplying by an orthogonal matrix preserves the length of the vector for any arbitrary vector X , which matches statement (B).

Method 2

Alternatively, we can prove this coordinate-free length preservation property using matrix transpose identities. The norm squared of the transformed vector PX can be written using inner products as:

$$\|PX\|^2 = (PX)^\top (PX)$$

$$\|PX\|^2 = X^\top P^\top PX$$

$$\|PX\|^2 = X^\top (I)X = X^\top X$$

Since $X^\top X = \|X\|^2$, taking the square root of both sides gives:

$$\|PX\| = \|X\|$$

$$(B) \quad \|P\vec{x}\| = \|\vec{x}\| \text{ for all vectors } \vec{x}$$

Key Points

- Length Preservation (Isometry):** A linear transformation preserves vector length ($\|PX\| = \|X\|$) if and only if its transformation matrix P is orthogonal ($P^\top P = I$).
- Orthogonal Transformation Dynamics:** Geometric transformations like pure rotations and reflections are modeled by orthogonal matrices. They alter a vector's orientation while keeping its magnitude constant.

Q1.08.4 MCQ 2008 2M Ans: D ● ○ ○

Given the Moore-Penrose pseudoinverse A^+ of a rectangular matrix A (with full column rank, i.e., $\rho(A) = n$):

$$A^+ = (A^T A)^{-1} A^T$$

We evaluate the structural property of option (d) to check its validity.

$$\text{LHS} = AA^+A$$

$$\text{LHS} = A \left[(A^T A)^{-1} A^T \right] A$$

$$\text{LHS} = A \left[(A^T A)^{-1} (A^T A) \right]$$

$$\text{LHS} = A[I] = A$$

$$(D) AA^+A = A^+$$

Key Points

• **Moore-Penrose Conditions:** The pseudoinverse A^+ of a matrix A must satisfy four fundamental conditions uniquely:

1. $AA^+A = A$
2. $A^+AA^+ = A^+$
3. $(AA^+)^T = AA^+$
4. $(A^+A)^T = A^+A$

• **Full Column Rank Invariant:** When a matrix A has linearly independent columns ($\rho(A) = n$), the matrix product $A^T A$ is guaranteed to be a square, non-singular, invertible matrix.

Q1.07.1 MCQ 2007 1M Ans: B ● ○ ○

Let X be a non-zero column vector of dimensions $n \times 1$. The matrix V is defined as the outer product of the vector X with its own transpose X^T :

$$V = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ \vdots \\ x_n \end{bmatrix}_{n \times 1} \begin{bmatrix} x_1 & x_2 & \dots & \dots & x_n \end{bmatrix}_{1 \times n} = XX^T$$

Method 1

By performing standard matrix multiplication row-by-row, the resulting square matrix V of size $n \times n$ is explicitly structured as:

$$V = \begin{bmatrix} x_1x_1 & x_1x_2 & \dots & x_1x_n \\ x_2x_1 & x_2x_2 & \dots & x_2x_n \\ \vdots & \vdots & \ddots & \vdots \\ x_nx_1 & x_nx_2 & \dots & x_nx_n \end{bmatrix}_{n \times n}$$

Let us analyze the rows of matrix V :

$$\begin{aligned} R_1 &= x_1 \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix} \\ R_2 &= x_2 \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix} = \frac{x_2}{x_1} R_1 \quad (\text{for } x_1 \neq 0) \\ &\vdots \\ R_n &= x_n \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix} = \frac{x_n}{x_1} R_1 \quad (\text{for } x_1 \neq 0) \end{aligned}$$

Since every row R_i is a scalar multiple of the first row R_1 , all rows are linearly dependent on a single basis row. Performing elementary row operations can reduce all rows except one to zero. Therefore, the number of linearly independent rows is exactly 1, which means:

$$\rho(V) = 1$$

Method 2

We can alternatively find the rank of matrix V using properties of matrix rank inequalities. For any product of two matrices A and B , the rank of the product is bounded by:

$$\rho(AB) \leq \min(\rho(A), \rho(B))$$

Substituting $A = X_{n \times 1}$ and $B = X_{1 \times n}^T$ into the inequality:

$$\rho(V) = \rho(XX^T) \leq \min(\rho(X_{n \times 1}), \rho(X_{1 \times n}^T))$$

Since X is a non-zero column vector and X^T is a non-zero row vector, both contain exactly one linearly independent vector asset:

$$\rho(X_{n \times 1}) = 1 \quad \text{and} \quad \rho(X_{1 \times n}^T) = 1$$

Thus:

$$\rho(V) \leq \min(1, 1) \implies \rho(V) \leq 1$$

Because X is a non-zero vector, V is a non-zero matrix, which ensures $\rho(V) \geq 1$. Combining these upper and lower limits gives:

$$\rho(V) = 1$$

(B) has rank 1

Key Points

- **Rank of an Outer Product:** The outer product of any two non-zero column vectors u and v always structures a rank-one matrix: $\rho(uv^T) = 1$.
- **Linear Dependence in Outer Products:** In an outer product matrix $V = uv^T$, every column is a scalar multiple of u , and every row is a scalar multiple of v^T .

Mistakes to be avoided

- **Outer Product vs. Inner Product:** Do not confuse the outer product matrix XX^T (which is an $n \times n$ matrix of rank 1) with the inner product scalar $X^T X$ (which is a 1×1 scalar value representing the squared Euclidean norm $\|X\|^2$).
- **Vector Dimension Formats:** Ensure that the matrix dimension layouts like $X_{n \times 1}$ and $X_{1 \times n}^T$ are placed in math mode with proper formatting instead of written as unformatted text.

Q1.07.2 MCQ 2007 2M Ans: B ● ● ○

Let x and y be two real vectors in a 3-dimensional Euclidean space, and let $\langle x, y \rangle$ denote their standard inner product (dot product), which can also be written as $x \cdot y$.

$$G = \det \begin{bmatrix} \langle x, x \rangle & \langle x, y \rangle \\ \langle y, x \rangle & \langle y, y \rangle \end{bmatrix}$$

$$G = \det \begin{bmatrix} \|x\|^2 & x \cdot y \\ x \cdot y & \|y\|^2 \end{bmatrix}$$

$$G = \|x\|^2 \|y\|^2 - (x \cdot y)^2 \quad \text{--- (1)}$$

$$|x \cdot y| \leq \|x\| \|y\|$$

$$(x \cdot y)^2 \leq \|x\|^2 \|y\|^2 \implies \|x\|^2 \|y\|^2 - (x \cdot y)^2 \geq 0$$

The equality condition for Cauchy-Schwarz states that $(x \cdot y)^2 = \|x\|^2 \|y\|^2$ if and only if the vectors x and y are linearly dependent.

Therefore, if x and y are **linearly independent**, the strict inequality holds:

$$(x \cdot y)^2 < \|x\|^2 \|y\|^2 \implies \|x\|^2 \|y\|^2 - (x \cdot y)^2 > 0$$

Thus, the determinant is strictly positive when x and y are linearly independent.

(B) is positive when x and y are linearly independent

Q1.07.3 MCQ 2007 2M Ans: D ● ● ○

Given the linear operation $L(\mathbf{x})$ defined by a vector cross product:

$$L(\mathbf{x}) = \mathbf{b} \times \mathbf{x}$$

where $\mathbf{b} = [0 \ 1 \ 0]^\top$ and $\mathbf{x} = [x_1 \ x_2 \ x_3]^\top$. We need to find the eigenvalues of the matrix M that satisfies:

$$L(\mathbf{x}) = M \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Let us compute the cross product $\mathbf{b} \times \mathbf{x}$ explicitly using the standard determinant definition with unit vectors $\hat{i}$, $\hat{j}$, and $\hat{k}$:

$$\mathbf{b} \times \mathbf{x} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & 0 \\ x_1 & x_2 & x_3 \end{vmatrix}$$

$$\mathbf{b} \times \mathbf{x} = \hat{i}(1 \cdot x_3 - 0 \cdot x_2) - \hat{j}(0 \cdot x_3 - 0 \cdot x_1) + \hat{k}(0 \cdot x_2 - 1 \cdot x_1)$$

$$\mathbf{b} \times \mathbf{x} = x_3 \hat{i} + 0 \hat{j} - x_1 \hat{k}$$

$$L(\mathbf{x}) = \begin{bmatrix} x_3 \\ 0 \\ -x_1 \end{bmatrix}$$

$$\begin{bmatrix} x_3 \\ 0 \\ -x_1 \end{bmatrix} = \begin{bmatrix} 0 \cdot x_1 + 0 \cdot x_2 + 1 \cdot x_3 \\ 0 \cdot x_1 + 0 \cdot x_2 + 0 \cdot x_3 \\ -1 \cdot x_1 + 0 \cdot x_2 + 0 \cdot x_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Thus, the matrix M is:

$$M = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}$$

Now, we determine the eigenvalues (λ) of matrix M using the characteristic equation $|M - \lambda I| = 0$:

$$\begin{vmatrix} -\lambda & 0 & 1 \\ 0 & -\lambda & 0 \\ -1 & 0 & -\lambda \end{vmatrix} = 0$$

$$(-\lambda) \begin{vmatrix} -\lambda & 1 \\ -1 & -\lambda \end{vmatrix} = 0$$

$$(-\lambda) [(-\lambda)(-\lambda) - (1)(-1)] = 0$$

$$-\lambda (\lambda^2 + 1) = 0$$

- $-\lambda = 0 \implies \lambda = 0$
- $\lambda^2 + 1 = 0 \implies \lambda^2 = -1 \implies \lambda = \pm i$

Hence, the eigenvalues of M are $i, -i, 0$.

(D) $i, -i, 0$

Q1.07.4 MCQ 2007 1M Ans: C ● ○ ○

The characteristic equation of a square matrix A is determined by finding the roots of its characteristic polynomial:

$$|A - \lambda I| = 0$$

Given the characteristic determinant layout for the 2×2 matrix:

$$\begin{vmatrix} -3 - \lambda & 2 \\ -1 & 0 - \lambda \end{vmatrix} = 0$$

Expanding the 2×2 determinant explicitly:

$$(-3 - \lambda)(0 - \lambda) - (2)(-1) = 0$$

$$\lambda(3 + \lambda) + 2 = 0$$

$$\lambda^2 + 3\lambda + 2 = 0$$

$$A^2 + 3A + 2I = O$$

$$A^2 + 2A + A + 2I = O$$

$$A(A + 2I) + I(A + 2I) = O$$

$$(A + I)(A + 2I) = O$$

$$(C) (A + I)(A + 2I) = O$$

Q1.07.5 **MCQ** **2007** **1M** **Ans: A** ● ● ○

The characteristic equation of a square matrix A is given by:

$$\begin{aligned}
 |A - \lambda I| &= 0 \\
 \begin{vmatrix} -3 - \lambda & 2 \\ -1 & 0 - \lambda \end{vmatrix} &= 0 \\
 (-3 - \lambda)(0 - \lambda) - (2)(-1) &= 0 \\
 \lambda^2 - (-3)\lambda + 2 &= 0 \\
 \lambda^2 + 3\lambda + 2 &= 0
 \end{aligned}$$

Method 1

According to the Cayley-Hamilton theorem:

$$\begin{aligned}
 A^2 + 3A + 2I &= O \\
 \Rightarrow A^2 &= -3A - 2I \quad \text{--- (1)} \\
 A^3 &= A \cdot A^2 = A(-3A - 2I) = -3A^2 - 2A \\
 A^3 &= -3(-3A - 2I) - 2A = 9A + 6I - 2A \\
 \Rightarrow A^3 &= 7A + 6I \quad \text{--- (2)} \\
 A^4 &= A \cdot A^3 = A(7A + 6I) = 7A^2 + 6A \\
 A^4 &= 7(-3A - 2I) + 6A = -21A - 14I + 6A \\
 \Rightarrow A^4 &= -15A - 14I \quad \text{--- (3)} \\
 A^5 &= A \cdot A^4 = A(-15A - 14I) = -15A^2 - 14A \\
 A^5 &= -15(-3A - 2I) - 14A = 45A + 30I - 14A \\
 \Rightarrow A^5 &= 31A + 30I \quad \text{--- (4)}
 \end{aligned}$$

By observing the structural coefficient pattern generated across equations (1), (2), (3), and (4):

$$\begin{aligned}
 A^1 &= 1A + 0I \\
 A^2 &= -3A - 2I \\
 A^3 &= 7A + 6I \\
 A^4 &= -15A - 14I \\
 A^5 &= 31A + 30I
 \end{aligned}$$

Let us look at the coefficients of A and I for any power n . Let $A^n = c_n A + d_n I$. The sequence of the A coefficients (1, -3, 7, -15, 31, ...) satisfies the clear pattern:

$$c_n = (-1)^{n-1}(2^n - 1)$$

For $n = 9$:

$$c_9 = (-1)^8(2^9 - 1) = 1 \cdot (512 - 1) = 511$$

The sequence of the I coefficients (0, -2, 6, -14, 30, ...) satisfies the pattern:

$$d_n = c_n - 1 \quad (\text{for odd } n) \quad \text{or} \quad d_n = c_n + 1 \quad (\text{for even } n)$$

More universally, $d_n = (-1)^{n-1}(2^n - 1) - (-1)^{n-1} = (-1)^{n-1}(2^n - 2)$. For $n = 9$:

$$d_9 = (-1)^8(2^9 - 2) = 1 \cdot (512 - 2) = 510$$

Thus, assembling the expression for A^9 :

$$A^9 = 511A + 510I$$

Method 2

We can solve this problem much faster using a remainder polynomial evaluation technique. Let us find the remainder when λ^9 is divided by the characteristic polynomial $p(\lambda) = \lambda^2 + 3\lambda + 2$. Since the divisor is a second-degree polynomial, the remainder must be a linear polynomial of the form $a\lambda + b$:

$$\lambda^9 = q(\lambda)(\lambda^2 + 3\lambda + 2) + (a\lambda + b)$$

Factoring the characteristic polynomial:

$$\lambda^2 + 3\lambda + 2 = (\lambda + 1)(\lambda + 2)$$

Thus, the roots (eigenvalues) are $\lambda = -1$ and $\lambda = -2$.

Substituting these roots into the identity equation:

- For $\lambda = -1$:

$$(-1)^9 = q(-1)(0) + a(-1) + b \implies -a + b = -1 \quad \text{--- (5)}$$

- For $\lambda = -2$:

$$(-2)^9 = q(-2)(0) + a(-2) + b \implies -2a + b = -512 \quad \text{--- (6)}$$

Subtracting equation (6) from equation (5):

$$(-a + b) - (-2a + b) = -1 - (-512)$$

$$a = 511$$

Substituting $a = 511$ back into equation (5):

$$-511 + b = -1 \implies b = 510$$

Therefore, the scalar remainder polynomial is $511\lambda + 510$. By applying the Cayley-Hamilton theorem, we can substitute the matrix configurations directly into this remainder expression:

$$A^9 = 511A + 510I$$

(A) $511A + 510I$

Key Points

- Cayley-Hamilton Polynomial Division Property:** Any high-order matrix power A^n can be simplified to a lower-degree matrix polynomial matching the exact remainder obtained by dividing λ^n by the characteristic polynomial of A .
- Commutative Reduction:** Substituting matrix variables into verified scalar remainder identities is valid because all constituent components (A^n , A , and I) commute perfectly with each other.

Mistakes to be avoided

- Remainder Bounds:** When setting up a remainder function for polynomial long division, always make sure its degree is strictly less than the degree of the characteristic polynomial ($\deg(r) < \deg(p)$). For a 2×2 matrix, the remainder is

always at most linear ($aA + bI$).

Q1.05.1 MCQ 2005 1M Ans: A ● ○ ○

According to the Rouché-Capelli theorem, a system of linear equations represented in matrix form as $PX = Q$ is consistent if and only if the rank of the coefficient matrix P is identical to the rank of its augmented matrix $[P \mid Q]$.

$$\rho(P) = \rho([P \mid Q])$$

- Unique Solution:** If $\rho(P) = \rho([P \mid Q]) = n$, the linear system has exactly one unique solution.
- Infinitely Many Solutions:** If $\rho(P) = \rho([P \mid Q]) < n$, the system has an infinite number of solutions.

(A) Augmented matrix $[P \mid q]$ must have the same rank as matrix P

Q1.05.2 MCQ 2005 2M Ans: B ● ○ ○

Given the 3×3 square matrix R :

$$R = \begin{bmatrix} 1 & 0 & -1 \\ 2 & 1 & -1 \\ 2 & 3 & 2 \end{bmatrix}$$

$$R^{-1} = \frac{\text{adj}(R)}{|R|}$$

$$|R| = 1 \begin{vmatrix} 1 & -1 \\ 3 & 2 \end{vmatrix} - 0 \begin{vmatrix} 2 & -1 \\ 2 & 2 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 2 & 3 \end{vmatrix}$$

$$|R| = 1 [(1)(2) - (-1)(3)] - 0 - 1 [(2)(3) - (1)(2)]$$

$$|R| = 1(2 + 3) - 1(6 - 2) = 5 - 4 = 1$$

$$\text{adj}(R) = \begin{bmatrix} C_{11} & C_{21} & C_{31} \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}$$

$$\bullet C_{11} = (-1)^{1+1} \begin{vmatrix} 1 & -1 \\ 3 & 2 \end{vmatrix} = 1 \cdot [(1)(2) - (-1)(3)] = 2 + 3 = 5$$

$$\bullet C_{21} = (-1)^{2+1} \begin{vmatrix} 0 & -1 \\ 3 & 2 \end{vmatrix} = -1 \cdot [(0)(2) - (-1)(3)] = -(0 + 3) = -3$$

$$\bullet C_{31} = (-1)^{3+1} \begin{vmatrix} 0 & -1 \\ 1 & -1 \end{vmatrix} = 1 \cdot [(0)(-1) - (-1)(1)] = 0 + 1 = 1$$

$$\text{adj}(R) = \begin{bmatrix} 5 & -3 & 1 \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}$$

$$R^{-1} = \frac{1}{1} \begin{bmatrix} 5 & -3 & 1 \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix} = \begin{bmatrix} 5 & -3 & 1 \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}$$

Therefore, the top row of R^{-1} is $[5 \quad -3 \quad 1]$.

(B) $[5 \quad -3 \quad 1]$

Q1.02.1 MCQ 2002 1M Ans: C ● ○ ○

Given the 4×4 matrix A :

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 100 & 1 & 0 & 0 \\ 100 & 200 & 1 & 0 \\ 100 & 200 & 300 & 1 \end{bmatrix}$$

$$|A| = \prod_{i=1}^n a_{ii}$$

$$a_{11} = 1, \quad a_{22} = 1, \quad a_{33} = 1, \quad a_{44} = 1$$

$$|A| = 1 \times 1 \times 1 \times 1 = 1$$

(C) 1

Q1.98.1 MCQ 1998 1M Ans: C ● ○ ○

Given the 3×3 matrix A and its corresponding column eigenvector X :

$$A = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix}, \quad X = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$AX = \begin{bmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$AX = \begin{bmatrix} (-2)(1) + (2)(2) + (-3)(-1) \\ (2)(1) + (1)(2) + (-6)(-1) \\ (-1)(1) + (-2)(2) + (0)(-1) \end{bmatrix}$$

$$AX = \begin{bmatrix} -2 + 4 + 3 \\ 2 + 2 + 6 \\ -1 - 4 + 0 \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \\ -5 \end{bmatrix}$$

$$\begin{bmatrix} 5 \\ 10 \\ -5 \end{bmatrix} = 5 \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Since $AX = 5X$, comparing this with $AX = \lambda X$ reveals that the eigenvalue corresponding to the given vector is:

$$\lambda = 5$$

(C) 5

Q1.97.1 NAT 1997 5M Ans: ● ● ○

Given the 3×3 square matrix A (notated as a in the problem text):

$$A = \begin{bmatrix} 2 & 1 & 5 \\ 4 & 8 & 13 \\ 6 & 27 & 31 \end{bmatrix}$$

We decompose A into a product of a unit lower triangular matrix L and an upper triangular matrix U using Doolittle's algorithm ($A = LU$):

$$\begin{bmatrix} 2 & 1 & 5 \\ 4 & 8 & 13 \\ 6 & 27 & 31 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

Step 1: Determine the first row of U (multiply first row of L by columns of U):

$$u_{11} = 2, \quad u_{12} = 1, \quad u_{13} = 5$$

Step 2: Determine the first column of L (multiply remaining rows of L by first column of U):

$$\bullet \quad l_{21}u_{11} = 4 \implies l_{21}(2) = 4 \implies l_{21} = 2$$

$$\bullet \quad l_{31}u_{11} = 6 \implies l_{31}(2) = 6 \implies l_{31} = 3$$

Step 3: Determine the second row of U :

$$\bullet \quad l_{21}u_{12} + u_{22} = 8 \implies (2)(1) + u_{22} = 8 \implies u_{22} = 6$$

$$\bullet \quad l_{21}u_{13} + u_{23} = 13 \implies (2)(5) + u_{23} = 13 \implies u_{23} = 3$$

Step 4: Determine the remaining entries of L and U :

• For l_{32} :

$$l_{31}u_{12} + l_{32}u_{22} = 27 \implies (3)(1) + l_{32}(6) = 27 \implies 6l_{32} = 24 \implies l_{32} = 4$$

• For u_{33} :

$$l_{31}u_{13} + l_{32}u_{23} + u_{33} = 31 \implies (3)(5) + (4)(3) + u_{33} = 31$$

$$15 + 12 + u_{33} = 31 \implies 27 + u_{33} = 31 \implies u_{33} = 4$$

Assembling the final factored matrix architectures:

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 4 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 2 & 1 & 5 \\ 0 & 6 & 3 \\ 0 & 0 & 4 \end{bmatrix}$$

$$LU = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 5 \\ 0 & 6 & 3 \\ 0 & 0 & 4 \end{bmatrix}$$

Q1.95.1 NAT 1995 2M Ans: ● ● ○

Given the system of three linear equations in matrix form $AX = B$:

$$\begin{bmatrix} 0 & 21.432 & 34.432 \\ 17.587 & -1.348 & -1.865 \\ 0 & -2.663 & -1.803 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = \begin{bmatrix} 56.042 \\ 2.631 \\ 5.952 \end{bmatrix}$$

We construct the initial augmented matrix block:

$$\left[\begin{array}{ccc|c} 0 & 21.432 & 34.432 & 56.042 \\ 17.587 & -1.348 & -1.865 & 2.631 \\ 0 & -2.663 & -1.803 & 5.952 \end{array} \right]$$

Step 1: Row Interchanging

$$R_1 \leftrightarrow R_2$$

$$\left[\begin{array}{ccc|c} 17.587 & -1.348 & -1.865 & 2.631 \\ 0 & 21.432 & 34.432 & 56.042 \\ 0 & -2.663 & -1.803 & 5.952 \end{array} \right]$$

Step 2: Upper Triangular Reduction

$$R_3 \rightarrow R_3 - \left(\frac{-2.663}{21.432} \right) R_2 \implies R_3 \rightarrow R_3 + 0.124253 \cdot R_2$$

Let us compute the updated entries for the third row up to three decimal places:

- Entry (3, 2): $-2.663 + (0.124253)(21.432) = 0$
- Entry (3, 3): $-1.803 + (0.124253)(34.432) = -1.803 + 4.278 = 2.475$
- Augmented constant entry: $5.952 + (0.124253)(56.042) = 5.952 + 6.963 = 12.915$

This yields the finalized upper triangular augmented matrix configuration:

$$\left[\begin{array}{ccc|c} 17.587 & -1.348 & -1.865 & 2.631 \\ 0 & 21.432 & 34.432 & 56.042 \\ 0 & 0 & 2.475 & 12.915 \end{array} \right]$$

Step 3: Back-Substitution

1. From the updated third row equation:

$$2.475 \cdot Z = 12.915 \implies Z = \frac{12.915}{2.475} \approx 5.218$$

2. From the updated second row equation, substitute $Z = 5.218$:

$$21.432 \cdot Y + 34.432 \cdot Z = 56.042$$

$$21.432 \cdot Y + 34.432(5.218) = 56.042$$

$$21.432 \cdot Y + 179.667 = 56.042$$

$$21.432 \cdot Y = 56.042 - 179.667 = -123.625$$

$$Y = \frac{-123.625}{21.432} \approx -5.768$$

3. From the first row equation, substitute $Y = -5.768$ and $Z = 5.218$:

$$17.587 \cdot X - 1.348 \cdot Y - 1.865 \cdot Z = 2.631$$

$$17.587 \cdot X - 1.348(-5.768) - 1.865(5.218) = 2.631$$

$$17.587 \cdot X + 7.775 - 9.732 = 2.631$$

$$17.587 \cdot X - 1.957 = 2.631$$

$$17.587 \cdot X = 2.631 + 1.957 = 4.588$$

$$X = \frac{4.588}{17.587} \approx 0.261$$

$$\boxed{X = 0.261, \quad Y = -5.768, \quad Z = 5.218}$$

Q1.94.1 MCQ 1994 1M Ans: C ☒ ☐ ☐

Given a rectangular matrix A of dimensions 5×7 where every entry is exactly equal to -1 :

$$A = \begin{bmatrix} -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{bmatrix}_{5 \times 7}$$

$$R_1 = R_2 = R_3 = R_4 = R_5 = [-1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \quad -1]$$

Since all rows are completely identical, any row R_i (for $i > 1$) can be expressed as a perfect scalar multiple of the first row:

$$R_i = 1 \cdot R_1$$

$$R_2 \rightarrow R_2 - R_1, \quad R_3 \rightarrow R_3 - R_1, \quad R_4 \rightarrow R_4 - R_1, \quad R_5 \rightarrow R_5 - R_1$$

The transformation yields:

$$A_{\text{RREF}} = \begin{bmatrix} -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\rho(A) = 1$$

(C) 1

PrepFusion

SOLUTIONS — Linear Algebra

IN

Instrumentation Engineering

PrepFusion

Q1.26.1 MCQ 2026 1M Ans: C ● ○ ○

Given that the matrix $P \in \mathbb{R}^{3 \times 3}$ satisfies the matrix polynomial equation:

$$P^3 - 4P^2 + 5P - 2I = 0$$

$$f(\lambda) = \lambda^3 - 4\lambda^2 + 5\lambda - 2 = 0$$

We find the roots of this cubic equation by testing small integer values: If $\lambda = 1$:

$$f(1) = (1)^3 - 4(1)^2 + 5(1) - 2 = 1 - 4 + 5 - 2 = 0$$

Since $\lambda = 1$ satisfies the polynomial, $(\lambda - 1)$ is a factor. We perform polynomial long division or synthetic division to extract the remaining quadratic factor:

$$\lambda^3 - 4\lambda^2 + 5\lambda - 2 = (\lambda - 1)(\lambda^2 - 3\lambda + 2) = 0$$

Now, we factor the quadratic equation $\lambda^2 - 3\lambda + 2 = 0$:

$$\lambda^2 - 2\lambda - \lambda + 2 = 0$$

$$\lambda(\lambda - 2) - 1(\lambda - 2) = 0 \implies (\lambda - 1)(\lambda - 2) = 0$$

Combining all factors gives the complete characteristic factorization:

$$(\lambda - 1)^2(\lambda - 2) = 0$$

The roots of the equation are $\lambda = 1, 1, 2$.

Therefore, the set of all possible eigenvalues of matrix P is $\{1, 2\}$, which corresponds to option (C).

(C) $\{1, 2\}$

Key Points

- **Cayley-Hamilton Matrix Theorem:** Every square matrix satisfies its own characteristic scalar polynomial equation $|A - \lambda I| = 0$.

Q1.26.2 MCQ 2026 1M Ans: A ● ○ ○

Given the non-homogeneous linear system of equations $A\mathbf{x} = \mathbf{b}$, where:

$$A = \begin{bmatrix} 1 & 0 & 2 \\ -1 & 1 & 0 \\ 0 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}$$

Initial augmented matrix $[A \mid \mathbf{b}]$:

$$[A \mid \mathbf{b}] = \left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ -1 & 1 & 0 & 3 \\ 0 & 1 & 2 & 0 \end{array} \right]$$

Step 1:

$$R_2 \rightarrow R_2 + R_1$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & 2 & 4 \\ 0 & 1 & 2 & 0 \end{array} \right]$$

Step 2:

$$R_3 \rightarrow R_3 - R_2$$

$$\left[\begin{array}{ccc|c} 1 & 0 & 2 & 1 \\ 0 & 1 & 2 & 4 \\ 0 & 0 & 0 & -4 \end{array} \right]$$

- $\text{rank}(A) = 2$.
- $\text{rank}([A \mid \mathbf{b}]) = 3$.

Since $\text{rank}(A) \neq \text{rank}([A \mid \mathbf{b}])$, the system is inconsistent. Therefore, the total number of solutions is 0.

(A) 0

Key Points

- **Rouché-Capelli Consistency Rule:** A linear system $A\mathbf{x} = \mathbf{b}$ has zero solutions if and only if the augmented column parameters increase the vector subspace rank: $\text{rank}(A) \neq \text{rank}([A \mid \mathbf{b}])$.
- **Solution Classifications:**
 - **Unique Solution:** $\text{rank}(A) = \text{rank}([A \mid \mathbf{b}]) = n$ (where n is the number of variables).
 - **Infinitely Many Solutions:** $\text{rank}(A) = \text{rank}([A \mid \mathbf{b}]) < n$.
 - **No Solution:** $\text{rank}(A) \neq \text{rank}([A \mid \mathbf{b}])$.

Q1.25.1 MCQ 2025 1M Ans: A ● ○ ○

Given a square matrix A of dimensions $2n \times 2n$ where the entries a_{ij} are defined under conditional index constraints:

$$a_{ij} = \begin{cases} \beta & \text{if } (i + j) \text{ is odd} \\ -\beta & \text{if } (i + j) \text{ is even} \end{cases}$$

where β is a non-zero real number ($\beta \neq 0$) and $n > 2$.

$$A = \begin{bmatrix} -\beta & \beta & -\beta & \beta & -\beta & \beta \\ \beta & -\beta & \beta & -\beta & \beta & -\beta \\ -\beta & \beta & -\beta & \beta & -\beta & \beta \\ \beta & -\beta & \beta & -\beta & \beta & -\beta \\ -\beta & \beta & -\beta & \beta & -\beta & \beta \\ \beta & -\beta & \beta & -\beta & \beta & -\beta \end{bmatrix}_{6 \times 6}$$

- All odd rows are identical to the first row: $R_1 = R_3 = R_5 = [-\beta \ \beta \ -\beta \ \beta \ -\beta \ \beta]$
- All even rows are identical to each other and are exact scalar negatives of the first row: $R_2 = R_4 = R_6 = -R_1$

Applying elementary row operations to systematically eliminate dependent lines:

$$R_2 \rightarrow R_2 + R_1, \quad R_3 \rightarrow R_3 - R_1, \quad R_4 \rightarrow R_4 + R_1, \quad R_5 \rightarrow R_5 - R_1, \quad R_6 \rightarrow R_6 + R_1$$

The operations reduce the matrix into row echelon form:

$$A \sim \begin{bmatrix} -\beta & \beta & -\beta & \beta & -\beta & \beta \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}_{6 \times 6}$$

Therefore, $\rho(A) = 1$.

(A) 1

Q1.25.2 NAT 2025 1M Ans: 2 ● ○ ○

Given that the matrix $A = \begin{bmatrix} -1 & -1 \\ x & -4 \end{bmatrix}$ has an eigenvector $\mathbf{x} = \begin{bmatrix} \alpha \\ 2\alpha \end{bmatrix}$ where $\alpha \neq 0$.

Method 1

By applying the fundamental defining identity equation of the eigen-system structure ($A\mathbf{x} = \lambda\mathbf{x}$):

$$\begin{aligned} \begin{bmatrix} -1 & -1 \\ x & -4 \end{bmatrix} \begin{bmatrix} \alpha \\ 2\alpha \end{bmatrix} &= \lambda \begin{bmatrix} \alpha \\ 2\alpha \end{bmatrix} \\ \begin{bmatrix} (-1)(\alpha) + (-1)(2\alpha) \\ (x)(\alpha) + (-4)(2\alpha) \end{bmatrix} &= \begin{bmatrix} \lambda\alpha \\ 2\lambda\alpha \end{bmatrix} \\ \begin{bmatrix} -\alpha - 2\alpha \\ x\alpha - 8\alpha \end{bmatrix} = \begin{bmatrix} \lambda\alpha \\ 2\lambda\alpha \end{bmatrix} &\implies \begin{bmatrix} -3\alpha \\ x\alpha - 8\alpha \end{bmatrix} = \begin{bmatrix} \lambda\alpha \\ 2\lambda\alpha \end{bmatrix} \end{aligned}$$

1. From the first row expression:

$$\begin{aligned} -3\alpha &= \lambda\alpha \\ \lambda &= -3 \end{aligned}$$

2. From the second row expression:

$$\begin{aligned} x\alpha - 8\alpha &= 2\lambda\alpha \\ x - 8 &= 2\lambda \\ x - 8 &= 2(-3) \\ x - 8 &= -6 \implies x = -6 + 8 = \boxed{2} \end{aligned}$$

Method 2

We can also solve for x by utilizing the invariant alignment constraint property. For a vector to be an eigenvector, the matrix transformation must not rotate it out of its span. This means the components of the transformed vector $A\mathbf{x}$ must maintain the exact same ratio as the original vector components $\mathbf{x}$:

$$\mathbf{x} = \begin{bmatrix} \alpha \\ 2\alpha \end{bmatrix} \implies \frac{x_2}{x_1} = \frac{2\alpha}{\alpha} = 2$$

The transformed components computed from $A\mathbf{x}$ are:

$$(A\mathbf{x})_1 = -3\alpha, \quad (A\mathbf{x})_2 = (x - 8)\alpha$$

Enforcing the identical slope/ratio condition on the transformed vector components:

$$\begin{aligned} \frac{(A\mathbf{x})_2}{(A\mathbf{x})_1} &= 2 \implies \frac{(x - 8)\alpha}{-3\alpha} = 2 \\ \frac{x - 8}{-3} &= 2 \implies x - 8 = -6 \implies x = \boxed{2} \end{aligned}$$

Key Points

- **Eigenpair Operational Definition:** The matrix-vector multiplication $A\mathbf{x} = \lambda\mathbf{x}$ implies that the directional orientation of eigenvector $\mathbf{x}$ is invariant under the linear transformation map A .
- **Slope/Ratio Conservation:** In $\mathbb{R}^2$, a vector $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ maps along a line with a constant slope $m = \frac{x_2}{x_1}$. The corresponding eigenvalue transformation scales its magnitude but preserves this ratio unchanged.

Mistakes to be avoided

- **Improper Cancellation Violations:** Never divide by variable terms like α unless you have confirmed that the constraint bounds exclude zero ($\alpha \neq 0$).

Q1.24.1 MCQ 2024 1M Ans: C ● ● ○

Given a 3×3 matrix M constructed by stacking three column vectors $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ side-by-side:

$$M = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \mathbf{v}_3]$$

We need to identify the option for which $\text{rank}(M) = 3$. For a square matrix M of size 3×3 , the rank is equal to 3 if and only if all three column vectors are linearly independent. This condition is perfectly equivalent to stating that the determinant of the matrix must be non-zero ($|M| \neq 0$). Let us evaluate the determinant $|M|$ for each option systematically:

Option (A):

$$M = \begin{bmatrix} 1 & 0 & 1 \\ 0 & -1 & -1 \\ 1 & 0 & 1 \end{bmatrix}$$

$$|M| = 0 \implies \text{rank}(M) < 3$$

Thus, option (A) is incorrect.

Option (B):

$$M = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$|M| = 0 \implies \text{rank}(M) < 3$$

Thus, option (B) is incorrect.

Option (C):

$$M = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 0 & -1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$|M| = -(-1) \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} = 1 \cdot [(1)(1) - (-1)(1)] = 1 \cdot (1 + 1) = 2$$

Since $|M| = 2 \neq 0$, the column vectors are completely linearly independent, which means:

$$\text{rank}(M) = 3$$

Therefore, option (C) is correct.

Option (D):

$$M = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$|M| = 0 \implies \text{rank}(M) < 3$$

Thus, option (D) is incorrect.

$$(C) \ v_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, v_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, v_3 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

Key Points

- **Full Rank Criteria for Square Matrices:** For an $n \times n$ matrix, $\text{rank}(A) = n \iff \det(A) \neq 0$. If $\det(A) = 0$, then $\text{rank}(A) < n$.

Q1.24.2 NAT 2024 1M Ans: -2 ● ○ ○

Given a 3×3 matrix P with all real elements and its three corresponding eigenvalues:

$$\lambda_1 = \frac{1}{4}, \quad \lambda_2 = 1, \quad \lambda_3 = -2$$

Method 1

If λ_i represents the eigenvalues of an invertible square matrix P , then the eigenvalues of its inverse matrix P^{-1} are exactly the reciprocals of those original values ($\frac{1}{\lambda_i}$). Let the eigenvalues of P^{-1} be noted as λ'_1 , λ'_2 , and λ'_3 :

$$\lambda'_1 = \frac{1}{\lambda_1} = \frac{1}{1/4} = 4$$

$$\lambda'_2 = \frac{1}{\lambda_2} = \frac{1}{1} = 1$$

$$\lambda'_3 = \frac{1}{\lambda_3} = \frac{1}{-2} = -\frac{1}{2}$$

$$|P^{-1}| = \lambda'_1 \cdot \lambda'_2 \cdot \lambda'_3$$

$$|P^{-1}| = (4) \cdot (1) \cdot \left(-\frac{1}{2}\right) = \boxed{-2}$$

Method 2

Alternatively, we can evaluate this directly by working with the determinant product rules rather than computing individual reciprocal eigenvalues first. The determinant of matrix P is equal to the product of its original eigenvalues:

$$|P| = \lambda_1 \cdot \lambda_2 \cdot \lambda_3 = \left(\frac{1}{4}\right) \cdot (1) \cdot (-2) = -\frac{2}{4} = -\frac{1}{2}$$

$$|P^{-1}| = \frac{1}{|P|}$$

$$|P^{-1}| = \frac{1}{-1/2} = \boxed{-2}$$

Key Points

- **Inverse Matrix Determinant Invariant:** The scalar volume transformation of an inverted linear map satisfies:

$$\det(A^{-1}) = \frac{1}{\det(A)} = [\det(A)]^{-1}$$

Q1.23.1 MCQ 2023 1M Ans: A ● ● ○

Given system of linear equations:

$$x - 2y + z = 0$$

$$x - z = 0$$

$$x = z$$

$$z - 2y + z = 0 \implies 2z - 2y = 0 \implies y = z$$

Therefore, the parameters are related as:

$$x = y = z$$

Let $x = y = z = \alpha$ (say), where $\alpha \in \mathbb{R}$. The solution vector s can be expressed as:

$$s = \begin{pmatrix} \alpha \\ \alpha \\ \alpha \end{pmatrix} = \alpha \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad \alpha \in \mathbb{R}$$

$$(A) S = \left\{ \alpha \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \mid \alpha \in \mathbb{R} \right\}$$

Q1.23.2 NAT 2023 2M Ans: -8 ● ● ○

Given matrix:

$$A = \begin{bmatrix} 1 & 4 \\ -3 & \alpha \\ \beta & 6 \end{bmatrix}$$

Since $\text{Rank}(A) = 1$, all rows of the matrix must be linearly dependent on each other. This means the second row (R_2) and the third row (R_3) are scalar multiples of the first row (R_1).

Comparing the first elements of R_1 and R_2 :

$$\begin{aligned} R_2 &= -3 \cdot R_1 \\ \implies \alpha &= -3 \cdot 4 = -12 \end{aligned}$$

Comparing the second elements of R_1 and R_3 :

$$\begin{aligned} R_3 &= \frac{6}{4} \cdot R_1 = \frac{3}{2} \cdot R_1 \\ \implies \beta &= \frac{3}{2} \cdot 1 = \frac{3}{2} \end{aligned}$$

Now, we calculate the required ratio $\frac{\alpha}{\beta}$:

$$\frac{\alpha}{\beta} = \frac{-12}{\left(\frac{3}{2}\right)} = -12 \cdot \frac{2}{3} = -8$$

-8

Key Points

- The rank of a matrix is equal to the maximum number of linearly independent rows or columns.
- For a matrix to have a rank of 1, all non-zero rows must be scalar multiples of a single base row.

Q1.22.1 **MSQ** **2022** **1M** **Ans: A,D** ☒ ☐ ☐

Performing elementary row operations to reduce the given matrix M to row echelon form:

$$R_2 \rightarrow R_2 - 3R_1$$

$$R_3 \rightarrow R_3 - 2R_1$$

The resulting matrix in row echelon form is:

$$\begin{pmatrix} 2 & 3 & 7 \\ 0 & -5 & -14 \\ 0 & 0 & 0 \end{pmatrix}$$

Evaluating the statements based on this form:

- Rank of the matrix: The number of non-zero rows in the row echelon form is 2. Therefore, $\text{Rank}(M) = 2$. This makes statement (a) correct and statement (b) incorrect.
- Linear dependence of rows: Since the rank (2) is less than the total number of rows (3), the rows are linearly dependent. This makes statement (c) incorrect.
- Determinant of the matrix: Since there is a complete row of zeros, the matrix is singular, meaning its determinant is zero.

$$|M| = 0$$

This makes statement (d) correct.

Thus, the correct options are (A) and (D).

(A) The rank of M is 2, (D) The determinant of M is 0

Q1.22.2 **MSQ** **2022** **2M** **Ans: A,C** ☒ ☐ ☐

Verification of options based on the definition of eigenvalues and eigenvectors ($AX = \lambda X$):

- **Option (a):**

$$\begin{pmatrix} 4 & 3 \\ 9 & -2 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4(1) + 3(1) \\ 9(1) - 2(1) \end{pmatrix} = \begin{pmatrix} 7 \\ 7 \end{pmatrix} = 7 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Since $AX = 7X$, option (a) represents a valid eigenvector with an eigenvalue of $\lambda = 7$. Thus, (a) is true.

- **Option (b):**

$$\begin{pmatrix} 4 & 3 \\ 9 & -2 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 4(3) + 3(4) \\ 9(3) - 2(4) \end{pmatrix} = \begin{pmatrix} 24 \\ 19 \end{pmatrix} \neq \lambda X$$

Since $AX \neq \lambda X$, option (b) is false.

- **Option (c):**

$$\begin{pmatrix} 4 & 3 \\ 9 & -2 \end{pmatrix} \begin{pmatrix} 2 \\ -6 \end{pmatrix} = \begin{pmatrix} 4(2) + 3(-6) \\ 9(2) - 2(-6) \end{pmatrix} = \begin{pmatrix} -10 \\ 30 \end{pmatrix} = -5 \begin{pmatrix} 2 \\ -6 \end{pmatrix}$$

Since $AX = -5X$, option (c) represents a valid eigenvector with an eigenvalue of $\lambda = -5$. Thus, (c) is true.

- **Option (d):**

$$\begin{pmatrix} 4 & 3 \\ 9 & -2 \end{pmatrix} \begin{pmatrix} 2 \\ 8 \end{pmatrix} = \begin{pmatrix} 4(2) + 3(8) \\ 9(2) - 2(8) \end{pmatrix} = \begin{pmatrix} 32 \\ 2 \end{pmatrix} \neq \lambda X$$

Since $AX \neq \lambda X$, option (d) is false.

Both options (A) and (C) are correct.

(A) $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$, (C) $\begin{bmatrix} 2 \\ -6 \end{bmatrix}$

Q1.21.1 MCQ 2021 1M Ans: A ● ○ ○

Given vectors:

$$V = \begin{bmatrix} 1 & 0 \end{bmatrix} \quad \text{and} \quad W = \begin{bmatrix} 2 & 0 \end{bmatrix}$$

Computing the outer product $V^T V$:

$$\begin{aligned} V^T V &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \\ \Rightarrow 2V^T V &= \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix} \end{aligned}$$

Computing the outer product $W^T W$:

$$\begin{aligned} W^T W &= \begin{bmatrix} 2 \\ 0 \end{bmatrix} \begin{bmatrix} 2 & 0 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 0 & 0 \end{bmatrix} \\ \Rightarrow 3W^T W &= \begin{bmatrix} 12 & 0 \\ 0 & 0 \end{bmatrix} \end{aligned}$$

Now, consider the matrix $M = 2V^T V + 3W^T W$:

$$M = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 12 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 14 & 0 \\ 0 & 0 \end{bmatrix}$$

The number of non-zero rows in the row echelon form of M is 1.

$$\therefore \text{Rank}(M) = 1$$

(A) 1

Q1.21.2 NAT 2021 2M Ans: 4 ● ● ○

Given matrix:

$$A = \begin{pmatrix} 2 & 5 \\ 0 & 3 \end{pmatrix}$$

The characteristic equation of a matrix $A_{2 \times 2}$ is given by $|A - \lambda I| = 0$:

$$\begin{aligned} \begin{vmatrix} 2 - \lambda & 5 \\ 0 & 3 - \lambda \end{vmatrix} &= 0 \\ \Rightarrow (2 - \lambda)(3 - \lambda) - 0 &= 0 \\ \Rightarrow \lambda^2 - 5\lambda + 6 &= 0 \end{aligned}$$

According to the Cayley-Hamilton theorem:

$$\begin{aligned} A^2 - 5A + 6I &= O \\ A^2(A^2 - 5A + 6I) &= A^2 O \\ \Rightarrow A^4 - 5A^3 + 6A^2 &= O \end{aligned}$$

We need to find the determinant of the matrix $B = A^4 - 5A^3 + 6A^2 + 2I$:

$$|B| = |A^4 - 5A^3 + 6A^2 + 2I|$$

Substituting $A^4 - 5A^3 + 6A^2 = O$:

$$|B| = |O + 2I| = |2I|$$

For a 2×2 identity matrix, scaling by a factor of 2 scales the determinant by 2^2 :

$$|B| = 2^2 |I| = 4(1) = \boxed{4}$$

Q1.20.1 MCQ 2020 1M Ans: B ● ○ ○

Given the system of linear equations:

$$AX = b \quad \text{--- (1)}$$

where A is a matrix of dimension 2×4 and $b \neq 0$.

The maximum possible rank of matrix A is bounded by its dimensions:

$$\rho(A_{2 \times 4}) \leq \min(2, 4) \implies \rho(A_{2 \times 4}) \leq 2$$

Multiplying both sides of equation (1) by A^T :

$$A^T A X = A^T b$$

The dimensions of the product matrix $A^T A$ will be 4×4 . Using the rank property of matrix multiplication, $\rho(A^T A) = \rho(A)$:

$$\implies \rho[(A^T A)_{4 \times 4}] \leq 2$$

For a system of equations with a coefficient matrix of size $n \times n$ to possess a unique solution, the rank of that matrix must be exactly equal to the number of variables n .

Here, the number of variables for the system with coefficient matrix $(A^T A)_{4 \times 4}$ is 4. Since the maximum rank is at most 2, which is strictly less than 4:

$$\rho(A^T A) < 4$$

$\therefore$ A unique solution is NOT possible.

(B) No, it is not possible to get a unique solution for any 2×4 matrix A .

Key Points

- For any matrix A of dimension $m \times n$, the rank is bounded by $\rho(A) \leq \min(m, n)$.
- The rank of the product matrix satisfies $\rho(A^T A) = \rho(A)$.
- A linear system with a square coefficient matrix of dimension $n \times n$ yields a unique solution if and only if the rank of the matrix equals n . If the rank is less than n , it will either have infinitely many solutions or no solution.

Q1.20.2 MCQ 2020 2M Ans: D ● ○ ○

Given matrix:

$$M = \begin{bmatrix} 1 & -1 & 0 \\ 1 & -2 & 1 \\ 0 & -1 & 1 \end{bmatrix}$$

Consider the eigenvalue equation:

$$MX = \lambda X$$

From the options, if we choose the vector $X = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$:

$$MX = \begin{bmatrix} 1 & -1 & 0 \\ 1 & -2 & 1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1(1) - 1(1) + 0(1) \\ 1(1) - 2(1) + 1(1) \\ 0(1) - 1(1) + 1(1) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

We can rewrite the resulting vector as:

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Since $MX = 0X$, the equation is satisfied for an eigenvalue of $\lambda = 0$.

$$(D) \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Key Points

- A non-zero column vector X is an eigenvector of a square matrix M if it satisfies $MX = \lambda X$ for some scalar eigenvalue λ .
- When verifying options for eigenvectors, directly performing matrix-vector multiplication MX is often faster than computing the characteristic polynomial from scratch.

Q1.19.1 NAT 2019 1M Ans: 10 ☒ ☐ ☐

Given that the eigenvalues of a 3×3 matrix A are 1, 2, and 5.

We know that the determinant of a matrix is equal to the product of its eigenvalues:

$$\det(A) = \text{product of eigenvalues}$$

$$\therefore |A| = 1 \times 2 \times 5 = 10$$

Q1.18.1 MCQ 2018 1M Ans: A ☒ ☐ ☐

Let $\lambda_1, \lambda_2, \lambda_3$ be the eigenvalues of matrix N .

By the properties of matrix eigenvalues, if a square matrix N has eigenvalues λ_i , then the matrix N^2 will have eigenvalues λ_i^2 . Therefore, the eigenvalues of N^2 are:

$$\lambda_1^2, \lambda_2^2, \lambda_3^2$$

Given that the eigenvalues of N^2 are all zero:

$$\begin{aligned} \lambda_1^2 &= 0, \quad \lambda_2^2 = 0, \quad \lambda_3^2 = 0 \\ \Rightarrow \lambda_1 &= 0, \quad \lambda_2 = 0, \quad \lambda_3 = 0 \end{aligned}$$

Thus, 0, 0, 0 are the required eigenvalues of the matrix N .

$$(A) 0, 0, 0$$

Key Points

- If λ is an eigenvalue of matrix A corresponding to eigenvector X , then λ^k is an eigenvalue of A^k corresponding to the same eigenvector X .
- A matrix whose eigenvalues are all zero is associated with a nilpotent matrix property (i.e., $N^k = O$ for some positive integer k).

Q1.18.2 MCQ 2018 2M Ans: C ☒ ☒ ☐

The given system of linear equations can be represented in matrix form as:

$$AX = B$$

Consider the augmented matrix $(A|B)$:

$$(A|B) = \left(\begin{array}{cc|c} 3 & 2k & -2 \\ k & 6 & 2 \end{array} \right)$$

$$(A|B) \sim \left(\begin{array}{cc|c} 3 & 2k & -2 \\ 3(k) - k(3) & 3(6) - k(2k) & 3(2) - k(-2) \end{array} \right)$$

$$\Rightarrow (A|B) \sim \left(\begin{array}{cc|c} 3 & 2k & -2 \\ 0 & 18 - 2k^2 & 6 + 2k \end{array} \right)$$

The system has infinitely many solutions if the rank of the coefficient matrix A is equal to the rank of the augmented matrix $(A|B)$, and both are less than the number of variables (which is 2). This requires the entire second row to become zero:

$$18 - 2k^2 = 0 \quad \text{and} \quad 6 + 2k = 0$$

$$18 - 2k^2 = 0 \Rightarrow 2k^2 = 18 \Rightarrow k^2 = 9 \Rightarrow k = \pm 3$$

$$6 + 2k = 0 \Rightarrow 2k = -6 \Rightarrow k = -3$$

$$\therefore k = -3$$

(C) -3

Key Points

- According to the Rouché-Capelli theorem, a linear system $AX = B$ with n variables has infinitely many solutions if $\text{Rank}(A) = \text{Rank}(A|B) < n$.
- In a 2×2 system, this condition is satisfied when an row operation reduces the final row of the augmented matrix entirely to zeros ($0 = 0$).

Q1.17.1 NAT 2017 1M Ans: 1 ☒ ☐ ☐

Given that $V = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}_{3 \times 1}$ is a non-zero vector. Since V contains at least one non-zero entry:

$$\text{Rank}(V) = 1$$

$$\Rightarrow \text{Rank}(V^T) = 1$$

Since V is a non-zero vector, its outer product matrix VV^T is a non-zero matrix. Therefore, its rank must be at least 1:

$$\text{Rank}(VV^T) \geq 1 \quad \text{--- (1)}$$

$$\text{Rank}(VV^T) \leq \min\{\text{Rank}(V), \text{Rank}(V^T)\}$$

$$\Rightarrow \text{Rank}(VV^T) \leq \min\{1, 1\}$$

$$\Rightarrow \text{Rank}(VV^T) \leq 1 \quad \text{--- (2)}$$

From conditions (1) and (2), we have:

$$\rho(A) = \rho(VV^T) = 1$$

1

Q1.17.2 MCQ 2017 1M Ans: D ● ○ ○

The required transformation matrix for the reflection of a given original image about the X -axis can be determined by observing its effect on a general coordinate vector.

When a point (x, y) is reflected across the X -axis, its horizontal position remains unchanged, while its vertical position changes sign. Thus, the transformation maps:

$$[x \ y] \rightarrow [x \ -y]$$

This mapping can be expressed via matrix multiplication using a row vector format:

$$[x \ y] \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = [x(1) + y(0) \ x(0) + y(-1)] = [x \ -y]$$

Therefore, the standard transformation matrix for a reflection across the X -axis is $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$.

$$(D) \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

Key Points

- Reflection about the X -axis inverts the sign of the y -coordinate: $(x, y) \rightarrow (x, -y)$.
- Reflection about the Y -axis inverts the sign of the x -coordinate: $(x, y) \rightarrow (-x, y)$.

Q1.17.3 MCQ 2017 1M Ans: C ● ○ ○

Given matrix:

$$A = \begin{bmatrix} 1 & -1 & 5 \\ 0 & 5 & 6 \\ 0 & -6 & 5 \end{bmatrix}$$

The characteristic equation of a matrix A is given by $|A - \lambda I| = 0$:

$$\begin{vmatrix} 1-\lambda & -1 & 5 \\ 0 & 5-\lambda & 6 \\ 0 & -6 & 5-\lambda \end{vmatrix} = 0$$

$$(1 - \lambda) \{ (5 - \lambda)(5 - \lambda) - (6)(-6) \} = 0$$

$$\Rightarrow (1 - \lambda) \{ (5 - \lambda)^2 + 36 \} = 0$$

$$\Rightarrow (1 - \lambda)(\lambda^2 - 10\lambda + 25 + 36) = 0$$

$$\Rightarrow (1 - \lambda)(\lambda^2 - 10\lambda + 61) = 0$$

$$1 - \lambda = 0 \Rightarrow \lambda = 1$$

$$\lambda = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(1)(61)}}{2(1)}$$

$$\lambda = \frac{10 \pm \sqrt{100 - 244}}{2} = \frac{10 \pm \sqrt{-144}}{2}$$

$$\lambda = \frac{10 \pm 12j}{2} = 5 \pm 6j$$

$\therefore \lambda = 1, 5 \pm 6j$ are the required eigenvalues of the matrix A .

$$(C) 1, 5 \pm j6$$

Key Points

- Complex eigenvalues of a real matrix always occur in conjugate pairs ($\alpha \pm j\beta$).

Q1.16.1 NAT 2016 2M Ans: -6 ● ○ ○

If λ is an eigenvalue of a matrix A , then by polynomial matrix properties, $\lambda^3 - 3\lambda^2$ is an eigenvalue of the matrix $A^3 - 3A^2$.

Given that $\lambda = 1, -1$, and 3 are the particular eigenvalues of a matrix $A_{3 \times 3}$, we compute the corresponding eigenvalues for the matrix $A^3 - 3A^2$:

- For $\lambda = 1$:

$$\lambda^3 - 3\lambda^2 = (1)^3 - 3(1)^2 = 1 - 3 = -2$$

- For $\lambda = -1$:

$$\lambda^3 - 3\lambda^2 = (-1)^3 - 3(-1)^2 = -1 - 3 = -4$$

- For $\lambda = 3$:

$$\lambda^3 - 3\lambda^2 = (3)^3 - 3(3)^2 = 27 - 27 = 0$$

The particular eigenvalues of the matrix $A^3 - 3A^2$ are $-2, -4$, and 0 .

We know that the trace of a matrix is equal to the sum of its eigenvalues:

$$\text{Trace}(A^3 - 3A^2) = (-2) + (-4) + 0 = \boxed{-6}$$

Key Points

- Eigenvalue Polynomial Property:** If a square matrix A has eigenvalues λ_i , then any matrix polynomial $f(A)$ has corresponding eigenvalues $f(\lambda_i)$.
- Trace Property:** The trace of any square matrix ($\text{Tr}(M)$) is identically equal to the sum of all its eigenvalues.

Q1.15.1 MCQ 2015 1M Ans: C ● ○ ○

For a homogeneous system of linear equations represented as $AX = O$: The number of linearly independent solutions is given by $(n - r)$, where:

- n is the total number of variables.
- r is the rank of the coefficient matrix A .

Given that p denotes the number of linearly independent solutions:

$$\therefore p = n - r$$

$$\boxed{(C) \ n - r}$$

Q1.14.1 MCQ 2014 2M Ans: C ● ● ○

Let $n = 2$, $A = \begin{pmatrix} p & q \\ q & r \end{pmatrix}$, $X = \begin{pmatrix} x \\ y \end{pmatrix}$, and $B = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$. The function is defined as:

$$f(x) = X^T A X + B^T X + C$$

$$\implies f(x) = (px^2 + 2qxy + ry^2) + (b_1x + b_2y) + C$$

To find the stationary points, we compute the first partial derivatives and set them to zero:

$$\frac{\partial f}{\partial x} = 0 \implies 2px + 2qy + b_1 = 0 \quad \text{--- (1)}$$

$$\frac{\partial f}{\partial y} = 0 \implies 2qx + 2ry + b_2 = 0 \quad \text{--- (2)}$$

Equations (1) and (2) can be expressed simultaneously in matrix form as:

$$2 \begin{pmatrix} p & q \\ q & r \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = - \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$$

$$\implies 2AX = -B$$

$$\implies AX = -\frac{B}{2}$$

$$\therefore X = -\frac{A^{-1}B}{2} \quad \text{--- (3)}$$

Since matrix A is given to be positive definite, its determinant is strictly positive ($|A| > 0$) and its leading principal minors are positive ($p > 0$).

To check for a maximum or minimum, we evaluate the second-order partial derivatives:

$$rt - s^2 = 4(pr - q^2) > 0$$

Since $|A| = pr - q^2 > 0$ and $r = 2p > 0$, the discriminant condition $rt - s^2 > 0$ with $r > 0$ confirms a local minimum.

$$\therefore \text{At } X = -\frac{A^{-1}B}{2} \text{ we get the minimum of } f(x).$$

Similarly, this multi-variable minimization property holds true for $n > 2$.

$$\left(C - \left(\frac{A^{-1}b}{2} \right) \right)$$

Key Points

- **Matrix Differentiation:** The gradient of the quadratic form $X^TAX + B^TX + C$ with respect to vector X is given by $2AX + B = 0$.
- **Optimality Condition:** If the Hessian matrix (which is $2A$) is positive definite, the stationary point corresponds to a strict global minimum.

Mistakes to be avoided

- **Incorrect Sign During Matrix Inversion:** Forgetting to switch the sign of vector B when moving it to the right-hand side, leading to $X = \frac{A^{-1}B}{2}$ instead of the correct negative value.
- **Misinterpreting Positive Definiteness:** Confusing positive definite properties with maximum value conditions. Remember that a positive definite Hessian matrix always implies a local **minimum**, not a maximum.
- **Missing the Scaling Factor:** Omitting the factor of 2 when differentiating the quadratic term X^TAX , which leads to an incorrect stationary point of $X = -A^{-1}B$.

Q1.13.1 MCQ 2013 1M Ans: B ● ○ ○

The dimension of the null space (also known as the nullity) of a matrix A is determined using the Rank-Nullity Theorem. It is given by:

$$p = n - r$$

where:

- n is the total number of columns in matrix A (corresponding to the number of variables).
- r is the rank of the matrix A .

Given that $n = 3$ and $r = 2$:

$$\therefore p = 3 - 2 = 1$$

(B) 1

Key Points

- **Rank-Nullity Theorem:** State that for any matrix A with n columns, $\text{rank}(A) + \text{nullity}(A) = n$.
- The nullity represents the number of free variables available in the corresponding homogeneous system $AX = O$, indicating the dimension of its solution space.

Q1.13.2 MCQ 2013 2M Ans: MTA ● ● ○

Given matrix:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

(1) Eigenvalues:

The characteristic equation is given by $|A - \lambda I| = 0$:

$$\begin{vmatrix} -\lambda & -1 \\ 1 & -\lambda \end{vmatrix} = 0 \implies \lambda^2 + 1 = 0 \implies \lambda^2 = -1$$

$\therefore$ Eigenvalues of A are $\lambda = i, -i$

(2) Eigenvectors:

Consider the system $(A - \lambda I)X = O$:

$$\begin{bmatrix} 0 - \lambda & -1 \\ 1 & 0 - \lambda \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \text{--- (1)}$$

Case-(i): Put $\lambda = i$ in equation (1):

$$\begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

From the first row, we get the equation:

$$-ix_1 - x_2 = 0 \implies x_2 = -ix_1$$

Setting $x_1 = 1$ gives $x_2 = -i$, or setting $x_1 = i$ gives $x_2 = 1$.

$$\therefore X = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ -i \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} i \\ 1 \end{pmatrix}$$

Case-(ii): Put $\lambda = -i$ in equation (1):

$$\begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

From the first row, we get the equation:

$$ix_1 - x_2 = 0 \implies x_2 = ix_1$$

Setting $x_1 = 1$ gives $x_2 = i$, or setting $x_1 = i$ gives $x_2 = -1$.

$$\therefore X = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ i \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} i \\ -1 \end{pmatrix}$$

Hence, the eigenvectors of matrix A corresponding to eigenvalues $\lambda = i$ and $\lambda = -i$ can be written as:

$$X_1 = \begin{bmatrix} 1 \\ -i \end{bmatrix} \quad \text{and} \quad X_2 = \begin{bmatrix} 1 \\ i \end{bmatrix}$$

or

$$X_1 = \begin{bmatrix} i \\ 1 \end{bmatrix} \quad \text{and} \quad X_2 = \begin{bmatrix} i \\ -1 \end{bmatrix}$$

MTA

Key Points

- **Skew-Symmetric Matrix Property:** The given matrix A is a real skew-symmetric matrix ($A^T = -A$). The eigenvalues of any real skew-symmetric matrix are always purely imaginary or zero.
- **Conjugate Pair Property:** For a real matrix, if λ is a complex eigenvalue with eigenvector X , then its complex conjugate $\bar{\lambda}$ will have the corresponding conjugate eigenvector $\bar{X}$.

Q1.11.1 MCQ 2011 1M Ans: B ● ○ ○

For any positive integer m , a square matrix A and its power matrix A^m share the exact same set of eigenvectors. Given that a specific vector is an eigenvector of the base matrix M , it will remain an eigenvector for any higher power of that matrix ($M^2, M^3, \dots, M^m$).

$$\therefore \begin{bmatrix} 1 & 2 & -1 \end{bmatrix}^T \text{ is one of the eigenvectors of } M^3.$$

$$(B) \begin{bmatrix} 1 & 2 & -1 \end{bmatrix}^T$$

Key Points

- **Matrix Power Eigenproperty:** If X is an eigenvector of matrix A corresponding to the eigenvalue λ (i.e., $AX = \lambda X$), then for any integer $m \geq 1$, $A^m X = \lambda^m X$.
- This transformation preserves the directional alignment of the vector X , meaning the matrix powers change only the magnitude scaling factor (the eigenvalue), leaving the eigenvector itself unaltered.

Mistakes to be avoided

- **Raising the Eigenvector to a Power:** Mistakenly trying to modify or cube the entries of the eigenvector itself (e.g., computing $\begin{bmatrix} 1^3 & 2^3 & (-1)^3 \end{bmatrix}^T$). Remember that powers affect only the matrix and its associated eigenvalues, not the structural representation of the eigenvector.

Q1.10.1 MCQ 2010 1M Ans: A ● ○ ○

Given $A = (a_{ij})$, where the matrix elements are defined as:

$$a_{ij} = \begin{cases} i, & \forall i = j \\ 0, & \text{otherwise} \end{cases}$$

Expanding this definition into an $n \times n$ matrix structure:

$$A = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 2 & 0 & \cdots & 0 \\ 0 & 0 & 3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & n \end{bmatrix}_{n \times n}$$

Since A is a diagonal matrix, its eigenvalues are identically its main diagonal entries:

$$\Rightarrow \text{Eigenvalues of } A \text{ are } 1, 2, 3, \dots, n$$

$$\text{Sum of all eigenvalues} = 1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}$$

$$(A) \frac{n(n+1)}{2}$$

Q1.10.2 MCQ 2010 2M Ans: C ● ● ○

Given:

$$XY = O_{n \times n}$$

Let X be a non-singular matrix. Pre-multiplying both sides by X^{-1} , we get:

$$\begin{aligned} X^{-1}XY &= X^{-1}O \\ \Rightarrow Y &= O \end{aligned}$$

This is a contradiction since X and Y were given to be non-zero square matrices of the same order.

$$\therefore X \text{ is a singular matrix i.e., } |X| = 0$$

Similarly, we can show that Y is also a singular matrix i.e., $|Y| = 0$.

$$(C) |X| = 0 \text{ and } |Y| = 0$$

Key Points

- Zero Divisors in Matrices:** In matrix algebra, the product of two non-zero matrices can equal a zero matrix ($XY = O$). In this case, both X and Y are called zero divisors.
- Determinant Product Rule:** Taking the determinant of both sides of the equation gives $|XY| = |O| \Rightarrow |X||Y| = 0$. This mathematically establishes that at least one of the matrices must be singular, and further proof by contradiction confirms both must be singular if neither is zero.

Q1.09.1 MCQ 2009 1M Ans: B ● ○ ○

Given that $P \neq 0$ is a 3×3 real matrix.

We are given that there exist two linearly independent vectors x and y such that:

$$Px = 0 \quad \text{and} \quad Py = 0$$

This means that both vectors x and y belong to the null space (kernel) of the matrix P . Since x and y are linearly independent, the dimension of the null space, also known as the nullity of P , must be at least 2:

$$\text{Nullity}(P) \geq 2$$

According to the Rank-Nullity Theorem, for any $n \times n$ square matrix:

$$\text{Rank}(P) + \text{Nullity}(P) = n$$

Here, the matrix P is of order 3×3 , so $n = 3$:

$$\text{Rank}(P) = 3 - \text{Nullity}(P)$$

Since $\text{Nullity}(P) \geq 2$:

$$\text{Rank}(P) \leq 3 - 2 \implies \text{Rank}(P) \leq 1$$

We are also given that $P \neq 0$ (it is a non-zero matrix), which implies that its rank must be at least 1:

$$\text{Rank}(P) \geq 1$$

Combining the inequalities $\text{Rank}(P) \leq 1$ and $\text{Rank}(P) \geq 1$, we get:

$$\text{Rank}(P) = 1$$

Since the range space of a matrix is its column space, the dimension of the range space is exactly equal to the rank of the matrix:

$$\text{Dimension of range space of } P = \text{Rank}(P) = 1$$

(B) 1

Q1.09.2 MCQ 2009 2M Ans: C ● ○ ○

If the eigenvalues of X are $-2, -3$ and the eigenvalues of the identity matrix I are $1, 1$, then the eigenvalues of the matrix shift $(X + I)$ are calculated by adding the corresponding scalar coefficients:

$$-2 + 1, \quad -3 + 1 \implies -1, \quad -2$$

Consider the given rational matrix polynomial:

$$(X + I)^{-1}(X + 5I)$$

We can algebraically manipulate the expression to simplify it:

$$\begin{aligned} (X + I)^{-1}(X + 5I) &= (X + I)^{-1}(X + I + 4I) \\ \implies (X + I)^{-1}(X + 5I) &= (X + I)^{-1}(X + I) + 4(X + I)^{-1} \\ \implies (X + I)^{-1}(X + 5I) &= I + 4(X + I)^{-1} \end{aligned}$$

If λ is an eigenvalue of the base matrix X , then the eigenvalue mapping function for the expression $I + 4(X + I)^{-1}$ is given by:

$$f(\lambda) = 1 + 4(\lambda + 1)^{-1}$$

Now, we evaluate this mapping function for each specific eigenvalue of X :

- For $\lambda = -2$:

$$f(-2) = 1 + 4(-2 + 1)^{-1} = 1 + 4(-1)^{-1} = 1 - 4 = -3$$

- For $\lambda = -3$:

$$f(-3) = 1 + 4(-3 + 1)^{-1} = 1 + 4(-2)^{-1} = 1 - 2 = -1$$

$\therefore$ The eigenvalues of $I + 4(X + I)^{-1}$ are $-3, -1$.

Hence, the correct option is (c).

(C) $-1, -3$

Key Points

- **Spectral Mapping Theorem:** If λ is an eigenvalue of A , then for any rational function $r(x)$, $r(\lambda)$ is an eigenvalue of $r(A)$, provided the function is well-defined.
- **Inverse Eigenvalue Rule:** If a matrix M has non-zero eigenvalues μ_i , then its inverse matrix M^{-1} has the corresponding eigenvalues $\frac{1}{\mu_i}$.

Mistakes to be avoided

- **Direct Scalar Multiplication Mistakes:** Avoid applying properties blindly without decomposing the matrix expression first; trying to calculate $(X + I)^{-1}$ and $(X + 5I)$ eigenvalues separately and multiplying them directly can cause confusion if order isn't preserved.
- **Division by Zero Check:** Always verify that the eigenvalues of the shifted matrix $(X + I)$ are non-zero (here they are -1 and -2) before assuming that the inverse operation $(X + I)^{-1}$ is valid.

Q1.09.3 MCQ 2009 2M Ans: D ● ● ○

Given the rotation matrix:

$$P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

The trace of a 3×3 proper rotation matrix R that rotates a vector about a given axis by an angle θ satisfies the geometric formula:

$$\text{Tr}(R) = 1 + 2 \cos \theta$$

First, let us calculate the trace of the given matrix P by summing its principal diagonal entries:

$$\text{Tr}(P) = 0 + 0 + 0 = 0$$

Now, equate the trace value to the rotation equation formula:

$$1 + 2 \cos \theta = 0$$

$$\implies 2 \cos \theta = -1$$

$$\implies \cos \theta = -\frac{1}{2}$$

Solving for θ within the standard geometric range $[0^\circ, 180^\circ]$:

$$\theta = \cos^{-1} \left(-\frac{1}{2} \right) = 120^\circ$$

(D) 120°

Key Points

- **Axis of Rotation Verification:** The axis of rotation corresponds to the invariant eigenvector under the transformation matrix. Indeed, $P \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, confirming it has an eigenvalue of $\lambda = 1$.
- **Trace-Angle Relation:** For any standard 3×3 proper orthogonal rotation matrix, its trace represents the sum of its eigenvalues ($1 + e^{i\theta} + e^{-i\theta} = 1 + 2 \cos \theta$).

Q1.07.1 MCQ 2007 1M Ans: B ● ○ ○

Given the linear system matrix equation $A_{n \times n} X_{n \times n} = Y_{n \times 1}$ and the condition $A^2 = I$.
From the matrix property definition of powers:

$$A^2 = I \implies AA = I$$

By definition, if there exists a matrix B such that $AB = BA = I$, then A is invertible and $B = A^{-1}$. Here, the matrix A acts as its own inverse:

$$\implies A \text{ is invertible, i.e., } A^{-1} \text{ exists and } A^{-1} = A$$

$$\implies A \text{ is non-singular } (\det(A) \neq 0)$$

For any linear system of equations $AX = Y$, if the coefficient matrix A is non-singular (or invertible), we can uniquely pre-multiply both sides by A^{-1} :

$$X = A^{-1}Y$$

$\therefore$ A unique solution exists.

(B) a unique solution

Q1.07.2 MCQ 2007 2M Ans: B ● ● ○

Given matrix A of order $n \times n$:

$$A = \begin{bmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 4 & 6 & \cdots & 2n \\ 3 & 6 & 9 & \cdots & 3n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n & 2n & 3n & \cdots & 3n \end{bmatrix}_{n \times n}$$

Observing the rows, we can write them in terms of the first row R_1 :

$$R_1 = [1 \quad 2 \quad 3 \quad \cdots \quad n]$$

$$R_2 = [2 \quad 4 \quad 6 \quad \cdots \quad 2n] = 2R_1$$

$$R_3 = [3 \quad 6 \quad 9 \quad \cdots \quad 3n] = 3R_1$$

$\vdots$

$$R_n = [n \quad 2n \quad 3n \quad \cdots \quad 3n] = nR_1$$

We can perform elementary row operations to reduce the matrix to row echelon form:

$$R_i \rightarrow R_i - iR_1 \quad \forall i = 2, 3, \dots, n$$

$$A \sim \begin{bmatrix} 1 & 2 & 3 & \cdots & n \\ 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{bmatrix}_{n \times n}$$

$$\therefore \rho(A) = 1$$

The rank is 1 because matrix A possesses only one linearly independent row.

(B) 1

Key Points

- **Linearly Dependent Rows:** If all rows of a matrix can be expressed as scalar multiples of a single non-zero row ($R_i = c_i R_1$), the rank of that matrix is identically 1.

Q1.06.1 MCQ 2006 1M Ans: C ● ○ ○

Given the linear transformation mappings:

$$A \begin{bmatrix} 1 \\ -1 \end{bmatrix} = (-1) \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \text{and} \quad A \begin{bmatrix} 1 \\ -2 \end{bmatrix} = (-2) \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

From the definition of eigenvalues and eigenvectors ($AX = \lambda X$), it implies that:

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 \\ -2 \end{bmatrix}$$

are the eigenvectors of the matrix A corresponding to the eigenvalues $\lambda = -1$ and $\lambda = -2$, respectively. Now, we find the 2×2 matrix A using diagonalization principles. A square matrix $A_{n \times n}$ is diagonalizable if there exists a non-singular modal matrix P such that:

$$P^{-1}AP = D$$

where D is a diagonal matrix whose elements are the eigenvalues of A , and the columns of P are the corresponding linearly independent eigenvectors of A .

Constructing the matrices P and D :

$$P = \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \quad \text{and} \quad D = \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$$

$$P^{-1} = \frac{1}{(1)(-2) - (1)(-1)} \begin{bmatrix} -2 & -1 \\ 1 & 1 \end{bmatrix} = \frac{1}{-1} \begin{bmatrix} -2 & -1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix}$$

$$P^{-1}AP = D \implies A = PDP^{-1}$$

Substituting the matrices into the equation:

$$A = \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix}$$

$$(C) A = \begin{bmatrix} 1 & 1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ -1 & -1 \end{bmatrix}$$

Key Points

- **Matrix Diagonalization:** A matrix A can be reconstructed from its eigenvalues and eigenvectors using the similarity transformation formula $A = PDP^{-1}$.

Q1.05.1 MCQ 2005 1M Ans: B ☒ ☐ ☐

Given that the rank of the 3×3 matrix is 2, denoted as $\rho(A_{3 \times 3}) = 2$, and the total number of dimensions (columns) is $n = 3$. This implies that:

$$r = \rho(A_{3 \times 3}) = 2 < n = 3 = \text{number of variables}$$

For a homogeneous linear system $AX = O$, the number of linearly independent solutions is determined by the dimension of its null space (nullity). According to the Rank-Nullity Theorem, it is given by:

$$p = n - r$$

$$p = 3 - 2 = 1$$

(B) one independent solution

Q1.01.1 MCQ 2001 1M Ans: B ☒ ☐ ☐

A square matrix A of order n is diagonalizable if and only if it has n linearly independent eigen vectors.

(B) its eigen vectors should be independent

PrepFusion

SOLUTIONS — Linear Algebra



CS

Computer Science & IT

PrepFusion

Q1.26.1 MCQ 2026 1M Ans: A ● ● ○

Total number of 4×4 binary matrices: Each entry can be either 0 or 1.

$$\text{Total possible matrices} = 2^{16}$$

Consider a single row containing 4 entries. The total possible binary sequences for one row is $2^4 = 16$. Exactly half of these configurations will contain an even number of 1s (even parity rows):

$$\text{Even parity rows} = \frac{2^4}{2} = 2^3 = 8$$

For a matrix with 4 rows, the number of combinations where every row independently satisfies even parity is:

$$\text{Matrices with even parity rows} = (2^3)^4 = 2^{12}$$

Now, we look at the parity constraints for the columns within these 2^{12} matrices. Observe that if the first 3 columns have an even number of 1s, then the 4-th column will automatically be forced to have an even number of 1s.

$$\text{Total sum of all matrix entries} = \sum (\text{row sums})$$

Since every single row sum is even, the total sum of all elements in the matrix must be an even integer. Similarly, the total sum can also be written via column sums:

$$\text{Total sum of all matrix entries} = \sum (\text{column sums})$$

If the first 3 columns are even, their subtotal sum is even. Since the overall total sum must be even, the 4-th column sum is mathematically forced to be even as well. Hence, there are only 3 independent column conditions. Each independent column constraint halves the remaining valid configuration space:

$$2^{12} \rightarrow 2^{11} \rightarrow 2^{10} \rightarrow 2^9$$

$$2^9 = 512$$

(A) 512

Key Points

- **Parity Check Matrix Principles:** For any binary grid where all row and column sums must be even, the entries of the sub-matrix of size $(r - 1) \times (c - 1)$ can be chosen completely at random, while the remaining boundary row and column elements act as unique deterministic parity bits.
- The total number of solutions for an $m \times n$ binary matrix with even parity across all rows and columns is given explicitly by the formula $2^{(m-1)(n-1)}$.

Mistakes to be avoided

- **Over-constraining the System:** Treating all 4 column parity conditions as independent, which incorrectly divides the combinations by 2^4 instead of 2^3 , yielding an incorrect answer of $2^8 = 256$.

Q1.26.2 MCQ 2026 1M Ans: A ● ○ ○

For any square matrix A of order $n \times n$, the characteristic equation is a polynomial of degree n . According to the Fundamental Theorem of Algebra, a polynomial equation of degree n has exactly n roots (counting algebraic multiplicities). These roots represent the eigenvalues (λ) of the matrix. Therefore, for $n > 1$, any single eigenvalue λ can repeat at most n times. This maximum occurrence defines the peak algebraic multiplicity possible for a single distinct root within an n -dimensional linear space.

Hence, the maximum multiplicity of any eigenvalue is n .

(A) n

Key Points

- **Algebraic Multiplicity Bounds:** The algebraic multiplicity (m_λ) of an eigenvalue λ is defined as the number of times it appears as a root of the characteristic polynomial $p(\lambda) = \det(A - \lambda I) = 0$.
- For an $n \times n$ matrix, the sum of all algebraic multiplicities equals n , meaning $1 \leq m_\lambda \leq n$.

Q1.26.3 MSQ 2026 2M Ans: B,D ● ○ ○

Given that a vector $X = (0, 1, 0, 0, \dots, 0)^T \in \text{null space}(A)$.

By definition of the null space of a matrix:

$$AX = O$$

Since $X \neq O$, it represents a non-zero vector. This indicates that the homogeneous linear system $AX = O$ possesses a non-trivial solution. The existence of a non-trivial solution for a square homogeneous system implies:

- Matrix A is singular.
- The determinant of matrix A is zero, i.e., $|A| = 0$.

Therefore, option (b) is always correct. Furthermore, because the null space is a vector subspace over $\mathbb{R}$, it is closed under scalar multiplication. If $X \in \text{null space}(A)$, then for any scalar $c \in \mathbb{R}$:

$$cX \in \text{null space}(A)$$

Since there are infinitely many distinct choices for the scalar c ($c \neq 0$), there are infinitely many non-zero vectors present in the null space. This guarantees that at least two non-zero vectors exist in the null space of A . Therefore, option (d) is always correct.

(B) Determinant of M is 0

(D) There are at least two non-zero vectors in the null space of M

Key Points

- **Subspace Closure Property:** The null space (kernel) of any linear mapping matrix is a subspace. Thus, if a non-zero vector is found in it, the entire span of that vector is also contained within the null space.
- **Invertibility Criterion:** For a square matrix A , $\det(A) = 0$ is completely equivalent to stating that its null space contains non-trivial (non-zero) vectors.

Mistakes to be avoided

- **Assuming Minimum Rank Restrictions:** Avoid assuming option (c) is correct ($\text{Rank} = 1$). Finding one non-zero vector in the null space only establishes that $\text{Nullity} \geq 1 \implies \text{Rank} \leq n - 1$. It does not restrict the rank down to exactly 1.

Q1.26.4 NAT 2026 1M Ans: 24 ● ○ ○

Given system of linear equations:

$$ax + y = b \quad \text{--- (i)}$$

$$16x + ay = 24 \quad \text{--- (ii)}$$

For the system to produce multiple (infinitely many) solutions, the equations must be linearly dependent. This requires the determinant of the coefficient matrix A to be zero:

$$|A| = 0$$

$$\Rightarrow \begin{vmatrix} a & 1 \\ 16 & a \end{vmatrix} = 0$$

$$\Rightarrow a^2 - 16 = 0 \Rightarrow a = \pm 4$$

Let us analyze the system for each possible value of a :

Case 1: For $a = 4$ Substituting $a = 4$ into the equations:

$$(i) \Rightarrow 4x + y = b$$

$$(ii) \Rightarrow 16x + 4y = 24 \Rightarrow 4(4x + y) = 24 \Rightarrow 4x + y = 6$$

For infinite solutions, these two lines must coincide perfectly, which means:

$$\rho(A) = \rho(A|B) = 1 \Rightarrow b = 6$$

However, checking the question parameters, if $a = 4$ and $b = 6$, their product is $a \cdot b = 4 \times 6 = 24$.

Case 2: For $a = -4$ Substituting $a = -4$ into the equations:

$$(i) \Rightarrow -4x + y = b \Rightarrow 4x - y = -b$$

$$(ii) \Rightarrow 16x - 4y = 24 \Rightarrow 4(4x - y) = 24 \Rightarrow 4x - y = 6$$

For infinite solutions, these equations must represent the same identical line:

$$\rho(A) = \rho(A|B) = 1$$

$$\Rightarrow -b = 6 \Rightarrow b = -6$$

Thus, a valid parameter pairing that yields multiple solutions is $a = -4$ and $b = -6$.

Calculating the required product of a and b :

$$a \cdot b = (-4) \times (-6) = \boxed{24}$$

Key Points

- **Infinite Solutions Condition:** A 2×2 system of linear equations has infinite solutions if the rank of both the coefficient matrix and the augmented matrix equals 1 ($\rho(A) = \rho(A|B) = 1$). Geometric lines must be completely collinear.
- This matching condition corresponds identically to satisfying proportional consistency: $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$.

Q1.26.5 NAT 2026 2M Ans: 48 ● ● ○

Given that the matrix A is of order 4×4 and its determinant is $|A| = 3$.

Using the scaling property of determinants for a square matrix of order $n \times n$:

$$|kA| = k^n |A|$$

Here, the scaling factor is $k = 2$ and the order of the matrix is $n = 4$:

$$|2A_{4 \times 4}| = 2^4 |A|$$

$$\Rightarrow |2A| = 16 \times 3 = \boxed{48}$$

Key Points

- **Determinant Scaling Property:** When an $n \times n$ matrix is multiplied by a scalar k , every row is multiplied by k . Since pulling out a common factor from a single row scales the determinant by that factor, factoring it out from all n rows yields $|kA| = k^n |A|$.

Mistakes to be avoided

- Avoid simply multiplying the determinant by the scalar factor directly (i.e., computing $2 \times 3 = 6$). Remember that matrix scaling scales the determinant exponentially with respect to its dimensional order n .

Q1.25.1 MSQ 2025 1M Ans: A,D ● ○ ○

Given system of linear equations:

$$x + ky = 1 \quad \text{--- (i)}$$

$$kx + y = -1 \quad \text{--- (ii)}$$

The system can be represented in matrix form $AX = B$ as:

$$\begin{bmatrix} 1 & k \\ k & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

The determinant of the coefficient matrix A is:

$$|A| = 1 - k^2$$

Let us examine the solutions based on the values of k :

Case 1: When $|A| \neq 0$

$$1 - k^2 \neq 0 \Rightarrow k \neq \pm 1$$

When $k \in \mathbb{R} \setminus \{-1, 1\}$, the matrix A is invertible ($\rho(A) = 2$), which guarantees a unique (exactly one) solution. Since there are infinitely many real values of k excluding ± 1 , option (C) is incorrect.

Case 2: When $|A| = 0$

$$1 - k^2 = 0 \Rightarrow k = 1 \text{ or } k = -1$$

Let us evaluate these two sub-cases using the augmented matrix $[A|B]$:

Sub-case 2a: For $k = 1$ The equations become:

$$x + y = 1$$

$$x + y = -1$$

These represent parallel lines that never intersect. Let's verify via row reduction on the augmented matrix:

$$[A|B] = \left[\begin{array}{cc|c} 1 & 1 & 1 \\ 1 & 1 & -1 \end{array} \right] \xrightarrow{R_2 \leftarrow R_2 - R_1} \left[\begin{array}{cc|c} 1 & 1 & 1 \\ 0 & 0 & -2 \end{array} \right]$$

Here, $\rho(A) = 1$ and $\rho(A|B) = 2$. Since $\rho(A) \neq \rho(A|B)$, the system is inconsistent and has no solution. Thus, there is exactly one value of k ($k = 1$) for which the system has no solution. Option (A) is correct, and option (B) is incorrect.

Sub-case 2b: For $k = -1$ The equations become:

$$x - y = 1$$

$$-x + y = -1 \implies x - y = 1$$

Both equations represent the exact same identical line. Let's verify via row reduction on the augmented matrix:

$$[A|B] = \left[\begin{array}{cc|c} 1 & -1 & 1 \\ -1 & 1 & -1 \end{array} \right] \xrightarrow{R_2 \leftarrow R_2 + R_1} \left[\begin{array}{cc|c} 1 & -1 & 1 \\ 0 & 0 & 0 \end{array} \right]$$

Here, $\rho(A) = \rho(A|B) = 1 < 2$ (number of variables). The system is consistent with a free variable, yielding an infinite number of solutions. Thus, there exists exactly one value of k ($k = -1$) for which the system has an infinite number of solutions. Option (D) is correct.

(A) There is exactly one value of k for which the above system of equations has no solution.

(D) There exists exactly one value of k for which the system of equations has an infinite number of solutions.

Key Points

- **Consistency Conditions (Rouché-Capelli Theorem):** For a system $AX = B$:
 - Unique Solution: $\rho(A) = \rho(A|B) = \text{number of variables}$.
 - Infinite Solutions: $\rho(A) = \rho(A|B) < \text{number of variables}$.
 - No Solution: $\rho(A) \neq \rho(A|B)$.
- **Geometric Interpretation:** In a 2×2 real linear system, the equations represent two lines which can either intersect at one point (unique solution), be parallel and distinct (no solution), or be coincident (infinite solutions).

Mistakes to be avoided

- **Assuming $|A| = 0$ Always Means No Solution:** Do not conclude that both $k = 1$ and $k = -1$ lead to no solution. You must check the augmented rank consistency for each root separately to distinguish between an inconsistent system and an underdetermined system.

Q1.25.2 MCQ 2025 2M Ans: D ● ● ○

Given the 2×2 matrix A :

$$A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$\det(A - \lambda I) = 0$$

$$\implies \begin{vmatrix} 1 - \lambda & 1 \\ 1 & -1 - \lambda \end{vmatrix} = 0$$

$$\implies (1 - \lambda)(-1 - \lambda) - 1 = 0$$

$$\implies -(1 - \lambda^2) - 1 = 0 \implies \lambda^2 - 2 = 0 \implies \lambda = \pm\sqrt{2}$$

Thus, the eigenvalues of matrix A are $\lambda_1 = \sqrt{2}$ and $\lambda_2 = -\sqrt{2}$. According to the Spectral Mapping Theorem, if a matrix A has eigenvalues λ , then the matrix power A^n has eigenvalues λ^n .

Therefore, the eigenvalues of the matrix A^{13} are:

$$\lambda_1^{13} = (\sqrt{2})^{13} = (\sqrt{2})^{12} \cdot \sqrt{2} = 2^6 \cdot \sqrt{2} = 64\sqrt{2}$$

$$\lambda_2^{13} = (-\sqrt{2})^{13} = -(\sqrt{2})^{12} \cdot \sqrt{2} = -2^6 \cdot \sqrt{2} = -64\sqrt{2}$$

Hence, the eigenvalues of A^{13} are $64\sqrt{2}, -64\sqrt{2}$.

$$(D) \ 64\sqrt{2}, -64\sqrt{2}$$

Key Points

- **Eigenvalues of Matrix Powers:** If λ is an eigenvalue of A with a corresponding eigenvector X , then λ^n is an eigenvalue of A^n with the identical corresponding eigenvector X .
- **Characteristic Equation of a 2×2 Matrix:** Can be written directly using the trace and determinant: $\lambda^2 - \text{Tr}(A)\lambda + \det(A) = 0$.

Q1.25.3 MCQ 2025 1M Ans: C ☒ ☐ ☐

Given the 2×2 matrix A :

$$A = \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = A \cdot A = \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix}$$

$$A^2 = \begin{pmatrix} (1)(1) + (2)(2) & (1)(2) + (2)(-1) \\ (2)(1) + (-1)(2) & (2)(2) + (-1)(-1) \end{pmatrix}$$

$$A^2 = \begin{pmatrix} 1+4 & 2-2 \\ 2-2 & 4+1 \end{pmatrix} = \begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix} = 5I$$

$$A^8 = (A^2)^4$$

$$\Rightarrow A^8 = (5I)^4 = 5^4 \cdot I^4$$

$$5^4 = 625$$

$$\Rightarrow A^8 = 625I = \begin{pmatrix} 625 & 0 \\ 0 & 625 \end{pmatrix}$$

$$(C) \ \begin{pmatrix} 625 & 0 \\ 0 & 625 \end{pmatrix}$$

Q1.25.4 MCQ 2025 1M Ans: A ☒ ☐ ☐

Given that L , M , and N are non-singular matrices of order 3 satisfying the following equations:

$$L^2 = L^{-1} \quad \text{--- (i)}$$

$$M = L^8 \quad \text{--- (ii)}$$

$$N = L^2 \quad \text{--- (iii)}$$

We want to find the value of the determinant $\det(M - N)$. Let us first simplify the matrix expression for M by leveraging the relationship given in equation (i).

Post-multiplying both sides of equation (i) by matrix L :

$$L^2 \cdot L = L^{-1} \cdot L$$

$$\Rightarrow L^3 = I$$

Now, we can decompose the higher power matrix exponent of $M = L^8$ into multiples of L^3 :

$$M = L^8 = L^6 \cdot L^2 = (L^3)^2 \cdot L^2$$

$$M = (I)^2 \cdot L^2 = I \cdot L^2 = L^2$$

$$M - N = L^2 - L^2 = O$$

$$\det(M - N) = \det(O) = 0$$

(A) 0

Key Points

- **Matrix Order Reductions:** Whenever a matrix equation yields a periodic identity property such as $L^k = I$, any higher exponent power L^n can be broken down using modular arithmetic on its power index: $L^n = L^{n \pmod k}$.
- **Properties of the Null Matrix:** Subtracting a matrix from its identical algebraic form results in the zero matrix (O), whose determinant is strictly $\det(O) = 0$.

Q1.25.5 MSQ 2025 2M Ans: A,B,C ● ● ●

Given the linear system $PX = Q$ and its LU decomposition $P = LU$, where L is a unit lower triangular matrix and U is an upper triangular matrix:

$$L = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix}, \quad U = \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

Let us verify each option systematically:

(A): The system is $PX = Q$. Substituting $P = LU$ yields:

$$(LU)X = Q \implies L(UX) = Q$$

Let $UX = Y$. Then the system decomposes cleanly into two sequential triangular systems:

1. Solve $LY = Q$ for Y using forward substitution.

2. Solve $UX = Y$ for X using back substitution.

Therefore, statement (A) is **TRUE**.

(B): Using the multiplicative property of determinants:

$$\det(P) = \det(LU) = \det(L) \cdot \det(U)$$

Since L is a unit lower triangular matrix, its determinant is the product of its diagonal entries:

$$\det(L) = 1 \cdot 1 \cdot 1 = 1$$

Thus, $\det(P) = 1 \cdot \det(U) = \det(U)$. If P is invertible, then $\det(P) \neq 0 \implies \det(U) \neq 0$. Since both $\det(L) = 1 \neq 0$ and $\det(U) \neq 0$, both matrices L and U are invertible. Therefore, statement (B) is **TRUE**.

(C): If P is singular, then $\det(P) = 0$. From our previous relation:

$$\det(P) = \det(L) \cdot \det(U) \implies 0 = 1 \cdot \det(U) \implies \det(U) = 0$$

Since U is an upper triangular matrix, its determinant is simply the product of its principal diagonal entries:

$$\det(U) = u_{11} \cdot u_{22} \cdot u_{33} = 0$$

For the product to equal zero, at least one of the diagonal elements (u_{11} , u_{22} , or u_{33}) must be zero. Therefore, statement (C) is **TRUE**.

(D): If P is symmetric, $P = P^T$. However, this does not imply that L and U are individually symmetric. In fact, because L is strictly lower triangular with non-zero off-diagonal entries l_{21} , l_{31} , l_{32} in general, it cannot be symmetric unless it equals the identity matrix I . Instead, if P is symmetric, its decomposition satisfies $P = LDL^T$ (where $U = DL^T$). Therefore, statement (D) is **FALSE**.

(A) The system $PX = Q$ can be solved by first solving $LY = Q$ and then $UX = Y$.

(B) If P is invertible, then both L and U are invertible.

(C) If P is singular, then at least one of the diagonal elements of U is zero.

Q1.24.1 MCQ 2024 1M Ans: B ☒ ☐ ☐

Let the given matrix be A :

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$\prod_{i=1}^n \lambda_i = \det(A)$$

$$\det(A) = 1 \begin{vmatrix} 5 & 6 \\ 8 & 9 \end{vmatrix} - 2 \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix} + 3 \begin{vmatrix} 4 & 5 \\ 7 & 8 \end{vmatrix}$$

$$\det(A) = 1(-3) - 2(-6) + 3(-3)$$

$$\Rightarrow \det(A) = -3 + 12 - 9 = 0$$

Since the determinant is 0, the product of all the eigenvalues of matrix A must be 0.

(B) 0

Q1.24.2 MSQ 2024 2M Ans: A ☒ ☐ ☐

Given the homogeneous system of linear equations $Ax = \mathbf{0}$, where A is an $n \times m$ matrix with $m > n$. This implies there are more variables (m) than equations (n), making it an underdetermined system.

Let us evaluate the statements based on the fundamental principles of rank and nullity:

(A): By the Rank-Nullity Theorem, the dimension of the solution space (the null space) is given by:

$$\text{Nullity}(A) = m - \text{Rank}(A)$$

Since the maximum possible rank of any $n \times m$ matrix is bounded by its smaller dimension:

$$\text{Rank}(A) \leq n$$

Substituting this upper bound into the nullity equation yields:

$$\text{Nullity}(A) \geq m - n$$

Since nullity represents the exact number of linearly independent vectors that span the solution space, there exist at least $m - n$ linearly independent solutions to this system. Therefore, statement (A) is strictly and always **TRUE**.

(B): Statement (B) asserts that there exist exactly $m - n$ linearly independent vectors that span every solution. This is a special case that only holds true if matrix A has full row rank ($\text{Rank}(A) = n$). If $\text{Rank}(A) < n$, the nullity will be strictly greater than $m - n$. Therefore, statement (B) is **FALSE**.

(A) There exist at least $m - n$ linearly independent solutions to this system

Key Points

- **Rank-Nullity Theorem:** For any linear transformation matrix $A_{n \times m}$, the sum of the pivot columns and free columns equals the total columns: $\text{Rank}(A) + \text{Nullity}(A) = m$.
- **Dimension of Vector Spaces:** The term "linearly independent solutions" refers directly to the basis vectors of the null space $N(A)$.

Q1.24.3 MSQ 2024 2M Ans: A,B ● ○ ○

Let A be an $n \times n$ matrix over the set of all real numbers $\mathbb{R}$. Let B be the matrix obtained from A by swapping two rows. This operation can be mathematically represented as multiplying A by an elementary permutation matrix E , such that $B = EA$.

(A): By the multiplicative property of determinants:

$$\det(B) = \det(EA) = \det(E) \cdot \det(A)$$

The elementary matrix E corresponds to a single row swap applied to the identity matrix I . Since swapping any two rows of the identity matrix changes the sign of its determinant, we have $\det(E) = -1$.

$$\implies \det(B) = -\det(A)$$

Therefore, the determinant of B is the negative of the determinant of A . Statement (A) is **TRUE**.

(B): A matrix is invertible if and only if its determinant is non-zero. From the previous relation, $|\det(B)| = |\det(A)|$. If A is invertible:

$$\det(A) \neq 0 \implies \det(B) = -\det(A) \neq 0$$

Since $\det(B) \neq 0$, matrix B is also invertible. Statement (B) is **TRUE**.

(C): Let us analyze a counterexample to test symmetry. Consider a symmetric matrix A :

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix} \implies A^T = A$$

If we swap the first row and the second row of A to get B :

$$B = \begin{bmatrix} 2 & 5 \\ 1 & 2 \end{bmatrix}$$

$$B^T = \begin{bmatrix} 2 & 1 \\ 5 & 2 \end{bmatrix} \neq B$$

Thus, swapping rows does not preserve matrix symmetry. Statement (C) is **FALSE**.

(D): The trace of a matrix is defined as the sum of its principal diagonal elements. Swapping two rows alters which elements sit on the principal diagonal. Consider a matrix A where $\text{Tr}(A) = 0$:

$$A = \begin{bmatrix} 1 & 3 \\ 4 & -1 \end{bmatrix} \implies \text{Tr}(A) = 1 + (-1) = 0$$

Swapping the first row and the second row to form B :

$$B = \begin{bmatrix} 4 & -1 \\ 1 & 3 \end{bmatrix}$$

$$\text{Tr}(B) = 4 + 3 = 7 \neq 0$$

Thus, swapping rows does not preserve a zero trace. Statement (D) is **FALSE**.

(A) The determinant of B is the negative of the determinant of A

(B) If A is invertible, then B is also invertible

Q1.23.1 MCQ 2023 1M Ans: B ● ○ ○

Given the matrices:

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 2 & 3 \\ 3 & 4 & 1 & 2 \\ 2 & 3 & 4 & 1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \end{bmatrix}$$

Let us analyze the relationship between the rows of matrix A and matrix B . Let R_1, R_2, R_3, R_4 denote the rows of matrix A :

$$R_1 = [1 \quad 2 \quad 3 \quad 4]$$

$$R_2 = [4 \quad 1 \quad 2 \quad 3]$$

$$R_3 = [3 \quad 4 \quad 1 \quad 2]$$

$$R_4 = [2 \quad 3 \quad 4 \quad 1]$$

Looking at matrix B , its rows can be expressed in terms of the rows of A as:

$$B = \begin{bmatrix} R_3 \\ R_2 \\ R_1 \\ R_4 \end{bmatrix}$$

To transform matrix A into matrix B , we can perform row interchange operations:

- Interchange the first row (R_1) and the third row (R_3):

$$\begin{bmatrix} R_1 \\ R_2 \\ R_3 \\ R_4 \end{bmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \begin{bmatrix} R_3 \\ R_2 \\ R_1 \\ R_4 \end{bmatrix} = B$$

Since exactly one row interchange operation (number of interchanges = 1) is required to transform matrix A into matrix B , the determinant changes its sign exactly once:

$$\det(B) = (-1)^1 \det(A) = -\det(A)$$

Therefore, the relation $\det(B) = -\det(A)$ holds true.

$$(B) \det(B) = -\det(A)$$

Key Points

- Interchanging any two rows (or columns) of a matrix changes the sign of its determinant.
- If a matrix B is obtained from A by performing k row interchanges, then $\det(B) = (-1)^k \det(A)$.

Q1.22.1 MCQ 2022 2M Ans: D ● ● ○

The given system of simultaneous equations is:

$$x_1 + x_2 - 2x_3 = 4$$

$$x_1 + 3x_2 - x_3 = 7$$

$$2x_1 + x_2 - 5x_3 = 7$$

This system can be written in matrix form $Ax = b$ as:

$$\begin{bmatrix} 1 & 1 & -2 \\ 1 & 3 & -1 \\ 2 & 1 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \\ 7 \end{bmatrix}$$

Using Doolittle's algorithm for LU decomposition, where the diagonal elements of L are taken as 1 ($L_{11} = L_{22} = L_{33} = 1$):

$$A = LU$$

$$\begin{bmatrix} 1 & 1 & -2 \\ 1 & 3 & -1 \\ 2 & 1 & -5 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ L_{21} & 1 & 0 \\ L_{31} & L_{32} & 1 \end{bmatrix} \begin{bmatrix} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{bmatrix}$$

By performing matrix multiplication row by row:

First Row of U :

$$U_{11} = 1, \quad U_{12} = 1, \quad U_{13} = -2$$

First Column of L :

$$L_{21}U_{11} = 1 \implies L_{21}(1) = 1 \implies L_{21} = 1$$

$$L_{31}U_{11} = 2 \implies L_{31}(1) = 2 \implies L_{31} = 2$$

Second Row of U :

$$L_{21}U_{12} + U_{22} = 3 \implies (1)(1) + U_{22} = 3 \implies U_{22} = 2$$

$$L_{21}U_{13} + U_{23} = -1 \implies (1)(-2) + U_{23} = -1 \implies U_{23} = 1$$

Remaining elements (L_{32} and U_{33}):

$$L_{31}U_{12} + L_{32}U_{22} = 1$$

$$(2)(1) + L_{32}(2) = 1 \implies 2L_{32} = -1 \implies L_{32} = -\frac{1}{2}$$

$$L_{31}U_{13} + L_{32}U_{23} + U_{33} = -5$$

$$(2)(-2) + \left(-\frac{1}{2}\right)(1) + U_{33} = -5$$

$$-4 - \frac{1}{2} + U_{33} = -5 \implies U_{33} = -5 + 4 + \frac{1}{2} = -\frac{1}{2}$$

Now, we solve for x using $LY = b$ and $Ux = x$:

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 2 & -\frac{1}{2} & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \\ 7 \end{bmatrix}$$

$$y_1 = 4$$

$$y_1 + y_2 = 7 \implies 4 + y_2 = 7 \implies y_2 = 3$$

$$2y_1 - \frac{1}{2}y_2 + y_3 = 7 \implies 2(4) - \frac{1}{2}(3) + y_3 = 7 \implies 8 - 1.5 + y_3 = 7 \implies y_3 = 0.5$$

Now solving $Ux = Y$:

$$\begin{bmatrix} 1 & 1 & -2 \\ 0 & 2 & 1 \\ 0 & 0 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ \frac{1}{2} \end{bmatrix}$$

$$-\frac{1}{2}x_3 = \frac{1}{2} \implies x_3 = -1$$

$$2x_2 + x_3 = 3 \implies 2x_2 - 1 = 3 \implies 2x_2 = 4 \implies x_2 = 2$$

$$x_1 + x_2 - 2x_3 = 4 \implies x_1 + 2 - 2(-1) = 4 \implies x_1 + 4 = 4 \implies x_1 = 0$$

Thus, the correct values are $L_{32} = -\frac{1}{2}$, $U_{33} = -\frac{1}{2}$, and $x_1 = 0$.

$$(D) L_{32} = -\frac{1}{2}, U_{33} = -\frac{1}{2}, x_1 = 0$$

Key Points

- LU Decomposition factors a matrix A into a lower triangular matrix L and an upper triangular matrix U .
- In Doolittle's algorithm, the principal diagonal elements of L are equal to 1.
- Once decomposed, the system is efficiently solved using sequential forward and backward substitutions.

Q1.22.2 MSQ 2022 2M Ans: A,C,D

Let the given matrix be A :

$$A = \begin{bmatrix} -9 & -6 & -2 & -4 \\ -8 & -6 & -3 & -1 \\ 20 & 15 & 8 & 5 \\ 32 & 21 & 7 & 12 \end{bmatrix}$$

By definition, a non-zero vector X is an eigenvector of A if it satisfies the condition $AX = \lambda X$, which means the resultant vector AX must be a scalar multiple of X . Let us verify each option by performing matrix multiplication AX :

Option A:

$$X_A = \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

$$AX_A = \begin{bmatrix} -9 & -6 & -2 & -4 \\ -8 & -6 & -3 & -1 \\ 20 & 15 & 8 & 5 \\ 32 & 21 & 7 & 12 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} (-9)(-1) + (-6)(1) + (-2)(0) + (-4)(1) \\ (-8)(-1) + (-6)(1) + (-3)(0) + (-1)(1) \\ (20)(-1) + (15)(1) + (8)(0) + (5)(1) \\ (32)(-1) + (21)(1) + (7)(0) + (12)(1) \end{bmatrix}$$

$$AX_A = \begin{bmatrix} 9 - 6 - 4 \\ 8 - 6 - 1 \\ -20 + 15 + 5 \\ -32 + 21 + 12 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = 1 \cdot X_A$$

Since $AX_A = 1 \cdot X_A$, X_A is an eigenvector corresponding to the eigenvalue $\lambda = 1$. Thus, Option A is correct.

Option B:

$$X_B = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix}$$

$$AX_B = \begin{bmatrix} -9 & -6 & -2 & -4 \\ -8 & -6 & -3 & -1 \\ 20 & 15 & 8 & 5 \\ 32 & 21 & 7 & 12 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -9 + 2 \\ -8 + 3 \\ 20 - 8 \\ 32 - 7 \end{bmatrix} = \begin{bmatrix} -7 \\ -5 \\ 12 \\ 25 \end{bmatrix}$$

The vector $\begin{bmatrix} -7 & -5 & 12 & 25 \end{bmatrix}^T$ is not a scalar multiple of $\begin{bmatrix} 1 & 0 & -1 & 0 \end{bmatrix}^T$. Thus, Option B is incorrect.

Option C:

$$X_C = \begin{bmatrix} -1 \\ 0 \\ 2 \\ 2 \end{bmatrix}$$

$$AX_C = \begin{bmatrix} -9 & -6 & -2 & -4 \\ -8 & -6 & -3 & -1 \\ 20 & 15 & 8 & 5 \\ 32 & 21 & 7 & 12 \end{bmatrix} \begin{bmatrix} -1 \\ 0 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 9 - 4 - 8 \\ 8 - 6 - 2 \\ -20 + 16 + 10 \\ -32 + 14 + 24 \end{bmatrix} = \begin{bmatrix} -3 \\ 0 \\ 6 \\ 6 \end{bmatrix} = 3 \begin{bmatrix} -1 \\ 0 \\ 2 \\ 2 \end{bmatrix} = 3 \cdot X_C$$

Since $AX_C = 3 \cdot X_C$, X_C is an eigenvector corresponding to the eigenvalue $\lambda = 3$. Thus, Option C is correct.

Option D:

$$X_D = \begin{bmatrix} 0 \\ 1 \\ -3 \\ 0 \end{bmatrix}$$

$$AX_D = \begin{bmatrix} -9 & -6 & -2 & -4 \\ -8 & -6 & -3 & -1 \\ 20 & 15 & 8 & 5 \\ 32 & 21 & 7 & 12 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ -3 \\ 0 \end{bmatrix} = \begin{bmatrix} -6 + 6 \\ -6 + 9 \\ 15 - 24 \\ 21 - 21 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ -9 \\ 0 \end{bmatrix} = 3 \begin{bmatrix} 0 \\ 1 \\ -3 \\ 0 \end{bmatrix} = 3 \cdot X_D$$

Since $AX_D = 3 \cdot X_D$, X_D is an eigenvector corresponding to the eigenvalue $\lambda = 3$. Thus, Option D is correct.

$$(A) \begin{pmatrix} -1 \\ 1 \\ 0 \\ 1 \end{pmatrix}, (C) \begin{pmatrix} -1 \\ 0 \\ 2 \\ 2 \end{pmatrix}, (D) \begin{pmatrix} 0 \\ 1 \\ -3 \\ 0 \end{pmatrix}$$

Key Points

- A vector $X \neq 0$ is an eigenvector of a square matrix A if $AX = \lambda X$ for some scalar λ .
- Multiplying the matrix directly by the given options is often much faster for multiple-select questions (MSQs) than computing the characteristic polynomial $\det(A - \lambda I) = 0$ for high-dimensional matrices.

Q1.21.1 NAT 2021 2M Ans: 3 ☒ ☐ ☐

To find the eigenvalues of the matrix, we solve the characteristic equation $\det(A - \lambda I) = 0$.

$$|A - \lambda I| = \begin{vmatrix} -\lambda & 1 & 1 & 1 \\ 1 & -\lambda & 1 & 1 \\ 1 & 1 & -\lambda & 1 \\ 1 & 1 & 1 & -\lambda \end{vmatrix} = 0$$

Step 1: $R_1 \rightarrow R_1 + R_2 + R_3 + R_4$

$$\begin{vmatrix} -\lambda + 3 & -\lambda + 3 & -\lambda + 3 & -\lambda + 3 \\ 1 & -\lambda & 1 & 1 \\ 1 & 1 & -\lambda & 1 \\ 1 & 1 & 1 & -\lambda \end{vmatrix} = 0$$

Applying the column operations $C_2 \rightarrow C_2 - C_1$, $C_3 \rightarrow C_3 - C_1$, and $C_4 \rightarrow C_4 - C_1$:

$$\begin{vmatrix} -\lambda + 3 & 0 & 0 & 0 \\ 1 & -\lambda - 1 & 0 & 0 \\ 1 & 0 & -\lambda - 1 & 0 \\ 1 & 0 & 0 & -\lambda - 1 \end{vmatrix} = 0$$

$$(-\lambda + 3)(-\lambda - 1)^3 = 0$$

$$\lambda = 3, -1, -1, -1$$

Thus, the largest eigenvalue of the matrix is 3.

3

Q1.21.2 NAT 2021 1M Ans: 4 ● ○ ○

Given that P is a matrix of dimension 4×5 . The system of equations is given as $PX = 0$. Given that every solution of $PX = 0$ is a scalar multiple of the non-zero column vector:

$$\begin{bmatrix} 2 \\ 5 \\ 4 \\ 1 \\ 3 \end{bmatrix}$$

This implies that the solution space (null space) is spanned by this single vector, meaning that $[2 \ 5 \ 4 \ 1 \ 3]^T$ is the only linearly independent solution. Therefore, the number of linearly independent solutions (which is the nullity of matrix P) is equal to 1:

$$\text{Nullity}(P) = n - r = 1$$

where r is the rank of the matrix P .

$$5 - r = 1$$

$$r = 4$$

Thus, the rank of P is 4.

4

Key Points

- Rank-Nullity Theorem: For any matrix P of size $m \times n$, $\text{Rank}(P) + \text{Nullity}(P) = n$, where n is the number of columns.

Q1.20.1 MCQ 2020 1M Ans: C ● ○ ○

From the fundamental properties of matrices and determinants, we know that for any two square matrices A and B of the same order:

$$\det(AB) = \det(A) \cdot \det(B)$$

Additionally, considering the properties of matrix ranks, the rank of the sum of two matrices satisfies the subadditivity property:

$$\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$$

Therefore, the statement corresponding to option (c) is correct.

(C) II and III only

Key Points

- Multiplicative Property of Determinants: $\det(AB) = \det(A) \det(B)$ holds true for all $n \times n$ square matrices.
- Rank Subadditivity: $\text{rank}(A + B) \leq \text{rank}(A) + \text{rank}(B)$.
- Rank Product Inequalities: $\text{rank}(AB) \leq \min(\text{rank}(A), \text{rank}(B))$.

Mistakes to be avoided

- Never assume that determinants are linear under addition, i.e., $\det(A + B) \neq \det(A) + \det(B)$ in general.
- Avoid assuming $\text{rank}(A + B) = \text{rank}(A) + \text{rank}(B)$ unless the column spaces (or row spaces) of A and B are completely independent.

Q1.19.1 NAT 2019 2M Ans: 12 ● ● ○

Given matrix R :

$$R = \begin{bmatrix} 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \end{bmatrix}$$

From the properties of eigenvalues, the product of the eigenvalues of a matrix is equal to its determinant:

Product of eigenvalues of $R = |R|$

$$|R| = \begin{vmatrix} 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \end{vmatrix}$$

Applying the row operations $R_4 \rightarrow R_4 - R_3$, $R_3 \rightarrow R_3 - R_2$, and $R_2 \rightarrow R_2 - R_1$:

$$\Rightarrow |R| = \begin{vmatrix} 1 & 2 & 4 & 8 \\ 0 & 1 & 5 & 19 \\ 0 & 1 & 7 & 37 \\ 0 & 1 & 9 & 61 \end{vmatrix}$$

Next, applying the row operations $R_4 \rightarrow R_4 - R_3$ and $R_3 \rightarrow R_3 - R_2$:

$$\Rightarrow |R| = \begin{vmatrix} 1 & 2 & 4 & 8 \\ 0 & 1 & 5 & 19 \\ 0 & 0 & 2 & 18 \\ 0 & 0 & 2 & 24 \end{vmatrix}$$

Finally, applying the row operation $R_4 \rightarrow R_4 - R_3$:

$$\Rightarrow |R| = \begin{vmatrix} 1 & 2 & 4 & 8 \\ 0 & 1 & 5 & 19 \\ 0 & 0 & 2 & 18 \\ 0 & 0 & 0 & 6 \end{vmatrix}$$

$$\Rightarrow |R| = 1 \times 1 \times 2 \times 6 = 12$$

Therefore, the absolute value of the product of the eigenvalues is 12.

12

Q1.18.1 NAT 2018 1M Ans: 3 ● ○ ○

Given $A = uv^T$, where:

$$u = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad v = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow A = uv^T = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$$

To find the eigenvalues of matrix A , consider the characteristic equation $|A - \lambda I| = 0$:

$$\Rightarrow \begin{vmatrix} 1 - \lambda & 1 \\ 2 & 2 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda)(2 - \lambda) - (1)(2) = 0$$

$$\Rightarrow 2 - \lambda - 2\lambda + \lambda^2 - 2 = 0$$

$$\Rightarrow \lambda^2 - 3\lambda = 0$$

$$\Rightarrow \lambda(\lambda - 3) = 0$$

This gives the eigenvalues:

$$\lambda = 0, 3$$

Therefore, the largest eigenvalue of A is 3.

3

Q1.18.2 MCQ 2018 2M Ans: D ● ● ○

Given that the matrix P has only one linearly independent eigenvector, it implies that P has a repeated eigenvalue with a geometric multiplicity less than its algebraic multiplicity.

A square matrix $A_{2 \times 2}$ is diagonalizable if and only if it possesses two linearly independent eigenvectors (i.e., its eigenvectors form a basis for $\mathbb{R}^2$).

Since P is a 2×2 matrix and has only one linearly independent eigenvector, it does not have a complete set of eigenvectors. Therefore, P cannot be diagonalized. Such a matrix is referred to as a defective matrix.

Thus, the statement corresponding to option (D) is correct.

(D) Only II and III are necessarily true

Q1.17.1 MCQ 2017 1M Ans: C ● ○ ○

Given that the vectors $a_1, a_2, \dots, a_n$ are linearly dependent, they form a column space where at least one vector can be written as a linear combination of the others. Consider the linear non-homogeneous system $AX = B$, where A is the coefficient matrix and $[A \mid B]$ represents the augmented matrix.

According to Rouché-Capelli theorem, the system is consistent if and only if:

$$\text{rank}(A) = \text{rank}([A \mid B]) = k$$

If this common rank k is strictly less than the number of unknowns (n), i.e., $k < n$:

The system possesses a degree of freedom equal to $n - k$.

There exist free variables which can take any arbitrary real value.

Therefore, the system has infinitely many solutions.

(C) infinitely many solutions

Q1.17.2 MCQ 2017 2M Ans: B ● ○ ○

Given that the rank of the real symmetric matrix $A_{n \times n}$ is 2. From the fundamental relation between the rank of a matrix and its eigenvalues, a matrix of rank r has at most r non-zero eigenvalues. Therefore, the remaining $n - 2$ eigenvalues must be zero. Let the eigenvalues of matrix A be:

$$\lambda_1, \lambda_2, 0, 0, \dots, 0$$

We are given the sum of the squares of all matrix entries:

$$\sum_{i=1}^n \sum_{j=1}^n A_{ij}^2 = 50 \quad \dots (1)$$

From matrix properties, the sum of the squares of all elements of any matrix is equal to the trace of AA^T . Since A is a real symmetric matrix ($A = A^T$), this simplifies to the trace of A^2 :

$$\sum_{i=1}^n \sum_{j=1}^n A_{ij}^2 = \text{Trace}(AA^T) = \text{Trace}(A^2)$$

The trace of A^2 is equal to the sum of the squares of the eigenvalues of A :

$$\text{Trace}(A^2) = \lambda_1^2 + \lambda_2^2 + 0^2 + \dots + 0^2 \quad \dots (2)$$

Equating equations (1) and (2):

$$\lambda_1^2 + \lambda_2^2 = 50$$

Let us test the statements based on this algebraic relation:

- Statement (I): If both λ_1^2 and λ_2^2 were strictly greater than 25, their sum would exceed 50. Therefore, at least one of λ_1^2 or λ_2^2 must be less than or equal to 25, which implies at least one eigenvalue must lie in the closed interval $[-5, 5]$. Thus, Statement (I) is true.
- Statement (II): Consider a counterexample where $\lambda_1 = \pm 5$ and $\lambda_2 = \pm 5$. In this scenario, $\lambda_1^2 + \lambda_2^2 = 25 + 25 = 50$, which completely satisfies the equation, but neither eigenvalue is strictly less than 5 or greater than 5 in magnitude. Thus, Statement (II) need not be true.

Hence, option (B) is the correct choice.

(B) (I) only

Key Points

- The sum of squares of all elements of a matrix is called the square of the Frobenius norm, satisfying $\sum A_{ij}^2 = \text{Trace}(AA^T)$.
- For a real symmetric matrix, $\text{Trace}(A^2) = \sum \lambda_i^2$.

Mistakes to be avoided

- Remember that $\text{Trace}(A^2) = \sum \lambda_i^2$ is perfectly straightforward here because A is symmetric. If A were a general non-symmetric matrix, $\sum A_{ij}^2$ would not simply equal $\sum \lambda_i^2$.

Q1.17.3 NAT 2017 1M Ans: 2 ● ○ ○

Given matrices P and Q :

$$P = \begin{bmatrix} 1 & 1 & -1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix}, \quad Q = \begin{bmatrix} -1 & -2 & -1 \\ 6 & 12 & 6 \\ 5 & 10 & 5 \end{bmatrix}$$

Let us consider the sum matrix $P + Q$:

$$P + Q = \begin{bmatrix} 1 + (-1) & 1 + (-2) & -1 + (-1) \\ 2 + 6 & -3 + 12 & 4 + 6 \\ 3 + 5 & -2 + 10 & 3 + 5 \end{bmatrix} = \begin{bmatrix} 0 & -1 & -2 \\ 8 & 9 & 10 \\ 8 & 8 & 8 \end{bmatrix}$$

$$|P + Q| = \begin{vmatrix} 0 & -1 & -2 \\ 8 & 9 & 10 \\ 8 & 8 & 8 \end{vmatrix}$$

$$|P + Q| = 0 - (-1)((8)(8) - (10)(8)) + (-2)((8)(8) - (9)(8))$$

$$|P + Q| = 1(64 - 80) - 2(64 - 72)$$

$$|P + Q| = -16 - 2(-8) = -16 + 16 = 0$$

Since $|P + Q| = 0$, the rank of $(P + Q)$ is strictly less than 3:

$$\text{Rank}(P + Q) < 3$$

Now, we check if there is any non-zero 2×2 minor. Consider the minor from the bottom-left corner:

$$\begin{vmatrix} 8 & 9 \\ 8 & 8 \end{vmatrix} = (8)(8) - (9)(8) = 64 - 72 = -8 \neq 0$$

Since there exists at least one 2×2 minor whose determinant is non-zero, the matrix $(P + Q)$ has 2 linearly independent rows.

Therefore, the rank of $(P + Q)$ is exactly 2.

2

Q1.17.4 NAT 2017 2M Ans: 5 ● ○ ○

The characteristic equation of the matrix M is given as:

$$\lambda^3 - 4\lambda^2 + a\lambda + 30 = 0 \quad \dots (1)$$

Since $\lambda = 2$ is an eigenvalue of the matrix M , it must satisfy the characteristic equation. Substituting $\lambda = 2$ into equation (1):

$$(2)^3 - 4(2)^2 + a(2) + 30 = 0$$

$$8 - 16 + 2a + 30 = 0$$

$$22 + 2a = 0 \implies a = -11$$

$$\lambda^3 - 4\lambda^2 - 11\lambda + 30 = 0$$

$$\implies (\lambda - 2)(\lambda^2 - 2\lambda - 15) = 0$$

$$\implies (\lambda - 2)(\lambda - 5)(\lambda + 3) = 0$$

$$\lambda = 2, -3, 5$$

$$\max(|\lambda|) = \max(|2|, |-3|, |5|) = \max(2, 3, 5) = 5$$

Therefore, the largest absolute value among the eigenvalues of M is 5.

5

Q1.16.1 NAT 2016 1M Ans: 15 ● ○ ○

Given that one of the eigenvalues of the real matrix P is complex:

$$\lambda_1 = 2 + \sqrt{-1} = 2 + i$$

Since the matrix has real coefficients, complex eigenvalues must occur in conjugate pairs. Therefore, the complex conjugate of λ_1 must also be an eigenvalue:

$$\lambda_2 = 2 - i$$

Let the third eigenvalue be given as $\lambda_3 = 3$.

From the fundamental properties of matrices, the determinant of a matrix is equal to the product of all its eigenvalues:

$$|P| = \lambda_1 \cdot \lambda_2 \cdot \lambda_3$$

$$\Rightarrow |P| = (2 + i)(2 - i)(3)$$

Using the algebraic identity $(a + ib)(a - ib) = a^2 + b^2$:

$$\Rightarrow |P| = (2^2 + 1^2)(3)$$

$$\Rightarrow |P| = (4 + 1)(3) = 5 \times 3 = 15$$

Therefore, the determinant of the matrix P is 15.

15

Key Points

- Complex Conjugate Root Theorem: For any matrix with real-valued entries, if $a + ib$ is an eigenvalue, then its complex conjugate $a - ib$ must also be an eigenvalue.

Q1.16.2 MCQ 2016 1M Ans: C ● ○ ○

Let us evaluate each of the given statements systematically by analyzing their properties and potential counterexamples:

- Statement I: Consider a system of 2 linear equations in 3 variables. An example of such a system is:

$$x - y + z = 1$$

$$-x + y - z = 2$$

Adding these two equations gives $0 = 3$, which is an impossibility. This demonstrates that a system with fewer equations than variables can be inconsistent and have no solution. Therefore, Statement I is false.

- Statement II: Consider a system of 3 linear equations in 2 variables. An example of such a system is:

$$x + y = 2$$

$$x - y = 0$$

$$3x + y = 4$$

From the first two equations, we easily find $x = 1$ and $y = 1$. Substituting these values into the third equation:

$$3(1) + (1) = 4$$

Since it satisfies all three equations, the system has a unique solution given by $x = 1$ and $y = 1$. This proves that a system with more equations than variables can still yield a unique solution. Therefore, Statement II is false.

- Statement III: Consider a system with 2 equations and 2 variables. An example of such a system is:

$$x + y = 2 \quad \text{and} \quad x - y = 0$$

Solving this linear system simultaneously gives a valid, well-defined solution:

$$x = 1 \quad \text{and} \quad y = 1$$

Therefore, Statement III is true.

Hence, the correct option is (c).

(C) Only III is true

Q1.16.3 NAT 2016 1M Ans: 0.124 to 0.126 ☒ ☐ ☐

Given that $\lambda = 1, 2, 4$ are the eigenvalues of a matrix $A_{3 \times 3}$.

From the properties of matrices, the determinant of a matrix is equal to the product of its eigenvalues:

$$|A| = \text{product of eigenvalues}$$

$$\Rightarrow |A| = 1 \times 2 \times 4 = 8$$

Since $|A| \neq 0$, the inverse matrix A^{-1} exists.

$$|(A^{-1})^T| = |A^{-1}|$$

$$\Rightarrow |(A^{-1})^T| = |A|^{-1}$$

$$\therefore |(A^{-1})^T| = \frac{1}{|A|} = \frac{1}{8} = 0.125$$

Therefore, the value of $|(A^{-1})^T|$ is 0.125.

0.125

Key Points

- For an $n \times n$ matrix A , $\det(A) = \prod_{i=1}^n \lambda_i$.
- $\det(A^T) = \det(A)$.
- $\det(A^{-1}) = \frac{1}{\det(A)}$.

Q1.16.4 NAT 2016 2M Ans: 1500 ☒ ☒ ☐

For multiplying four matrices A, B, C , and D , there are 5 possible ways to place the parentheses: 1. $A(B(CD))$ 2. $A((BC)D)$ 3. $((AB)C)D$ 4. $(A(BC))D$ 5. $(AB)(CD)$

The total number of scalar multiplications required for each corresponding case evaluates to 1750, 1500, 3500, 2000, and 3000 respectively.

Comparing these values, the minimum number of scalar multiplications needed is 1500.

1500

Key Points

- Matrix multiplication is associative, meaning $(AB)C = A(BC)$, but the grouping order heavily impacts the computational cost (number of scalar multiplications).
- For multiplying two matrices of dimensions $p \times q$ and $q \times r$, the number of scalar multiplications required is $p \times q \times r$.
- The total number of valid parenthesizations for n matrices is given by the $(n - 1)^{\text{th}}$ Catalan number: $C_{n-1} = \frac{1}{n} \binom{2n-2}{n-1}$. For 4 matrices, $C_3 = 5$.

Q1.15.1 MCQ 2015 2M Ans: D ● ○ ○

Given that the matrix $A = \begin{bmatrix} 1 & 4 \\ b & a \end{bmatrix}$ has eigenvalues -1 and 7 . From the essential properties of eigenvalues:

- Sum of Eigenvalues Property: The sum of the eigenvalues is equal to the trace of the matrix (the sum of its principal diagonal elements):

$$\text{Sum of the eigenvalues of } A = \text{sum of diagonal elements of } A$$

$$\Rightarrow 1 + a = -1 + 7$$

$$\Rightarrow 1 + a = 6 \Rightarrow a = 5$$

- Product of Eigenvalues Property: The product of the eigenvalues is equal to the determinant of the matrix:

$$|A| = \text{Product of the eigenvalues}$$

$$\Rightarrow (1)(a) - (4)(b) = (-1) \times 7$$

$$\Rightarrow a - 4b = -7$$

Substituting the value of $a = 5$ into the determinant equation:

$$5 - 4b = -7$$

$$\Rightarrow -4b = -12 \Rightarrow b = 3$$

Therefore, the parameters are $a = 5$ and $b = 3$, corresponding to option (d).

$$(D) \ a = 5, b = 3$$

Q1.15.2 MCQ 2015 1M Ans: B ● ○ ○

Let the given matrix be A :

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 2 & 1 \end{bmatrix}$$

The eigenvectors $X = \begin{bmatrix} x & y & z \end{bmatrix}^T$ of matrix A corresponding to the eigenvalue $\lambda = 1$ are determined by solving the homogeneous linear system $(A - \lambda I)X = O$.

Substituting $\lambda = 1$:

$$\begin{bmatrix} 1-1 & -1 & 2 \\ 0 & 1-1 & 0 \\ 1 & 2 & 1-1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & -1 & 2 \\ 0 & 0 & 0 \\ 1 & 2 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$-y + 2z = 0 \quad \dots (1)$$

$$x + 2y = 0 \quad \dots (2)$$

Let $z = \alpha$ (where $\alpha \neq 0$).

From equation (1):

$$y = 2z = 2\alpha$$

$$x = -2y = -2(2\alpha) = -4\alpha$$

$$\therefore X = \begin{bmatrix} -4\alpha \\ 2\alpha \\ \alpha \end{bmatrix} = \alpha \begin{bmatrix} -4 \\ 2 \\ 1 \end{bmatrix}, \quad \text{where } \alpha \neq 0$$

$$(B) \{ \alpha(-4, 2, 1) \mid \alpha \neq 0, \alpha \in \mathbb{R} \}$$

Q1.15.3 NAT 2015 1M Ans: 6 ☒ ☐ ☐

The characteristic equation of a square matrix A is given by $|A - \lambda I| = 0$.

$$\Rightarrow \begin{vmatrix} 4 - \lambda & 5 \\ 2 & 1 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (4 - \lambda)(1 - \lambda) - (5)(2) = 0$$

$$\Rightarrow 4 - 4\lambda - \lambda + \lambda^2 - 10 = 0$$

$$\Rightarrow \lambda^2 - 5\lambda - 6 = 0$$

$$\Rightarrow (\lambda - 6)(\lambda + 1) = 0$$

$$\lambda = 6, -1$$

Comparing the two values, the larger eigenvalue is 6.

6

Q1.15.4 NAT 2015 2M Ans: 0 ☒ ☒ ☐

Let the given matrix be A :

$$A = \begin{bmatrix} 3 & 4 & 45 \\ 7 & 9 & 105 \\ 13 & 2 & 195 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 3 & 4 & 45 \\ 7 & 9 & 105 \\ 13 & 2 & 195 \end{vmatrix}$$

From the properties of determinants, the value remains unchanged after performing elementary row or column operations of the form $R_i \rightarrow R_i + kR_j$ or $C_i \rightarrow C_i + kC_j$.

$$C_3 = \begin{bmatrix} 15 \times 3 \\ 15 \times 7 \\ 15 \times 13 \end{bmatrix} = 15 \begin{bmatrix} 3 \\ 7 \\ 13 \end{bmatrix} = 15C_1$$

Since Column 3 is a direct linear multiple of Column 1 ($C_3 = 15C_1$), the two columns are linearly dependent. By the property of determinants, if any two rows or columns of a square matrix are identical or proportional, the value of the determinant is zero.

$$\therefore |A| = 0$$

Key Points

- If any two rows or columns of a square matrix are proportional (i.e., $C_i = kC_j$), its determinant is identically zero.

Q1.15.5 MCQ 2015 2M Ans: C ● ● ●

If the given homogeneous system of linear equations has a non-trivial solution, the determinant of its coefficient matrix must be equal to zero:

$$\begin{vmatrix} p & q & r \\ q & r & p \\ r & p & q \end{vmatrix} = 0$$

Applying the column operation $C_1 \rightarrow C_1 + C_2 + C_3$:

$$\Rightarrow \begin{vmatrix} p+q+r & q & r \\ p+q+r & r & p \\ p+q+r & p & q \end{vmatrix} = 0$$

$$\Rightarrow (p+q+r) \begin{vmatrix} 1 & q & r \\ 1 & r & p \\ 1 & p & q \end{vmatrix} = 0$$

Applying the elementary row operations $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$\Rightarrow (p+q+r) \begin{vmatrix} 1 & q & r \\ 0 & r-q & p-r \\ 0 & p-q & q-r \end{vmatrix} = 0$$

Expanding the determinant along the first column:

$$\Rightarrow (p+q+r) [(r-q)(q-r) - (p-q)(p-r)] = 0$$

$$\Rightarrow (p+q+r) [-(r-q)^2 - (p^2 - pr - pq + qr)] = 0$$

$$\Rightarrow -(p+q+r)(p^2 + q^2 + r^2 - pq - qr - pr) = 0$$

Multiplying and dividing by 2, we can rewrite the expression inside the brackets as a sum of perfect squares:

$$\Rightarrow -\frac{1}{2}(p+q+r)[(p-q)^2 + (q-r)^2 + (r-p)^2] = 0$$

For this product to equal zero, either of the two terms must be zero:

$$p+q+r=0 \quad \text{or} \quad (p-q)^2 + (q-r)^2 + (r-p)^2 = 0$$

Since a sum of real squared terms can only be zero if each individual term is zero simultaneously ($p-q=0$, $q-r=0$, $r-p=0$), we get:

$$\therefore p+q+r=0 \quad \text{or} \quad p=q=r$$

Thus, the condition corresponding to option (C) is correct.

$$(C) \quad p+q+r=0 \text{ or } p=q=r$$

Key Points

- Circulant Determinant Identity: The determinant of a circulant matrix is given by:

$$\begin{vmatrix} p & q & r \\ q & r & p \\ r & p & q \end{vmatrix} = -(p+q+r)(p^2+q^2+r^2-pq-qr-pr)$$

Q1.14.1 NAT 2014 1M Ans: 0 ● ○ ○

From the fundamental spectral properties of real symmetric matrices, the eigenvectors corresponding to distinct eigenvalues are mutually orthogonal. A symmetric positive definite matrix is a special class of real symmetric matrices whose eigenvalues are strictly positive.

Let X_1 and X_2 be two eigenvectors corresponding to distinct eigenvalues λ_1 and λ_2 ($\lambda_1 \neq \lambda_2$) for a symmetric matrix A :

$$AX_1 = \lambda_1 X_1 \quad \text{and} \quad AX_2 = \lambda_2 X_2$$

Since the vectors are orthogonal, their inner product (dot product) satisfies the following relation:

$$\langle X_1, X_2 \rangle = X_1 \cdot X_2 = X_1^T X_2 = 0$$

$\therefore$ Dot product between any two eigenvectors corresponding to distinct eigenvalues is zero.

0

Q1.14.2 NAT 2014 2M Ans: 6 ● ● ○

Let the given matrix be A . The characteristic equation of a matrix A is given by $|A - \lambda I| = 0$:

$$\Rightarrow \begin{vmatrix} 1-\lambda & 0 & 0 & 0 & 1 \\ 0 & 1-\lambda & 1 & 1 & 0 \\ 0 & 1 & 1-\lambda & 1 & 0 \\ 0 & 1 & 1 & 1-\lambda & 0 \\ 1 & 0 & 0 & 0 & 1-\lambda \end{vmatrix} = 0$$

Expanding the 5×5 determinant along the 1st row:

$$(1-\lambda) \begin{vmatrix} 1 & 1 & 1 & 0 \\ 1 & 1-\lambda & 1 & 0 \\ 1 & 1 & 1-\lambda & 0 \\ 0 & 0 & 0 & 1-\lambda \end{vmatrix} + 1 \begin{vmatrix} 0 & 1-\lambda & 1 & 1 \\ 0 & 1 & 1-\lambda & 1 \\ 0 & 1 & 1 & 1-\lambda \\ 1 & 0 & 0 & 0 \end{vmatrix} = 0$$

Expanding each of these sub-determinants further:

$$\Rightarrow (1-\lambda)^2 \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{vmatrix} = 0$$

Factoring out the common 3×3 determinant:

$$\Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{vmatrix} [(1-\lambda)^2 - 1] = 0$$

Evaluating the remaining terms yields the characteristic polynomial:

$$\Rightarrow \lambda^3(\lambda - 2)(\lambda - 3) = 0$$

Solving for the roots, the eigenvalues of the matrix A are:

$$\lambda = 0, 0, 0, 2, 3$$

The question asks for the product of the non-zero eigenvalues of the matrix:

$$\text{Product} = (2) \times (3) = 6$$

6

Q1.14.3 NAT 2014 1M Ans: 2 ● ○ ○

From Grassmann's dimension theorem for vector subspaces, the relation between the dimensions of the sum and intersection of two subspaces V_1 and V_2 is given by:

$$\dim(V_1 + V_2) = \dim(V_1) + \dim(V_2) - \dim(V_1 \cap V_2)$$

To find the minimum possible value of $\dim(V_1 \cap V_2)$, we rearrange the equation to isolate the intersection term:

$$\dim(V_1 \cap V_2) = \dim(V_1) + \dim(V_2) - \dim(V_1 + V_2)$$

For $\dim(V_1 \cap V_2)$ to be minimized, the term $\dim(V_1 + V_2)$ must be maximized:

$$\text{Minimum value of } \dim(V_1 \cap V_2) = \dim(V_1) + \dim(V_2) - \max \text{ of } \dim(V_1 + V_2)$$

Assuming the subspaces belong to a ambient vector space V of dimension 6 (such that $\max(\dim(V_1 + V_2)) = 6$), and given that $\dim(V_1) = 4$ and $\dim(V_2) = 4$:

$$= 4 + 4 - 6 = 2$$

Therefore, the minimum value of $\dim(V_1 \cap V_2)$ is 2.

2

Key Points

- $\dim(V_1 + V_2) + \dim(V_1 \cap V_2) = \dim(V_1) + \dim(V_2)$.

Q1.14.4 MCQ 2014 1M Ans: A ● ○ ○

From the fundamental properties of matrix algebra, the determinant of a square matrix A is equal to the product of all its eigenvalues:

$$\det(A) = \lambda_1 \cdot \lambda_2 \dots \lambda_n$$

If it is given that the determinant of the matrix is negative ($\det(A) < 0$):

- The product of the eigenvalues must result in a negative real number.
- For a product of real numbers to be negative, there must be an odd number of negative factors.

Therefore, there must exist at least one eigenvalue that is strictly negative.

Thus, the statement corresponding to option (A) is correct.

(A) If the trace of the matrix is positive and the determinant of the matrix is negative, at least one of its eigenvalues is negative.

Q1.13.1 MCQ 2013 1M Ans: A ● ○ ○

Consider the first determinant Δ_1 :

$$\Delta_1 = \begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix}$$

Applying the elementary column operations $C_2 \rightarrow C_2 + C_1$ and $C_3 \rightarrow C_3 + C_1$:

$$\Rightarrow \Delta_1 = \begin{vmatrix} 1 & x+1 & x^2+1 \\ 1 & y+1 & y^2+1 \\ 1 & z+1 & z^2+1 \end{vmatrix}$$

From option (a), consider the second determinant Δ_2 :

$$\Delta_2 = \begin{vmatrix} 1 & x(x+1) & x+1 \\ 1 & y(y+1) & y+1 \\ 1 & z(z+1) & z+1 \end{vmatrix} = \begin{vmatrix} 1 & x^2+x & x+1 \\ 1 & y^2+y & y+1 \\ 1 & z^2+z & z+1 \end{vmatrix}$$

Interchanging columns $C_2 \leftrightarrow C_3$ introduces a negative sign according to determinant properties:

$$\Rightarrow \Delta_2 = (-1) \begin{vmatrix} 1 & x+1 & x^2+x \\ 1 & y+1 & y^2+y \\ 1 & z+1 & z^2+z \end{vmatrix}$$

Comparing the final forms, we find that $\Delta_1 \neq \Delta_2$. Thus, the unequal relation proposed by option (a) is correct.

$$(A) \begin{vmatrix} 1 & x(x+1) & x+1 \\ 1 & y(y+1) & y+1 \\ 1 & z(z+1) & z+1 \end{vmatrix}$$

Q1.12.1 MCQ 2012 1M Ans: D ● ○ ○

Given the elements of the 2×2 matrix A , we can construct it as follows:

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

To find the eigenvalues of matrix A , we set up the characteristic equation $\det(A - \lambda I) = 0$:

$$\begin{vmatrix} 1-\lambda & 1 \\ 1 & -1-\lambda \end{vmatrix} = 0$$

$$(1-\lambda)(-1-\lambda) - (1)(1) = 0$$

$$\Rightarrow -(1-\lambda^2) - 1 = 0$$

$$\Rightarrow \lambda^2 - 1 - 1 = 0 \Rightarrow \lambda^2 = 2$$

$$\lambda = \pm\sqrt{2}$$

According to spectral matrix properties, if a matrix A has eigenvalues λ , then the powered matrix A^n will have eigenvalues λ^n . Therefore, the eigenvalues of A^{19} are:

$$\lambda_{19} = (\pm\sqrt{2})^{19}$$

$$(\sqrt{2})^{19} = (\sqrt{2})^{18} \cdot \sqrt{2} = (2^{1/2})^{18} \cdot \sqrt{2} = 2^9 \cdot \sqrt{2} = 512\sqrt{2}$$

$$\lambda_{19} = 512\sqrt{2} \quad \text{and} \quad -512\sqrt{2}$$

Let us alternative verify by computing A^2 :

$$A^2 = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1+1 & 1-1 \\ 1-1 & 1+1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = 2I$$

Using this direct matrix relationship for A^{19} :

$$A^{19} = (A^2)^9 \cdot A = (2I)^9 \cdot A = 2^9 I \cdot A = 512A$$

$$A^{19} = 512 \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 512 & 512 \\ 512 & -512 \end{bmatrix}$$

The eigenvalues of $512A$ are simply 512 times the eigenvalues of A :

$$\lambda_{A^{19}} = 512 \cdot (\pm\sqrt{2}) = \pm 512\sqrt{2}$$

Therefore, matching the derived mathematical evaluation, option (d) is the exact correct numerical value.

(D) $512\sqrt{2}$ and $-512\sqrt{2}$

Q1.11.1 MCQ 2011 2M Ans: C ● ● ○

Given the dimensions of the four matrices M_1, M_2, M_3 , and M_4 :

- $M_1 : p \times q = 10 \times 100$
- $M_2 : q \times r = 100 \times 20$
- $M_3 : r \times s = 20 \times 5$
- $M_4 : s \times t = 5 \times 80$

There are 5 possible ways to parenthesize the product $M_1 \times M_2 \times M_3 \times M_4$. Let us evaluate the total number of scalar multiplications required for each configuration:

- Case 1: $(M_1 \times (M_2 \times (M_3 \times M_4)))$
 - Cost of $(M_3 \times M_4) = r \times s \times t = 20 \times 5 \times 80 = 8000$, resulting matrix is 20×80 .
 - Cost of $(M_2 \times (M_3 M_4)) = q \times r \times t = 100 \times 20 \times 80 = 160000$, resulting matrix is 100×80 .
 - Cost of $(M_1 \times (M_2(M_3 M_4))) = p \times q \times t = 10 \times 100 \times 80 = 80000$.
 - Total cost = $8000 + 160000 + 80000 = 248000$.
- Case 2: $(M_1 \times ((M_2 \times M_3) \times M_4))$
 - Cost of $(M_2 \times M_3) = q \times r \times s = 100 \times 20 \times 5 = 10000$, resulting matrix is 100×5 .
 - Cost of $((M_2 M_3) \times M_4) = q \times s \times t = 100 \times 5 \times 80 = 40000$, resulting matrix is 100×80 .
 - Cost of $(M_1 \times ((M_2 M_3) M_4)) = p \times q \times t = 10 \times 100 \times 80 = 80000$.
 - Total cost = $10000 + 40000 + 80000 = 130000$.
- Case 3: $((M_1 \times M_2) \times M_3) \times M_4$
 - Given formula: $pqr + prs + pst$

- Total cost = $(10 \times 100 \times 20) + (10 \times 20 \times 5) + (10 \times 5 \times 80) = 20000 + 1000 + 4000 = 25000$.
- Case 4: $((M_1 \times (M_2 \times M_3)) \times M_4)$
 - Cost of $(M_2 \times M_3) = q \times r \times s = 100 \times 20 \times 5 = 10000$, resulting matrix is 100×5 .
 - Cost of $(M_1 \times (M_2 M_3)) = p \times q \times s = 10 \times 100 \times 5 = 5000$, resulting matrix is 10×5 .
 - Cost of $((M_1(M_2 M_3)) \times M_4) = p \times s \times t = 10 \times 5 \times 80 = 4000$.
 - Total cost = $10000 + 5000 + 4000 = 19000$.
- Case 5: $((M_1 \times M_2) \times (M_3 \times M_4))$
 - Given formula: $pqr + rst + prt$
 - Total cost = $(10 \times 100 \times 20) + (20 \times 5 \times 80) + (10 \times 20 \times 80) = 20000 + 8000 + 16000 = 44000$.

Comparing the values, the absolute mathematical minimum is 19000, which corresponds to Case 4. Since 19000 is listed as option C, it is the correct choice.

(C) 19000

Key Points

- The scalar multiplication count for multiplying a $m \times n$ matrix by a $n \times p$ matrix is given by $m \times n \times p$.

Q1.10.1 MCQ 2010 2M Ans: D ● ○ ○

Given that the eigenvalues of the matrix $A = \begin{bmatrix} 2 & 3 \\ x & y \end{bmatrix}$ are 4 and 8.

From the key properties of eigenvalues:

Sum of the eigenvalues = $\text{tr}(A)$

$$\Rightarrow 4 + 8 = 2 + y$$

$$\Rightarrow 12 = 2 + y \Rightarrow y = 10$$

Product of eigenvalues = $\det(A)$

$$\Rightarrow (4)(8) = (2)(y) - (3)(x)$$

$$\Rightarrow 32 = 2y - 3x$$

$$32 = 2(10) - 3x$$

$$\Rightarrow 32 = 20 - 3x$$

$$\Rightarrow 3x = 20 - 32$$

$$\Rightarrow 3x = -12 \Rightarrow x = -4$$

Therefore, the calculated values are $x = -4$ and $y = 10$, which corresponds to option (D).

(D) $x = -4, y = 10$

Q1.08.1 **MCQ** **2008** **1M** **Ans: MTA** ● ○ ○

Given the system of linear equations in matrix form $AX = B$:

$$\begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 1 & 4 & \alpha \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$$

We can solve this problem using two different approaches.

Method 1

Consider the augmented matrix $[A \mid B]$:

$$[A \mid B] = \left[\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 1 & 2 & 3 & 2 \\ 1 & 4 & \alpha & 4 \end{array} \right]$$

Applying elementary row operations $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$\Rightarrow [A \mid B] \sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 3 & \alpha - 2 & 3 \end{array} \right]$$

Applying row operation $R_3 \rightarrow R_3 - 3R_2$:

$$\Rightarrow [A \mid B] \sim \left[\begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & \alpha - 5 & 0 \end{array} \right]$$

To have a unique solution, the rank of the coefficient matrix A must equal the rank of the augmented matrix $[A \mid B]$, which must equal the number of variables (3). Therefore, the pivot element in the third row must be non-zero:

$$\alpha - 5 \neq 0 \implies \alpha \neq 5$$

Hence, for all values of α except $\alpha = 5$, the system will have a unique solution.

Method 2

To have a unique solution, the determinant of the coefficient matrix A must be non-singular (non-zero):

$$\text{i.e., } \begin{vmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 1 & 4 & \alpha \end{vmatrix} \neq 0$$

Expanding along the first row:

$$1(2\alpha - 12) - 1(\alpha - 3) + 2(4 - 2) \neq 0$$

$$2\alpha - 12 - \alpha + 3 + 4 \neq 0$$

$$\alpha - 5 \neq 0$$

$$\therefore \alpha \neq 5$$

MTA

Q1.08.2 MCQ 2008 2M Ans: A ● ○ ○

Let us find the eigenvalues for each of the given matrices systematically by setting up their characteristic equations $\det(M - \lambda I) = 0$:

- Matrix A:

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

Since A is a diagonal matrix, its eigenvalues are simply its main diagonal entries:

$$\lambda = 1, 0$$

Matrix A has an eigenvalue of 1.

- Matrix B:

$$B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Since B is an upper triangular matrix, its eigenvalues are its main diagonal entries:

$$\lambda = 0, 0$$

Matrix B does not have an eigenvalue of 1.

- Matrix C:

$$C = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

The characteristic equation is given by:

$$\det(C - \lambda I) = \begin{vmatrix} 1 - \lambda & -1 \\ 1 & 1 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda)^2 - (-1)(1) = 0$$

$$\Rightarrow (1 - \lambda)^2 + 1 = 0 \Rightarrow (1 - \lambda)^2 = -1$$

$$\Rightarrow 1 - \lambda = \pm i \Rightarrow \lambda = 1 + i, 1 - i$$

Matrix C has complex eigenvalues, not the real eigenvalue 1.

- Matrix D:

$$D = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\lambda = -1, -1$$

Matrix D does not have an eigenvalue of 1.

Comparing all four matrices, only one matrix (A) has an eigenvalue equal to 1. This corresponds to option (A).

(A) one

Key Points

- Triangular and Diagonal Matrices:** For any upper triangular, lower triangular, or diagonal matrix, the eigenvalues are identically equal to the elements located along the principal diagonal.
- Complex Conjugate Pairs:** If a real matrix produces a complex root, its eigenvalues must appear as a conjugate pair ($\lambda = a \pm ib$).

Q1.07.1 **MCQ** **2007** **2M** **Ans: C** ● ○ ○

Let the given 8×8 block matrix be M :

$$M = \begin{bmatrix} A & I \\ I & A \end{bmatrix}$$

Let λ be an eigenvalue of matrix A corresponding to an eigenvector v , such that $Av = \lambda v$. We want to find an eigenvalue μ and a corresponding block eigenvector $X = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ for the block matrix M :

$$MX = \mu X$$

$$\Rightarrow \begin{bmatrix} A & I \\ I & A \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \mu \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$$

This expands into a system of two block vector equations:

$$Av_1 + Iv_2 = \mu v_1 \quad \dots (1)$$

$$Iv_1 + Av_2 = \mu v_2 \quad \dots (2)$$

Let $v_1 = v$ and $v_2 = kv$ for some scalar constant k . Substituting into equations (1) and (2):

$$Av + kv = \mu v \Rightarrow (\lambda + k)v = \mu v$$

$$v + kAv = \mu kv \Rightarrow (1 + k\lambda)v = \mu kv$$

Since $v \neq 0$, we can cancel v to obtain two scalar equations:

$$\lambda + k = \mu$$

$$1 + k\lambda = \mu k$$

Substituting $\mu = \lambda + k$ into the second equation:

$$1 + k\lambda = (\lambda + k)k$$

$$1 + k\lambda = k\lambda + k^2 \Rightarrow k^2 = 1 \Rightarrow k = \pm 1$$

Now, substituting $k = \pm 1$ back into the expression for μ :

- For $k = 1$: $\mu = \lambda + 1$
- For $k = -1$: $\mu = \lambda - 1$

Therefore, if λ is an eigenvalue of A , then the eigenvalues of the block matrix M are given by $\lambda + 1$ and $\lambda - 1$.

Given that the eigenvalues of A are $\{-5, -2, 1, 4\}$, let us compute the complete set of eigenvalues for M :

- From $\lambda = -5$: $\mu = -5 + 1 = -4$ or $\mu = -5 - 1 = -6$
- From $\lambda = -2$: $\mu = -2 + 1 = -1$ or $\mu = -2 - 1 = -3$
- From $\lambda = 1$: $\mu = 1 + 1 = 2$ or $\mu = 1 - 1 = 0$
- From $\lambda = 4$: $\mu = 4 + 1 = 5$ or $\mu = 4 - 1 = 3$

The complete set of eigenvalues of M is $\{-6, -4, -3, -1, 0, 2, 3, 5\}$. Therefore, 2 is an eigenvalue of the block matrix, which corresponds to option (c).

(C) 2

Q1.04.1 MCQ 2004 1M Ans: C ● ○ ○

The total number of elements in an $n \times n$ matrix is n^2 .

In a symmetric matrix, the elements satisfy the condition $a_{ij} = a_{ji}$. Because of this symmetry, the elements below the main diagonal are completely determined by the elements above the main diagonal. Therefore, to construct a unique symmetric matrix, we only need to choose values for the elements on the principal diagonal and the elements lying strictly above it.

Let us find the total count of these independent choices:

- Number of diagonal elements: n
- Total number of non-diagonal elements: $n^2 - n$
- Number of elements strictly above the diagonal:

$$\frac{n^2 - n}{2}$$

Summing these up gives the total number of independent element positions that can be chosen:

$$\text{Total independent elements} = n + \frac{n^2 - n}{2} = \frac{2n + n^2 - n}{2} = \frac{n^2 + n}{2} = \frac{n(n + 1)}{2}$$

Each of these independent positions can be filled with either 0 or 1 (a total of 2 choices per element position). Therefore, the total number of ways to form different symmetric matrices is:

$$\text{Total matrices} = 2^{\frac{n(n+1)}{2}} = 2^{\frac{n^2+n}{2}}$$

$$(C) \ 2^{\frac{n^2+n}{2}}$$

Key Points

- For a symmetric matrix of order n , the number of independent elements is $\frac{n(n+1)}{2}$.
- If a matrix contains k independent positions and each position can accept m distinct values, the total number of unique matrices that can be generated is m^k .

Q1.04.2 MCQ 2004 1M Ans: B ● ○ ○

Given the matrix equation:

$$ABCD = I$$

We can solve for B^{-1} using properties of matrix inversion.

Method 1

Taking the inverse on both sides of the equation:

$$\begin{aligned} \Rightarrow (ABCD)^{-1} &= I^{-1} \\ \Rightarrow D^{-1}C^{-1}B^{-1}A^{-1} &= I \\ \Rightarrow D(D^{-1}C^{-1}B^{-1}A^{-1})A &= DIA \\ \Rightarrow (DD^{-1})C^{-1}B^{-1}(A^{-1}A) &= DA \\ \Rightarrow IC^{-1}B^{-1}I &= DA \\ \Rightarrow C^{-1}B^{-1} &= DA \end{aligned}$$

Now, multiplying by C on both sides:

$$\Rightarrow C(C^{-1}B^{-1}) = CDA$$

$$\Rightarrow (CC^{-1})B^{-1} = CDA$$

$$\Rightarrow IB^{-1} = CDA$$

$$\therefore B^{-1} = CDA$$

$$\boxed{(B) CDA}$$

Method 2

$$ABCD = I$$

$$\Rightarrow A^{-1}(ABCD) = A^{-1}I$$

$$\Rightarrow BCD = A^{-1}$$

$$\Rightarrow (BCD)D^{-1} = A^{-1}D^{-1}$$

$$\Rightarrow BC = A^{-1}D^{-1}$$

Now, take the inverse of both sides:

$$\Rightarrow (BC)^{-1} = (A^{-1}D^{-1})^{-1}$$

$$\Rightarrow C^{-1}B^{-1} = (D^{-1})^{-1}(A^{-1})^{-1}$$

$$\Rightarrow C^{-1}B^{-1} = DA$$

$$\Rightarrow C(C^{-1}B^{-1}) = CDA$$

$$\therefore B^{-1} = CDA$$

$$\boxed{(B) CDA}$$

Key Points

- For any invertible matrices X and Y , $(XY)^{-1} = Y^{-1}X^{-1}$. This rule extends to multiple matrices: $(XYZ)^{-1} = Z^{-1}Y^{-1}X^{-1}$.
- Multiplying any matrix by the identity matrix I leaves the matrix unchanged ($AI = IA = A$).

Q1.01.1 MCQ 2001 1M Ans: A ● ○ ○

Let us evaluate each statement by constructing simple counterexamples using 2×2 matrices ($n = 2$):

- Statement S_1 : The sum of two singular $n \times n$ matrices may be non-singular. Let us choose two distinct singular matrices A and B ($\det(A) = 0$ and $\det(B) = 0$):

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \Rightarrow \det(A) = 0$$

$$B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \Rightarrow \det(B) = 0$$

Computing their sum $A + B$:

$$A + B = \begin{bmatrix} 1+0 & 0+0 \\ 0+0 & 0+1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

Evaluating the determinant of the sum matrix:

$$\det(A + B) = \det(I) = 1 \neq 0$$

Since the resulting matrix is non-singular, Statement S_1 is true.

- Statement S_2 : The sum of two $n \times n$ non-singular matrices may be singular. Let us choose a non-singular matrix C ($\det(C) \neq 0$) and its additive inverse $-C$:

$$C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \implies \det(C) = 1 \neq 0$$

$$-C = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = -I \implies \det(-C) = 1 \neq 0$$

Computing their sum $C + (-C)$:

$$C + (-C) = \begin{bmatrix} 1-1 & 0+0 \\ 0+0 & 1-1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O$$

Evaluating the determinant of the sum matrix:

$$\det(C + (-C)) = \det(O) = 0$$

Since the resulting zero matrix is singular, Statement S_2 is true.

Therefore, both statements S_1 and S_2 are true, which corresponds to option (A).

(A) S_1 and S_2 are both true

Q1.00.1 MCQ 2000 1M Ans: A ● ○ ○

Given an $n \times n$ array v defined by the relation $v[i, j] = i - j$ for all indices $1 \leq i \leq n$ and $1 \leq j \leq n$.

We can solve for the sum of the elements using two distinct methods.

Method 1

Let us write out the matrix structural visualization of the array v :

$$v = \begin{bmatrix} 1-1 & 1-2 & 1-3 & \dots & 1-n \\ 2-1 & 2-2 & 2-3 & \dots & 2-n \\ 3-1 & 3-2 & 3-3 & \dots & 3-n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n-1 & n-2 & n-3 & \dots & n-n \end{bmatrix} = \begin{bmatrix} 0 & -1 & -2 & \dots & -(n-1) \\ 1 & 0 & -1 & \dots & -(n-2) \\ 2 & 1 & 0 & \dots & -(n-3) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n-1 & n-2 & n-3 & \dots & 0 \end{bmatrix}$$

From the structural pattern, we observe:

- The principal diagonal elements ($i = j$) are all exactly equal to 0.
- For any off-diagonal entry located at position (i, j) , its corresponding symmetric element across the principal diagonal at position (j, i) is:

$$v[j, i] = j - i = -(i - j) = -v[i, j]$$

Since $v[j, i] + v[i, j] = 0$, every positive element in the lower-triangular portion of the matrix has a perfectly matched, opposite-signed negative element in the upper-triangular portion. Consequently, when summing all elements of the array together, the opposing terms cancel out pairwise, yielding a net sum of 0.

Method 2

We can alternatively calculate the total sum mathematically using double summation indices:

$$\text{Total Sum} = \sum_{i=1}^n \sum_{j=1}^n v[i, j] = \sum_{i=1}^n \sum_{j=1}^n (i - j)$$

Distributing the inner summation across subtraction:

$$= \sum_{i=1}^n \left(\sum_{j=1}^n i - \sum_{j=1}^n j \right)$$

Evaluating the inner terms over the index j :

- Since i is independent of j , $\sum_{j=1}^n i = n \cdot i$.
- Using the arithmetic series identity, $\sum_{j=1}^n j = \frac{n(n+1)}{2}$.

Substituting these expressions back into the outer summation:

$$\begin{aligned} &= \sum_{i=1}^n \left(n \cdot i - \frac{n(n+1)}{2} \right) \\ &= n \sum_{i=1}^n i - \sum_{i=1}^n \frac{n(n+1)}{2} \end{aligned}$$

Evaluating the summation over the index i :

$$= n \cdot \left[\frac{n(n+1)}{2} \right] - n \cdot \left[\frac{n(n+1)}{2} \right] = 0$$

Thus, the total sum of the elements is 0, which matches option (A).

(A) 0

Key Points

- Skew-Symmetric Properties: A matrix A satisfying $a_{ji} = -a_{ij}$ is classified as skew-symmetric. The sum of all elements in any skew-symmetric matrix is inherently 0.
- Summation Splitting Laws: Linear summation components can be segmented individually using the distribution properties: $\sum (x_i \pm y_i) = \sum x_i \pm \sum y_i$.

Mistakes to be avoided

- When evaluating $\sum_{j=1}^n i$, remember that the row variable i acts as a constant relative to j , so it scales as $n \cdot i$, not as $\frac{n(n+1)}{2}$.

Q1.98.1 MCQ 1998 2M Ans: B ● ○ ○

Given the determinant Δ :

$$\Delta = \begin{vmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix}$$

Applying the row operation $R_1 \rightarrow R_1 - R_2$:

$$\Rightarrow \Delta = \begin{vmatrix} 0 & a-b & bc-ca \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix}$$

We can rewrite the third entry of the first row by factoring out $-c$:

$$bc - ca = -c(a - b)$$

Substituting this back into the determinant:

$$\Rightarrow \Delta = \begin{vmatrix} 0 & a-b & -c(a-b) \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix}$$

Taking $(a - b)$ as a common factor from the first row (R_1):

$$\Rightarrow \Delta = (a - b) \begin{vmatrix} 0 & 1 & -c \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix}$$

Since the determinant can be cleanly factored as a product containing $(a - b)$, it is explicitly clear that $(a - b)$ is a factor of Δ . This matches option (b).

(B) a-b

Q1.97.1 MCQ 1997 1M Ans: C ☒ ☐ ☐

Given an n -rowed square matrix A :

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

The matrix I_{12} is an elementary matrix obtained by interchanging the first and second rows of the n -rowed identity matrix I :

$$I_{12} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}$$

When a matrix A is post-multiplied by an elementary matrix (i.e., the multiplication is of the form AE), the operation performs a **column operation** on A matching the row operation used to create E .

Let us explicitly compute the product $B = AI_{12}$:

$$B = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}$$

Thus, the resulting matrix is:

$$AI_{12} = \begin{bmatrix} a_{12} & a_{11} & a_{13} & \dots & a_{1n} \\ a_{22} & a_{21} & a_{23} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n2} & a_{n1} & a_{n3} & \dots & a_{nn} \end{bmatrix}$$

Now let us analyze the specific properties of the first column of this product matrix AI_{12} : The first column consists of the elements $[a_{12} \ a_{22} \ \dots \ a_{n2}]'$, which is identically the second column of A . This corresponds perfectly to option (C).

(C) column is the same as the second column of A

Q1.96.1 NAT 1996 5M Ans: * ● ● ●

Given the matrices $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ and $B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$.

We are given the following matrix equations:

$$CD = I \quad \dots (1)$$

$$C = A \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad \dots (2)$$

$$AB = I \quad \dots (3)$$

From equation (1), we can express D as the inverse of C :

$$D = C^{-1}$$

$$\Rightarrow D = \left(A \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right)^{-1}$$

$$\Rightarrow D = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} A^{-1}$$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} = \frac{1}{(1)(1) - (0)(1)} \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$$

Substituting this inverse back into the expression for D :

$$\Rightarrow D = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} A^{-1}$$

From equation (3), since $AB = I$, it implies that $B = A^{-1}$. Substituting B for A^{-1} :

$$\Rightarrow D = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} B$$

$$\Rightarrow D = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$$

$$\therefore D = \begin{bmatrix} (1)(b_{11}) + (0)(b_{21}) & (1)(b_{12}) + (0)(b_{22}) \\ (-1)(b_{11}) + (1)(b_{21}) & (-1)(b_{12}) + (1)(b_{22}) \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ -b_{11} + b_{21} & -b_{12} + b_{22} \end{bmatrix}$$

$$\begin{bmatrix} b_{11} & b_{12} \\ -b_{11} + b_{21} & -b_{12} + b_{22} \end{bmatrix}$$

Key Points

- For a matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ with $\det(M) \neq 0$, the inverse is $M^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$.

Q1.96.2 MCQ 1996 2M Ans: A ● ● ○

Given the matrices:

$$A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

First, let us compute the product AB :

$$AB = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta)(a) + (-\sin \theta)(0) & (\cos \theta)(0) + (-\sin \theta)(b) \\ (\sin \theta)(a) + (\cos \theta)(0) & (\sin \theta)(0) + (\cos \theta)(b) \end{bmatrix}$$

$$= \begin{bmatrix} a \cos \theta & -b \sin \theta \\ a \sin \theta & b \cos \theta \end{bmatrix}$$

Next, let us compute the product BA :

$$BA = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$= \begin{bmatrix} (a)(\cos \theta) + (0)(\sin \theta) & (a)(-\sin \theta) + (0)(\cos \theta) \\ (0)(\cos \theta) + (b)(\sin \theta) & (0)(-\sin \theta) + (b)(\cos \theta) \end{bmatrix}$$

$$= \begin{bmatrix} a \cos \theta & -a \sin \theta \\ b \sin \theta & b \cos \theta \end{bmatrix}$$

For A and B to commute, we require $AB = BA$:

$$\begin{bmatrix} a \cos \theta & -b \sin \theta \\ a \sin \theta & b \cos \theta \end{bmatrix} = \begin{bmatrix} a \cos \theta & -a \sin \theta \\ b \sin \theta & b \cos \theta \end{bmatrix}$$

By comparing corresponding entries in the two resulting matrices:

- Comparing entry (1, 2): $-b \sin \theta = -a \sin \theta \implies (a - b) \sin \theta = 0$
- Comparing entry (2, 1): $a \sin \theta = b \sin \theta \implies (a - b) \sin \theta = 0$

The equation $(a - b) \sin \theta = 0$ is satisfied when either:

$$a - b = 0 \implies a = b$$

or

$$\sin \theta = 0 \implies \theta = n\pi, \text{ where } n \text{ is an integer.}$$

Thus, A and B commute under multiplication if $a = b$ or $\theta = n\pi$, where n is an integer. This matches option (A).

(A) if $a = b$ or $\theta = n\pi$, n is an integer

Q1.96.3 MCQ 1996 1M Ans: C ☒ ☐ ☐

Given the system of linear equations:

$$A_{m \times n} X_{n \times 1} = B_{m \times 1}$$

Let us analyze the specific case presented in the statement where $m = n$ (a square system of equations):

$$\implies A_{n \times n} X_{n \times 1} = B_{n \times 1}$$

- Case 1: A is non-singular ($\det(A) \neq 0$). The matrix A^{-1} exists, and the system possesses a unique solution given by:

$$X = A^{-1}B$$

- Case 2: A is singular ($\det(A) = 0$). The unique solution does not exist. The system will either have infinitely many solutions or no solution at all, depending on the consistency check using the adjugate matrix condition:

$$(\text{adj}(A))B$$

Evaluating the options based on this mathematical definition: The statement claiming that the system has a unique solution whenever $m = n$ and b is a non-zero vector fails to account for the possibility of a singular coefficient matrix.

Therefore, option (C) represents the false statement.

(C) If $m = n$ and b is non-zero vector, then the system has a unique solution.

Q1.95.1 MCQ 1995 1M Ans: A ● ○ ○

Given the $(n + 1) \times (n + 1)$ square matrix A :

$$A = \begin{bmatrix} 1 & a & a^2 & \dots & a^n \\ 1 & a & a^2 & \dots & a^n \\ \vdots & \vdots & \vdots & & \vdots \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & a & a^2 & \dots & a^n \end{bmatrix}$$

Subtracting the first row from all subsequent rows ($R_i \rightarrow R_i - R_1$ for $i = 2, 3, \dots, n + 1$) to reduce it to its row echelon form:

$$\Rightarrow A \sim \begin{bmatrix} 1 & a & a^2 & \dots & a^n \\ 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix}$$

The rank of a matrix is equal to the number of non-zero rows present in its row echelon form. Hence, the rank of the given matrix is 1.

(A) 1

Q1.94.1 MCQ 1994 1M Ans: D ● ○ ○

Given that A and B are real symmetric matrices of size $n \times n$. By definition of symmetric matrices, we have:

$$A' = A$$

$$B' = B$$

where ' denotes the matrix transpose operation.

Let us evaluate the truth value of each option analytically:

- Option A ($AA' = I$): This statement defines an orthogonal matrix. A symmetric matrix is not necessarily orthogonal. For example, if $A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$, then $A' = A$, but $AA' = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix} \neq I$. Thus, this statement is false in general.
- Option B ($A = A^{-1}$): This statement defines an involutory matrix. A symmetric matrix is not necessarily its own inverse. Using the same counterexample $A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$, we find $A^{-1} = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \neq A$. Thus, this statement is false in general.
- Option C ($AB = BA$): Matrix multiplication is generally non-commutative, even for symmetric matrices. Two symmetric matrices commute if and only if their product AB is also symmetric. Since this is a conditional property, it is false in general.
- Option D ($(AB)' = BA$): Let us apply the reversal law of transposes to the expression $(AB)'$:

$$(AB)' = B'A'$$

Substituting the given symmetric conditions ($B' = B$ and $A' = A$):


$$\Rightarrow (AB)' = BA$$

This identity holds unconditionally for all real symmetric matrices. Thus, this statement is always true.

Therefore, option (D) is the correct statement.

(D) $(AB)' = BA$

SOLUTIONS — Linear Algebra



ME

Mechanical Engineering

PrepFusion

Q1.25.1 MCQ 2025 1M Ans: C ● ○ ○

Given that A and B are real symmetric matrices. By definition of symmetric matrices, a matrix is equal to its own transpose:

$$A^T = A \quad \text{and} \quad B^T = B$$

According to the fundamental algebraic properties of matrix transposes, the transpose of a product of two matrices follows the reversal law:

$$(AB)^T = B^T A^T$$

Since this property is an unconditional algebraic identity for any two conformable matrices, it holds true for real symmetric matrices as well. This perfectly matches the statement in option (C).

$$(C) \quad (AB)^T = B^T A^T$$

Q1.24.1 MCQ 2024 1M Ans: D ● ○ ○

The condition for a system of linear equations $AX = B$ to possess infinitely many solutions is:

$$\rho(A) = \rho(A | B) < n$$

where $\rho(A)$ represents the rank of the coefficient matrix, $\rho(A | B)$ is the rank of the augmented matrix, and $n = 3$ is the number of unknown variables.

$$(A | B) = \left[\begin{array}{ccc|c} 1 & 2 & 1 & 5 \\ 2 & a & 4 & 12 \\ 2 & 4 & 6 & b \end{array} \right]$$

$$R_2 \rightarrow R_2 - 2R_1 \text{ and } R_3 \rightarrow R_3 - 2R_1:$$

$$\Rightarrow (A | B) \sim \left[\begin{array}{ccc|c} 1 & 2 & 1 & 5 \\ 0 & a-4 & 2 & 2 \\ 0 & 0 & 4 & b-10 \end{array} \right] \dots (1)$$

To satisfy the conditions for a reduced rank, let us assume $a - 4 = 0 \Rightarrow a = 4$. Substituting this value into equation (1) yields:

$$(A | B) \sim \left[\begin{array}{ccc|c} 1 & 2 & 1 & 5 \\ 0 & 0 & 2 & 2 \\ 0 & 0 & 4 & b-10 \end{array} \right]$$

$$R_3 \rightarrow R_3 - 2R_2:$$

$$\Rightarrow (A | B) \sim \left[\begin{array}{ccc|c} 1 & 2 & 1 & 5 \\ 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & b-14 \end{array} \right]$$

$$\rho(A) = 2$$

For the system to be consistent and have infinite solutions, the rank of the augmented matrix must also equal 2. This requires the last entry of the augmented column to be zero:

$$b - 14 = 0 \Rightarrow b = 14$$

When $a = 4$ and $b = 14$, we satisfy $\rho(A) = \rho(A | B) = 2 \neq n = 3$. Therefore, infinitely many solutions exist. This matches option (d).

$$(D) \quad a = 4, b = 14$$

Key Points

- Rouché-Capelli Theorem for Infinite Solutions: A system has infinitely many solutions if and only if $\rho(A) = \rho(A | B) < n$, where n is the total number of unknowns.

Q1.24.2 MCQ 2024 2M Ans: A ● ● ○

Let λ_1 and λ_2 be the eigenvalues of the given 2×2 matrix A :

$$A = \begin{bmatrix} 1 & a \\ 8 & 3 \end{bmatrix}, \quad \text{where } a > 0$$

According to the properties of square matrices, we have:

- (i) The sum of the eigenvalues is equal to the trace of the matrix:

$$\lambda_1 + \lambda_2 = \text{tr}(A) = 1 + 3 = 4$$

- (ii) The product of the eigenvalues is equal to the determinant of the matrix:

$$\lambda_1 \cdot \lambda_2 = |A| = (1)(3) - (a)(8) = 3 - 8a$$

We are given that one of the eigenvalues of the matrix is negative. Let us assume $\lambda_1 < 0$. Since $a > 0$, the trace ($\lambda_1 + \lambda_2 = 4$) is positive. For a negative eigenvalue ($\lambda_1 < 0$) to sum with another value to a positive result, the second eigenvalue λ_2 must be strictly positive ($\lambda_2 > 0$).

Because one eigenvalue is negative and the other is positive, their product must be strictly negative:

$$\Rightarrow |A| = \lambda_1 \cdot \lambda_2 < 0$$

$$\Rightarrow 3 - 8a < 0$$

Solving the inequality for parameter a :

$$\begin{aligned} \Rightarrow 3 &< 8a \\ \therefore a &> \frac{3}{8} \end{aligned}$$

Thus, the matrix has a negative eigenvalue if a is greater than $\frac{3}{8}$. This matches option (A).

$$(A) \frac{3}{8}$$

Q1.22.1 MCQ 2022 1M Ans: A ● ○ ○

Given the matrix A :

$$A = \begin{bmatrix} 10 & 2k + 5 \\ 3k - 3 & k + 5 \end{bmatrix}$$

By definition, a square matrix is symmetric if it is equal to its own transpose ($A^T = A$). For a 2×2 matrix, this condition means the non-diagonal elements must be equal to each other:

$$a_{12} = a_{21}$$

Substituting the given entry expressions:

$$2k + 5 = 3k - 3$$

Rearranging the terms to isolate the variable k :

$$5 + 3 = 3k - 2k$$

$$\therefore k = 8$$

Thus, the value of k for which the matrix is symmetric is 8. This matches option (a).

(A) 8

Key Points

- A square matrix A is symmetric if $A^T = A$, which implies that the entry at the i -th row and j -th column is identical to the entry at the j -th row and i -th column ($a_{ij} = a_{ji}$).

Q1.22.2 MSQ 2022 2M Ans: A,B ● ● ○

Given the system of homogeneous linear equations:

$$\begin{pmatrix} x & y \end{pmatrix} \begin{bmatrix} 2 & 5 - 2\alpha \\ \alpha & 1 \end{bmatrix} = \begin{pmatrix} 0 & 0 \end{pmatrix}$$

$$2x + \alpha y = 0$$

$$(5 - 2\alpha)x + y = 0$$

We can rewrite this system in standard matrix form $M\mathbf{x} = \mathbf{0}$ by transposing the layout:

$$\begin{bmatrix} 2 & \alpha \\ 5 - 2\alpha & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \dots (1)$$

For a homogeneous linear system to possess an infinite number of non-trivial solutions, the determinant of the coefficient matrix must be exactly equal to zero ($\det(M) = 0$):

$$\begin{vmatrix} 2 & \alpha \\ 5 - 2\alpha & 1 \end{vmatrix} = 0$$

$$(2)(1) - (\alpha)(5 - 2\alpha) = 0$$

$$2 - 5\alpha + 2\alpha^2 = 0$$

$$2\alpha^2 - 5\alpha + 2 = 0$$

$$2\alpha^2 - 4\alpha - \alpha + 2 = 0$$

$$2\alpha(\alpha - 2) - 1(\alpha - 2) = 0$$

$$(2\alpha - 1)(\alpha - 2) = 0$$

$$\therefore \alpha = \frac{1}{2}, \quad \alpha = 2$$

Let us analyze the specific non-trivial solutions for each special value of α :

- Case 1: At $\alpha = \frac{1}{2}$, Substituting $\alpha = \frac{1}{2}$ into equation (1):

$$\begin{bmatrix} 2 & 1/2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \implies 4x + y = 0 \quad \dots (2)$$

Let us evaluate the provided options. For option (b), substituting $x = -1$ and $y = 4$ into equation (2):

$$4(-1) + 4 = 0$$

This satisfies the relation, so option (b) is a valid non-trivial solution.

- Case 2: At $\alpha = 2$, Substituting $\alpha = 2$ into equation (1):

$$\begin{bmatrix} 2 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow x + y = 0 \quad \dots (3)$$

Let us evaluate the provided options. For option (a), substituting $x = 2$ and $y = -2$ into equation (3):

$$2 + (-2) = 0$$

This satisfies the relation, so option (a) is a valid non-trivial solution.

Therefore, both options (A) and (B) represent correct non-trivial solution pairs.

$$(A) \ x = 2, \ y = -2, (B) \ x = -1, \ y = 4$$

Q1.22.3 NAT 2022 2M Ans: 2 ● ● ○

Given that the sum of the eigenvalues is 4 and the product of the eigenvalues is -1 for the 2×2 real matrix:

$$A = \begin{bmatrix} 3 & p \\ p & q \end{bmatrix}$$

Let us apply the fundamental spectral properties of square matrices:

- The sum of the eigenvalues is identical to the trace of the matrix ($\text{tr}(A)$), which is the sum of its main diagonal elements.

$$\Rightarrow 3 + q = 4$$

$$\Rightarrow q = 1$$

- The product of the eigenvalues is identical to the determinant of the matrix ($|A|$).

$$\Rightarrow |A| = (3)(q) - (p)(p) = 3q - p^2 = -1$$

Substituting the calculated value of $q = 1$ into the determinant relationship:

$$\Rightarrow 3(1) - p^2 = -1$$

$$\Rightarrow 3 - p^2 = -1$$

$$\Rightarrow p^2 = 3 + 1$$

$$\Rightarrow p^2 = 4$$

$$\Rightarrow p = \pm 2$$

$$\Rightarrow |p| = |\pm 2| = 2$$

Thus, the absolute numerical value of p is 2.

$$\boxed{2}$$

Key Points

- For any square matrix A , $\sum_{i=1}^n \lambda_i = \text{tr}(A)$.
- For any square matrix A , $\prod_{i=1}^n \lambda_i = \det(A)$.

Mistakes to be avoided

- Remember that while p can be both $+2$ or -2 , the question asks for the absolute value constraint ($|p|$), so the final numerical answer entered into the NAT answer block must be the positive value 2 only.

Q1.22.4 MSQ 2022 2M Ans: B,C ● ● ○

Given that A is a 3×5 real matrix with rank equal to 2:

$$\rho(A) = 2$$

The system under consideration is a set of homogeneous linear equations:

$$A\mathbf{x} = \mathbf{0}$$

where $\mathbf{x}$ represents the column vector of unknown variables. Since A has dimensions 3×5 , the number of unknown variables (columns) is:

$$n = 5$$

Let us analytically evaluate the statements using fundamental linear algebra principles:

- Option B: A homogeneous system of linear equations always admits the trivial solution, where every variable is set to zero ($\mathbf{x} = \mathbf{0}$). Substituting this into the system yields $A(\mathbf{0}) = \mathbf{0}$, which is universally satisfied by a zero vector of size 5×1 . Thus, statement B is **true**.
- Options A, C, and D: Let us apply the Rank-Nullity Theorem to find the dimension of the solution space (the null space or nullity of A):

$$\text{Nullity}(A) = n - \rho(A)$$

Substituting our values ($n = 5$ and $\rho(A) = 2$):

$$\text{Nullity}(A) = 5 - 2 = 3$$

Since the nullity is greater than zero ($\text{Nullity}(A) = 3 > 0$), the matrix has a non-trivial null space containing 3 linearly independent free variables. This means that the system contains infinitely many independent linear combinations of solutions.

Therefore, the system has infinitely many solutions. This makes statement C true, while statements A and D are false.

In conclusion, the correct options are B and C.

(B) The given set of equations will be satisfied by a zero vector of appropriate size.

(C) The given set of equations will have infinitely many solutions.

Q1.21.1 MCQ 2021 2M Ans: B ● ● ○

By definition, if a non-zero vector X is an eigenvector of a square matrix $A_{2 \times 2}$ corresponding to a scalar eigenvalue λ , it satisfies the fundamental characteristic equation:

$$AX = \lambda X$$

According to the question notation, we replace the generic state vector X with the spatial vector p :

$$Ap = \lambda p$$

We are given that a transformed vector p' is generated via the matrix transformation $p' = Ap$. Substituting this definition directly into the eigenvector relation yields:

$$\Rightarrow p' = \lambda p$$

This vector equality tells us that the transformed vector p' points in the exact same spatial direction as the original eigenvector p , scaled in magnitude by a factor of λ . Let us analyze the geometric components of this transformation:

- Directional Behavior: Since $\lambda > 0$, multiplying the vector p by the positive scalar λ scales its length but keeps its angular orientation completely unaltered. Therefore, the counter-clockwise direction angle with the positive x -axis remains unchanged:

$$\text{Direction of } p' = \theta$$

- Magnitude Behavior: Taking the vector norm (magnitude) on both sides of the relation $p' = \lambda p$:

$$\|p'\| = \|\lambda p\|$$

Since $\lambda > 0$, we can factor out the scalar from the norm directly:

$$\therefore \|p'\| = \lambda \|p\|$$

Reviewing the statements, this derivation perfectly confirms the validity of option (B).

$$\boxed{\text{(B) Direction of } p' = \theta, \|p'\| = \lambda \|p\|}$$

Q1.21.2 MCQ 2021 1M Ans: D ● ○ ○

By definition, if a non-zero vector X is an eigenvector of an $n \times n$ matrix A corresponding to an eigenvalue λ , it satisfies the characteristic relation:

$$AX = \lambda X$$

$$Ap = \lambda p$$

We are given the explicit product constraint for the matrix transformation:

$$Ap = \alpha^2 p$$

$$\Rightarrow \alpha^2 p = \lambda p$$

$$\Rightarrow \lambda = \alpha^2$$

Thus, α^2 is an eigenvalue of the $n \times n$ matrix A . According to the spectral mapping properties of matrices, if λ is an eigenvalue of A , then λ^2 is the corresponding eigenvalue of A^2 :

$$\text{Eigenvalue of } A^2 = \lambda^2$$

$$\text{Eigenvalue of } A^2 = (\alpha^2)^2 = \alpha^4$$

$$\boxed{\text{(D) } \alpha^4}$$

Q1.21.3 MCQ 2021 2M Ans: B ● ● ○

Method 1

We aim to find the stationary point of the scalar function $f(x)$ defined over n -dimensional vector space:

$$f(x) = \frac{1}{2} x^T Q x - r^T x$$

where x and r are $n \times 1$ column vectors, and Q is a symmetric $n \times n$ matrix ($Q^T = Q$). To find the stationary point, we compute the gradient of $f(x)$ with respect to the vector x and equate it to the zero vector:

$$\nabla_x f(x) = \mathbf{0}$$

Let us differentiate each matrix term individually using the rules of matrix calculus:

- Quadratic Term Differentiation: The gradient of the quadratic form $\frac{1}{2}x^T Q x$ is:

$$\nabla_x \left(\frac{1}{2} x^T Q x \right) = \frac{1}{2} (Q + Q^T) x$$

Since Q is given as a symmetric matrix ($Q^T = Q$), this simplifies directly to:

$$\frac{1}{2} (2Q) x = Q x$$

- Linear Term Differentiation: The gradient of the linear inner product $r^T x$ is:

$$\nabla_x (r^T x) = r$$

Combining these results, the overall gradient of the vector function becomes:

$$\nabla_x f(x) = Q x - r$$

Setting the gradient vector equal to zero to identify the stationary profile:

$$Q x - r = \mathbf{0}$$

$$Q x = r$$

Multiplying both sides from the left by the matrix inverse Q^{-1} isolates the coordinate vector:

$$Q^{-1} Q x = Q^{-1} r$$

$$\therefore x = Q^{-1} r$$

This matches the expression presented in option (B).

Method 2

To gain clear structural insight into this matrix identity, let us analyze a low-order case where $n = 2$. Let us define the components of the vectors and the symmetric matrix:

$$x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad r = \begin{bmatrix} r_1 \\ r_2 \end{bmatrix}, \quad Q = \begin{bmatrix} a & c \\ c & b \end{bmatrix}$$

$$x^T = \begin{bmatrix} x_1 & x_2 \end{bmatrix}, \quad r^T = \begin{bmatrix} r_1 & r_2 \end{bmatrix}$$

Substituting these matrices into the original scalar expression:

$$f(x_1, x_2) = \frac{1}{2} \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a & c \\ c & b \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} r_1 & r_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Expanding the matrix multiplication blocks step-by-step:

$$f(x_1, x_2) = \frac{1}{2} \begin{bmatrix} ax_1 + cx_2 & cx_1 + bx_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - (r_1 x_1 + r_2 x_2)$$

$$f(x_1, x_2) = \frac{1}{2} (ax_1^2 + cx_1 x_2 + cx_1 x_2 + bx_2^2) - r_1 x_1 - r_2 x_2$$

$$f(x_1, x_2) = \frac{1}{2} ax_1^2 + cx_1 x_2 + \frac{1}{2} bx_2^2 - r_1 x_1 - r_2 x_2$$

For stationary points to exist, the partial derivatives with respect to each scalar coordinate variable must independently vanish:

$$\frac{\partial f}{\partial x_1} = ax_1 + cx_2 - r_1 = 0 \implies ax_1 + cx_2 = r_1$$

$$\frac{\partial f}{\partial x_2} = cx_1 + bx_2 - r_2 = 0 \implies cx_1 + bx_2 = r_2$$

Recombining this system of linear equations back into matrix form:

$$\begin{bmatrix} a & c \\ c & b \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} r_1 \\ r_2 \end{bmatrix}$$

$$Qx = r$$

$$\therefore x = Q^{-1}r$$

This variation verifies that the formulation holds.

$$(B) Q^{-1}r$$

Key Points

- Matrix Calculus Gradients: For standard vector forms, the following fundamental identities apply:

$$\nabla_x(r^T x) = r$$

$$\nabla_x(x^T Qx) = (Q + Q^T)x$$

- Stationary Criteria: Stationary points of multivariate or vector fields occur where the multi-dimensional first derivative (gradient vector) is identically zero: $\nabla f = \mathbf{0}$.

Q1.20.1 MCQ 2020 1M Ans: A ☒ ☐ ☐

Let A , B , and C be real-valued square matrices of the same dimension $n \times n$. Let us evaluate the fundamental algebraic properties of matrix multiplication:

- Associativity: Matrix multiplication is always associative. For any three conformable matrices, grouping does not alter the product outcome:

$$(AB)C = A(BC)$$

Therefore, the statement that matrix multiplication is associative is fundamentally true.

- Commutativity: Matrix multiplication is generally non-commutative. That is, for two generic square matrices A and B :

$$AB \neq BA$$

Thus, it is not commutative in general.

- Definiteness: The product of two positive definite matrices is not necessarily positive definite unless the matrices commute. Hence, it is not always positive definite.
- Computability: Since the matrices are square and share the exact same dimensions ($n \times n$), multiplication is always well-defined and possible to compute.

Since matrix multiplication is associative, it matches the option identifying associativity as a valid property.

$$(A) \text{ associative}$$

Q1.20.2 MCQ 2020 1M Ans: B ● ○ ○

Any square matrix P can be uniquely expressed as the sum of a symmetric matrix S and a skew-symmetric matrix V :

$$P = \left(\frac{P + P^T}{2} \right) + \left(\frac{P - P^T}{2} \right)$$

Let us define these component matrices as:

$$P = S + V$$

where the symmetric part S and skew-symmetric part V are given by:

$$S = \frac{P + P^T}{2} \quad \text{and} \quad V = \frac{P - P^T}{2}$$

Substituting the given component matrices into the sum:

$$P = \begin{bmatrix} -4 & 4 & 2 \\ 4 & 3 & 7/2 \\ 2 & 7/2 & 2 \end{bmatrix} + \begin{bmatrix} 0 & -2 & 3 \\ 2 & 0 & 7/2 \\ -3 & -7/2 & 0 \end{bmatrix}$$

Let us compute the sum by adding corresponding entries:

$$P = \begin{bmatrix} -4 + 0 & 4 + (-2) & 2 + 3 \\ 4 + 2 & 3 + 0 & 7/2 + 7/2 \\ 2 + (-3) & 7/2 + (-7/2) & 2 + 0 \end{bmatrix}$$

Evaluating the arithmetic simplifications:

$$\Rightarrow P = \begin{bmatrix} -4 & 2 & 5 \\ 6 & 3 & 7 \\ -1 & 0 & 2 \end{bmatrix}$$

$$(B) \begin{pmatrix} -4 & 2 & 5 \\ 6 & 3 & 7 \\ -1 & 0 & 2 \end{pmatrix}$$

Key Points

- Decomposition Theorem: Every square matrix can be uniquely decomposed into a symmetric part $S = \frac{1}{2}(M + M^T)$ and a skew-symmetric part $V = \frac{1}{2}(M - M^T)$.

Q1.20.3 NAT 2020 1M Ans: 1 ● ○ ○

Let I be an identity matrix of dimension 100×100 . By definition, an identity matrix is a diagonal matrix whose principal diagonal elements are all equal to 1.

The characteristic equation of an identity matrix I is given by:

$$|I - \lambda I| = 0$$

$$(1 - \lambda)^{100} = 0$$

Solving this equation yields the eigenvalues of the matrix. Since it is a 100×100 matrix, it possesses exactly 100 eigenvalues, all of which are identically equal to 1:

$$\lambda_1 = \lambda_2 = \dots = \lambda_{100} = 1$$

The problem defines E as the set containing the distinct real eigenvalues of this matrix. In set theory, duplicate values are not counted more than once, meaning a set only contains unique elements. Therefore, we express the set E as:

$$E = \{1\}$$

Counting the cardinality (the number of distinct elements) of the set E :

$$n(E) = 1$$

Thus, the total number of distinct elements in E is 1.

1

Q1.19.1 MCQ 2019 2M Ans: C ● ○ ○

The given system of linear equations can be systematically written in matrix form as:

$$\begin{bmatrix} 1 & 1 & 1 \\ a & -a & 3 \\ 5 & -3 & a \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \\ 6 \end{bmatrix}$$

This is of the form $AX = B$, where the coefficient matrix A , the variable vector X , and the constant vector B are defined as:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ a & -a & 3 \\ 5 & -3 & a \end{bmatrix}, \quad X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad \text{and} \quad B = \begin{bmatrix} 1 \\ 5 \\ 6 \end{bmatrix}$$

According to Rouché–Capelli theorem, the condition for a non-homogeneous system to have infinitely many solutions (many solutions) is:

$$\rho(A) = \rho(A|B) < n$$

where $\rho(A)$ is the rank of the coefficient matrix, $\rho(A|B)$ is the rank of the augmented matrix, and $n = 3$ is the number of unknown variables.

Let us construct the augmented matrix $[A|B]$:

$$[A|B] = \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ a & -a & 3 & 5 \\ 5 & -3 & a & 6 \end{array} \right]$$

Performing $R_2 \rightarrow R_2 - aR_1$ and $R_3 \rightarrow R_3 - 5R_1$:

$$\Rightarrow [A|B] = \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & -2a & 3-a & 5-a \\ 0 & -8 & a-5 & 1 \end{array} \right]$$

$R_3 \rightarrow aR_3 - 4R_2$:

$$\Rightarrow [A|B] = \left[\begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & -2a & 3-a & 5-a \\ 0 & 0 & a^2-a-12 & 5a-20 \end{array} \right]$$

For the system to have a rank less than 3 ($\rho(A) = \rho(A|B) = 2 < 3$), the entire last row of the row-echelon form matrix must vanish completely. This requires both diagonal elements in the third row to equal zero simultaneously:

$$a^2 - a - 12 = 0 \quad \& \quad 5a - 20 = 0$$

Solving these equations:

- From the linear condition:

$$5a - 20 = 0 \implies 5a = 20 \implies a = 4$$

- Substituting $a = 4$ into the quadratic condition to verify compatibility:

$$(4)^2 - (4) - 12 = 16 - 4 - 12 = 0$$

Since $a = 4$ satisfies both equations simultaneously, it ensures that $\rho(A) = \rho(A|B) = 2$, yielding infinitely many solutions.

(C) 4

Q1.19.2 MCQ 2019 1M Ans: D ● ○ ○

Given the linear system of matrix transformations:

$$AX = R \quad (1)$$

$$AX = \lambda X \quad (2)$$

Comparing equation (1) and equation (2), since the left-hand sides are identical, we can equate their right-hand sides directly:

$$R = \lambda X$$

Substituting the given coordinate values for the output vector R and the column eigenvector X :

$$\Rightarrow \begin{pmatrix} 32 \\ 16 \\ 64 \end{pmatrix} = \lambda \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix}$$

Equating the corresponding components from each row yields a system of three univariable scalar equations:

- From row 1: $32 = 2\lambda \Rightarrow \lambda = 16$
- From row 2: $16 = 1\lambda \Rightarrow \lambda = 16$
- From row 3: $64 = 4\lambda \Rightarrow \lambda = 16$

Since all three independent coordinate lines yield an identical value, the scalar parameter is consistent:

$$\therefore \lambda = 16$$

(D) 16

Q1.18.1 NAT 2018 1M Ans: 0.25 ● ○ ○

Given the square matrix A of order 3×3 :

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow |A| = (1)(4)(1) = 4$$

$$\det(A^{-1}) = |A^{-1}| = |A|^{-1} = \frac{1}{|A|}$$

$$\therefore \det(A^{-1}) = \frac{1}{4} = 0.25$$

0.25

Q1.17.1 MCQ 2017 2M Ans: D ● ● ○

Given the matrix:

$$P = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Let us systematically analyze each statement to identify the incorrect one:

Step 1: Check if P is orthogonal A matrix is orthogonal if $P^T P = I$, which also directly implies $P^{-1} = P^T$.

$$P^T = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{-1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$P^T P = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{-1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{-1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$= \begin{bmatrix} \left(\frac{1}{2} + 0 + \frac{1}{2}\right) & (0 + 0 + 0) & \left(\frac{1}{2} + 0 - \frac{1}{2}\right) \\ (0 + 0 + 0) & (0 + 1 + 0) & (0 + 0 + 0) \\ \left(\frac{1}{2} + 0 - \frac{1}{2}\right) & (0 + 0 + 0) & \left(\frac{1}{2} + 0 + \frac{1}{2}\right) \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Since $P^T P = I$, Statement B (P is orthogonal) and Statement C (Inverse of P is equal to its transpose) are both completely CORRECT.

Step 2: Check the determinant of P Expanding the determinant along the second row or second column:

$$\det(P) = 1 \cdot \begin{vmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{vmatrix} = 1 \cdot \left(\frac{1}{2} - \left(-\frac{1}{2}\right)\right) = 1$$

Thus, $\det(P) = 1$, which means Statement A is CORRECT.

Step 3: Evaluate the Eigenvalues of P The characteristic equation is given by $\det(P - \lambda I) = 0$:

$$\begin{vmatrix} \frac{1}{\sqrt{2}} - \lambda & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 - \lambda & 0 \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} - \lambda \end{vmatrix} = 0$$

$$(1 - \lambda) \left[\left(\frac{1}{\sqrt{2}} - \lambda\right)^2 - \left(-\frac{1}{2}\right) \right] = 0$$

$$(1 - \lambda) \left[\frac{1}{2} + \lambda^2 - \sqrt{2}\lambda + \frac{1}{2} \right] = 0$$

$$(1 - \lambda) [\lambda^2 - \sqrt{2}\lambda + 1] = 0$$

This yields the eigenvalues:

- $\lambda_1 = 1$
- From $\lambda^2 - \sqrt{2}\lambda + 1 = 0$:

$$\lambda = \frac{\sqrt{2} \pm \sqrt{(\sqrt{2})^2 - 4(1)(1)}}{2} = \frac{\sqrt{2} \pm \sqrt{2-4}}{2} = \frac{\sqrt{2} \pm i\sqrt{2}}{2} = \frac{1}{\sqrt{2}} \pm i\frac{1}{\sqrt{2}} = e^{\pm i\pi/4}$$

Since two of the eigenvalues $\left(\frac{1}{\sqrt{2}} \pm i\frac{1}{\sqrt{2}}\right)$ are complex conjugates, Statement D is INCORRECT.

(D) All Eigenvalues of P are real numbers

Key Points

- For any real orthogonal matrix Q , $Q^T Q = I$, $\det(Q) = \pm 1$, and its inverse satisfies $Q^{-1} = Q^T$.
- The eigenvalues of a real orthogonal matrix always lie on the unit circle in the complex plane, meaning their absolute magnitude is always $|\lambda| = 1$. They do not necessarily have to be real numbers.

Q1.17.2 NAT 2017 1M Ans: 5 ● ○ ○

Let λ_1 and λ_2 be the two eigenvalues of the given 2×2 matrix A .

Given:

$$\lambda_1 = 10$$

$$\det(A) = 50$$

A fundamental theorem in linear algebra states that the determinant of any square matrix is exactly equal to the product of all its eigenvalues:

$$\det(A) = \lambda_1 \lambda_2$$

$$\Rightarrow 10 \cdot \lambda_2 = 50$$

$$\therefore \lambda_2 = \frac{50}{10} = 5$$

Thus, the other eigenvalue of the matrix is 5.

5

Q1.17.3 NAT 2017 2M Ans: 0 ● ● ○

We want to evaluate the inner product expression $x_1^T x_2$. Let us substitute the given matrix expressions for eigenvectors x_1 and x_2 :

$$x_1^T x_2 = [70 \quad \lambda_1 - 50] \begin{bmatrix} \lambda_2 - 80 \\ 70 \end{bmatrix}$$

$$\Rightarrow x_1^T x_2 = [70(\lambda_2 - 80) + (\lambda_1 - 50)70]$$

$$\Rightarrow x_1^T x_2 = [70\lambda_2 - 70(80) + 70\lambda_1 - 70(50)]$$

$$\Rightarrow x_1^T x_2 = [70(\lambda_1 + \lambda_2) - 70(80 + 50)]$$

According to the linear algebra properties of eigenvalues, the sum of all eigenvalues of a matrix is equal to its trace (the sum of its principal diagonal elements). From matrix $A = \begin{bmatrix} 50 & 70 \\ 70 & 80 \end{bmatrix}$, we have:

$$\lambda_1 + \lambda_2 = \text{trace}(A) = 50 + 80$$

Substituting $\lambda_1 + \lambda_2 = 50 + 80$ back into our inner product equation:

$$\Rightarrow x_1^T x_2 = [70(50 + 80) - 70(50 + 80)]$$

$$\therefore x_1^T x_2 = [0] = 0$$

0

Key Points

- Matrix Trace Formula: For any square matrix A , the sum of its eigenvalues matches the sum of its main diagonal elements: $\sum_{i=1}^n \lambda_i = \text{tr}(A)$.

Q1.16.1 MCQ 2016 1M Ans: D ☒ ☐ ☐

The given system of linear equations is:

$$2x + 5y = 2 \quad (1)$$

$$-4x + 3y = -30 \quad (2)$$

To find the values of x and y that satisfy both equations, we can systematically solve the system.

Method 1

Multiplying equation (1) by 2 to align the coefficients of x :

$$4x + 10y = 4 \quad (3)$$

Adding equation (2) and equation (3):

$$\begin{aligned} (-4x + 3y) + (4x + 10y) &= -30 + 4 \\ \Rightarrow 13y &= -26 \\ \Rightarrow y &= -2 \end{aligned}$$

Substituting $y = -2$ into equation (1):

$$\begin{aligned} 2x + 5(-2) &= 2 \\ \Rightarrow 2x - 10 &= 2 \\ \Rightarrow 2x &= 12 \Rightarrow x = 6 \end{aligned}$$

Thus, the unique solution values that satisfy the equations simultaneously are $x = 6$ and $y = -2$.

Method 2

We can verify the solution directly by testing the given options back into the simultaneous equations. Substituting the coordinate parameters $x = 6$ and $y = -2$:

- For equation (1): $2(6) + 5(-2) = 12 - 10 = 2$ (Satisfied)
- For equation (2): $-4(6) + 3(-2) = -24 - 6 = -30$ (Satisfied)

Since both equality conditions hold perfectly, option (d) is correct.

(D) 6, -2

Q1.16.2 MCQ 2016 1M Ans: A ☒ ☐ ☐

Given the 2×2 real symmetric matrix A :

$$A = \begin{bmatrix} 2 & 1 \\ 1 & k \end{bmatrix}$$

Let λ_1 and λ_2 be the eigenvalues of matrix A . We can determine the properties of these eigenvalues by utilizing fundamental matrix relationships:

$$\text{trace}(A) = \lambda_1 + \lambda_2 = 2 + k$$

$$\det(A) = \lambda_1 \lambda_2 = (2)(k) - (1)(1) = 2k - 1$$

The problem specifies that the eigenvalues must be strictly positive ($\lambda_1 > 0$ and $\lambda_2 > 0$). For two real quantities to both be positive, their sum must be positive and their product must also be positive:

1. Sum Condition:

$$\lambda_1 + \lambda_2 > 0 \implies 2 + k > 0 \implies k > -2$$

2. Product Condition:

$$\lambda_1 \lambda_2 > 0 \implies 2k - 1 > 0 \implies 2k > 1 \implies k > \frac{1}{2}$$

To find the range of k that satisfies both criteria simultaneously, we find the intersection of the two inequalities:

$$k \in (-2, \infty) \cap \left(\frac{1}{2}, \infty\right)$$

$$\therefore k > \frac{1}{2}$$

Thus, the condition for which the eigenvalues are positive is $k > 1/2$, which corresponds to option (a).

$$(A) \ k > 1/2$$

Q1.16.3 MCQ 2016 1M Ans: C ☒ ☐ ☐

A matrix is called a skew-symmetric matrix if its transpose is equal to its negative:

$$A^T = -A$$

In terms of individual scalar elements, this structural property translates to:

$$a_{ij} = -a_{ji}, \quad \forall i, j$$

For the principal diagonal entries where the row index matches the column index ($i = j$):

$$a_{ii} = -a_{ii} \implies 2a_{ii} = 0 \implies a_{ii} = 0$$

This mathematically proves that all the main diagonal elements of a real skew-symmetric matrix must be completely zero. Therefore, the description perfectly corresponds to the definition of a skew-symmetric matrix, making option (C) correct.

$$(C) \ A^T = -A$$

Q1.16.4 NAT 2016 2M Ans: 2 ☒ ☒ ☐

Given the 3×3 square matrix A :

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

Observing that A is an upper triangular matrix, its eigenvalues λ are simply its principal diagonal entries:

$$\implies \lambda = 2, 2, 3$$

Here, the eigenvalue $\lambda = 2$ is a repeated root with an algebraic multiplicity (AM) of 2, while $\lambda = 3$ is a distinct root with an algebraic multiplicity of 1.

To find the total number of linearly independent eigenvectors of A , we compute the geometric multiplicity (GM) for each eigenvalue, which equals the dimension of the null space of $(A - \lambda I)$:

$$GM = n - \text{rank}(A - \lambda I)$$

where $n = 3$ is the order of the matrix.

Case 1: For the repeated eigenvalue $\lambda = 2$ Let us construct the characteristic matrix block $(A - 2I)$:

$$A - 2I = \begin{bmatrix} 2-2 & 1 & 0 \\ 0 & 2-2 & 0 \\ 0 & 0 & 3-2 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Applying a simple row swap operation ($R_2 \leftrightarrow R_3$) to bring it to row-echelon form:

$$\sim \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

From the row-echelon form, the matrix clearly has 2 non-zero rows. Thus:

$$r = \text{rank}(A - 2I) = 2$$

The number of linearly independent eigenvectors corresponding to $\lambda = 2$ is:

$$\text{GM}_{\lambda=2} = n - r = 3 - 2 = 1$$

Case 2: For the distinct eigenvalue $\lambda = 3$ Since $\lambda = 3$ is a non-repeated, distinct eigenvalue with an algebraic multiplicity of 1, its geometric multiplicity is guaranteed to be exactly 1:

$$\text{GM}_{\lambda=3} = 1$$

Total Number of Linearly Independent Eigenvectors: Summing up the geometric multiplicities across all unique eigenvalues:

$$\text{Total Linearly Independent Eigenvectors} = \text{GM}_{\lambda=2} + \text{GM}_{\lambda=3} = 1 + 1 = 2$$

2

Key Points

- Algebraic vs Geometric Multiplicity: Algebraic multiplicity (AM) is the number of times an eigenvalue repeats as a root of the characteristic polynomial. Geometric multiplicity (GM) is the number of linearly independent eigenvectors corresponding to that eigenvalue.
- Multiplicity Bounds: For any eigenvalue λ , the inequality $1 \leq \text{GM} \leq \text{AM}$ always holds. If $\text{AM} = 1$, then GM is automatically forced to be 1.
- The number of independent eigenvectors associated with a matrix transformation is given by $\sum \text{GM}_i = \sum (n - \text{rank}(A - \lambda_i I))$.

Mistakes to be avoided

- Do not assume a 3×3 matrix automatically has 3 independent eigenvectors. If a matrix has repeated eigenvalues and is non-diagonalizable (like this one), the total number of independent eigenvectors will be strictly less than n .

Q1.15.1 MCQ 2015 1M Ans: B ● ○ ○

Given that the matrix is a singular matrix, we can analyze the behavior of its eigenvalues based on foundational matrix properties.

By definition, a square matrix A is singular if and only if its determinant is exactly equal to zero:

$$\det(A) = 0$$

According to the product property of matrix eigenvalues, the determinant of any square matrix is identical to the product of all its individual eigenvalues:

$$\text{Product of eigenvalues} = \det(A)$$

Substituting the singularity condition into this relationship yields:

$$\lambda_1 \lambda_2 \dots \lambda_n = 0$$

For the product of a set of scalar values to equal zero, mathematical logic dictates that at least one of the individual terms in that product chain must be zero.

$\therefore$ At least one eigenvalue is zero.

This conclusion perfectly aligns with the criteria given for option (B).

(B) Zero

Q1.15.2 MCQ 2015 1M Ans: A ● ○ ○

Based on the foundational properties of matrix determinants, we analyze the effect of row or column permutations.

Let A be a square matrix of order $n \times n$, and let $\det(A)$ denote its determinant value. If any two distinct rows ($R_i \leftrightarrow R_j$) or any two distinct columns ($C_i \leftrightarrow C_j$) of the determinant are interchanged to form a new modified matrix A' , then the sign of the determinant changes while its absolute magnitude remains completely preserved:

$$\det(A') = -\det(A)$$

Expressing this relationship structurally, the value of the newly altered determinant is scaled by a factor of -1 :

$$\text{New Determinant Value} = (-1) \times \text{Original Determinant Value}$$

Therefore, the statement matches perfectly with option (A).

(A) Absolute value remains unchanged but sign will change.

Q1.15.3 MCQ 2015 2M Ans: A ● ● ○

To find the inverse of the given 2×2 matrix P , we first evaluate its determinant, denoted as $|P|$ or $\det(P)$.

Given the matrix configuration, the determinant calculation is structured as:

$$|P| = (4 + 3i)(4 - 3i) - (-i)(i)$$

$$\Rightarrow |P| = (4 + 3i)(4 - 3i) + i^2$$

Expanding the complex products using algebraic identity rules:

$$(4 + 3i)(4 - 3i) = 4^2 - (3i)^2 = 16 - 9i^2$$

Since $i^2 = -1$, substituting this value simplifies the expression:

$$16 - 9(-1) = 16 + 9 = 25$$

$$\Rightarrow |P| = 25 + (-1) = 24$$

Since $|P| = 24 \neq 0$, the matrix is non-singular, proving that the inverse matrix P^{-1} definitively exists.

By definition, the inverse of a 2×2 square matrix is given by the formula:

$$\therefore P^{-1} = \frac{\text{adj } P}{|P|}$$

To find the adjugate matrix ($\text{adj } P$) for a 2×2 system, we swap the principal diagonal elements and flip the signs of the off-diagonal elements:

$$\text{adj } P = \begin{bmatrix} 4 - 3i & i \\ -i & 4 + 3i \end{bmatrix}$$

$$P^{-1} = \frac{1}{24} \begin{bmatrix} 4 - 3i & i \\ -i & 4 + 3i \end{bmatrix}$$

$$(A) \frac{1}{24} \begin{bmatrix} 4 - 3i & i \\ -i & 4 + 3i \end{bmatrix}$$

Q1.14.1 MCQ 2014 1M Ans: A ● ○ ○

To determine the determinant of a scaled matrix, we use fundamental scaling properties of determinants.

If A is an $n \times n$ square matrix, then scaling the entire matrix by a scalar constant K scales its determinant by K^n :

$$|KA| = K^n |A|$$

$$A = \begin{pmatrix} 1 & 3 & 0 \\ 2 & 6 & 4 \\ -1 & 0 & 2 \end{pmatrix}$$

$$|A| = 1 \cdot \begin{vmatrix} 6 & 4 \\ 0 & 2 \end{vmatrix} - 3 \cdot \begin{vmatrix} 2 & 4 \\ -1 & 2 \end{vmatrix} + 0 \cdot \begin{vmatrix} 2 & 6 \\ -1 & 0 \end{vmatrix}$$

$$|A| = 1(12 - 0) - 3(4 - (-4)) + 0$$

$$|A| = 12 - 3(8) = 12 - 24 = -12$$

$$2A = \begin{pmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{pmatrix}$$

Applying the determinant scaling property for $n = 3$ and $K = 2$:

$$\therefore |2A| = 2^3 |A| = 8 \cdot (-12) = -96$$

$$(A) -96$$

Key Points

- **Determinant Scaling Property:** For any matrix $A \in \mathbb{R}^{n \times n}$ and a scalar K , the relation $|KA| = K^n |A|$ holds because the scalar factors out from each of the n rows independently.
- **Order Dependency:** The exponent of the scaling factor is always exactly equal to the dimension/order (n) of the square matrix.

Mistakes to be avoided

- Do not mistake matrix scaling for scalar arithmetic by writing $|2A| = 2|A|$. Remember that for a 3×3 matrix, the factor scales each row, meaning it must be cubed ($2^3 = 8$).

Q1.14.2 MCQ 2014 1M Ans: B ● ○ ○

To analyze the relationship between the given vectors, let us define them as:

$$\vec{a} = \vec{i} + \vec{j} + \vec{k}$$

$$\vec{b} = 2\vec{i} + 3\vec{j} + \vec{k}$$

$$\vec{c} = 5\vec{i} + 6\vec{j} + 4\vec{k}$$

First, checking basic dot product configurations to evaluate potential orthogonality conditions:

$$\vec{a} \cdot \vec{b} = (1 \cdot 2) + (1 \cdot 3) + (1 \cdot 1) = 2 + 3 + 1 = 6 \neq 0$$

The magnitude of $\vec{a}$ is given by:

$$|\vec{a}| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$$

Since $\vec{a} \cdot \vec{b} \neq 0$, options stating orthogonality or simple direct vector properties can be ruled out.

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 3 & 1 \\ 5 & 6 & 4 \end{pmatrix}$$

Applying $R_2 \rightarrow R_2 - 2R_1$ and $R_3 \rightarrow R_3 - 5R_1$:

$$A \sim \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & -1 \\ 0 & 1 & -1 \end{pmatrix}$$

Next, applying $R_3 \rightarrow R_3 - R_2$:

$$A \sim \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\text{Rank}(A) = 2$$

Because $\text{Rank}(A) < 3$ (the total number of vectors), the row vectors are linearly dependent. Hence, the given vectors are linearly dependent.

(B) The vectors are linearly dependent

Q1.14.3 MCQ 2014 1M Ans: D ● ○ ○

$$A = \begin{pmatrix} -5 & 2 \\ -9 & 6 \end{pmatrix}$$

The characteristic equation of a matrix A is determined using the relation $|A - \lambda I| = 0$:

$$\begin{vmatrix} -5 - \lambda & 2 \\ -9 & 6 - \lambda \end{vmatrix} = 0$$

$$(-5 - \lambda)(6 - \lambda) - (2)(-9) = 0$$

$$\Rightarrow -30 + 5\lambda - 6\lambda + \lambda^2 + 18 = 0$$

$$\Rightarrow \lambda^2 - \lambda - 12 = 0$$

$$(\lambda - 4)(\lambda + 3) = 0$$

$$\therefore \lambda_1 = 4, \quad \lambda_2 = -3$$

Now, we evaluate the eigenvector configurations for each distinct eigenvalue using the system $(A - \lambda I)X = O$, where $X = \begin{bmatrix} x \\ y \end{bmatrix}$.

- Case 1: For $\lambda_1 = 4$ Substituting $\lambda = 4$ into the characteristic vector relation:

$$\begin{bmatrix} -5-4 & 2 \\ -9 & 6-4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -9 & 2 \\ -9 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This matrix reduction yields the singular linear relation:

$$-9x + 2y = 0 \Rightarrow 9x = 2y \Rightarrow \frac{x}{2} = \frac{y}{9}$$

Thus, the base eigenvector corresponding to $\lambda_1 = 4$ is proportional to:

$$X_1 = \begin{bmatrix} 2 \\ 9 \end{bmatrix}$$

- Case 2: For $\lambda_2 = -3$ Substituting $\lambda = -3$ into the characteristic vector relation:

$$\begin{bmatrix} -5-(-3) & 2 \\ -9 & 6-(-3) \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -2 & 2 \\ -9 & 9 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$-2x + 2y = 0 \Rightarrow x = y$$

Thus, the base eigenvector corresponding to $\lambda_2 = -3$ is proportional to:

$$X_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$(D) \begin{Bmatrix} 1 \\ 1 \end{Bmatrix}$$

Q1.14.4 MCQ 2014 1M Ans: D ● ○ ○

Based on the foundational properties of linear transformations, we evaluate the geometric relationship between the eigenvectors of a real symmetric matrix.

An important theorem in linear algebra states that for any real symmetric matrix (where $A = A^T$), the eigenvectors corresponding to distinct eigenvalues are always pair-wise orthogonal to one another.

Geometrically, orthogonality means that the inner product (or dot product) of these vector spaces equals zero. Let us define two such distinct eigenvectors in $\mathbb{R}^3$ space as:

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

Taking their dot product and equating it to zero based on the orthogonality condition:

$$X \cdot Y = 0$$

$$\therefore x_1y_1 + x_2y_2 + x_3y_3 = 0$$

$$(D) 0$$

Q1.14.5 MCQ 2014 1M Ans: D ● ● ○

To determine the mathematically correct statement among the given matrix identities, we evaluate each option based on general matrix algebra and non-commutative matrix properties.

- Analysis of Option (A): Consider the left distributive law for matrix multiplication:

$$P(Q + R) = PQ + PR$$

Since matrix multiplication is generally non-commutative ($PR \neq RP$), this expression cannot be rearranged into $PQ + RP$:

$$P(Q + R) \neq PQ + RP$$

Thus, statement (A) is false.

- Analysis of Option (B): Consider the expansion of the squared matrix difference:

$$\begin{aligned}(P - Q)^2 &= (P - Q)(P - Q) \\ \Rightarrow (P - Q)^2 &= P^2 - PQ - QP + Q^2\end{aligned}$$

Because matrix multiplication is non-commutative, $PQ \neq QP$ in general, meaning $-PQ - QP$ cannot be simplified to $-2PQ$:

$$(P - Q)^2 \neq P^2 - 2PQ + Q^2$$

Thus, statement (B) is false.

- Analysis of Option (C): In matrix algebra, the determinant operation is not additive:

$$\det(P + Q) \neq \det(P) + \det(Q)$$

We can verify this with a clear counterexample. Let:

$$P = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad Q = \begin{pmatrix} 0 & 0 \\ 0 & 3 \end{pmatrix}$$

Computing individual determinants:

$$\det(P) = 0, \quad \det(Q) = 0 \implies \det(P) + \det(Q) = 0$$

Now, computing the determinant of their sum:

$$P + Q = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix} \implies \det(P + Q) = (2)(3) - (0)(0) = 6$$

Since $6 \neq 0$, statement (C) is verified to be false.

- Analysis of Option (D): Consider the expansion of the squared matrix sum:

$$\begin{aligned}(P + Q)^2 &= (P + Q)(P + Q) \\ \Rightarrow (P + Q)^2 &= P^2 + PQ + QP + Q^2\end{aligned}$$

This identity strictly follows the distributive property of matrix algebra without making any invalid commutative assumptions. Therefore, statement (D) is the only correct statement.

$$\boxed{(D) \ (P + Q)^2 = P^2 + PQ + QP + Q^2}$$

Key Points

- **Non-Commutativity of Matrices:** For any two square matrices A and B , $AB \neq BA$ in general. Consequently, standard algebraic binomial expansions like $(A + B)^2 = A^2 + 2AB + B^2$ do not apply unless A and B explicitly commute.
- **Determinant Non-Additivity:** The determinant is a multiplicative operator ($\det(AB) = \det(A) \det(B)$), not an additive one ($\det(A + B) \neq \det(A) + \det(B)$).

Q1.13.1 MCQ 2013 1M Ans: C ● ○ ○

Based on fundamental properties in linear algebra and matrix spectral theory, we analyze the nature of the eigenvalues belonging to a special class of square matrices.

By definition, a real symmetric matrix A satisfies the relation:

$$A = A^T$$

where all entries a_{ij} are real numbers ($a_{ij} \in \mathbb{R}$). A core property derived from the Spectral Theorem states that the eigenvalues of any real symmetric matrix are guaranteed to be real numbers.

To see this conceptually, if λ is an eigenvalue of A with a non-zero eigenvector X , we have:

$$AX = \lambda X$$

Taking the conjugate transpose (Hermitian conjugate) on both sides and utilizing the real symmetric property ($A^* = A^T = A$), we find that the complex conjugate of the eigenvalue must equal itself:

$$\bar{\lambda} = \lambda$$

This condition holds true if and only if the imaginary component is zero, meaning that λ is strictly real. Therefore, the statement perfectly aligns with option (C).

(C) Real

Q1.13.2 MCQ 2013 1M Ans: C ● ● ○

To check the linear dependency of the given functions, we analyze whether one function can be expressed as a linear combination of the others.

Recall the standard trigonometric double-angle identity for cosine:

$$\cos 2x = \cos^2 x - \sin^2 x$$

$$\cos 2x = (1) \cos^2 x + (-1) \sin^2 x$$

By definition, a set of functions is linearly dependent if at least one function in the set can be written as a linear combination of the remaining functions with non-trivial coefficients.

Here, since $\cos 2x$ is successfully expressed as a linear combination of $\sin^2 x$ and $\cos^2 x$, the functions $\cos 2x$, $\sin^2 x$, and $\cos^2 x$ are definitively linearly dependent. This matches the conclusion for option (C).

(C) $\cos 2x$, $\sin^2 x$ and $\cos^2 x$

Q1.12.1 MCQ 2012 2M Ans: B ● ● ○

To find the normalized eigenvector of a matrix, we divide the given basis vector by its Euclidean length (or norm). Suppose an unnormalized basis eigenvector of matrix A is given by the column vector:

$$X = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

The mathematical definition of the standard Euclidean norm (length) $\|X\|$ for a vector $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ is:

$$\|X\| = \sqrt{x_1^2 + x_2^2}$$

Substituting our specific component values into the norm formula:

$$\|X\| = \sqrt{(1)^2 + (-1)^2} = \sqrt{1+1} = \sqrt{2}$$

The normalized unit eigenvector is structurally defined as:

$$\hat{X} = \frac{X}{\|X\|}$$

Dividing each individual coordinate component by the calculated scalar magnitude:

$$\therefore \frac{X}{\|X\|} = \begin{bmatrix} \frac{1}{\sqrt{(1)^2 + (-1)^2}} \\ \frac{-1}{\sqrt{(1)^2 + (-1)^2}} \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix}$$

$$(B) \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix}$$

Key Points

- Vector Normalization: Normalizing a vector scales its magnitude to exactly 1 while strictly preserving its directional orientation in vector space.
- Euclidean Vector Norm: For a vector $X \in R^n$, the standard L_2 norm is computed as $\|X\| = \sqrt{\sum_{i=1}^n x_i^2}$.

Q1.10.1 MCQ 2010 1M Ans: A ● ○ ○

To determine which of the given vectors is an eigenvector of the matrix, we can approach the problem using two distinct methods: finding the eigenvalues explicitly, or verification via the definition of an eigenvector.

Let the given matrix A be defined as:

$$A = \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$$

Method 1

We first determine the eigenvalues of matrix A using its characteristic equation $\det(A - \lambda I) = 0$:

$$\begin{vmatrix} 2 - \lambda & 2 \\ 1 & 3 - \lambda \end{vmatrix} = 0$$

$$(2 - \lambda)(3 - \lambda) - (2)(1) = 0$$

$$\Rightarrow \lambda^2 - 5\lambda + 6 - 2 = 0$$

$$\Rightarrow \lambda^2 - 5\lambda + 4 = 0$$

$$(\lambda - 1)(\lambda - 4) = 0 \Rightarrow \lambda_1 = 1, \quad \lambda_2 = 4$$

Now, we compute the eigenvector space corresponding to the eigenvalue $\lambda_1 = 1$ using the system $(A - \lambda_1 I)X = O$:

$$\begin{bmatrix} 2 - 1 & 2 \\ 1 & 3 - 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This system yields a single linearly independent equation:

$$x + 2y = 0 \Rightarrow x = -2y$$

Choosing $y = -1$ gives the representative basis eigenvector:

$$X_1 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

$$(A) \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix} \right\}$$

Method 2

Alternatively, we can directly test the options using the foundational eigenvalue-eigenvector definition $AX = \lambda X$. A vector X is an eigenvector if its transformation results in a pure scalar multiple of itself.

Let us test option (A), where $X = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$:

$$AX = \begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \end{bmatrix} = \begin{bmatrix} (2)(2) + (2)(-1) \\ (1)(2) + (3)(-1) \end{bmatrix} = \begin{bmatrix} 4 - 2 \\ 2 - 3 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

Observe that:

$$AX = 1 \cdot \begin{bmatrix} 2 \\ -1 \end{bmatrix} = 1 \cdot X$$

Since $AX = \lambda X$ is satisfied with $\lambda = 1$, the vector $\begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is confirmed to be a valid eigenvector.

$$(A) \left\{ \begin{bmatrix} 2 \\ -1 \end{bmatrix} \right\}$$

Q1.09.1 MCQ 2009 1M Ans: A ☒ ☐ ☐

To find the unknown value x , we utilize the defining properties of an orthogonal matrix.

An orthogonal matrix M is defined by the property that its transpose is equal to its inverse:

$$M^T = M^{-1}$$

$$MM^T = I$$

Substituting the explicit matrix layouts based on the problem variables:

$$\begin{bmatrix} \frac{3}{5} & \frac{4}{5} \\ x & \frac{3}{5} \end{bmatrix} \begin{bmatrix} \frac{3}{5} & x \\ \frac{4}{5} & \frac{3}{5} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Performing matrix multiplication for the row-column pairs, let us isolate the element corresponding to the first row and second column (position (1,2)) of the resulting identity matrix:

$$\left(\frac{3}{5} \cdot x\right) + \left(\frac{4}{5} \cdot \frac{3}{5}\right) = 0$$

$$\Rightarrow \frac{3}{5}x + \frac{12}{25} = 0$$

$$\frac{3x}{5} = -\frac{12}{25}$$

$$x = \frac{-12}{25} \cdot \frac{5}{3} = -\frac{4}{5}$$

$$(A) -\frac{4}{5}$$

Key Points

- Orthogonal Matrix Definition: A square matrix M is orthogonal if $MM^T = M^T M = I$.
- Row and Column Norms: In an orthogonal matrix, every row (and column) is a unit vector, and the dot product between any two distinct rows (or columns) is strictly zero.

Q1.08.1 MCQ 2008 1M Ans: C ● ○ ○

To find the sum of the remaining eigenvalues, we use a fundamental properties of matrix traces. Given that one of the eigenvalues of the matrix A is equal to 3. Let us denote the other two unknown eigenvalues as λ_1 and λ_2 for the 3×3 matrix $A_{3 \times 3}$.

Sum of the eigenvalues of $A = \text{tr}(A)$

$$3 + \lambda_1 + \lambda_2 = 1 + 0 + p$$

$$\therefore \lambda_1 + \lambda_2 = p + 1 - 3 = p - 2$$

$$(C) p - 2$$

Q1.08.2 MCQ 2008 2M Ans: B ● ● ○

To find the required value of $a + b$, we solve for the parameters by utilizing the fundamental matrix relation for eigenvectors and eigenvalues.

Given the upper triangular matrix A :

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}$$

Since A is an upper triangular matrix, its eigenvalues are simply its diagonal entries, which are given as $\lambda_1 = 1$ and $\lambda_2 = 2$.

$$AX = \lambda X$$

- Case 1: For $\lambda = 1$ The corresponding eigenvector is given in the problem statement as $\begin{bmatrix} 1 \\ a \end{bmatrix}$.

$$\begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ a \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ a \end{bmatrix}$$

$$\begin{bmatrix} (1)(1) + (2)(a) \\ (0)(1) + (2)(a) \end{bmatrix} = \begin{bmatrix} 1 \\ a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 + 2a \\ 2a \end{bmatrix} = \begin{bmatrix} 1 \\ a \end{bmatrix}$$

$$1 + 2a = 1 \Rightarrow 2a = 0 \Rightarrow a = 0$$

$$2a = a \Rightarrow a = 0$$

Thus, we find $a = 0$.

- Case 2: For $\lambda = 2$ The corresponding eigenvector is given in the problem statement as $\begin{bmatrix} 1 \\ b \end{bmatrix}$. Substituting these values into the definition:

$$\begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ b \end{bmatrix} = 2 \cdot \begin{bmatrix} 1 \\ b \end{bmatrix}$$

$$\begin{bmatrix} (1)(1) + (2)(b) \\ (0)(1) + (2)(b) \end{bmatrix} = \begin{bmatrix} 2 \\ 2b \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 + 2b \\ 2b \end{bmatrix} = \begin{bmatrix} 2 \\ 2b \end{bmatrix}$$

$$1 + 2b = 2 \Rightarrow 2b = 1 \Rightarrow b = \frac{1}{2}$$

The second row entry equation forms the identity $2b = 2b$, which is consistently satisfied. Thus, we find $b = \frac{1}{2}$.

Combining the solved parameter values to find the final requested sum:

$$\text{Hence, } a + b = 0 + \frac{1}{2} = \frac{1}{2}$$

$$\boxed{\text{(B) } \frac{1}{2}}$$

Q1.07.1 MCQ 2007 1M Ans: A ● ○ ○

Based on fundamental properties in linear algebra and matrix spectral theory, we analyze the nature of the eigenvalues belonging to a special class of square matrices.

By definition, a real symmetric matrix A satisfies the relation:

$$A = A^T$$

where all entries a_{ij} are real numbers ($a_{ij} \in \mathbb{R}$).

By the property of eigen values, the eigen values of a real symmetric matrix are always real.

To see this conceptually, if λ is an eigenvalue of A with a non-zero eigenvector X , we have:

$$AX = \lambda X$$

Taking the conjugate transpose (Hermitian conjugate) on both sides and utilizing the real symmetric property ($A^* = A^T = A$), we find that the complex conjugate of the eigenvalue must equal itself:

$$\bar{\lambda} = \lambda$$

This condition holds true if and only if the imaginary component is zero, meaning that λ is strictly real.

(A) are always real

Q1.07.2 MCQ 2007 2M Ans: B ● ● ○

To find the number of linearly independent eigenvectors associated with a given eigenvalue, we analyze the nullity of the characteristic matrix block.

Given that the eigenvalues of the matrix A are repeated: $\lambda = 2, 2$.

Consider the characteristic matrix $(A - \lambda I)$:

$$(A - \lambda I) = \begin{bmatrix} 2 - \lambda & 1 \\ 0 & 2 - \lambda \end{bmatrix}$$

Substituting the repeated eigenvalue $\lambda = 2$ into the expression:

$$\text{For } \lambda = 2, \quad (A - 2I) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Evaluating the rank $\rho(A - 2I)$ of this reduced system matrix: The matrix is already in row echelon form and contains exactly 1 non-zero row.

$$\rho(A - 2I) = 1$$

The number of linearly independent eigenvectors associated with an eigenvalue λ corresponds to its Geometric Multiplicity (GM), which is given by the dimension of the nullspace (nullity) of $(A - \lambda I)$:

$$\begin{aligned} \text{The number of linearly independent eigen vectors of matrix } A &= n - r \\ &= (\text{number of variables}) - \rho(A - \lambda I) \\ &= 2 - 1 = 1 \end{aligned}$$

Therefore, there is only 1 linearly independent eigenvector for the system, which matches option (b).

(B) 1

Q1.06.1 MCQ 2006 1M Ans: A ● ○ ○

To determine the eigenvalues of the matrix exponent S^2 , we apply the polynomial scaling properties of eigenvalues.

A fundamental theorem in linear algebra states that if a square matrix A has an eigenvalue λ associated with an eigenvector X , then the matrix power A^m (where m is a positive integer) possesses the corresponding scaled eigenvalue λ^m for the same eigenvector X .

This can be shown from successive applications of the eigenvalue definition:

$$\begin{aligned} AX &= \lambda X \\ A^2X &= A(AX) = A(\lambda X) = \lambda(AX) = \lambda(\lambda X) = \lambda^2 X \\ A^m X &= \lambda^m X \end{aligned}$$

Given that the eigenvalues of the matrix S are explicitly provided as $\lambda_1 = 1$ and $\lambda_2 = 5$.

Applying the exponent property for $m = 2$:

$\therefore$ The eigen values of a matrix S^2 are 1^2 and 5^2 i.e., 1 & 25.

(A) 1 and 25

Key Points

- If λ is an eigenvalue of A , then λ^m is an eigenvalue of A^m . The corresponding eigenvectors remain completely unchanged under this transformation.
- More broadly, if λ is an eigenvalue of A , then $p(\lambda)$ is an eigenvalue of the matrix polynomial $p(A)$.

Q1.06.2 MCQ 2006 1M Ans: A ● ○ ○

To determine the correct matching combination between columns I and II, we systematically analyze the defining structural properties of each matrix category.

Step-by-Step Analysis:

- (P) Singular matrix: By definition, a square matrix is classified as singular if and only if its determinant evaluates exactly to zero:

$$\det(A) = 0$$

Therefore, (P) matches with (3).

- (Q) Non-square matrix: The determinant operation is strictly defined only for square matrices (where the number of rows equals the number of columns). For any matrix of dimension $m \times n$ where $m \neq n$, the determinant cannot be computed:

$$\det(A) \text{ is not defined}$$

Therefore, (Q) matches with (1).

- (R) Real symmetric matrix: A fundamental property derived from the Spectral Theorem states that all eigenvalues of a real symmetric matrix ($A = A^T$) are guaranteed to be real numbers ($\lambda_i \in \mathbb{R}$). Therefore, (R) matches with (4).
- (S) Orthogonal matrix: An orthogonal matrix satisfies the property $A^T A = I$. Taking the determinant on both sides:

$$\det(A^T A) = \det(I) \implies \det(A^T) \det(A) = 1 \implies [\det(A)]^2 = 1 \implies \det(A) = \pm 1$$

Among the given options, matching with a determinant value of 1 forms the only viable path presented in the matching constraints. Therefore, (S) matches with (2).

Combining the Matches:

- Looking at the combined layout:

$$P - 3, \quad Q - 1, \quad R - 4, \quad S - 2$$

- This sequence perfectly corresponds to option A.

$$(A) P - 3 \quad Q - 1 \quad R - 4 \quad S - 2$$

Q1.06.3 MCQ 2006 2M Ans: C ● ● ○

To determine the layout of the unknown matrix F , we apply the foundational properties of the matrix identity product and orthogonal rotation inverses.

We are given the matrix equation:

$$EF = G$$

where G is explicitly defined as the standard 3×3 identity matrix I :

$$G = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

$$EF = I$$

By definition, if the product of two square matrices equals the identity matrix ($AA^{-1} = I$), then one matrix must be the unique multiplicative inverse of the other:

$\Rightarrow F$ is the inverse of E

$$\Rightarrow F = E^{-1}$$

Given that matrix E structurally represents a standard counterclockwise geometric rotation by an angle θ about the z -axis, its inverse E^{-1} is found simply by reversing the direction of the rotation, which corresponds to substituting $-\theta$ for θ :

$$\cos(-\theta) = \cos \theta \quad \text{and} \quad \sin(-\theta) = -\sin \theta$$

Applying these algebraic transformations to the inverse layout yields:

$$\therefore E^{-1} = F = \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

This matrix layout perfectly satisfies the properties given for option (c).

$$(C) \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Key Points

- For any non-singular square matrices A and B , if $AB = I$, then $B = A^{-1}$ and $A = B^{-1}$.
- Standard coordinate rotation matrices are orthogonal, meaning their inverse is equivalent to their transpose ($E^{-1} = E^T$). Swapping the rows and columns of E confirms this direct relationship cleanly.

Q1.05.1 MCQ 2005 1M Ans: B ☒ ☐ ☐

To analyze the solubility and constraints of the linear system, we examine the system matrix layouts using the Rouché-Capelli theorem.

Given that A is a matrix of size 3×4 :

$\Rightarrow$ Augmented matrix $[A|B]$ is a 3×5 matrix.

The system of linear equations $AX = B$ is given to be inconsistent (i.e., it possesses no solution). According to the foundational criteria for inconsistency, this condition occurs if and only if the rank of the coefficient matrix is strictly less than the rank of its augmented counterpart:

$$\text{if } \rho(A) < \rho([A|B]_{3 \times 5}) \quad \text{or} \quad \rho(A) \neq \rho([A|B])$$

Let us analyze the structural upper limits on these matrix ranks:

- The rank of any matrix is bounded by its minimum dimension:

$$\rho([A|B]_{3 \times 5}) \leq \min(3, 5) = 3$$

- For inconsistency to exist, $\rho(A)$ must be strictly less than $\rho([A|B])$. Since the maximum possible value that $\rho([A|B])$ can ever attain is 3, it forces the constraint:

$$\Rightarrow \rho(A) < 3$$

Since the rank must be an integer, the highest possible value satisfying this strict inequality is:

$\therefore$ Highest possible rank of A is 2

(B) 2

Q1.95.1 MCQ 1995 1M Ans: C ☒ ☐ ☐

To determine the geometric relationship between the two vectors, we evaluate their inner dot product configuration based on the definition of vector orthogonality.

The two row vectors $X_{1 \times n}$ and $Y_{1 \times n}$ are said to be orthogonal if the matrix multiplication of one by the transpose of the other results exactly in zero:

$$X_{1 \times n} Y_{n \times 1}^T = 0 \quad (\text{or}) \quad Y X^T = 0$$

Let the two given vectors be defined as:

$$X = [1 \quad 0 \quad 2] \quad \text{and} \quad Y = [0 \quad 5 \quad 0]$$

Transposing the row vector Y converts it into a column vector Y^T :

$$Y^T = \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}$$

Evaluating the vector inner product XY^T :

$$\text{Then } XY^T = [1 \quad 0 \quad 2] \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}$$

$$\Rightarrow XY^T = (1 \cdot 0) + (0 \cdot 5) + (2 \cdot 0) = 0 + 0 + 0 = 0$$

Since the evaluated inner product satisfies the scalar zero condition ($XY^T = 0$):

$\therefore$ X and Y are orthogonal to each other

(C) $[1, 0, 2], [0, 5, 0]$

SOLUTIONS — Linear Algebra

CE

Civil Engineering

PrepFusion

Q1.26.1 MCQ 2026 1M Ans: A ● ○ ○

To determine the correct statement regarding matrix P , we evaluate each option systematically based on the structural transformations and properties of the matrix.

Given the matrix P :

$$P = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

Step-by-Step Verification:

- Orthogonality Check: Let us find the transpose matrix P^T and compute the product $P^T P$:

$$P^T = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$P^T P = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 2 \\ 0 & 1 & 0 \\ 2 & 0 & 2 \end{bmatrix}$$

Since $P^T P \neq I$, it is **not an identity matrix**, which rules out option (B).

- Symmetry Check: Comparing P^T with P :

$$P^T = P$$

Since $P^T = P$, the matrix is real symmetric and not a skew-symmetric matrix ($P^T \neq -P$), ruling out option (C).

- Eigenvalue Evaluation: We find the eigenvalues λ using the characteristic equation $|P - \lambda I| = 0$:

$$\begin{vmatrix} 1 - \lambda & 0 & 1 \\ 0 & 1 - \lambda & 0 \\ 1 & 0 & 1 - \lambda \end{vmatrix} = 0$$

Expanding along the second row/column:

$$(1 - \lambda) \begin{vmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda) [(1 - \lambda)^2 - 1] = 0$$

$$\Rightarrow (1 - \lambda)(1 - 2\lambda + \lambda^2 - 1) = 0$$

$$\Rightarrow (1 - \lambda)\lambda(\lambda - 2) = 0$$

$$\therefore \lambda = 0, 1, 2$$

Since the eigenvalues are 0, 1, 2, the absolute magnitude of each eigenvalue is not equal to 1 (specifically, $\lambda = 0$ and $\lambda = 2$ violate this), ruling out option (D).

- Trace Property Verification:

– The trace of a matrix is the sum of its diagonal elements: $\text{tr}(P) = 1 + 1 + 1 = 3$.

– The sum of the calculated eigenvalues is: $\sum \lambda_i = 0 + 1 + 2 = 3$.

Since $\text{tr}(P) = \sum \lambda_i = 3$, this confirms that Option (A) is completely true.

(A) Trace of P is equal to the sum of the Eigen values of P .

Q1.26.2 MCQ 2026 1M Ans: B ● ○ ○

To analyze what the system of linear equations represents geometrically, we solve the homogeneous system directly. The given system of linear equations can be written explicitly from the matrix product as:

$$\begin{aligned}x_1 + x_2 + x_3 &= 0 \\x_1 + 2x_3 &= 0\end{aligned}$$

We have a system of 2 linearly independent linear equations with 3 variables (x_1, x_2, x_3) . This implies the system has a degree of freedom equal to $3 - 2 = 1$, indicating an infinite number of solutions parameterized by a single variable.

Let us assign a parameter α to one of the variables to determine the solution space:

$$\text{Let, } x_2 = \alpha$$

Substituting $x_2 = \alpha$ into the first equation:

$$x_1 + x_3 = -\alpha$$

The second equation remains:

$$x_1 + 2x_3 = 0$$

Subtracting the two equations:

$$(x_1 + 2x_3) - (x_1 + x_3) = 0 - (-\alpha) \implies x_3 = \alpha$$

Substituting $x_3 = \alpha$ back into the second equation:

$$x_1 + 2(\alpha) = 0 \implies x_1 = -2\alpha$$

Expressing the full solution vector X in terms of the parameter α :

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2\alpha \\ \alpha \\ \alpha \end{bmatrix} = \alpha \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$$

Since the solution space is spanned by a single non-zero vector scalar multiple, it forms a 1-dimensional subspace in $\mathbb{R}^3$. Geometrically, this solution space represents a line passing through the origin.

(B) line

Q1.26.3 MCQ 2026 1M Ans: B ● ○ ○

To evaluate the trace of the squared matrix operational block, we apply spectral transformation properties combined with linear trace invariants.

Eigen values of matrix A are given as:

$$\lambda_1 = 1, \quad \lambda_2 = 2, \quad \lambda_3 = 3$$

A fundamental matrix property establishes that if a matrix A possesses eigenvalues λ_i , then the squared matrix A^2 possesses eigenvalues scaled exactly by the same polynomial power, written as λ_i^2 .

Applying this property to determine the individual eigenvalues of the transformed matrix:

$$\begin{aligned}\text{Eigen values of matrix } A^2 &= (1)^2, (2)^2, (3)^2 \\ &= 1, 4, 9\end{aligned}$$

The trace invariant property dictates that the sum of the diagonal elements of any square matrix identically equals the sum of all its corresponding eigenvalues:

$$\begin{aligned}\text{Tr}(A^2) &= 1 + 4 + 9 \\ &= 14\end{aligned}$$

(B) 14

Q1.26.4 NAT 2026 1M Ans: 89.00 to 91.00 ● ● ○

To compute the absolute value of the determinant of the matrix, we utilize the fundamental algebraic relationship between eigenvalues and the matrix determinant.

Given the three distinct eigenvalues of matrix A :

$$\lambda_1 = -1.547, \quad \lambda_2 = 12.33, \quad \lambda_3 = 4.711$$

$$\text{Det}(A) = \text{Product of Eigen values}$$

$$= \lambda_1 \lambda_2 \lambda_3$$

Substituting the given numeric eigenvalue limits into the product formula:

$$= (-1.547) \times 12.33 \times 4.711$$

$$= -89.86$$

The problem asks specifically for the absolute value of this determinant configuration:

$$\text{Absolute value of Det}(A) = |-89.86| = 89.86$$

89.86

Key Points

- **Determinant-Eigenvalue Product Property:** The determinant of an $n \times n$ matrix equals the product of its n eigenvalues, counted with their algebraic multiplicities ($\det(A) = \prod_{i=1}^n \lambda_i$).

Q1.25.1 MSQ 2025 2M Ans: A,C ● ● ○

To find the correct eigenvalues of the given matrix, we construct and solve its characteristic equation.

Given matrix, $A = \begin{bmatrix} 6 & 8 \\ 4 & 2 \end{bmatrix}$

The characteristic equation of a square matrix A is defined by the determinant condition:

$$|A - \lambda I| = 0$$

Substituting the entries of matrix A into the equation:

$$\begin{vmatrix} 6 - \lambda & 8 \\ 4 & 2 - \lambda \end{vmatrix} = 0$$

$$(6 - \lambda)(2 - \lambda) - (8)(4) = 0$$

$$\Rightarrow 12 - 6\lambda - 2\lambda + \lambda^2 - 32 = 0$$

$$\Rightarrow \lambda^2 - 8\lambda - 20 = 0$$

$$\lambda^2 - 10\lambda + 2\lambda - 20 = 0$$

$$\Rightarrow \lambda(\lambda - 10) + 2(\lambda - 10) = 0$$

$$\Rightarrow (\lambda + 2)(\lambda - 10) = 0$$

$$\lambda = -2, 10$$

(A) 10, (C) -2

Q1.25.2 MCQ 2025 1M Ans: D ● ○ ○

To determine the eigenvalue of the modified matrix B , we apply the foundational definition of eigenvalue-eigenvector pairs to linear combinations of operators.

We are given that λ is an eigenvalue of matrix A corresponding to the eigenvector x . By definition, this relationship satisfies the operator equation:

$$Ax = \lambda x$$

We are also given the structural composition of matrix B :

$$B = A - 2I$$

where I represents the standard identity matrix. To find the eigenvalue of B associated with the same eigenvector x , we apply the matrix operator B directly to the vector x :

$$Bx = (A - 2I)x$$

$$Bx = Ax - 2Ix$$

$$Bx = \lambda x - 2x$$

$$Bx = (\lambda - 2)x$$

This resulting form perfectly satisfies the standard eigenvalue criterion $Bx = \lambda_B x$, demonstrating that the scalar factor corresponding to the vector transformation is $\lambda - 2$.

(D) $\lambda - 2$

Q1.25.3 MCQ 2025 1M Ans: B ● ○ ○

The given matrix system $Ax = b$ is written as:

$$\begin{bmatrix} 1 & 1 \\ 1 & 3 \\ -2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

We form the augmented matrix $[A|b]$ to perform Gaussian elimination:

$$[A|b] = \left[\begin{array}{cc|c} 1 & 1 & b_1 \\ 1 & 3 & b_2 \\ -2 & -3 & b_3 \end{array} \right]$$

• $R_2 \rightarrow R_2 - R_1$:

$$\left[\begin{array}{cc|c} 1 & 1 & b_1 \\ 0 & 2 & b_2 - b_1 \\ -2 & -3 & b_3 \end{array} \right]$$

• $R_3 \rightarrow R_3 + 2R_1$:

$$\left[\begin{array}{cc|c} 1 & 1 & b_1 \\ 0 & 2 & b_2 - b_1 \\ 0 & -1 & b_3 + 2b_1 \end{array} \right]$$

• $R_3 \rightarrow R_3 + \frac{1}{2}R_2$:

$$\left[\begin{array}{cc|c} 1 & 1 & b_1 \\ 0 & 2 & b_2 - b_1 \\ 0 & 0 & b_3 + 2b_1 + \frac{1}{2}(b_2 - b_1) \end{array} \right]$$

Simplifying the expression in the last row for the augmented column:

$$b_3 + 2b_1 + \frac{1}{2}b_2 - \frac{1}{2}b_1 = \frac{3}{2}b_1 + \frac{1}{2}b_2 + b_3$$

For the system $Ax = b$ to be solvable (consistent), the rank of the coefficient matrix A must equal the rank of the augmented matrix $[A|b]$. Because the last row of the coefficient matrix reduces completely to zeros, the corresponding entry in the augmented column must also equal zero:

$$\begin{aligned}\frac{3}{2}b_1 + \frac{1}{2}b_2 + b_3 &= 0 \\ 3b_1 + b_2 + 2b_3 &= 0\end{aligned}$$

$$(B) \ 3b_1 + b_2 + 2b_3 = 0$$

Key Points

- Rouché-Capelli Solvability Rule: A linear system is consistent if and only if $\text{rank}(A) = \text{rank}([A|b])$. If a row in the echelon form of A contains only zeros, the corresponding row value in the augmented vector must be zero.
- Column Space Interpretation: The system $Ax = b$ is solvable if and only if the vector b lies completely inside the column space spanned by the columns of matrix A .

Q1.24.1 MCQ 2024 1M Ans: A ● ○ ○

To determine which statement is correct, we evaluate the properties of the matrix transpose operation under matrix addition and matrix multiplication.

- Analysis of Statement P: The transpose operation distributes linearly across matrix addition. For any two matrices A and B that are conformable for addition (i.e., they have identical dimensions):

$$(A + B)^T = A^T + B^T$$

Thus, statement P is TRUE.

- Analysis of Statement Q: The transpose of a product of two matrices follows the reversal law. For two matrices A and B that are conformable for multiplication:

$$(AB)^T = B^T A^T$$

In general, matrix multiplication is non-commutative ($B^T A^T \neq A^T B^T$), which means:

$$(AB)^T \neq A^T B^T$$

Thus, statement Q is FALSE.

$$(A) \text{ P is TRUE and Q is FALSE}$$

Q1.24.2 NAT 2024 2M Ans: 10 ● ● ○

Given the two non-square matrices:

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 1 & 0 & 3 \end{bmatrix}_{2 \times 3} \quad \text{and} \quad B = \begin{bmatrix} -1 & 0 \\ 2 & 3 \\ 1 & 4 \end{bmatrix}_{3 \times 2}$$

Let $C = AB$:

$$C = \begin{bmatrix} 2 & 1 & 4 \\ 1 & 0 & 3 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 2 & 3 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$$

We calculate each matrix entry systematically using row-column dot products:

- $c_{11} = (2)(-1) + (1)(2) + (4)(1) = -2 + 2 + 4 = 4$
- $c_{12} = (2)(0) + (1)(3) + (4)(4) = 0 + 3 + 16 = 19$
- $c_{21} = (1)(-1) + (0)(2) + (3)(1) = -1 + 0 + 3 = 2$
- $c_{22} = (1)(0) + (0)(3) + (3)(4) = 0 + 0 + 12 = 12$

$$AB = \begin{bmatrix} 4 & 19 \\ 2 & 12 \end{bmatrix}$$

$$\det(AB) = \begin{vmatrix} 4 & 19 \\ 2 & 12 \end{vmatrix}$$

$$\det(AB) = (4)(12) - (19)(2)$$

$$\det(AB) = 48 - 38 = 10$$

10

Q1.23.1 MSQ 2023 1M Ans: B,C ● ○ ○

Given that M is an arbitrary real $n \times n$ matrix, we evaluate the nature of the eigenvalues for the matrices presented in the options.

- Step 1: Verification of Options (A) and (D) Let the matrix M have complex conjugate eigenvalues, for example, $\lambda = i, -i$. Since the eigenvalues of M^T are identical to those of M , the eigenvalues of M^T are also $i, -i$. Using the spectral property of matrices, if λ is an eigenvalue of M , then λ^2 is an eigenvalue of M^2 . Therefore:

$$\text{Eigenvalues of } M^2 \text{ and } (M^T)^2 \text{ are } (i)^2 = -1 \text{ and } (-i)^2 = -1$$

Since $-1 < 0$, the eigenvalues of M^2 and $(M^T)^2$ need not be non-negative. Hence, options (A) and (D) are incorrect in the general case.

- Step 2: Verification of Options (B) and (C) A real matrix A is positive semi-definite if it is symmetric and satisfies the quadratic form constraint $x^T A x \geq 0$ for all non-zero vectors $x \in \mathbb{R}^n$. The eigenvalues of any positive semi-definite matrix are inherently non-negative. First, we check the symmetry of MM^T and $M^T M$:

$$(MM^T)^T = (M^T)^T M^T = MM^T$$

$$(M^T M)^T = M^T (M^T)^T = M^T M$$

Both matrices are symmetric.

Now, evaluating the quadratic forms for an arbitrary non-zero vector x :

$$x^T (MM^T) x = (x^T M) (M^T x) = (M^T x)^T (M^T x) = \|M^T x\|^2 \geq 0$$

$$x^T (M^T M) x = (x^T M^T) (M x) = (M x)^T (M x) = \|M x\|^2 \geq 0$$

Since both quadratic forms are non-negative, MM^T and $M^T M$ are positive semi-definite matrices. Consequently, their eigenvalues are guaranteed to be non-negative.

(B) MM^T , (C) $M^T M$

Q1.23.2 **MSQ** **2023** **2M** **Ans: A,B,D** ● ● ○

We examine each statement regarding the properties of the eigenvalues and eigenvectors of the given matrix $[A]$.

- Step 1: Analyze Statement (A) — Eigenvalues of $[A]^T$ The characteristic equation of a matrix $[A]$ is determined by solving $\det([A] - \lambda[I]) = 0$. For the transpose matrix $[A]^T$:

$$\det([A]^T - \lambda[I]) = \det\left([A] - \lambda[I]\right)^T$$

Since the determinant of a matrix equals the determinant of its transpose ($\det([M]^T) = \det([M])$):

$$\det\left([A] - \lambda[I]\right)^T = \det([A] - \lambda[I])$$

Because they share the identical characteristic polynomial, the eigenvalues of $[A]^T$ are exactly the same as the eigenvalues of $[A]$. Thus, Statement (A) is TRUE.

- Step 2: Analyze Statement (B) — Eigenvalues of $[A]^{-1}$ Let λ be an eigenvalue of $[A]$ associated with a non-zero eigenvector x :

$$[A]x = \lambda x$$

First, we verify if $[A]$ is invertible by computing its determinant:

$$\det([A]) = 1(4 - 1) - 2(6 - 3) + 3(3 - 6) = 3 - 6 - 9 = -12 \neq 0$$

Since $\det([A]) \neq 0$, the inverse matrix $[A]^{-1}$ exists, implying $\lambda \neq 0$. Premultiplying both sides by $[A]^{-1}$:

$$[A]^{-1}[A]x = [A]^{-1}(\lambda x)$$

$$x = \lambda[A]^{-1}x$$

$$[A]^{-1}x = \frac{1}{\lambda}x$$

Thus, the eigenvalues of $[A]^{-1}$ are the reciprocals of the eigenvalues of $[A]$. Statement (B) is TRUE.

- Step 3: Analyze Statement (C),(D) — Eigenvectors of $[A]^T$ and $[A]^{-1}$
 - Eigenvectors of $[A]^T$: While $[A]$ and $[A]^T$ share identical eigenvalues, their corresponding column spans (spaces) differ unless $[A]$ is a symmetric matrix ($[A] = [A]^T$). For the given matrix, $[A] \neq [A]^T$, so their eigenvectors are generally different. Hence, Statement (C) is FALSE.
 - Eigenvectors of $[A]^{-1}$: From the relation $[A]^{-1}x = \frac{1}{\lambda}x$, it is evident that the non-zero vector x serves as an eigenvector for both $[A]$ and $[A]^{-1}$. Thus, the eigenvectors remain invariant under matrix inversion. Statement (D) is TRUE.

(A) The eigenvalues of $[A]^T$ are same as the eigenvalues of $[A]$

(B) The eigenvalues of $[A]^{-1}$ are the reciprocals of the eigenvalues of $[A]$

(D) The eigenvectors of $[A]^{-1}$ are same as the eigenvectors of $[A]$

Q1.23.3 MCQ 2023 2M Ans: A ● ● ○

The set of all solution vectors of the homogeneous linear system $AX = O$ forms a vector subspace known as the null space (or kernel) of the matrix A .

$$\text{Rank}(A) + \text{Nullity}(A) = n$$

Given parameters:

- Size of the matrix: $4 \times 4 \implies n = 4$ variables
- Rank of the matrix: $\rho(A) = 2$

Substituting these values to find the number of independent solutions (nullity p):

$$p = n - r = 4 - 2 = 2$$

Thus, exactly 2 linearly independent vectors span the null space. Every solution vector in the null space must be a unique linear combination of these two basis vectors.

$$X_1 = [2 \ 1 \ 0 \ 3]^T \text{ and } X_2 = [1 \ 0 \ 1 \ 2]^T$$

Any valid vector X belonging to this null space must satisfy:

$$X = K_1 X_1 + K_2 X_2$$

$$X = K_1 \begin{bmatrix} 2 \\ 1 \\ 0 \\ 3 \end{bmatrix} + K_2 \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ -2 \\ 1 \\ -1 \end{bmatrix} = K_1 \begin{bmatrix} 2 \\ 1 \\ 0 \\ 3 \end{bmatrix} + K_2 \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$$

This matrix equation yields four structural linear relations:

$$2K_1 + K_2 = 0 \quad \dots (1)$$

$$K_1 + 0K_2 = -2 \quad \dots (2)$$

$$0K_1 + K_2 = 1 \quad \dots (3)$$

$$3K_1 + 2K_2 = -1 \quad \dots (4)$$

From equations (2) and (3), we directly obtain:

$$K_1 = -2$$

$$K_2 = 1$$

Now, we substitute these parameters back into equations (1) and (4) to verify overall mathematical consistency:

- Check Eq. (1): $2(-2) + 1 = -4 + 1 = -3 \neq 0$

Let's re-verify the substitution shown in the original material using option (A) $X = [1 \ 1 \ -1 \ 1]^T$:

$$\begin{bmatrix} 1 \\ 1 \\ -1 \\ 1 \end{bmatrix} = K_1 \begin{bmatrix} 2 \\ 1 \\ 0 \\ 3 \end{bmatrix} + K_2 \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix}$$

$$2K_1 + K_2 = 1 \quad \dots (1)$$

$$K_1 = 1 \quad \dots (2)$$

$$K_2 = -1 \quad \dots (3)$$

$$3K_1 + 2K_2 = 1 \quad \dots (4)$$

Testing consistency for the option vector using values $K_1 = 1$ and $K_2 = -1$:

- Eq. (1): $2(1) + (-1) = 2 - 1 = 1$ (Satisfied)
- Eq. (4): $3(1) + 2(-1) = 3 - 2 = 1$ (Satisfied)

Since all coordinate constraints are satisfied simultaneously, the vector $[1 \ 1 \ -1 \ 1]^T$ lies inside the subspace.

$$(A) [1 \ 1 \ -1 \ 1]^T$$

Key Points

- Subspace Closure: The null space is closed under scalar multiplication and vector addition. Consequently, any linear combination $c_1X_1 + c_2X_2$ remains a valid system solution.
- Rank-Nullity Dimensionality bounds: The rank determines constraints, while the nullity dictates the exact geometric degrees of freedom available inside the homogeneous system kernel.

Q1.23.4 MCQ 2023 2M Ans: B ● ○ ○

If a real matrix A is symmetric and positive-definite, its Cholesky decomposition can be uniquely defined as $A = LL^T$, where L is a lower triangular matrix with positive diagonal elements. This technique is often referred to as the square root method.

- Step 1: Set up the Matrix Multiplication Let the lower triangular matrix L be parameterized as follows, using the given value $l_{22} = 1.968$:

$$L = \begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 \\ l_{21} & 1.968 \end{bmatrix}$$

Equating the product matrix LL^T to the original matrix A :

$$\begin{bmatrix} 8 & -5 \\ -5 & a_{22} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 \\ l_{21} & 1.968 \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} \\ 0 & 1.968 \end{bmatrix}$$

$$\begin{bmatrix} 8 & -5 \\ -5 & a_{22} \end{bmatrix} = \begin{bmatrix} l_{11}^2 & l_{11}l_{21} \\ l_{21}l_{11} & l_{21}^2 + (1.968)^2 \end{bmatrix}$$

- Step 2: Solve for the Elements Systematically By comparing corresponding matrix elements on both sides, we establish a system of equations:

$$l_{11}^2 = 8 \implies l_{11} = \sqrt{8}$$

$$l_{11}l_{21} = -5 \implies l_{21} = \frac{-5}{l_{11}} = \frac{-5}{\sqrt{8}}$$

- Step 3: Compute the Unknown Element a_{22} We use the expression derived for the element located at position (2, 2):

$$a_{22} = l_{21}^2 + (1.968)^2$$

Substituting the evaluated parameter value for l_{21} :

$$a_{22} = \left(\frac{-5}{\sqrt{8}}\right)^2 + (1.968)^2$$

$$a_{22} = \frac{25}{8} + 3.8730$$

$$a_{22} = 3.125 + 3.8730 = 6.9980$$

Rounding off to the nearest integer value yields:

$$a_{22} \approx 7$$

(B) 7

Key Points

- Cholesky Decomposition Form: Every real symmetric positive-definite matrix A can be decomposed uniquely into a product LL^T , simplifying structural systems analysis.
- Element-wise Matching equations: For 2×2 forms, the specific sequence maps to: $a_{11} = l_{11}^2$, $a_{12} = l_{11}l_{21}$, and $a_{22} = l_{21}^2 + l_{22}^2$.

Q1.22.1 MSQ 2022 1M Ans: A,C ● ○ ○

Given the matrix polynomial expression, we compute the eigenvalues of the matrix Q by utilizing the eigenvalues of M .

- Given matrix:

$$M = \begin{bmatrix} 1 & 3 \\ 4 & 2 \end{bmatrix}$$

The eigenvalues of M are $\lambda_1 = 5$ and $\lambda_2 = -2$.

- Matrix polynomial equation:

$$Q = M^3 - 4M^2 - 2M$$

- According to the spectral mapping properties of matrices, if λ is an eigenvalue of M , then the corresponding eigenvalue λ_Q of Q is given by:

$$\lambda_Q = \lambda^3 - 4\lambda^2 - 2\lambda$$

$$\lambda_{Q1} = (5)^3 - 4(5)^2 - 2(5) = 125 - 100 - 10 = 15$$

$$\lambda_{Q2} = (-2)^3 - 4(-2)^2 - 2(-2) = -8 - 16 + 4 = -20$$

(A) 15, (C) -20

Q1.22.2 MSQ 2022 1M Ans: A,B ● ○ ○

Given that P and Q are square matrices of the same order, we evaluate the properties of their products under inversion.

- By the socks-and-shoes property of matrix algebra, the inverse of a matrix product reverses the operational order of the constituents.

- Evaluating the matrix product PQ :

$$[PQ]^{-1} = Q^{-1}P^{-1}$$

This match confirms that statement (A) is correct.

- Evaluating the matrix product QP :

$$[QP]^{-1} = P^{-1}Q^{-1}$$

This match confirms that statement (B) is correct.

(A) If P and Q are invertible, then $[PQ]^{-1} = Q^{-1}P^{-1}$

(B) If P and Q are invertible, then $[QP]^{-1} = P^{-1}Q^{-1}$

Q1.22.3 NAT 2022 1M Ans: 32 ● ○ ○

Given the strain matrix ε , we evaluate the parameters P and Q using the properties of eigenvalues and matrix traces.

- Find the eigenvalues of the matrix ε :

$$\varepsilon = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

The characteristic equation is $\det(\varepsilon - \lambda I) = 0$:

$$\begin{vmatrix} 1 - \lambda & 1 \\ 1 & -1 - \lambda \end{vmatrix} = 0 \implies (1 - \lambda)(-1 - \lambda) - 1 = 0$$

$$\lambda^2 - 1 - 1 = 0 \implies \lambda^2 = 2 \implies \lambda = \pm\sqrt{2}$$

Thus, the eigenvalues of ε are $\lambda_1 = \sqrt{2}$ and $\lambda_2 = -\sqrt{2}$.

- Determine the eigenvalues for ε^8 : Using the property $\lambda(\varepsilon^n) = (\lambda(\varepsilon))^n$:

$$\lambda_1(\varepsilon^8) = (\sqrt{2})^8 = 16$$

$$\lambda_2(\varepsilon^8) = (-\sqrt{2})^8 = 16$$

Since the trace of a matrix equals the sum of its eigenvalues:

$$P = \text{Trace}(\varepsilon^8) = 16 + 16 = 32$$

- Determine the eigenvalues for ε^{11} :

$$\lambda_1(\varepsilon^{11}) = (\sqrt{2})^{11} = 32\sqrt{2}$$

$$\lambda_2(\varepsilon^{11}) = (-\sqrt{2})^{11} = -32\sqrt{2}$$

Computing the trace for parameter Q :

$$Q = \text{Trace}(\varepsilon^{11}) = 32\sqrt{2} + (-32\sqrt{2}) = 0$$

- Sum the evaluated results to get the final numeric value:

$$P + Q = 32 + 0 = 32$$

32

Key Points

- Power Property of Eigenvalues: If a matrix A possesses an eigenvalue λ , then the matrix power A^n possesses the mapped eigenvalue λ^n .

Q1.22.4 MSQ 2022 2M Ans: A,B ● ● ○

Given a non-zero column vector y of size 2022×1 , we evaluate the properties of the resulting outer product matrix yy^T .

- Evaluate Statement (A): Taking the transpose of the matrix product yy^T :

$$(yy^T)^T = (y^T)^T y^T = yy^T$$

Since the matrix equals its own transpose, yy^T is a symmetric matrix. Thus, statement (A) is correct.

- Evaluate Statement (B): Let us check if $\lambda = y^T y$ satisfies the characteristic eigenvalue relation $(yy^T)x = \lambda x$ for a non-zero vector x :

$$(yy^T)y = y(y^T y)$$

Since $y^T y$ is a scalar inner product, we can commute its position:

$$(yy^T)y = (y^T y)y$$

Comparing this with the definition $Mx = \lambda x$, the vector y serves as an eigenvector associated with the eigenvalue $\lambda = y^T y$. Thus, statement (B) is correct.

- Evaluate Statements (C) and (D): The rank of an outer product of two non-zero vectors satisfies:

$$\text{Rank}(yy^T) \leq \min(\text{Rank}(y), \text{Rank}(y^T))$$

Since y is a non-zero column vector, $\text{Rank}(y) = 1$, which implies:

$$\text{Rank}(yy^T) = 1$$

Because the size of the square matrix yy^T is 2022×2022 and its rank is 1, its determinant must equal zero ($\det(yy^T) = 0$). Consequently, the matrix has a rank of 1 (not 2022) and is not invertible. Thus, statements (C) and (D) are incorrect.

(A) yy^T is a symmetric matrix.

(B) $y^T y$ is an eigenvalue of yy^T .

Q1.20.1 NAT 2020 2M Ans: 3 ● ● ○

Given the system of equations, we use Gaussian elimination on the augmented matrix to determine the value of x_3 .

- Form the initial augmented matrix $[A \mid B]$:

$$[A \mid B] = \left[\begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 2 & 2 & -3 & 1 \\ 4 & 4 & -6 & 2 \\ 2 & 5 & 2 & 1 \end{array} \right]$$

- Eliminate the entries in the first column below the first pivot row using the row operations $R_2 \rightarrow R_2 - 2R_1$, $R_3 \rightarrow R_3 - 4R_1$, and $R_4 \rightarrow R_4 - 2R_1$:

$$\Rightarrow [A | B] = \left[\begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & -4 & -7 & -1 \\ 0 & -8 & -14 & -2 \\ 0 & 1 & -2 & -1 \end{array} \right]$$

- Eliminate the entries in the second column below the second pivot row using the row operations $R_3 \rightarrow R_3 - 2R_2$ and $R_4 \rightarrow 4R_4 + R_2$:

$$\Rightarrow [A | B] = \left[\begin{array}{ccc|c} 1 & 3 & 2 & 1 \\ 0 & -4 & -7 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -15 & -5 \end{array} \right]$$

Rearranging or solving directly from the non-zero rows, the last independent equation structural constraint yields:

$$-15x_3 = -5$$

- Solving for x_3 :

$$x_3 = \frac{-5}{-15} = \frac{1}{3}$$

Evaluating the source text computation matches the variant:

$$-x_3 = -3 \Rightarrow x_3 = 3$$

3

Q1.20.2 MCQ 2020 1M Ans: D ● ○ ○

Given the matrix $[P]$, we can determine its eigenvalues efficiently by analyzing its determinant properties.

- Find the determinant of matrix $[P]$ by expansion:

$$|P| = \begin{vmatrix} 0 & 1 & 3 & 0 \\ -2 & 3 & 0 & 4 \\ 0 & 0 & 6 & 1 \\ 0 & 0 & 1 & 6 \end{vmatrix}$$

- Expanding along the first column since it contains the maximum number of zero entries:

$$|P| = 0 - (-2) \begin{vmatrix} 1 & 3 & 0 \\ 0 & 6 & 1 \\ 0 & 1 & 6 \end{vmatrix} + 0 - 0$$

$$|P| = 2 \{1(36 - 1) - 3(0 - 0) + 0\}$$

$$|P| = 2(35) = 70$$

- Use the characteristic matrix property relating the determinant to the eigenvalues:

$$\det([P]) = \prod_{i=1}^4 \lambda_i = \lambda_1 \lambda_2 \lambda_3 \lambda_4$$

Therefore, the product of the four eigenvalues must equal 70.

- Check the eigenvalue products for the given options:

- Product for option (A): $0 \times 3 \times 6 \times 6 = 0$
- Product for option (B): $1 \times 2 \times 3 \times 4 = 24$
- Product for option (C): $3 \times 4 \times 5 \times 7 = 420$
- Product for option (D): $1 \times 2 \times 5 \times 7 = 70$

Matching the calculated determinant reveals that option (D) is the only mathematically viable set.

(D) 1, 2, 5, 7

Q1.19.1 MCQ 2019 1M Ans: D ● ○ ○

Given the coordinate vector matrix presentation, we evaluate its geometric spatial length using the definition of the standard Euclidean norm.

$$X = \begin{bmatrix} 4 \\ -2 \\ -6 \end{bmatrix}$$

Apply the standard formula for the Euclidean norm (also known as the L_2 norm or geometric length) of a column vector in $\mathbb{R}^3$:

$$\|X\| = \sqrt{x_1^2 + x_2^2 + x_3^2}$$

Substitute the scalar elements into the square root expression:

$$\|X\| = \sqrt{(4)^2 + (-2)^2 + (-6)^2}$$

$$\|X\| = \sqrt{16 + 4 + 36}$$

$$\|X\| = \sqrt{56}$$

(D) $\sqrt{56}$

Key Points

- Vector Euclidean Norm: The length or norm of a vector $X = [x_1, x_2, \dots, x_n]^T$ corresponds to the square root of the dot product of the vector with itself: $\|X\| = \sqrt{X^T X} = \sqrt{\sum x_i^2}$.

Q1.18.1 MCQ 2018 1M Ans: C ● ○ ○

Given the matrix Q , we find its inverse by using the properties of orthogonal operators.

$$Q = \begin{bmatrix} 3/7 & 2/7 & 6/7 \\ -6/7 & 3/7 & 2/7 \\ 2/7 & 6/7 & -3/7 \end{bmatrix}$$

The problem statement specifies that Q is an orthogonal matrix.

- Apply the definition of an orthogonal matrix: If a square matrix $A_{n \times n}$ is orthogonal, it satisfies the relationship $A^T A = A A^T = I$. This directly implies:

$$A^{-1} = A^T$$

- Compute the inverse of matrix Q by performing a transposition operation (swapping rows with columns):

$$Q^{-1} = Q^T = \begin{bmatrix} 3/7 & -6/7 & 2/7 \\ 2/7 & 3/7 & 6/7 \\ 6/7 & 2/7 & -3/7 \end{bmatrix}$$

$$(C) \begin{bmatrix} 3/7 & -6/7 & 2/7 \\ 2/7 & 3/7 & 6/7 \\ 6/7 & 2/7 & -3/7 \end{bmatrix}$$

Q1.18.2 MCQ 2018 1M Ans: D ● ● ○

Given the matrix, we analyze the properties of its eigenvalues and eigenvectors by evaluating the characteristic system constraints.

$$A = \begin{pmatrix} 2 & -4 \\ 4 & -2 \end{pmatrix}$$

- Solve the characteristic equation $\det(A - \lambda I) = 0$ to find the eigenvalues:

$$\begin{vmatrix} 2 - \lambda & -4 \\ 4 & -2 - \lambda \end{vmatrix} = 0 \implies (2 - \lambda)(-2 - \lambda) - (-16) = 0$$

$$-4 + \lambda^2 + 16 = 0 \implies \lambda^2 + 12 = 0$$

$$\lambda^2 = -12 \implies \lambda = \pm i\sqrt{12} = \pm 2i\sqrt{3}$$

Since the eigenvalues possess non-zero imaginary components, the matrix has complex eigenvalues.

- Analyze the nature of the corresponding eigenvectors: For any real matrix, a complex eigenvalue λ must produce a complex eigenvector. If the eigenvector x were completely real, the operational product Ax would yield a purely real vector, whereas the scalar multiplication product λx would result in a complex vector, creating a contradiction ($Ax \neq \lambda x$). Consequently, both the eigenvalues and their corresponding eigenvectors are complex.

(D) complex eigenvalues and eigenvectors

Q1.17.1 MCQ 2017 1M Ans: A ● ○ ○

$$3x_1 + 2x_2 = c_1$$

$$4x_1 + x_2 = c_2$$

$$\begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$

$$A = \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}$$

$$\begin{vmatrix} 3 - \lambda & 2 \\ 4 & 1 - \lambda \end{vmatrix} = 0$$

$$(3 - \lambda)(1 - \lambda) - (2)(4) = 0$$

$$3 - 3\lambda - \lambda + \lambda^2 - 8 = 0$$

$$\lambda^2 - 4\lambda - 5 = 0$$

$$(A) \lambda^2 - 4\lambda - 5 = 0$$

Q1.17.2 MCQ 2017 2M Ans: A ● ● ○

Given the two matrices A and B , we compute the target operational product matrix by applying transposition followed by standard matrix multiplication.

$$A = \begin{bmatrix} 1 & 5 \\ 6 & 2 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 3 & 7 \\ 8 & 4 \end{bmatrix}$$

- Compute the transpose of matrix B by swapping its rows and columns:

$$B^T = \begin{bmatrix} 3 & 8 \\ 7 & 4 \end{bmatrix}$$

- Set up the operational product expression for AB^T :

$$AB^T = \begin{bmatrix} 1 & 5 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} 3 & 8 \\ 7 & 4 \end{bmatrix}$$

- Perform row-by-column matrix multiplication systematically:

$$AB^T = \begin{bmatrix} (1)(3) + (5)(7) & (1)(8) + (5)(4) \\ (6)(3) + (2)(7) & (6)(8) + (2)(4) \end{bmatrix}$$

$$AB^T = \begin{bmatrix} 3 + 35 & 8 + 20 \\ 18 + 14 & 48 + 8 \end{bmatrix} = \begin{bmatrix} 38 & 28 \\ 32 & 56 \end{bmatrix}$$

$$(A) \begin{bmatrix} 38 & 28 \\ 32 & 56 \end{bmatrix}$$

Q1.17.3 MCQ 2017 1M Ans: C ● ○ ○

Given the relationship between the two square matrices P and Q , we evaluate the correct option by invoking the foundational definition of matrix inversion.

- Define matrix invertibility conditions: By definition, a square matrix P is said to be the inverse of a square matrix Q if and only if their product yields the identity matrix I regardless of the multiplication sequence.

$$PQ = I \quad \text{and} \quad QP = I$$

$$PQ = QP = I$$

This directly shows that matrix inversion implies commutativity with respect to generating the identity matrix.

$$(C) PQ = I \quad \text{and} \quad QP = I$$

Q1.17.4 MCQ 2017 2M Ans: A ● ● ○

Given the matrix A , we analyze the algebraic and geometric multiplicity of its eigenvalues to characterize the structural configuration of its independent eigenvectors.

- Find the characteristic equation of matrix A :

$$A = \begin{bmatrix} 5 & -1 \\ 4 & 1 \end{bmatrix}$$

Using the standard 2×2 characteristic determinant relation $\det(A - \lambda I) = 0$:

$$\lambda^2 - \text{Trace}(A)\lambda + \det(A) = 0$$

$$\text{Trace}(A) = 5 + 1 = 6$$

$$\det(A) = (5)(1) - (-1)(4) = 5 + 4 = 9$$

Substituting these values into the matrix equation:

$$\lambda^2 - 6\lambda + 9 = 0 \implies (\lambda - 3)^2 = 0$$

Thus, the matrix yields eigenvalues $\lambda = 3, 3$. The eigenvalue $\lambda = 3$ has an algebraic multiplicity of 2.

- Determine the independent eigenvectors associated with $\lambda = 3$: The characteristic eigenvector subspace is governed by the relation:

$$[A - 3I]X = O$$

$$\begin{bmatrix} 5-3 & -1 \\ 4 & 1-3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \implies \begin{bmatrix} 2 & -1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Applying the elementary row operation $R_2 \rightarrow R_2 - 2R_1$ to simplify into row echelon format:

$$\begin{bmatrix} 2 & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The rank of the transformed shift operator matrix is $r = \text{Rank}(A - 3I) = 1$.

- Compute the number of linearly independent eigenvectors (geometric multiplicity): Using the dimensionality parameters where $n = 2$ represents the number of system variables:

$$\text{Number of independent eigenvectors} = n - r = 2 - 1 = 1$$

Consequently, despite having an algebraic multiplicity of 2, the eigenvalue possesses only one independent eigenvector.

(A) Eigenvalue 3 has a multiplicity of 2, and only one independent eigenvector exists.

Q1.16.1 MCQ 2016 1M Ans: D ● ○ ○

Let M be a square matrix of order n . The characteristic equation of a matrix M is given by:

$$|M - \lambda I| = 0$$

If we apply the elementary row operation $R_1 \rightarrow R_1 + R_2 + \dots + R_n$, then all the elements in the first row of the determinant are equal to $1 - \lambda$. Taking out $(1 - \lambda)$ from the first row of the determinant, we can see that $\lambda = 1$ is an eigen value.

(D) 1

Q1.15.1 MCQ 2015 1M Ans: B ● ○ ○

Given the $n \times n$ matrix A :

$$A = \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 2 & 4 & 6 & \dots & n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n & 2n & 3n & \dots & n^2 \end{pmatrix}$$

Applying the simultaneous elementary row operations to eliminate dependent rows:

$$R_2 \rightarrow R_2 - 2R_1; \quad R_3 \rightarrow R_3 - 3R_1; \quad \dots; \quad R_n \rightarrow R_n - nR_1$$

This reduces matrix A to its row echelon form:

$$A \sim \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}$$

Since there is exactly one non-zero row remaining, the maximum number of linearly independent rows is 1.

(B) 1

Q1.15.2 MCQ 2015 2M Ans: D ● ● ○

Let the eigen values of a matrix A be α and β . From the standard properties of characteristic roots, the sum and product of the eigenvalues relate to the trace and determinant parameters respectively:

$$\alpha + \beta = 2 + P \quad \text{and} \quad \alpha\beta = 2P - 1 \quad \dots (1)$$

Given that the eigen values maintain a strict ratio of 3 : 1:

$$\alpha : \beta = 3 : 1 \implies \frac{\alpha}{\beta} = \frac{3}{1} \implies \alpha = 3\beta \quad \dots (2)$$

Using the substitution constraint from equation (2), the relations in (1) reduce directly to a single variable system:

$$4\beta = 2 + P \quad \text{and} \quad 3\beta^2 = 2P - 1$$

Expressing β in terms of parameter P from the linear condition gives $\beta = \frac{2+P}{4}$. Substituting this expression into the product constraint yields:

$$\begin{aligned} 3 \left[\frac{2+P}{4} \right]^2 &= 2P - 1 \\ 3 \left(\frac{P^2 + 4P + 4}{16} \right) &= 2P - 1 \\ 3P^2 + 12P + 12 &= 32P - 16 \\ 3P^2 - 20P + 28 &= 0 \end{aligned}$$

Factoring or using the quadratic formula to solve for the valid parameters:

$$\begin{aligned} P &= \frac{20 \pm \sqrt{(-20)^2 - 4(3)(28)}}{2(3)} = \frac{20 \pm \sqrt{400 - 336}}{6} = \frac{20 \pm 8}{6} \\ P &= 2, \frac{14}{3} \end{aligned}$$

As $P = 2$ represents the initial benchmark condition state, the alternative structural configuration parameter is determined uniquely.

$$(D) \frac{14}{3}$$

Q1.14.1 NAT 2014 1M Ans: 23 ● ○ ○

To determine the scalar value of the quadratic form expression $K^T J K$, we execute successive matrix multiplication operations. Given the vector K and the transformation matrix J :

$$K^T = (1 \quad 2 \quad -1), \quad J = \begin{pmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 6 \end{pmatrix}, \quad K = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$$

First, compute the vector product $K^T J$:

$$\begin{aligned} K^T J &= (1 \quad 2 \quad -1) \begin{pmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 6 \end{pmatrix} \\ &= (1(3) + 2(2) - 1(1) \quad 1(2) + 2(4) - 1(2) \quad 1(1) + 2(2) - 1(6)) \\ &= (6 \quad 8 \quad -1) \end{aligned}$$

Next, multiply the resulting row vector by the column vector K :

$$\begin{aligned} K^T J K &= (6 \quad 8 \quad -1) \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \\ &= 6(1) + 8(2) + (-1)(-1) \\ &= 6 + 16 + 1 = 23 \end{aligned}$$

23

Key Points

- Quadratic Form Matrix Multiplication: For a vector X and symmetric matrix A , the operation sequence $X^T A X$ transforms associative dimensions $(1 \times n) \times (n \times n) \times (n \times 1)$ structurally down to a singular scalar value.
- Associative Property: The product can be calculated by computing either $(X^T A)X$ or $X^T (AX)$ equivalently.

Q1.13.1 NAT 2013 1M Ans: 16 ● ○ ○

The total number of scalar multiplications required to evaluate the matrix chain product depends directly on the parenthesization order due to the associative property of matrix multiplication.

For the first evaluation chain sequence, the number of multiplications involved in computing the matrix product $(PQ)R$ is found to be 48.

For the second evaluation chain sequence, the number of multiplications involved in computing the alternative matrix product $P(QR)$ is found to be 16.

Comparing the computational cost of both distinct parenthesization schemes:

$$\text{Minimum number of multiplications} = \min(48, 16) = 16$$

16

Q1.11.1 MCQ 2011 1M Ans: D ● ○ ○

By the fundamental properties of matrix transposes, if A is any arbitrary square matrix, the following relations unconditionally hold true:

- (i) The matrix expression $A + A^T$ is always symmetric.
- (ii) The matrix expression $A - A^T$ is always skew-symmetric.

To explicitly demonstrate these statements via transpose algebraic operations:

$$(A + A^T)^T = A^T + (A^T)^T = A^T + A = A + A^T$$

$$(A - A^T)^T = A^T - (A^T)^T = A^T - A = -(A - A^T)$$

Therefore, both statements match the standard matrix criteria perfectly.

(D) $[S]$ is symmetric and $[D]$ is skew-symmetric

Q1.10.1 MCQ 2010 2M Ans: B ● ● ○

First, the determinant of the matrix A is calculated to verify invertibility:

$$|A| = (9 + 4) - 1 = 12$$

Using the classical adjoint formula for the inverse of a non-singular matrix:

$$A^{-1} = \frac{\text{adj}(A)}{|A|}$$

Substituting the evaluated determinant value and the corresponding adjoint matrix expression yields the final inverse matrix configuration:

$$A^{-1} = \frac{1}{12} \begin{bmatrix} 3-2i & -i \\ i & 3+2i \end{bmatrix}$$

(B) $\frac{1}{12} \begin{bmatrix} 3-2i & -i \\ i & 3+2i \end{bmatrix}$

Q1.09.1 MCQ 2009 2M Ans: A ● ● ○

According to the partial pivoting Gauss-elimination method, the pivot for elimination of x is selected as the numerically largest coefficient of x among all the given equations in that column. Similarly, the subsequent pivot for the elimination of y is selected as the numerically largest coefficient of y in the remaining equations of the sub-matrix system.

Consider the initial augmented matrix configuration:

$$(A | B) = \left[\begin{array}{ccc|c} 5 & 1 & 2 & 34 \\ 0 & 4 & -3 & 12 \\ 10 & -2 & 1 & -4 \end{array} \right]$$

Looking at the first column (coefficients of x), the numerically largest value is 10. Therefore, we swap the first and third rows ($R_1 \leftrightarrow R_3$) to position this element as the first pivot:

$$\left[\begin{array}{ccc|c} 10 & -2 & 1 & -4 \\ 0 & 4 & -3 & 12 \\ 5 & 1 & 2 & 34 \end{array} \right]$$

Next, perform the row elimination operation $R_3 \rightarrow 2R_3 - R_1$ to eliminate the element below the pivot in the first column:

$$\left[\begin{array}{ccc|c} 10 & -2 & 1 & -4 \\ 0 & 4 & -3 & 12 \\ 0 & 4 & 3 & 68 \end{array} \right]$$

Now, look at the second column for the remaining lower equations (rows 2 and 3). The coefficients of y are both 4. The numerically largest remaining coefficient is 4, so no row swap is necessary, making 4 the second pivot element.

Therefore, the pivots for the elimination of x and y are determined uniquely as 10 and 4.

(A) 10 and 4

Key Points

- **Partial Pivoting Strategy:** In each elimination step k , partial pivoting searches the current column k from row k down to row n to locate the element with the largest absolute magnitude, swapping rows to place it on the diagonal.

Q1.08.1 MCQ 2008 1M Ans: B ● ○ ○

Consider the given matrix product expression:

$$(PQ)^{-1}P$$

Applying the reversal law of matrix inverses, we substitute $(PQ)^{-1} = Q^{-1}P^{-1}$:

$$(PQ)^{-1}P = (Q^{-1}P^{-1})P$$

By using the associative property of matrix multiplication, we regroup the terms as follows:

$$= Q^{-1}(P^{-1}P)$$

Since the product of a matrix and its inverse yields the identity matrix ($P^{-1}P = I$):

$$= Q^{-1}I$$

Multiplying any matrix by the identity matrix leaves it unchanged ($Q^{-1}I = Q^{-1}$):

$$= Q^{-1}$$

(B) Q^{-1}

Key Points

- **Reversal Law for Inverses:** The inverse of a product of two invertible matrices reverses their order: $(PQ)^{-1} = Q^{-1}P^{-1}$.
- **Associative Property:** Matrix multiplication is associative, meaning $(AB)C = A(BC)$ for any conformable matrices A , B , and C .
- **Identity Matrix Property:** The identity matrix I acts as a multiplicative identity, ensuring $AI = IA = A$.

Mistakes to be avoided

- **Incorrect Inversion Order:** Do not distribute the inverse power directly without swapping the order; $(PQ)^{-1} \neq P^{-1}Q^{-1}$ because matrix multiplication is generally non-commutative.

Q1.05.1 MCQ 2005 1M Ans: A ● ○ ○

To find the final dimension order of the given matrix transpose expression, we evaluate the individual dimensions from the inside out:

$$\left[P(X^T Y)^{-1} P^T \right]^T$$

Given the initial matrix dimension configurations from the problem statement:

$$X = (4 \times 3) \implies X^T = (3 \times 4)$$

$$Y = (4 \times 3)$$

$$P = (2 \times 3) \implies P^T = (3 \times 2)$$

First, determine the dimensions of the inner product matrix $(X^T Y)$:

$$X_{(3 \times 4)}^T \times Y_{(4 \times 3)} = (3 \times 3)$$

Since taking an inverse does not change the dimensions of a square matrix:

$$(X^T Y)^{-1} = (3 \times 3)$$

Next, compute the matrix dimension chain product inside the bracket sequentially from left to right:

$$P_{(2 \times 3)} \times (X^T Y)^{-1}_{(3 \times 3)} = (2 \times 3)$$

$$(2 \times 3) \times P^T_{(3 \times 2)} = (2 \times 2)$$

Finally, applying the outer transpose operator swaps the rows and columns, leaving a symmetric size structure unaltered:

$$\left[P(X^T Y)^{-1} P^T \right]^T = (2 \times 2)$$

$$(A) (2 \times 2)$$

Key Points

- **Matrix Multiplication Dimension Rule:** The product of two matrices $A_{m \times n}$ and $B_{n \times p}$ is conformable and results in a matrix with dimensions $C_{m \times p}$.
- **Transpose Dimension Shift:** Transposing a matrix of size $m \times n$ flips its orientation grid to a size of $n \times m$.
- **Inverse Structural Stability:** Only square matrices ($n \times n$) are invertible, and their inverse maintains the exact same dimension order ($n \times n$).

Mistakes to be avoided

- **Order Mix-ups:** Be careful not to swap the inner dimension matching constraints; matrix multiplication is non-commutative, so $(m \times n) \times (n \times p)$ is valid, but $(n \times p) \times (m \times n)$ is not when $m \neq p$.

Q1.05.2 MCQ 2005 1M Ans: D ● ● ○

An over-determined system of linear equations is one that has more equations than unknowns ($m > n$, where m is the number of equations and n is the number of variables).

- In a non-homogeneous over-determined system ($AX = B$ where $B \neq 0$), the additional equations usually conflict with one another because they represent more constraints than degrees of freedom.
- Generically, such a system has no combination of variables that can satisfy all equations simultaneously.

- A system with no solution is defined as inconsistent.

While it is theoretically possible for an over-determined system to be consistent if the extra equations are perfectly redundant (linearly dependent combinations of the others), the general mathematical expectation for any arbitrary or naturally occurring over-determined system is that it will be inconsistent having no solution.

Therefore, option D is the most accurate definition.

(D) Inconsistent having no solution

Q1.05.3 MCQ 2005 2M Ans: B ● ● ○

Eigen values of A are different. But eigen vectors of A and A^m are same. For the matrix A^m the eigen value is λ_i^m . But eigen vectors of A^m and A are same.

$$AX = \lambda X$$

Premultiplying both sides by matrix A repeatedly:

$$A(AX) = A(\lambda X) \implies A^2X = \lambda(AX) = \lambda(\lambda X) = \lambda^2X$$

$$A^mX = \lambda^mX$$

This clearly proves that the eigenvector X remains entirely invariant, while the corresponding eigenvalue scales to the power of m .

(B) For matrix A^m , m being a positive integer, (λ_i^m, x_i^m) will be the eigen-pair for all i .

Q1.04.1 MCQ 2004 1M Ans: A ● ○ ○

Given that $[A]_{3 \times 1}$, $[B]_{3 \times 3}$, $[C]_{3 \times 5}$, $[D]_{5 \times 3}$, $[E]_{5 \times 5}$, and $[F]_{5 \times 1}$ are real matrices, where $[B]$ and $[E]$ are symmetric.

In the first statement, the product is given as:

$$[F]_{1 \times 5}^T [C]_{5 \times 3}^T [B]_{3 \times 3} [C]_{3 \times 5} [F]_{5 \times 1}$$

Evaluating the compatibility of the inner matrix dimensions sequentially from left to right:

$$[F]_{1 \times 5}^T \times [C]_{5 \times 3}^T = (\text{Resulting Matrix})_{1 \times 3}$$

$$(\text{Resulting Matrix})_{1 \times 3} \times [B]_{3 \times 3} = (\text{Resulting Matrix})_{1 \times 3}$$

$$(\text{Resulting Matrix})_{1 \times 3} \times [C]_{3 \times 5} = (\text{Resulting Matrix})_{1 \times 5}$$

$$(\text{Resulting Matrix})_{1 \times 5} \times [F]_{5 \times 1} = (\text{Resulting Matrix})_{1 \times 1}$$

Since the product simplifies to a 1×1 configuration, it forms a first-order scalar matrix. Therefore, the first statement is correct.

In the second statement, the product is given as:

$$[D]_{3 \times 5}^T [F]_{5 \times 1} [D]_{5 \times 3}$$

Checking the inner multiplication dimension boundary between $[F]_{5 \times 1}$ and $[D]_{5 \times 3}$: the number of columns of the first matrix (1) does not equal the number of rows of the second matrix (5). Hence, the matrix product $[D]_{3 \times 5}^T [F]_{5 \times 1} [D]_{5 \times 3}$ is not defined, making the second statement wrong.

(A) Statement I is true but II is false

Q1.01.1 MCQ 2001 2M Ans: A ● ● ○

Given the matrices $[P]$ and $[Q]$:

$$[P] = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} \quad \text{and} \quad [Q] = \begin{bmatrix} 4 & 8 \\ 9 & 2 \end{bmatrix}$$

First, find the transpose of matrix $[Q]$ by interchanging its rows and columns:

$$[Q]^T = \begin{bmatrix} 4 & 9 \\ 8 & 2 \end{bmatrix}$$

Now, compute the matrix product $[P][Q]^T$:

$$\begin{aligned} [P][Q]^T &= \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 4 & 9 \\ 8 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2(4) + 3(8) & 2(9) + 3(2) \\ 4(4) + 5(8) & 4(9) + 5(2) \end{bmatrix} \\ &= \begin{bmatrix} 8 + 24 & 18 + 6 \\ 16 + 40 & 36 + 10 \end{bmatrix} = \begin{bmatrix} 32 & 24 \\ 56 & 46 \end{bmatrix} \end{aligned}$$

$$(A) \begin{bmatrix} 32 & 24 \\ 56 & 46 \end{bmatrix}$$

Q1.00.1 MCQ 2000 1M Ans: B ● ○ ○

By the definition of rank of a matrix A , the first statement is correct (i.e., true).

If A is an $n \times n$ matrix and $|A_{n \times n}| \neq 0$ then $\rho(A_{n \times n}) = n$. Therefore, the second statement is also correct (i.e., true).

Hence, option (B) is correct.

(B) Both the statements are true.

Q1.99.1 MCQ 1999 1M Ans: C ● ○ ○

By the properties of determinant of matrices, we have:

$$|kA_{n \times n}| = k^n |A_{n \times n}|$$

Therefore, matching the structural scaling condition parameters directly:

$$\alpha = k^n \quad \text{where } k \text{ is a scalar}$$

$$(C) k^n$$

Q1.99.2 MCQ 1999 1M Ans: B ● ● ●

If $A = [a_{11}]_{1 \times 1}$ then:

$$|A| = |a_{11}| = a_{11}$$

If $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ then:

$$|A| = a_{11}a_{22} - a_{12}a_{21}$$

If $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ then:

$$|A| = a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} + a_{12}a_{31}a_{23} - a_{12}a_{21}a_{33} + a_{13}a_{22}a_{32} - a_{13}a_{31}a_{22}$$

Therefore, looking at the pattern of the total number of terms in the permutation expansions: The number of terms in expansion of 1st order determinant is 1!. The number of terms in expansion of 2nd order determinant is 2!. The number of terms in expansion of 3rd order determinant is 3!.

Following this mathematical induction pattern, the number of terms in expansion of an n^{th} order determinant is $n!$.

(B) $n!$

Key Points

- **Leibniz Formula for Determinants:** The determinant of an $n \times n$ matrix is defined via the Leibniz formula as a sum over all possible permutations σ of the symmetric group S_n :

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n a_{i, \sigma(i)}$$

- **Permutations Count:** Since the symmetric group S_n contains exactly $n!$ distinct permutation elements, the full algebraic expansion of an unconstrained n^{th} -order determinant consists of exactly $n!$ terms.

Q1.99.3 MCQ 1999 2M Ans: B ● ● ○

Given the determinant equation:

$$\begin{vmatrix} 2 & 1 & 1 \\ 1 & 1 & -1 \\ y & x^2 & x \end{vmatrix} = 0$$

Expanding the determinant along the first row:

$$2 \begin{vmatrix} 1 & -1 \\ x^2 & x \end{vmatrix} - 1 \begin{vmatrix} 1 & -1 \\ y & x \end{vmatrix} + 1 \begin{vmatrix} 1 & 1 \\ y & x^2 \end{vmatrix} = 0$$

$$2(x + x^2) - 1(x + y) + 1(x^2 - y) = 0$$

Multiplying through and expanding the individual terms:

$$2x + 2x^2 - x - y + x^2 - y = 0$$

$$3x^2 + x - 2y = 0$$

Rearranging the terms to isolate the variable y yields the standard conic equation:

$$2y = 3x^2 + x$$

Since the expression contains a quadratic term in x and a linear term in y , it represents a parabola. This matches the structural coordinate points configuration given in option (b), as the parabola $2y = 3x^2 + x$ passes smoothly through the verification coordinates $(0, 0)$, $(-1, 1)$, and $(1, 2)$.

(B) $(0, 0), (-1, 1), (1, 2)$

Q1.98.1 MCQ 1998 1M Ans: C ● ○ ○

Consider the given matrix relationship equation:

$$AS = AT$$

By the left cancellation law of matrix multiplication, we have:

$$S = T$$

The cancellation of matrix A from both sides of the equation is mathematically valid if and only if there exists a unique inverse matrix A^{-1} such that pre-multiplying the system holds:

$$A^{-1}(AS) = A^{-1}(AT)$$

$$(A^{-1}A)S = (A^{-1}A)T$$

$$IS = IT \implies S = T$$

Therefore, the above cancellation is possible only if matrix A is non-singular, meaning its inverse exists.

(C) A is non-singular

Q1.98.2 MCQ 1998 2M Ans: D ● ● ○

Quadratic form (Q.F): A 2nd degree homogenous polynomial in two variables x_1 and x_2 is called a quadratic form (Q.F) (and it can be generalized to any number of variables with degree 2).

- Ex (1): $ax^2 + 2hxy + by^2$
- Ex (2): $ax_1x_1 + 2hx_1x_2 + bx_2x_2$

Matrix notation of Q.F:- A quadratic form in two variables x_1 and x_2 of the form $q(x_1, x_2) = a_{11}x_1^2 + a_{12}x_1x_2 + a_{21}x_2x_1 + a_{22}x_2^2$ can be expressed in matrix form as follows:

$$q(x_1, x_2) = a_{11}x_1^2 + a_{12}x_1x_2 + a_{21}x_2x_1 + a_{22}x_2^2 = X^TAX$$

where $X = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and symmetric matrix $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$.

Here, a_{11} = the coefficient of x_1^2 , $[a_{12} = a_{21} = \frac{1}{2}(\text{coefficient of } x_1x_2 \text{ (or) } x_2x_1)]$ and a_{22} = the coefficient of x_2^2 .

Given that $Q(x_1, x_2) = 4x_1x_2 - 5x_2x_2$

$$\implies Q = (0)x_1^2 + (4)x_1x_2 + (-5)x_2^2$$

Here, a_{11} = the coefficient of $x_1^2 = 0$ $[a_{12} = a_{21} = \frac{1}{2}(\text{the coefficient of } x_1x_2) = \frac{1}{2}(4) = 2]$ and a_{22} = the coefficient of $x_2^2 = -5$.

$\therefore$ The symmetric matrix is:

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 2 & -5 \end{bmatrix}$$

(D) $\begin{bmatrix} 0 & 2 \\ 2 & -5 \end{bmatrix}$

Q1.97.1 MCQ 1997 1M Ans: A ● ○ ○

The product matrix (BA) exists, only if, number of columns in B is equal to number of rows in A .

To state this mathematically, let the dimensions of the matrices be given by:

$$B = [b_{ij}]_{m \times n} \quad \text{and} \quad A = [a_{ij}]_{p \times q}$$

The matrix multiplication operation $(B \times A)$ is conformable and physically defined if and only if the inner adjacent dimension indices match identically:

$$n = p$$

When this criteria is satisfied, the resulting product matrix fields a final structural dimension order of $m \times q$.

$\therefore$ Option (A) is correct.

(A) only if A has as many rows as B has columns



SOLUTIONS — Linear Algebra



CH

Chemical Engineering

PrepFusion

Q1.26.1 MCQ 2026 1M Ans: B ● ○ ○

Let the given matrix be denoted as A :

$$A = \begin{bmatrix} -3 & 4 \\ 4 & 3 \end{bmatrix}$$

We can find the eigenvalues by using the properties of the matrix trace and determinant.

- The sum of eigenvalues equals the trace of the matrix:

$$\lambda_1 + \lambda_2 = -3 + 3 = 0$$

- The product of eigenvalues equals the determinant of the matrix:

$$\lambda_1 \cdot \lambda_2 = (-3)(3) - (4)(4) = -9 - 16 = -25$$

Evaluating the given options:

- For option (A), the sum is 0 and the product is -49 .
- For option (B), the sum is 0 and the product is -25 .

Therefore, the eigenvalues are -5 and 5 .

(B) $-5, 5$

Key Points

- The sum of the eigenvalues of a matrix is equal to the sum of the elements on its main diagonal.
- The product of the eigenvalues of a matrix is equal to its determinant.

Q1.24.1 MSQ 2024 1M Ans: A,C ● ○ ○

For a linear homogeneous system of equations $Ax = 0$, where A is an $n \times n$ square matrix:

- The system always admits the trivial solution $x = 0$.
- For a non-trivial solution ($x \neq 0$) to exist, the columns of matrix A must be linearly dependent.
- This structural dependency implies that matrix A must be singular, meaning its determinant value reduces exactly to zero:

$$\text{Determinant of } A = 0$$

- Consequently, the rank r of the matrix (the maximum number of linearly independent rows or columns) must be strictly less than the total dimension count n :

$$r < n$$

Evaluating the conditions presented in the options:

- Option (A) is satisfied since the determinant must equal 0.
- Option (C) is satisfied since the rank r must be less than n .

Therefore, the conditions satisfied are given by options (A) and (C).

(A) Determinant of $A = 0$, (C) $r < n$

Q1.24.2 NAT 2024 1M Ans: -2 ● ○ ○

Given the matrix A :

$$A = \begin{bmatrix} -5 & a \\ -2 & -2 \end{bmatrix}$$

The eigenvalues of the matrix are $\lambda_1 = -1$ and $\lambda_2 = -6$. We can determine the value of the constant a by leveraging the property that the product of the eigenvalues of a matrix is equal to its determinant.

Evaluating the product of the given eigenvalues:

$$\lambda_1 \cdot \lambda_2 = (-1) \cdot (-6) = 6$$

Next, computing the determinant of matrix A :

$$\det(A) = (-5)(-2) - (a)(-2) = 10 + 2a$$

Setting the determinant equal to the product of the eigenvalues gives:

$$10 + 2a = 6$$

$$2a = 6 - 10$$

$$2a = -4$$

$$a = -2$$

Therefore, the value of the constant a is -2 .

$$\boxed{-2}$$

Key Points

- The determinant of a square matrix is identically equal to the product of its eigenvalues.
- The trace of a square matrix corresponds exactly to the sum of its eigenvalues.

Q1.23.1 NAT 2023 2M Ans: 4160 ● ● ○

Given the matrix M :

$$M = \begin{bmatrix} 10 & 6 \\ 6 & 10 \end{bmatrix}$$

First, we determine the eigenvalues of matrix M by establishing its characteristic equation $\det(M - \lambda I) = 0$:

$$\det \begin{bmatrix} 10 - \lambda & 6 \\ 6 & 10 - \lambda \end{bmatrix} = 0$$

$$(10 - \lambda)^2 - 36 = 0$$

$$(10 - \lambda)^2 = 36$$

$$10 - \lambda = \pm 6$$

$$\lambda_1 = 10 - 6 = 4$$

$$\lambda_2 = 10 + 6 = 16$$

If a matrix has eigenvalues λ , then the matrix raised to the power of n (M^n) carries eigenvalues equal to λ^n . Therefore, the eigenvalues of the cubed matrix M^3 are:

$$\lambda_1^3 = 4^3 = 64$$

$$\lambda_2^3 = 16^3 = 4096$$

The total sum of all the eigenvalues belonging to M^3 is computed as:

$$\text{Sum} = 64 + 4096 = 4160$$

Therefore, the sum of all eigenvalues of M^3 is 4160.

4160

Key Points

- If λ is an eigenvalue of M , then λ^n represents the corresponding eigenvalue for the matrix power M^n .
- The total sum of all eigenvalues belonging to any square matrix equals the trace value of that specific matrix.

Q1.22.1 MCQ 2022 1M Ans: B ☒ ☐ ☐

Given the matrix A :

$$A = \begin{bmatrix} x & 1 & 3 \\ y & 2 & 6 \\ 3 & 5 & 7 \end{bmatrix}$$

$$\det(A) = x \begin{vmatrix} 2 & 6 \\ 5 & 7 \end{vmatrix} - 1 \begin{vmatrix} y & 6 \\ 3 & 7 \end{vmatrix} + 3 \begin{vmatrix} y & 2 \\ 3 & 5 \end{vmatrix} = 0$$

Computing the individual 2×2 determinants:

$$x(2 \cdot 7 - 6 \cdot 5) - 1(y \cdot 7 - 6 \cdot 3) + 3(y \cdot 5 - 2 \cdot 3) = 0$$

$$x(14 - 30) - 1(7y - 18) + 3(5y - 6) = 0$$

$$-16x - 7y + 18 + 15y - 18 = 0$$

$$-16x + 8y = 0$$

$$8y = 16x$$

$$y = 2x$$

Substituting the given options into the derived relation $y = 2x$:

- For option (A) $(1, 1)$, $1 \neq 2(1)$.
- For option (B) $(1, 2)$, $2 = 2(1)$, which satisfies the relation.

Therefore, the correct ordered pair is $(1, 2)$.

(B) $(1, 2)$

Q1.21.1 NAT 2021 1M Ans: 3 ☒ ☐ ☐

Let M be the given 4×4 matrix composed of row vectors A , B , C , and D :

$$M = \begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \\ d_1 & d_2 & d_3 & d_4 \end{bmatrix} = \begin{bmatrix} A \\ B \\ C \\ D \end{bmatrix}$$

The rank of a matrix equals the maximum number of linearly independent row vectors it contains. We evaluate the dependency structure of the rows: Since B is not a scalar multiple of A , the set $\{A, B\}$ consists of 2 linearly independent

vectors. Since C is linearly independent of both A and B , the set $\{A, B, C\}$ forms a collection of 3 linearly independent vectors.

The fourth row vector D is explicitly given as a linear combination of the preceding rows:

$$D = 3A + 2B + C$$

Because row D depends entirely on rows A , B , and C , it does not contribute to the count of independent rows. Performing the row operation $R_4 \rightarrow R_4 - 3R_1 - 2R_2 - R_3$ reduces the fourth row completely to a zero row:

$$\begin{bmatrix} A \\ B \\ C \\ 0 \end{bmatrix}$$

Consequently, the number of linearly independent rows in the matrix is exactly 3. Therefore, the rank of the matrix is 3.

3

Q1.21.2 MCQ 2021 2M Ans: D ● ● ○

Given the description of the elements a_{ij} for the square matrix A of size $n \times n$:

$$a_{ij} = \begin{cases} i \times j, & \text{if } i \geq j \\ 0, & \text{if } i < j \end{cases}$$

Let us write out the structural format of the matrix A :

$$A = \begin{bmatrix} 1 \times 1 & 0 & 0 & \dots & 0 \\ 2 \times 1 & 2 \times 2 & 0 & \dots & 0 \\ 3 \times 1 & 3 \times 2 & 3 \times 3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ n \times 1 & n \times 2 & n \times 3 & \dots & n \times n \end{bmatrix}$$

Observing the structure, all elements above the main diagonal are equal to zero ($a_{ij} = 0$ for $i < j$). Therefore, A is a lower triangular matrix.

The determinant of a triangular matrix is equal to the product of its main diagonal elements:

$$\det(A) = a_{11} \cdot a_{22} \cdot a_{33} \dots a_{nn}$$

$$\det(A) = (1 \times 1) \cdot (2 \times 2) \cdot (3 \times 3) \dots (n \times n)$$

Rearranging the terms yields:

$$\det(A) = (1 \cdot 2 \cdot 3 \dots n) \cdot (1 \cdot 2 \cdot 3 \dots n)$$

$$\det(A) = (n!) \cdot (n!) = (n!)^2$$

Therefore, the determinant of A is $(n!)^2$.

(D) $(n!)^2$

Q1.19.1 MCQ 2019 1M Ans: C ● ○ ○

Consider a system of n homogeneous linear equations containing n unknowns represented in matrix form as:

$$Ax = 0$$

Here, A represents the $n \times n$ square coefficient matrix, x is the $n \times 1$ column vector of unknowns, and 0 is the $n \times 1$ null vector. A homogeneous linear system always yields the trivial solution $x = 0$. For the system to possess non-trivial solutions, the columns of the coefficient matrix A must be linearly dependent. This geometric restriction requires that A must be a singular matrix, meaning its determinant must equal zero:

$$\det(A) = 0$$

If $\det(A) \neq 0$, the matrix is invertible, and multiplying both sides by A^{-1} isolates the unique trivial solution $x = 0$. Therefore, non-trivial solutions exist if and only if the determinant is exactly 0.

(C) 0

Key Points

- A square homogeneous system $Ax = 0$ yields non-trivial solutions if and only if the matrix A is singular ($\det(A) = 0$).
- A vanishing determinant corresponds identically to a reduced rank ($r < n$), which introduces free variables into the system layout.

Mistakes to be avoided

- Do not assume that a homogeneous system can ever be inconsistent; it always has at least the trivial solution.

Q1.18.1 NAT 2018 2M Ans: 1 ● ● ○

Given the matrix A :

$$A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

We want to find the value of $\det(A^T A)$. This can be solved efficiently using two different methods.

Method 1

We can use the multiplicative properties of determinants. For any square matrices A and B of the same size, $\det(AB) = \det(A) \det(B)$. Furthermore, the determinant of a matrix is equal to the determinant of its transpose, $\det(A^T) = \det(A)$.

Applying these properties gives:

$$\det(A^T A) = \det(A^T) \cdot \det(A) = \det(A) \cdot \det(A) = [\det(A)]^2$$

Now, we evaluate the determinant of the 2×2 matrix A :

$$\det(A) = (\cos \theta)(\cos \theta) - (-\sin \theta)(\sin \theta)$$

$$\det(A) = \cos^2 \theta + \sin^2 \theta = 1$$

Substituting this value back into the determinant expression yields:

$$\det(A^T A) = (1)^2 = 1$$

Method 2

Alternatively, we can compute the matrix product $A^T A$ directly. First, we write the transpose matrix A^T :

$$A^T = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

Now, multiplying A^T and A :

$$A^T A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

$$A^T A = \begin{pmatrix} \cos^2 \theta + \sin^2 \theta & -\cos \theta \sin \theta + \sin \theta \cos \theta \\ -\sin \theta \cos \theta + \cos \theta \sin \theta & \sin^2 \theta + \cos^2 \theta \end{pmatrix}$$

$$A^T A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$$

Since $A^T A$ forms an identity matrix I , its determinant is identically equal to 1:

$$\det(A^T A) = \det(I) = 1$$

Therefore, the value of $\det(A^T A)$ is 1.

1

Key Points

- The determinant of a product of matrices satisfies $\det(AB) = \det(A) \det(B)$.
- Transposition preserves the determinant value of a square matrix, meaning $\det(A^T) = \det(A)$.
- A matrix that satisfies $A^T A = I$ is classified as an orthogonal matrix, and its determinant is always ± 1 .

Q1.16.1 NAT 2016 2M Ans: 15 ● ● ○

Given the matrix representation of simultaneous linear equations:

$$\begin{bmatrix} 0 & 0 & 0 & 4 & 13 \\ 2 & 5 & 5 & 2 & 10 \\ 0 & 0 & 2 & 5 & 3 \\ 0 & 0 & 0 & 4 & 5 \\ 2 & 3 & 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 46 \\ 161 \\ 61 \\ 30 \\ 81 \end{bmatrix}$$

We isolate equations that contain fewer variables to systematically solve for the unknowns. Extracting the first and fourth equations: From row 1:

$$4x_4 + 13x_5 = 46 \quad \text{--- (Equation 1)}$$

From row 4:

$$4x_4 + 5x_5 = 30 \quad \text{--- (Equation 2)}$$

Subtracting Equation 2 from Equation 1 eliminates the x_4 term:

$$(4x_4 + 13x_5) - (4x_4 + 5x_5) = 46 - 30$$

$$8x_5 = 16$$

$$x_5 = 2$$

Substituting $x_5 = 2$ back into Equation 2 to determine x_4 :

$$4x_4 + 5(2) = 30$$

$$4x_4 + 10 = 30$$

$$4x_4 = 20$$

$$x_4 = 5$$

Next, we extract the third equation from row 3, which contains x_3 , x_4 , and x_5 :

$$2x_3 + 5x_4 + 3x_5 = 61$$

Substituting the known values $x_4 = 5$ and $x_5 = 2$ into this equation:

$$2x_3 + 5(5) + 3(2) = 61$$

$$2x_3 = 30$$

$$x_3 = 15$$

$$\boxed{15}$$

Q1.15.1 MCQ 2015 1M Ans: D ☒ ☐ ☐

Let the given column vectors form the columns of a matrix A :

$$A = \begin{bmatrix} 1 & x & 3 \\ 2 & 6 & 4 \\ 1 & x & 2 \end{bmatrix}$$

A set of vectors is linearly dependent if and only if the determinant of the matrix formed by these vectors equals zero. Evaluating the determinant of matrix A by expanding along the first row:

$$\det(A) = 1 \begin{vmatrix} 6 & 4 \\ x & 2 \end{vmatrix} - x \begin{vmatrix} 2 & 4 \\ 1 & 2 \end{vmatrix} + 3 \begin{vmatrix} 2 & 6 \\ 1 & x \end{vmatrix} = 0$$

Computing the individual 2×2 determinants:

$$1(6 \cdot 2 - 4 \cdot x) - x(2 \cdot 2 - 4 \cdot 1) + 3(2 \cdot x - 6 \cdot 1) = 0$$

$$1(12 - 4x) - x(4 - 4) + 3(2x - 6) = 0$$

$$12 - 4x - 0 + 6x - 18 = 0$$

$$2x = 6$$

$$x = 3$$

Therefore, the set of vectors is linearly dependent when x is equal to 3.

$$\boxed{(D) 3}$$

Q1.15.2 NAT 2015 1M Ans: 7 ☒ ☐ ☐

Let the given matrix be denoted as A and the eigenvector as x :

$$A = \begin{pmatrix} 4 & 3 \\ 3 & 4 \end{pmatrix}, \quad x = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

By definition, if x is an eigenvector of a matrix A , it satisfies the fundamental relation $Ax = \lambda x$, where λ represents the corresponding eigenvalue.

Multiplying the matrix A by the vector x :

$$Ax = \begin{pmatrix} 4 & 3 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 4 \cdot 1 + 3 \cdot 1 \\ 3 \cdot 1 + 4 \cdot 1 \end{pmatrix}$$

$$Ax = \begin{pmatrix} 7 \\ 7 \end{pmatrix}$$

Factoring out the common scalar value from the resulting vector:

$$Ax = 7 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Comparing this outcome directly with the eigenvalue relation $Ax = \lambda x$:

$$\lambda = 7$$

7

Q1.13.1 MCQ 2013 1M Ans: B ● ○ ○

We analyze each statement individually to determine its truth value:

- Statement P: The eigenvalues of a symmetric matrix are real. This statement is true. A real symmetric matrix satisfies $A = A^T$, which mathematically guarantees that all its roots of the characteristic polynomial lie entirely on the real line.
- Statement Q: The value of the determinant of an orthogonal matrix can only be +1. This statement is false. An orthogonal matrix satisfies $A^T A = I$. Taking the determinant on both sides yields $\det(A^T) \det(A) = \det(I) \implies [\det(A)]^2 = 1$, which gives $\det(A) = \pm 1$. Thus, it can be +1 or -1.
- Statement R: The transpose of a square matrix A has the same eigenvalues as those of A . This statement is true. The eigenvalues are determined by the characteristic equation $\det(A - \lambda I) = 0$. Since transposition preserves the determinant value of a matrix, $\det(A - \lambda I)^T = \det(A^T - \lambda I) = 0$, meaning both matrices share identical characteristic roots.
- Statement S: The inverse of an $n \times n$ matrix exists if and only if the rank is less than n . This statement is false. An $n \times n$ square matrix is invertible if and only if it has full rank, meaning its rank equals n . If the rank is less than n , the matrix is singular and its inverse does not exist.

Since statements P and R are true, the correct selection corresponds to option (B).

(B) P and R only

Q1.12.1 MCQ 2012 1M Ans: D ● ○ ○

Let the given matrix be denoted as A :

$$A = \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} = 4I$$

where I is the 2×2 identity matrix. By definition, a non-zero column vector x is an eigenvector of a matrix A if it satisfies the condition $Ax = \lambda x$ for a scalar value λ . Let us multiply matrix A by any non-zero vector x :

$$Ax = (4I)x = 4(Ix) = 4x$$

Since this fundamental relation holds true for every non-zero vector in the vector space, any non-zero vector serves as a valid eigenvector of this scalar matrix with a corresponding eigenvalue of $\lambda = 4$.

However, by formal mathematical definition, the null vector (or zero vector) is strictly excluded from being a valid eigenvector because it satisfies $A \cdot 0 = \lambda \cdot 0$ trivially for any scalar value λ , which provides no structural transformation data. Evaluating the choices presented in the options:

- Options (A), (B), and (C) represent non-zero column vectors, making them valid eigenvectors.
- Option (D) is the null vector $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, which cannot be an eigenvector.

Therefore, the vector that is not a valid eigenvector corresponds to option (D).

$$(D) \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Q1.11.1 MCQ 2011 1M Ans: D ● ○ ○

Given the eigenvalues $\lambda_1 = -1$ and $\lambda_2 = 3$ for a real 2×2 matrix R . The corresponding eigenvectors are:

$$V_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

We are given that the modal matrix P is constructed using the eigenvectors as its column vectors:

$$P = [V_1 \quad V_2]$$

According to the matrix diagonalization theorem, if a square matrix R has linearly independent eigenvectors that form the columns of matrix P , then the similarity transformation matrix $P^{-1}RP$ results in a diagonal matrix D :

$$P^{-1}RP = D = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$$

Here, the diagonal elements of D correspond precisely to the eigenvalues arranged in the exact same column order as their respective eigenvectors in P . Since V_1 occupies the first column and V_2 occupies the second column of P , the diagonal matrix is configured as:

$$P^{-1}RP = \begin{pmatrix} -1 & 0 \\ 0 & 3 \end{pmatrix}$$

$$(D) \begin{pmatrix} -1 & 0 \\ 0 & 3 \end{pmatrix}$$

Q1.10.1 MCQ 2010 1M Ans: A ● ● ○

Given the definition of the elements a_{ij} for the real square matrix A of size $n \times n$:

$$a_{ij} = \begin{cases} i, & \text{if } i = j \\ 0, & \text{otherwise} \end{cases}$$

Let us explicitly display the structure of matrix A :

$$A = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 2 & 0 & \dots & 0 \\ 0 & 0 & 3 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & n \end{bmatrix}$$

Observing the structure, A is a diagonal matrix. For any diagonal matrix, the eigenvalues are simply the elements located on the main diagonal. Therefore, the eigenvalues of A are:

$$\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 3, \dots, \lambda_n = n$$

The problem asks for the summation of all n eigenvalues, which matches the sum of the first n natural numbers:

$$\text{Sum} = \sum_{i=1}^n \lambda_i = 1 + 2 + 3 + \cdots + n$$

Using the standard arithmetic series summation formula:

$$\text{Sum} = \frac{n(n+1)}{2}$$

Therefore, the summation of all n eigenvalues is $n(n+1)/2$.

$$(A) \ n(n+1)/2$$

Key Points

- The eigenvalues of a diagonal matrix are exactly equal to its main diagonal entries.
- The sum of the first n natural numbers is evaluated using the identity $\sum i = n(n+1)/2$.

Q1.10.2 MCQ 2010 2M Ans: C ● ● ○

Given that X and Y are non-zero square matrices of size $n \times n$ satisfying the matrix product equation:

$$XY = 0_{n \times n}$$

$$|XY| = |0_{n \times n}|$$

Using the multiplicative property of determinants, $|XY| = |X||Y|$, and knowing that the determinant of a zero matrix is 0:

$$|X| \cdot |Y| = 0$$

This scalar equation implies that at least one of the determinants must be equal to zero, meaning $|X| = 0$ or $|Y| = 0$ (or both). Let us check the possibilities through contradiction:

Suppose $|X| \neq 0$. This means matrix X is non-singular and its inverse X^{-1} exists. Pre-multiplying both sides of the initial equation $XY = 0$ by X^{-1} gives:

$$X^{-1}(XY) = X^{-1} \cdot 0$$

$$(X^{-1}X)Y = 0$$

$$I \cdot Y = 0 \implies Y = 0$$

However, this contradicts the given condition that Y is a non-zero matrix. Therefore, $|X|$ cannot be non-zero, which forces:

$$|X| = 0$$

Similarly, suppose $|Y| \neq 0$. This means matrix Y is non-singular and its inverse Y^{-1} exists. Post-multiplying both sides of the initial equation $XY = 0$ by Y^{-1} gives:

$$(XY)Y^{-1} = 0 \cdot Y^{-1}$$

$$X(YY^{-1}) = 0$$

$$X \cdot I = 0 \implies X = 0$$

This contradicts the given condition that X is a non-zero matrix. Therefore, $|Y|$ cannot be non-zero, which forces:

$$|Y| = 0$$

Since both assumptions lead to a contradiction, both matrices must simultaneously be singular:

$$|X| = 0 \text{ and } |Y| = 0$$

Evaluating the choices presented in the options shows that this satisfies option (C).

$$(C) \ |X| = 0 \text{ and } |Y| = 0$$

Q1.09.1 MCQ 2009 1M Ans: C ● ○ ○

Consider the given system of n linear homogeneous equations in n unknowns:

$$Ax = 0$$

where A is an $n \times n$ square matrix, x is the vector of unknowns, and 0 is the null vector.

A homogeneous linear system always contains the trivial solution $x = 0$. For non-trivial solutions ($x \neq 0$) to exist, the coefficient matrix A must be singular, which means its determinant must drop to zero:

$$\det(A) = 0$$

According to matrix rank properties, when the determinant of a square matrix of size $n \times n$ is zero, the maximum number of linearly independent rows or columns is strictly less than n . Therefore, the rank of matrix A must satisfy:

$$\text{rank of } A < n$$

If the rank equals n , the matrix is non-singular, and the system possesses only the unique trivial solution. Thus, a non-trivial solution exists only if the rank of A is less than n .

(C) rank of $A < n$

Q1.09.2 MCQ 2009 2M Ans: A ● ● ○

Given the eigenvalues of matrix A are $\lambda_1 = 5$ and $\lambda_2 = -1$.

We need to find the eigenvalues of the matrix expression:

$$B = -2A + 3I$$

According to the properties of matrix polynomials, if λ is an eigenvalue of matrix A , then the corresponding eigenvalue λ_B of a linear matrix combination $B = c_1A + c_2I$ is given by substituting λ into the polynomial expression:

$$\lambda_B = c_1\lambda + c_2$$

Let us map each eigenvalue of matrix A to its corresponding value for matrix B : For $\lambda_1 = 5$:

$$\lambda_{B1} = -2(5) + 3 = -10 + 3 = -7$$

For $\lambda_2 = -1$:

$$\lambda_{B2} = -2(-1) + 3 = 2 + 3 = 5$$

The resulting pair of eigenvalues for the matrix expression is -7 and 5 .

(A) -7 and 5

Q1.07.1 MCQ 2007 2M Ans: B ● ● ○

Given that A and B are two 3×3 matrices such that $B \neq 0$ and:

$$A = \begin{bmatrix} -2 & 4 & 6 \\ 1 & 2 & 1 \\ 0 & 4 & 4 \end{bmatrix}, \quad AB = 0$$

First, let us calculate the determinant of matrix A to evaluate its rank structure:

$$\det(A) = -2(2 \cdot 4 - 1 \cdot 4) - 4(1 \cdot 4 - 1 \cdot 0) + 6(1 \cdot 4 - 2 \cdot 0)$$

$$\det(A) = -2(8 - 4) - 4(4) + 6(4)$$

$$\det(A) = -2(4) - 16 + 24 = -8 - 16 + 24 = 0$$

Since $\det(A) = 0$, matrix A is singular, meaning its rank is strictly less than its dimension: $\text{rank}(A) < 3$. Let us examine a 2×2 submatrix to determine if the rank is 2:

$$\begin{vmatrix} -2 & 4 \\ 1 & 2 \end{vmatrix} = -4 - 4 = -8 \neq 0$$

Since there is a non-zero 2×2 minor, the rank of matrix A is exactly 2:

$$\text{rank}(A) = 2$$

According to Sylvester's rank inequality for the product of two matrices A and B of size $n \times n$:

$$\text{rank}(A) + \text{rank}(B) - n \leq \text{rank}(AB)$$

Substituting the known conditions $\text{rank}(A) = 2$, $n = 3$, and $\text{rank}(AB) = \text{rank}(0) = 0$:

$$2 + \text{rank}(B) - 3 \leq 0$$

$$\text{rank}(B) - 1 \leq 0$$

$$\text{rank}(B) \leq 1$$

Since we are given that $B \neq 0$, the rank of B must be at least 1. Therefore, $\text{rank}(B) = 1$. Let us evaluate this condition against the choices provided: The constraint $\text{rank}(B) = 1$ satisfies the inequality condition $\text{rank}(B) < 3$ presented in option (B).

Therefore, the rank of matrix B satisfies $r < 3$.

$$(B) \ r < 3$$

Q1.05.1 MCQ 2005 2M Ans: D ● ● ○

Given the 2×2 matrix A :

$$A = \begin{pmatrix} 1 & 4 \\ a & 2 \end{pmatrix}$$

We determine the characteristic equation of matrix A via $\det(A - \lambda I) = 0$:

$$\det \begin{pmatrix} 1 - \lambda & 4 \\ a & 2 - \lambda \end{pmatrix} = 0$$

$$(1 - \lambda)(2 - \lambda) - 4a = 0$$

$$\lambda^2 - 3\lambda + 2 - 4a = 0$$

The roots of this quadratic equation represent the eigenvalues λ . For the eigenvalues to be real numbers, the discriminant D of the quadratic equation must be greater than or equal to zero:

$$D = (-3)^2 - 4(1)(2 - 4a) \geq 0$$

$$9 - 8 + 16a \geq 0$$

$$1 + 16a \geq 0 \implies a \geq -\frac{1}{16}$$

Next, for the eigenvalues to be non-negative ($\lambda \geq 0$), both their sum and their product must be non-negative. The sum of the eigenvalues is given by the trace of the matrix:

$$\lambda_1 + \lambda_2 = 3 > 0$$

This condition is automatically satisfied. The product of the eigenvalues is given by the determinant of the matrix, which must also be greater than or equal to zero:

$$\lambda_1 \cdot \lambda_2 = \det(A) = 2 - 4a \geq 0$$

$$2 \geq 4a \implies a \leq \frac{2}{4} \implies a \leq \frac{1}{2}$$

Combining the constraints established from both the real root condition and the non-negative root condition gives:

$$-\frac{1}{16} \leq a \leq \frac{1}{2}$$

$$(D) \left(-\frac{1}{16}\right) \leq a \leq \left(\frac{1}{2}\right)$$

Key Points

- For a quadratic characteristic equation $\lambda^2 - \text{Trace}(A)\lambda + \det(A) = 0$, eigenvalues are real if the discriminant satisfies $D \geq 0$.
- Real eigenvalues are non-negative if and only if both the trace and the determinant are non-negative.

Mistakes to be avoided

- Do not overlook checking the determinant non-negativity constraint after validating the discriminant bounds, as real roots can still be negative if the product condition is skipped.

Q1.02.1 NAT 2002 5M Ans: * ● ● ●

Given matrix A :

$$A = \begin{bmatrix} 0.1 & 0.5 & 0 \\ 0.8 & 0 & 0.4 \\ 0.1 & 0.5 & 0.6 \end{bmatrix}$$

We solve each part of the problem systematically:

- Part a.** The characteristic equation is defined by $\det(A - \lambda I) = 0$:

$$\det \begin{bmatrix} 0.1 - \lambda & 0.5 & 0 \\ 0.8 & -\lambda & 0.4 \\ 0.1 & 0.5 & 0.6 - \lambda \end{bmatrix} = 0$$

Expanding the determinant along the first row:

$$(0.1 - \lambda)[- \lambda(0.6 - \lambda) - 0.2] - 0.5[0.8(0.6 - \lambda) - 0.04] = 0$$

$$(0.1 - \lambda)(\lambda^2 - 0.6\lambda - 0.2) - 0.5(0.48 - 0.8\lambda - 0.04) = 0$$

$$0.1\lambda^2 - 0.06\lambda - 0.02 - \lambda^3 + 0.6\lambda^2 + 0.2\lambda - 0.22 + 0.4\lambda = 0$$

$$-\lambda^3 + 0.7\lambda^2 + 0.54\lambda - 0.24 = 0 \implies \lambda^3 - 0.7\lambda^2 - 0.54\lambda + 0.24 = 0$$

- Part b.** We are given the condition $Ax = x$ for some vector x . By definition of the eigenvalue equation $Ax = \lambda x$, this implies that one eigenvalue of the matrix corresponds precisely to:

$$\lambda_1 = 1$$

Alternatively, summing the elements of each column in matrix A :

- Column 1: $0.1 + 0.8 + 0.1 = 1.0$
- Column 2: $0.5 + 0 + 0.5 = 1.0$
- Column 3: $0 + 0.4 + 0.6 = 1.0$

Since all column sums are identically equal to 1, $\lambda = 1$ is guaranteed to be an eigenvalue.

- **Part c.** To find the remaining two eigenvalues, we use the trace and determinant matrix properties. The trace of matrix A represents the sum of all eigenvalues:

$$\lambda_1 + \lambda_2 + \lambda_3 = \text{Trace}(A) = 0.1 + 0 + 0.6 = 0.7$$

Substituting $\lambda_1 = 1$:

$$1 + \lambda_2 + \lambda_3 = 0.7 \implies \lambda_2 + \lambda_3 = -0.3$$

The product of all eigenvalues represents the determinant value:

$$\lambda_1 \cdot \lambda_2 \cdot \lambda_3 = \det(A) = -0.24$$

Substituting $\lambda_1 = 1$:

$$\lambda_2 \cdot \lambda_3 = -0.24$$

We form a quadratic equation to solve for λ_2 and λ_3 :

$$\lambda^2 - (\lambda_2 + \lambda_3)\lambda + (\lambda_2\lambda_3) = 0$$

$$\lambda^2 + 0.3\lambda - 0.24 = 0$$

$$(\lambda + 0.6)(\lambda - 0.4) = 0$$

Solving this equation yields the other two eigenvalues:

$$\lambda_2 = -0.6, \quad \lambda_3 = 0.4$$

a. $\lambda^3 - 0.7\lambda^2 - 0.54\lambda + 0.24 = 0$, b. $\lambda = 1$, c. -0.6 and 0.4

SOLUTIONS — Linear Algebra

PI

Production & Industrial Engineering

PrepFusion

Q1.26.1 MCQ 2026 2M Ans: C ● ● ○

We analyze the four given 3×3 matrices to map each one to its corresponding mathematical property:

- **Matrix P:** Notice that the elements across the main diagonal are mirror images:

$$P_{12} = P_{21} = 17, \quad P_{13} = P_{31} = 5, \quad P_{23} = P_{32} = -12$$

Since $P = P^T$, it is a **Symmetric** matrix (Matches 3).

- **Matrix Q:** The trace of a matrix is the sum of its main diagonal elements. Here:

$$\text{Trace} = 0 + 0 + 0 = 0$$

Thus, it is **Trace free** (Matches 4).

- **Matrix S:** All elements below the main diagonal are exactly zero. This defines an Upper **Triangular** matrix (Matches 2).
- **Matrix R:** By elimination, this must be singular. Let's verify by checking its determinant $|R|$. Expanding along the first column:

$$|R| = (2/3) \times [(2/3)(-2/3) - (-1/3)(4/3)] - 0 + 0$$

$$|R| = (2/3) \times [-4/9 + 4/9] = (2/3) \times 0 = 0$$

Since the determinant is zero, it is indeed a **Singular** matrix (Matches 1).

The correct pairing is P-3; Q-4; R-1; S-2.

(C) P-3; Q-4; R-1; S-2

Key Points

- A matrix is symmetric if it equals its transpose ($A = A^T$), meaning entries are symmetric relative to the main diagonal.
- The trace of a square matrix represents the sum of its main diagonal entries; a matrix with a trace of zero is trace-free.
- A matrix is singular if its determinant equals zero, which indicates that its rows or columns are linearly dependent.

Q1.25.1 MCQ 2025 1M Ans: A ● ○ ○

Let

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

The characteristic equation is $|A - \lambda I| = 0$

$$\begin{vmatrix} -\lambda & -1 \\ 1 & -\lambda \end{vmatrix} = 0$$

$$\lambda^2 + 1 = 0$$

$$\Rightarrow \lambda^2 = -1 = i^2$$

$$\lambda = i, -i$$

$$\lambda = \sqrt{-1} \text{ and } -\sqrt{-1}$$

(A) $-\sqrt{-1}$ and $\sqrt{-1}$

Key Points

- The characteristic equation of a square matrix A is given by $\det(A - \lambda I) = 0$, where λ represents the eigenvalues.
- For a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, the characteristic equation can also be expressed as $\lambda^2 - \text{trace}(A)\lambda + \det(A) = 0$.

Q1.24.1

MCQ

2024

2M

Ans: C



Method 1

Given that

$$A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix}$$

$$\Rightarrow \text{the characteristic equation is } \lambda^2 + (-a^2 - bc) = 0$$

$$\Rightarrow \lambda^2 - a^2 - bc = 0$$

$$\Rightarrow \lambda^2 - (a^2 + bc) = 0$$

By Cayley-Hamilton theorem, we have

$$A^2 - (a^2 + bc)I = 0$$

Given that $A^2 = I$:

$$\Rightarrow I - a^2I - bcI = 0$$

$$\Rightarrow (1 - a^2 - bc)I = 0$$

$$\therefore 1 - a^2 - bc = 0$$

Method 2

Consider $A^2 = I$

$$\Rightarrow A \cdot A = I$$

$$\Rightarrow \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \begin{pmatrix} a & b \\ c & -a \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} a^2 + bc & ab - ab \\ ac - ac & bc + a^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} a^2 + bc & 0 \\ 0 & bc + a^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\Rightarrow a^2 + bc = 1$$

$$\therefore 1 - a^2 - bc = 0$$

$$\boxed{(C) \ 1 - a^2 - bc = 0}$$

Key Points

- Cayley-Hamilton Theorem: Every square matrix satisfies its own characteristic equation.
- Matrix Equality: Two matrices are equal if and only if their corresponding elements are equal.

Q1.23.1 MCQ 2023 1M Ans: MTA

Given matrix:

$$A = \begin{bmatrix} 1 & -1 & 4 \\ 3 & 2 & -1 \\ 2 & 1 & -1 \end{bmatrix}$$

Every square matrix A can be uniquely expressed as the sum of a symmetric matrix P and a skew-symmetric matrix B :

$$A = P + B$$

where the skew-symmetric part B is defined as:

$$B = \frac{A - A^T}{2}$$

First, determine the transpose matrix A^T :

$$A^T = \begin{bmatrix} 1 & 3 & 2 \\ -1 & 2 & 1 \\ 4 & -1 & -1 \end{bmatrix}$$

Now, compute the matrix $A - A^T$:

$$A - A^T = \begin{bmatrix} 1-1 & -1-3 & 4-2 \\ 3-(-1) & 2-2 & -1-1 \\ 2-4 & 1-(-1) & -1-(-1) \end{bmatrix} = \begin{bmatrix} 0 & -4 & 2 \\ 4 & 0 & -2 \\ -2 & 2 & 0 \end{bmatrix}$$

Dividing by 2 to obtain the skew-symmetric matrix B :

$$B = \frac{1}{2} \begin{bmatrix} 0 & -4 & 2 \\ 4 & 0 & -2 \\ -2 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -2 & 1 \\ 2 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix}$$

Comparing this with the given matrix template for B , the element B_{13} located at the first row and third column is:

$$B_{13} = 1$$

Since the value 1 is not present among the given options (A, B, C, D), this question contains an error in its options and is classified as Marks to All (MTA).

MTA

Q1.23.2 MCQ 2023 1M Ans: A

Given coordinates of the original point:

$$P(x, y) = P(10, 15)$$

Coordinates of the translated point:

$$P'(x', y') = P(15, 25)$$

The translation relation along the x -axis is:

$$x' = x + t_x$$

$$15 = 10 + t_x$$

$$\Rightarrow t_x = 5$$

The translation relation along the y -axis is:

$$y' = y + t_y$$

$$25 = 15 + t_y$$

$$\Rightarrow t_y = 10$$

Using the homogeneous translation matrix equation $[P'] = [T][P]$:

$$\begin{aligned} \begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} &= \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix} = \begin{bmatrix} x + 5 \\ y + 10 \\ 1 \end{bmatrix} \\ \begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} &= \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 10 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \end{aligned}$$

Comparing with $P' = T \cdot P$, the translation matrix T is:

$$T = \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 10 \\ 0 & 0 & 1 \end{bmatrix}$$

$$(A) \begin{bmatrix} 1 & 0 & 5 \\ 0 & 1 & 10 \\ 0 & 0 & 1 \end{bmatrix}$$

Key Points

- 2D Translation Matrix: The textbook structure for a 2D translation matrix in a homogeneous coordinate system is defined as:

$$T = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

where t_x and t_y represent the translation distances along the x and y axes respectively.

Q1.23.3 NAT 2023 2M Ans: 1 ● ● ○

Given matrix

$$A = \begin{bmatrix} 4 & 2 \\ 3 & 3 \end{bmatrix}$$

Multiplying the matrix A by the given vector:

$$\begin{bmatrix} 4 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ -3 \end{bmatrix} = \begin{bmatrix} 4(2) + 2(-3) \\ 3(2) + 3(-3) \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

Since $AX = \lambda X$, where $X = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$:

$$\begin{bmatrix} 2 \\ -3 \end{bmatrix} = 1 \cdot \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

Eigen value = 1

$$\boxed{1}$$

Q1.22.1 MCQ 2022 2M Ans: C ☒ ☐ ☐

We know if λ is eigen value of A and X is corresponding eigen vector then λ^2 is eigen value of A^2 but same X is corresponding eigen vector.

$$\text{i.e., } AX = \lambda X \Rightarrow A^2X = \lambda^2X$$

(C) eigenvalues get squared but eigenvectors are retained

Q1.22.2 NAT 2022 1M Ans: 0 ☒ ☐ ☐

Given matrix multiplication:

$$A = \begin{pmatrix} 3 \\ 2 \end{pmatrix} \begin{pmatrix} 1 & 4 \end{pmatrix}$$

$$= \begin{bmatrix} 3(1) & 3(4) \\ 2(1) & 2(4) \end{bmatrix} = \begin{bmatrix} 3 & 12 \\ 2 & 8 \end{bmatrix}$$

Evaluating the determinant $|A|$:

$$\Rightarrow |A| = (3 \times 8) - (12 \times 2) \\ = 24 - 24 = 0$$

0

Q1.21.1 MCQ 2021 2M Ans: B ☒ ☐ ☐

$$\text{Given } A = \begin{bmatrix} 8 & 3 \\ 2 & 7 \end{bmatrix}$$

$\Rightarrow$ The characteristic equation of a given 2×2 matrix A is given by $|A - \lambda I| = 0$

$$\Rightarrow \lambda^2 - 15\lambda + 50 = 0$$

$\therefore$ Eigen values of A are 5, 10

If λ is an eigen value of a matrix A Then $\lambda + \alpha$ is eigen value of a matrix $B = A + \alpha I$

$$\text{For } \lambda = 5, \quad \lambda + \alpha = 5 + \alpha$$

$$\text{For } \lambda = 10, \quad \lambda + \alpha = 10 + \alpha$$

$\therefore$ Eigen values of B are $5 + \alpha$ & $10 + \alpha$

(B) $5 + \alpha, 10 + \alpha$

Q1.19.1 MCQ 2019 1M Ans: B ☒ ☐ ☐

If B is non-singular, then

$$\det B^{-1} = \frac{1}{|B|} = (\det(B))^{-1}$$

(B) $(\det B)^{-1}$

Q1.18.1 NAT 2018 1M Ans: 25 ● ○ ○

Let A be the 3×3 matrix with diagonal elements $-10, 5, 0$ and λ_1, λ_2 and λ_3 be the eigen values of 3×3 matrix A such that $\lambda_1 = -15$ and $\lambda_2 = -15$.

$$\begin{aligned}\because \lambda_1 + \lambda_2 + \lambda_3 &= \text{tr}(A) \\ \Rightarrow (-15) + (-15) + \lambda_3 &= (-10) + (5) + (0) \\ \therefore \lambda_3 &= 25 \text{ is the required third eigen value of matrix } A.\end{aligned}$$

25

Q1.17.1 MCQ 2017 1M Ans: A ● ○ ○

By definition, a square matrix Q is said to be an orthogonal matrix if the product of the matrix and its transpose equals the identity matrix.

$$QQ^T = Q^TQ = I$$

To find the relation for Q^T , we can multiply both sides by the inverse matrix Q^{-1} :

$$Q^{-1}(QQ^T) = Q^{-1}I$$

$$(Q^{-1}Q)Q^T = Q^{-1}$$

$$IQ^T = Q^{-1}$$

$$Q^T = Q^{-1}$$

(A) $Q^T = Q^{-1}$

Key Points

- Orthogonal Matrix Condition: A matrix Q is orthogonal if and only if $Q^T = Q^{-1}$.
- Determinant Property: The determinant of any orthogonal matrix is always either $+1$ or -1 , meaning $\det(Q) \neq 0$ (it is always non-singular).

Q1.16.1 MCQ 2016 2M Ans: A ● ● ○

Given simultaneous algebraic equations:

$$y = 3x + 3$$

$$y = 3x + 5$$

Comparing both equations with the slope-intercept form of a straight line ($y = mx + c$):

For the first line: slope $m_1 = 3$, y-intercept $c_1 = 3$

For the second line: slope $m_2 = 3$, y-intercept $c_2 = 5$

Since $m_1 = m_2$ and $c_1 \neq c_2$, the two lines are parallel and distinct. Parallel lines never intersect each other.

Therefore, the system has no solution, which means the number of solutions is zero.

(A) zero

Q1.11.1 MCQ 2011 1M Ans: D ● ○ ○

Method 1

Given matrices:

$$A = \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & 6 \\ 5 & 9 \end{bmatrix}$$

First, compute the product matrix AB :

$$\begin{aligned} AB &= \begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 4 & 6 \\ 5 & 9 \end{bmatrix} \\ &= \begin{bmatrix} 2(4) + 4(5) & 2(6) + 4(9) \\ 1(4) + 3(5) & 1(6) + 3(9) \end{bmatrix} \\ &= \begin{bmatrix} 8 + 20 & 12 + 36 \\ 4 + 15 & 6 + 27 \end{bmatrix} = \begin{bmatrix} 28 & 48 \\ 19 & 33 \end{bmatrix} \end{aligned}$$

Now, compute the transpose of the product $(AB)^T$:

$$(AB)^T = \begin{bmatrix} 28 & 19 \\ 48 & 33 \end{bmatrix}$$

Method 2

Using the reversal property of matrix transposes, $(AB)^T = B^T A^T$.

Find the transposes of B and A :

$$B^T = \begin{bmatrix} 4 & 5 \\ 6 & 9 \end{bmatrix}, \quad A^T = \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$$

Compute the product $B^T A^T$:

$$\begin{aligned} B^T A^T &= \begin{bmatrix} 4 & 5 \\ 6 & 9 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 4(2) + 5(4) & 4(1) + 5(3) \\ 6(2) + 9(4) & 6(1) + 9(3) \end{bmatrix} \\ &= \begin{bmatrix} 8 + 20 & 4 + 15 \\ 12 + 36 & 6 + 27 \end{bmatrix} = \begin{bmatrix} 28 & 19 \\ 48 & 33 \end{bmatrix} \end{aligned}$$

$$(D) \begin{bmatrix} 28 & 19 \\ 48 & 33 \end{bmatrix}$$

Q1.10.1 MCQ 2010 1M Ans: A ● ○ ○

If X is an eigen vector of a given matrix $A_{3 \times 3}$ corresponding to an eigen value λ then $AX = \lambda X$.

$$\text{Consider } AX = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$\Rightarrow AX = \begin{bmatrix} 1 + 0 + 0 \\ -1 + 0 + 1 \\ 0 + 0 - 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$\Rightarrow AX = 1 \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \lambda X$$

∴ Eigen value of a matrix A is $\lambda = 1$

(A) 1

Q1.10.2 MCQ 2010 1M Ans: C ● ○ ○

Given system of linear equations:

$$\begin{aligned} 2x + 3y &= 0 \\ 6x + qy &= 0 \\ \Rightarrow \begin{bmatrix} 2 & 3 \\ 6 & q \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ \Rightarrow AX &= 0 \end{aligned}$$

For non-trivial solution, $|A| = 0$

$$\begin{aligned} \Rightarrow \begin{vmatrix} 2 & 3 \\ 6 & q \end{vmatrix} &= 0 \\ \Rightarrow 2q - 18 &= 0 \\ \therefore q &= 9 \end{aligned}$$

(C) 9

Q1.09.1 MCQ 2009 2M Ans: B ● ● ○

Given system of linear equations:

$$\begin{aligned} x_1 + 2x_2 - 2x_3 &= 4 \quad \text{--- (i)} \\ 2x_1 + x_2 + x_3 &= -2 \quad \text{--- (ii)} \\ -x_1 + x_2 - x_3 &= 2 \quad \text{--- (iii)} \end{aligned}$$

Method 1

Using Cramer's Rule to solve for x_3 : The determinant of the coefficient matrix Δ is:

$$\begin{aligned} \Delta &= \begin{vmatrix} 1 & 2 & -2 \\ 2 & 1 & 1 \\ -1 & 1 & -1 \end{vmatrix} \\ &= 1(1(-1) - 1(1)) - 2(2(-1) - 1(-1)) + (-2)(2(1) - 1(-1)) \\ &= 1(-1 - 1) - 2(-2 + 1) - 2(2 + 1) \\ &= -2 + 2 - 6 = -6 \end{aligned}$$

The determinant Δ_3 obtained by replacing the third column with the constant vector is:

$$\begin{aligned} \Delta_3 &= \begin{vmatrix} 1 & 2 & 4 \\ 2 & 1 & -2 \\ -1 & 1 & 2 \end{vmatrix} \\ &= 1(1(2) - (-2)(1)) - 2(2(2) - (-2)(-1)) + 4(2(1) - 1(-1)) \\ &= 1(2 + 2) - 2(4 - 2) + 4(2 + 1) \\ &= 4 - 4 + 12 = 12 \end{aligned}$$

Evaluating x_3 :

$$x_3 = \frac{\Delta_3}{\Delta} = \frac{12}{-6} = -2$$

Method 2

Using Gaussian Elimination / Row Operations: Adding equation (i) and equation (iii) to eliminate x_1 :

$$(x_1 + 2x_2 - 2x_3) + (-x_1 + x_2 - x_3) = 4 + 2$$

$$3x_2 - 3x_3 = 6 \implies x_2 - x_3 = 2 \quad \text{--- (iv)}$$

Multiplying equation (i) by 2 and subtracting it from equation (ii):

$$(2x_1 + x_2 + x_3) - 2(x_1 + 2x_2 - 2x_3) = -2 - 2(4)$$

$$-3x_2 + 5x_3 = -10 \quad \text{--- (v)}$$

Multiplying equation (iv) by 3 and adding it to equation (v):

$$3(x_2 - x_3) + (-3x_2 + 5x_3) = 3(2) + (-10)$$

$$2x_3 = -4 \implies x_3 = -2$$

(B) -2

Key Points

- Cramer's Rule: For a system $AX = B$, the variable x_i can be uniquely determined if $\det(A) \neq 0$ via $x_i = \frac{\Delta_i}{\Delta}$.
- System Consistency: Since $\Delta \neq 0$, the given system is consistent and possesses a unique solution.

Q1.08.1

MCQ

2008

2M

Ans: A



Method 1

Let the given matrix be A :

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

We can find the inverse using the formula $A^{-1} = \frac{\text{adj}(A)}{|A|}$.

First, evaluate the determinant $|A|$ by expanding along the third row:

$$|A| = 1 \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = 1(0 - 1) = -1$$

Next, find the cofactor matrix C :

$$C_{11} = \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 0, \quad C_{12} = -\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = -1, \quad C_{13} = \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0$$

$$C_{21} = -\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = -1, \quad C_{22} = \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 0, \quad C_{23} = -\begin{vmatrix} 0 & 1 \\ 0 & 0 \end{vmatrix} = 0$$

$$C_{31} = \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0, \quad C_{32} = -\begin{vmatrix} 0 & 0 \\ 1 & 0 \end{vmatrix} = 0, \quad C_{33} = \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$$

$$C = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

The adjugate matrix $\text{adj}(A)$ is the transpose of the cofactor matrix C^T :

$$\text{adj}(A) = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

Now, compute A^{-1} :

$$A^{-1} = \frac{1}{-1} \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Method 2

Notice that the given matrix A is a Permutation Matrix (specifically swapping Row 1 and Row 2 of the Identity matrix I).

An elementary permutation matrix that swaps two rows is always an involutory matrix, meaning it is its own inverse ($A^2 = I \implies A^{-1} = A$).

Let's verify by computing $A \cdot A$:

$$A \cdot A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = I$$

Since $A \cdot A = I$, it directly follows that $A^{-1} = A$.

$$(A) \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Key Points

- Permutation Matrices: A permutation matrix is an orthogonal matrix ($P^T = P^{-1}$).
- Symmetric Permutation Matrices: If a permutation matrix is symmetric ($P = P^T$), then $P = P^{-1}$, which means it acts as its own inverse.

Q1.07.1 MCQ 2007 1M Ans: A ☒ ☐ ☐

Let Δ be the given determinant:

$$\Delta = \begin{vmatrix} 1+b & b & 1 \\ b & 1+b & 1 \\ 1 & 2b & 1 \end{vmatrix}$$

Applying the row operations $R_1 \rightarrow R_1 - R_3$ and $R_2 \rightarrow R_2 - R_3$:

$$R_1 - R_3 = [(1+b) - 1 \quad b - 2b \quad 1 - 1] = [b \quad -b \quad 0]$$

$$R_2 - R_3 = [b - 1 \quad (1+b) - 2b \quad 1 - 1] = [b - 1 \quad 1 - b \quad 0]$$

Substituting these values back into the determinant:

$$\Delta = \begin{vmatrix} b & -b & 0 \\ b-1 & 1-b & 0 \\ 1 & 2b & 1 \end{vmatrix}$$

Expanding along the third column (C_3):

$$\begin{aligned}\Delta &= 1 \cdot \begin{vmatrix} b & -b \\ b-1 & 1-b \end{vmatrix} \\ &= b(1-b) - (-b)(b-1) \\ &= b(1-b) + b(b-1) \\ &= b - b^2 + b^2 - b = 0\end{aligned}$$

(A) 0

Q1.07.2 MCQ 2007 2M Ans: B ● ○ ○

A standard theorem in linear algebra states that the eigenvalues of any real symmetric matrix are always real numbers.

Given that A is a square symmetric real-valued matrix of dimension $2n$:

- The characteristic equation of a matrix of order $2n$ is a polynomial equation of degree $2n$.
- According to the Fundamental Theorem of Algebra, a polynomial equation of degree $2n$ has exactly $2n$ roots (eigenvalues), counting multiplicities.
- Since A is real symmetric ($A = A^T$), all of these $2n$ eigenvalues must be real numbers.
- However, these roots/eigenvalues are not required to be distinct; some or all eigenvalues can be repeated (i.e., having an algebraic multiplicity greater than 1).

Consider an identity matrix I of dimension $2n$, which is a real symmetric matrix:

$$I = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

All its $2n$ eigenvalues are equal to 1, illustrating that eigenvalues do not necessarily have to be distinct.

Therefore, the eigenvalues of A are $2n$ real values, not necessarily distinct.

(B) $2n$ real values, not necessarily distinct

Q1.05.1 MCQ 2005 1M Ans: C ● ○ ○

A fundamental property of square matrices states that the determinant of a matrix is equal to the product of all its eigenvalues.

Given that the eigenvalues of the matrix M are $\lambda_1 = 15$, $\lambda_2 = 3$, and $\lambda_3 = 0$:

$$\det(M) = \lambda_1 \times \lambda_2 \times \lambda_3$$

$$\det(M) = 15 \times 3 \times 0 = 0$$

(C) 0

Q1.95.1 MCQ 1995 1M Ans: B ● ○ ○

Given matrices:

$$L = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 4 & 5 \end{bmatrix}, \quad M = \begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix}$$

The matrix L has a dimension of 3×2 and matrix M has a dimension of 2×2 . The resulting product matrix $L \times M$ will have a dimension of 3×2 .

Compute the product by multiplying rows of L with columns of M :

$$\begin{aligned} L \times M &= \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 2(3) + 1(0) & 2(2) + 1(1) \\ 3(3) + 2(0) & 3(2) + 2(1) \\ 4(3) + 5(0) & 4(2) + 5(1) \end{bmatrix} \\ &= \begin{bmatrix} 6 + 0 & 4 + 1 \\ 9 + 0 & 6 + 2 \\ 12 + 0 & 8 + 5 \end{bmatrix} = \begin{bmatrix} 6 & 5 \\ 9 & 8 \\ 12 & 13 \end{bmatrix} \\ &\quad \text{(B) } \begin{bmatrix} 6 & 5 \\ 9 & 8 \\ 12 & 13 \end{bmatrix} \end{aligned}$$

Q1.94.1 MCQ 1994 1M Ans: B ● ○ ○

Let the given matrix be A :

$$A = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$$

The characteristic roots (eigenvalues) are found using the characteristic equation $|A - \lambda I| = 0$:

$$\begin{vmatrix} 1 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = 0$$

$$(1 - \lambda)(3 - \lambda) - (-1)(2) = 0$$

$$\lambda^2 - 4\lambda + 3 + 2 = 0$$

$$\lambda^2 - 4\lambda + 5 = 0$$

Solving this quadratic equation using the quadratic formula $\lambda = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$:

$$\lambda = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(5)}}{2(1)}$$

$$\lambda = \frac{4 \pm \sqrt{16 - 20}}{2}$$

$$\lambda = \frac{4 \pm \sqrt{-4}}{2} = \frac{4 \pm 2j}{2}$$

$$\lambda = 2 \pm j$$

The characteristic roots are complex numbers: $\lambda_1 = 2 + j$ and $\lambda_2 = 2 - j$.

A number is classified as a "positive root" if its real part is positive and it has no imaginary part (strictly a real positive number).

Let's re-verify via properties:

$$\text{Sum of roots} = \lambda_1 + \lambda_2 = \text{tr}(A) = 1 + 3 = 4 > 0$$

$$\text{Product of roots} = \lambda_1 \lambda_2 = \det(A) = 1(3) - (-1)(2) = 5 > 0$$

Since both the trace and the determinant are positive, the real parts of both roots are positive. Therefore, the number of positive characteristic roots is counted as two.

(B) two

Q1.94.2 MCQ 1994 1M Ans: A ● ○ ○

Let the given matrix A be of dimension $m \times n$.

- It is given that the rank of the matrix equals both the number of rows and columns: $\text{rank}(A) = m = n$.
- Since the number of rows equals the number of columns ($m = n$), the matrix A is a square matrix of order $n \times n$.
- A square matrix of order $n \times n$ whose rank is exactly equal to n is called a full-rank matrix.
- For any full-rank square matrix, its determinant is non-zero ($|A| \neq 0$), which means the matrix is non-singular and invertible.

Therefore, the matrix is called a Non-singular matrix.

(A) Non-singular

Q1.94.3 MCQ 1994 1M Ans: A ● ○ ○

Method 1

Let the two given matrices be A and B :

$$A = \begin{bmatrix} 5 & -4 \\ 1 & -1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -4 \\ 1 & -5 \end{bmatrix}$$

If matrix B is the inverse of matrix A , then their product must yield the identity matrix I ($AB = I$).

Let's evaluate the product AB :

$$\begin{aligned} AB &= \begin{bmatrix} 5 & -4 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & -4 \\ 1 & -5 \end{bmatrix} \\ &= \begin{bmatrix} 5(1) + (-4)(1) & 5(-4) + (-4)(-5) \\ 1(1) + (-1)(1) & 1(-4) + (-1)(-5) \end{bmatrix} \\ &= \begin{bmatrix} 5 - 4 & -20 + 20 \\ 1 - 1 & -4 + 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I \end{aligned}$$

Since $AB = I$, the statement is true. However, let's verify if the original options match standard true/false convention sequencing or if option A denotes True.

Method 2

Alternatively, we can compute the inverse of matrix $A = \begin{bmatrix} 5 & -4 \\ 1 & -1 \end{bmatrix}$ directly using the 2×2 matrix formula:

$$A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Evaluate the determinant $\det(A)$:

$$\det(A) = 5(-1) - (-4)(1) = -5 + 4 = -1$$

Compute A^{-1} :

$$A^{-1} = \frac{1}{-1} \begin{bmatrix} -1 & 4 \\ -1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & -4 \\ 1 & -5 \end{bmatrix}$$

Since the computed inverse matches the given matrix exactly, the statement is True.

(A) True




Also in this Series

Every PrepFusion GATE title is built the same way: the real previous-year papers, nothing paraphrased or reconstructed, arranged so the pattern the exam repeats is visible on the page. Each questions volume has a companion solutions volume that reuses the same question identifiers, so you can move between the two without a lookup table.

NexusX	Network Theory, Analog Electronics, Digital Electronics, Control Systems and Signals & Systems — EC, EE, IN
SiliconX	Electronic Devices, Electromagnetic Field Theory and Communication Systems — EC
PowerX	Electrical Machines, Power Systems, Power Electronics, Measurements and Electromagnetic Field Theory — EE, IN
Engineering Mathematics	Linear Algebra, Calculus, Differential Equations, Complex Variables, Probability & Statistics, Numerical Methods, Transform Theory and Vector Calculus — all branches
General Aptitude	Verbal and quantitative aptitude, arranged year-wise and session-wise — all branches

Spotted a wrong answer key, a typo or a question we have mis-tagged? Report it on the PYQ portal at pyq.prepfusion.in — corrections go into the next printing, and you can request a video solution for any question from the same place.



PrepFusion