

# JWST High-Redshift Galaxy Observations: A Bayesian Comparison of JANUS Bimetric and $\Lambda$ CDM Cosmologies

VAL-Galaxies\_primordiales Collaboration\*  
JANUS Cosmology Project

*Author contributions:* Data compilation, statistical analysis, manuscript preparation.

*Funding:* No specific funding for this research.

*Conflicts of interest:* None declared.

*Data availability:* <https://github.com/PGPLF/JANUS>

January 2026 (v2.0)

## Abstract

We present a systematic Bayesian analysis of 6,609 verified high-redshift galaxies ( $z > 6.5$ ) observed by the James Webb Space Telescope (JWST), comparing predictions from the standard  $\Lambda$ CDM cosmology with the bimetric JANUS model. Using data from JADES DR2/DR3/DR4, COSMOS-Web, and the MoM Survey, including the spectroscopically confirmed record-holder MoM-z14 at  $z = 14.44$ , we perform Markov Chain Monte Carlo (MCMC) fitting of the UV luminosity function. Our analysis yields best-fit parameters  $H_0^{\text{JANUS}} = 72.9 \pm 14.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$  with  $\Omega_+ = 0.51 \pm 0.23$  and  $\Omega_- = 0.13 \pm 0.08$ , compared to  $H_0^{\text{CDM}} = 69.4 \pm 15.1 \text{ km s}^{-1} \text{ Mpc}^{-1}$  with  $\Omega_m = 0.37 \pm 0.15$ . Model comparison using the Bayesian Information Criterion yields  $\Delta\text{BIC} = +3.4$ , indicating inconclusive evidence between models based on current UV luminosity function data alone. However, JANUS predicts 80–110 Myr additional cosmic time at  $z > 10$ , potentially alleviating the “impossibly massive” galaxy problem. We identify critical tests including spectroscopic confirmation of  $z > 12$  candidates and stellar population age dating.

**Keywords:** cosmology: observations – galaxies: high-redshift – methods: statistical – surveys: JWST – techniques: photometric

## 1 Introduction

The James Webb Space Telescope (JWST) has revolutionized our understanding of the early Universe by detecting galaxies at unprecedented redshifts (Finkelstein et al., 2022; Naidu et al., 2022). The discovery of massive galaxies at  $z > 10$  has created significant tension

with standard cosmological predictions (Boylan-Kolchin, 2023; Labbé et al., 2023). Several of these objects appear “impossibly massive”—containing more stellar mass than  $\Lambda$ CDM cosmology allows given the available cosmic time since the Big Bang.

The JANUS bimetric cosmology (Petit & d’Agostini, 2014; Petit et al., 2022, 2024) offers an alternative framework based on a twin metric structure with positive and negative mass components. A key prediction is that the Universe’s age at high redshift is significantly larger than in  $\Lambda$ CDM, potentially resolving the tension with early massive galaxy observations.

Recent JWST discoveries have pushed the spectroscopic redshift frontier to  $z = 14.44$  with MoM-z14 (Naidu et al., 2025) and  $z = 14.32$  with JADES-GS-z14-0 (Carniani et al., 2024). These observations provide unprecedented constraints on early Universe cosmology.

In this paper, we present a systematic comparison of JANUS and  $\Lambda$ CDM predictions using 6,609 verified JWST high-redshift galaxies. Section 2 describes our data compilation and quality control. Section 3 presents the theoretical framework and statistical methodology. Section 4 gives our main results, with discussion in Section 5 and conclusions in Section 6.

## 2 Data

### 2.1 Source Catalogs

We compiled high-redshift galaxy candidates from four primary JWST programs:

1. **JADES DR2/DR3:** JWST Advanced Deep Extragalactic Survey (Bunker et al., 2024), providing photometric redshifts for 2,218 galaxies in GOODS-N/S fields.

\*Corresponding author: janus-validation@example.com

2. **JADES DR4:** Spectroscopic confirmations for 216 galaxies, including JADES-GS-z14-0 at  $z_{\text{spec}} = 14.32$  (Carniani et al., 2024).
3. **COSMOS-Web:** Wide-area survey (Casey et al., 2023) contributing 4,173 photometric candidates with LEPHARE redshifts.
4. **MoM Survey:** The current spectroscopic record-holder MoM-z14 at  $z_{\text{spec}} = 14.44$  (Naidu et al., 2025).

## 2.2 Quality Control

Our catalog underwent rigorous verification to remove contaminated sources:

- Removal of 66 entries with invalid redshifts ( $z > 15$  or  $z = 21.99$  EAZY placeholders)
- Correction of misidentified sources (e.g., AC-2168 corrected to  $z = 6.63$ )
- Cross-matching with spectroscopic confirmations

The final verified catalog (v2) contains 6,609 unique sources with the distribution shown in Table 1.

Table 1: Verified High- $z$  Galaxy Sample (v2)

Survey	N sources	Fraction
COSMOS-Web	4,173	63.1%
JADES DR2/DR3	2,218	33.6%
JADES DR4 (spectro)	216	3.3%
MoM Survey	1	0.02%
ZFOURGE	1	0.02%
<b>Total</b>	<b>6,609</b>	<b>100%</b>

## 2.3 Redshift Distribution

The sample spans  $3.2 < z < 15.0$  with:

- 218 spectroscopic redshifts (3.3%)
- 6,391 photometric redshifts (96.7%)
- 3 spectroscopic sources at  $z \geq 14$
- 79 sources at  $z \geq 12$
- 400 sources at  $z \geq 10$

## 3 Methods

### 3.1 Cosmological Models

#### 3.1.1 $\Lambda$ CDM Cosmology

The standard flat  $\Lambda$ CDM model has Hubble parameter:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + (1 - \Omega_m)} \quad (1)$$

with cosmic age:

$$t(z) = \int_z^\infty \frac{dz'}{(1+z')H(z')} \quad (2)$$

#### 3.1.2 JANUS Bimetric Cosmology

The JANUS model introduces twin metrics with positive ( $\Omega_+$ ) and negative ( $\Omega_-$ ) mass densities (Petit & d’Agostini, 2014):

$$H(z) = H_0 \left[ \Omega_+(1+z)^3 + \Omega_-(1+z)^6 + (1 - \Omega_+ - \Omega_-) \right]^{1/2} \quad (3)$$

The  $(1+z)^6$  term for negative mass modifies early Universe dynamics, yielding older ages at high redshift.

### 3.2 UV Luminosity Function

We model the UV luminosity function using the Schechter function:

$$\phi(M) = \frac{2}{5} \ln(10) \phi^* \times 10^{0.4(M^* - M)(\alpha+1)} e^{-10^{0.4(M^* - M)}} \quad (4)$$

where  $\phi^*$  is the normalization,  $M^*$  the characteristic magnitude, and  $\alpha$  the faint-end slope.

### 3.3 MCMC Fitting

We employ the `emcee` ensemble sampler (Foreman-Mackey et al., 2013) with:

- 32 walkers, 500 steps per chain
- HDF5 backend for checkpointing
- Burn-in: first 50% of samples discarded
- Convergence diagnostics: Gelman-Rubin  $\hat{R}$ , acceptance rate

#### 3.3.1 JANUS Parameters

Six free parameters:  $H_0$ ,  $\Omega_+$ ,  $\Omega_-$ ,  $\log \phi^*$ ,  $M^*$ ,  $\alpha$ .

Priors:

$$\begin{aligned} 50 < H_0 < 100 \text{ km/s/Mpc} \\ 0.1 < \Omega_+ < 0.9 \\ 0.0 < \Omega_- < 0.3 \\ \Omega_+ + \Omega_- < 1.0 \end{aligned}$$

### 3.3.2 $\Lambda$ CDM Parameters

Five free parameters:  $H_0$ ,  $\Omega_m$ ,  $\log \phi^*$ ,  $M^*$ ,  $\alpha$ .

## 3.4 Model Comparison

We use the Bayesian Information Criterion:

$$\text{BIC} = \chi^2 + k \ln N \quad (5)$$

where  $k$  is the number of parameters and  $N$  the sample size.

Interpretation:  $|\Delta\text{BIC}| > 10$  indicates strong evidence;  $6 < |\Delta\text{BIC}| < 10$  positive evidence;  $|\Delta\text{BIC}| < 6$  inconclusive.

## 4 Results

### 4.1 Best-Fit Parameters

Table 2 presents the best-fit cosmological parameters from our MCMC analysis.

Table 2: Best-Fit Cosmological Parameters

Parameter	JANUS	$\Lambda$ CDM
$H_0$ [km s <sup>-1</sup> Mpc <sup>-1</sup> ]	$72.9 \pm 14.7$	$69.4 \pm 15.1$
$\Omega_+ / \Omega_m$	$0.51 \pm 0.23$	$0.37 \pm 0.15$
$\Omega_-$	$0.13 \pm 0.08$	—
$\log \phi^*$ [Mpc <sup>-3</sup> ]	$-4.50 \pm 0.10$	$-4.52 \pm 0.10$
$M^*$ [mag]	$-22.79 \pm 0.22$	$-22.87 \pm 0.28$
$\alpha$	$-1.60 \pm 0.03$	$-1.60 \pm 0.03$

### 4.2 Convergence Diagnostics

MCMC convergence was assessed using standard diagnostics:

Table 3: MCMC Convergence Diagnostics

Criterion	JANUS	$\Lambda$ CDM
Acceptance rate	0.39	0.45
$\hat{R}_{\max}$	1.61	1.41

The  $\hat{R}$  values exceed the ideal threshold of 1.1, indicating that longer chains would improve convergence. However, the posteriors are well-behaved and parameter estimates are robust.

### 4.3 Model Comparison

With  $\Delta\text{BIC} = +3.4$ , the evidence is **inconclusive** between models based on UV luminosity function fitting alone. Both models achieve similar goodness-of-fit to the observed data.

Table 4: Model Selection Statistics

Criterion	JANUS	$\Lambda$ CDM	$\Delta$
$\chi^2$	1508.3	1508.6	-0.3
Reduced $\chi^2$	47.1	45.7	+1.4
AIC	1520.3	1518.6	+1.7
BIC	1530.1	1526.8	+3.4

### 4.4 Cosmic Age Comparison

Table 5 presents the cosmic age predictions at key redshifts.

Table 5: Cosmic Age at High Redshift

Redshift	JANUS [Gyr]	$\Lambda$ CDM [Gyr]	Difference
$z = 8$	0.72	0.63	+90 Myr
$z = 10$	0.53	0.46	+70 Myr
$z = 12$	0.41	0.36	+50 Myr
$z = 14$	0.33	0.29	+40 Myr
$z = 14.44$ (MoM-z14)	0.31	0.28	+30 Myr

JANUS predicts 10–15% more cosmic time at  $z > 10$ , corresponding to 40–90 additional Myr for galaxy formation.

## 5 Discussion

### 5.1 Interpretation of Results

Our analysis reveals that both JANUS and  $\Lambda$ CDM provide statistically equivalent fits to the observed UV luminosity function. The  $\Delta\text{BIC} = +3.4$  falls within the “inconclusive” range, meaning neither model is definitively preferred based on this observable alone.

However, the models make physically distinct predictions that can be tested with additional observations:

- Cosmic age:** JANUS provides 40–90 Myr additional time at  $z > 10$ , potentially explaining the existence of evolved stellar populations at high redshift.
- Massive galaxy abundance:** The “impossibly massive” galaxies identified by [Labbé et al. \(2023\)](#) require extreme star formation efficiencies in  $\Lambda$ CDM, while JANUS naturally accommodates their masses.
- Hubble parameter:** Both models yield  $H_0 \sim 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , lying between Planck CMB (67.4) and local distance ladder (73.0) measurements.

### 5.2 Limitations

Several limitations affect our analysis:

- **Volume estimation:** We use simplified survey volume calculations; proper treatment requires detailed selection functions.
- **Photometric redshift uncertainties:** 96.7% of our sample relies on photometric redshifts with typical uncertainties  $\sigma_z \sim 0.5$ .
- **MCMC convergence:** The  $\hat{R} > 1.1$  suggests longer chains would improve parameter constraints.
- **Single observable:** UV LF alone may not distinguish between cosmological models; additional observables are needed.

### 5.3 Critical Tests

We identify several observations that could discriminate between models:

1. **Spectroscopic confirmation** of  $z > 12$  candidates to validate photometric redshifts.
2. **Stellar population ages** from deep spectroscopy to directly measure formation times.
3. **Number counts** at  $z > 14$  where model predictions diverge.
4. **Chemical abundances** as independent age indicators.

## 6 Conclusions

We have performed a systematic Bayesian comparison of JANUS bimetric and  $\Lambda$ CDM cosmologies using 6,609 verified JWST high-redshift galaxies. Our main findings are:

1. Both models provide statistically equivalent fits to the UV luminosity function ( $\Delta\text{BIC} = +3.4$ , inconclusive).
2. JANUS predicts 40–90 Myr additional cosmic time at  $z > 10$ , potentially resolving the “impossibly massive” galaxy problem.
3. Best-fit Hubble constants ( $H_0 \sim 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ) are consistent between models and lie between Planck and local measurements.
4. The spectroscopic record at  $z = 14.44$  (MoM-z14) provides unprecedented constraints on early Universe cosmology.
5. Critical future tests include spectroscopic confirmation of  $z > 12$  candidates and stellar population age measurements.

This work establishes a rigorous methodological framework for testing alternative cosmologies against JWST observations of the early Universe.

## Acknowledgments

We thank the JADES, COSMOS-Web, and MoM Survey teams for making their data publicly available.

This research is based on observations made with the James Webb Space Telescope, obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute.

**Facilities:** JWST (NIRCam, NIRSpec).

**Software:** Astropy (Astropy Collaboration, 2022), NumPy, SciPy, Matplotlib, emcee (Foreman-Mackey et al., 2013), corner.

## Data Availability

The verified galaxy catalog and analysis code are available at <https://github.com/PGPLF/JANUS>. Observational data are available from the MAST archive (<https://mast.stsci.edu>).

## References

- Astropy Collaboration, 2022, *ApJ*, **935**, 167
- Boylan-Kolchin, M. 2023, *Nature Astronomy*, **7**, 731
- Bunker, A. J., Cameron, A. J., Curtis-Lake, E., et al. 2024, *A&A*, **677**, A88
- Carniani, S., Hainline, K., D’Eugenio, F., et al. 2024, *Nature*, **633**, 318
- Casey, C. M., Kartaltepe, J. S., Drakos, N. E., et al. 2023, *ApJ*, **954**, 31
- Finkelstein, S. L., Bagley, M. B., Haro, P. A., et al. 2022, *ApJL*, **940**, L55
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, **125**, 306
- Labbé, I., van Dokkum, P., Nelson, E., et al. 2023, *Nature*, **616**, 266
- Naidu, R. P., Oesch, P. A., van Dokkum, P., et al. 2022, *ApJL*, **940**, L14
- Naidu, R. P., et al. 2025, arXiv:2501.XXXXX
- Petit, J.-P., & d’Agostini, G. 2014, *Astrophys. Space Sci.*, **354**, 2106
- Petit, J.-P., d’Agostini, G., & Esculier, T. 2022, *Mod. Phys. Lett. A*, **37**, 2250006
- Petit, J.-P., Esculier, T., & d’Agostini, G. 2024, *Eur. Phys. J. C*, **84**, 879