

The Effect of Skooma Exposure on Count Choculitis

A fictional randomized crossover study for document extraction testing

Abstract


Background: Count Choculitis is a seasonal public-health nuisance characterized by nocturnal cough, velvet rash, and recurrent cocoa fever. We evaluated whether carefully titrated skooma changed symptom burden over four observation weeks in a deliberately synthetic population.

Methods: Ninety-six volunteer counts were assigned to low-dose skooma syrup or matched placebo tonic. Outcomes were fabricated but internally consistent so that sentence, page, coordinate, table, and figure extractors can be evaluated against predictable content.

Table 1. Baseline characteristics of fictional participants.

Characteristic	Skooma (n=48)	Placebo (n=48)
Mean age, centuries	3.4	3.5
Female, n (%)	21 (44)	22 (46)
Prior cocoa fever, n (%)	39 (81)	38 (79)
Baseline symptom index	62.1	61.8

Figure 1. Participant flow from screening through randomized allocation.





Results

The skooma arm showed a larger fictional reduction in the Count Choculitis symptom index than placebo. The most consistent changes were observed for night cough and velvet rash; cocoa fever varied by village.

Table 2. Change in Count Choculitis outcomes after four weeks.

Outcome	Skooma mean change	Placebo mean change	Difference
Total symptom index	-18.4	-6.2	-12.2
Night cough episodes	-7.1	-2.4	-4.7
Velvet rash score	-4.8	-1.9	-2.9
Cocoa fever days	-1.2	-0.9	-0.3

Figure 2. Dose-response pattern for fictional skooma exposure quaFigure 3. Forest-style summary of fabricated effect estimates.



Discussion

This fake study should not be interpreted clinically. Its purpose is to provide a compact PDF fixture with coherent prose, numbered tables, numbered figures, captions, and vector geometry suitable for testing GROBID and Azure Document Intelligence extraction pipelines.

Funding: None. Conflicts: All characters and conditions are synthetic.