

# JOIN



Telegram  
@PuneEngineers

For more Subjects

<https://www.studymedia.in/fe/notes>



SCAN ME





Sinhgad Institutes

# Partial Differentiation Introduction & Definition

Subject- Engg.Mathematcs-I

Faculty- Prof.Ms.S.A.Gurav

Sinhgad College Of Engineering

Other Subjects: <https://www.studymedia.in/fe/notes>



Sinhgad Institutes

## CONTENT OF THE TOPIC

- 1.Introduction to Partial Derivative, Definition, Physical Meaning and basic examples.
2. Properties of Partial Derivative and Higher order derivatives.
- 3.Derivative of Composite function
- 4.Variable to be treated as constant.
- 5.Homogeneous Function and Euler's Theorem.
- 6.Partial Derivative of Implicit function& Total Derivative.



Sinhgad Institutes

# PARTIAL DERIVATIVES

**Introduction** – Partial Derivative is used for functions with more than one independent variable.

If the function  $Z = f(x, y)$  be a function of two variables  $x, y$  then the rate of change of  $z$  with respect to  $x$ , keeping  $y$  const. is called partial derivative of  $z$  with respect to  $x$  & it is denoted by  $\frac{\partial z}{\partial x}$ .

Similarly the rate of change of  $z$  with respect to  $y$ , keeping  $x$  const. is called partial derivative of  $z$  with respect to  $y$  & it is denoted by  $\frac{\partial z}{\partial y}$ .

Thus for  $Z = f(x, y)$   $\frac{\partial z}{\partial x}$ ,  $\frac{\partial z}{\partial y}$  are called First Order Partial derivatives with respect to  $x$  &  $y$  respectively.

Other Subjects: <https://www.studymedia.in/fe/notes>



Sinhgad Institutes

Thus 
$$\frac{\partial z}{\partial x} = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

and 
$$\frac{\partial z}{\partial y} = \lim_{k \rightarrow 0} \frac{f(x, y+k) - f(x, y)}{k}$$

In general  $\frac{\partial z}{\partial x}$   $\frac{\partial z}{\partial y}$  are function of both the variables x

and y so we may obtain higher derivatives

Other Subjects: <https://www.studymedia.in/fe/notes>



Sinhgad Institutes

# GEOMETRICAL INTERPRETATION

Geometrically  $z=f(x, y)$  represents a surface  $S$ , so if  $y=k$  (constant) i.e. a plane parallel to  $XZ$  plane thus


$z=f(x, y)$  and  $y=k$  (constant) together represents a curve  $C$  which is the section of  $S$  by  $y=k$  plane.

Thus  $\frac{\partial z}{\partial x}$  represents the slope of tangent to  $C$  at  $(x, k, z)$

Similarly  $\frac{\partial z}{\partial y}$  represents the slope of tangent drawn to the curve of intersection of  $z=f(x, y)$  &  $x=k$ .



Sinhgad Institutes



If  $f(a,b)=c$  then pt  $P(a,b,c)$  lies on  $S$

$y=b$  intersects surface  $S$  in  $C_1$

$T_1$  is the slope of tangent to curve  $C_1$  i.e.  $\frac{\partial z}{\partial x}$

Similarly  $T_2$  is the slope of tangent to curve  $C_2$  i.e.  $\frac{\partial z}{\partial y}$



Sinhgad Institutes

# Examples on Direct Partial Derivatives

Q1. If  $x = r \cos \theta$ ,  $y = r \sin \theta$  show that

$$\left( \frac{\partial r}{\partial x} \right)^2 + \left( \frac{\partial r}{\partial y} \right)^2 = 1$$

Sol  $x = r \cos \theta$ ,  $y = r \sin \theta$  then

$$(x)^2 + (y)^2 = r^2 \quad \text{therefore}$$

$$2r \frac{\partial r}{\partial x} = 2x \quad \& \quad 2r \frac{\partial r}{\partial y} = 2y$$

$$\text{so } \frac{\partial r}{\partial x} = \frac{x}{r} = \frac{r \cos \theta}{r} = \cos \theta \quad \& \quad \frac{\partial r}{\partial y} = \frac{y}{r} = \frac{r \sin \theta}{r} = \sin \theta$$

$$\text{thus } \left( \frac{\partial r}{\partial x} \right)^2 + \left( \frac{\partial r}{\partial y} \right)^2 = \cos^2 \theta + \sin^2 \theta = 1$$





Sinhgad Institutes

# Particular Example

1) If  $z^3 - zx - y = 4$  then find  $\frac{\partial z}{\partial x}$  &  $\frac{\partial z}{\partial y}$

Solution : Given  $z^3 - zx - y = 4$ ----(1)

Differentiate eq(1) partially w.r.t. x keeping y as constant

$$3z^2 \frac{\partial z}{\partial x} - z - x \frac{\partial z}{\partial x} = 0$$

$$(3z^2 - x) \frac{\partial z}{\partial x} = z$$

$$\frac{\partial z}{\partial x} = \frac{z}{3z^2 - x}$$

Also, differentiate eq (1) partially w.r.t. y, keeping x as constant



Sinhgad Institutes

$$\left( 3z^2 \frac{\partial z}{\partial y} - x \frac{\partial z}{\partial y} - 1 \right) = 0$$

$$(3z^2 - x) \frac{\partial z}{\partial y} = 1$$

$$\frac{\partial z}{\partial y} = \frac{1}{(3z^2 - x)}$$

3) 1) If  $u = \frac{1}{\sqrt{x^2 + y^2 + z^2}}$  then find the value of

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$$



Sinhgad Institutes

Solution:

$$\frac{\partial u}{\partial x} = \frac{\partial}{\partial x} [(x^2 + y^2 + z^2)^{-1/2}]$$

$$\frac{\partial u}{\partial x} = \frac{-1}{2} \frac{2x}{(x^2 + y^2 + z^2)^{3/2}}$$

$$\frac{\partial u}{\partial x} = \frac{-x}{(x^2 + y^2 + z^2)^{3/2}}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial u}{\partial x} \right) = \frac{\partial}{\partial x} \left( \frac{-x}{(x^2 + y^2 + z^2)^{3/2}} \right)$$



Sinhgad Institutes

$$\frac{\partial^2 u}{\partial x^2} = (-x) \left( \frac{-3}{2} \right) (x^2 + y^2 + z^2)^{-5/2} + (-1)(x^2 + y^2 + z^2)^{-3/2}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{3x^2}{(x^2 + y^2 + z^2)^{5/2}} - \frac{1}{(x^2 + y^2 + z^2)^{3/2}}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{2x^2 - y^2 - z^2}{(x^2 + y^2 + z^2)^{5/2}}$$

Similarly

$$\frac{\partial^2 u}{\partial y^2} = \frac{-x^2 + 2y^2 - z^2}{(x^2 + y^2 + z^2)^{5/2}}$$



Sinhgad Institutes

$$\frac{\partial^2 u}{\partial z^2} = \frac{-x^2 - y^2 + 2z^2}{(x^2 + y^2 + z^2)^{5/2}}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} = \frac{2x^2 - y^2 - z^2 - x^2 + 2y^2 - z^2 - x^2 - y^2 + 2z^2}{(x^2 + y^2 + z^2)^{5/2}} = 0$$

3) If  $u = \log(\tan x + \tan y + \tan z)$  then prove that  $\sin 2x \frac{\partial u}{\partial x} + \sin 2y \frac{\partial u}{\partial y} + \sin 2z \frac{\partial u}{\partial z} = 2$



Sinhgad Institutes

Solution : Given  $u = \log(\tan x + \tan y + \tan z)$

$$\frac{\partial u}{\partial x} = \frac{\sec^2 x}{\tan x + \tan y + \tan z}$$

Similarly

$$\frac{\partial u}{\partial y} = \frac{\sec^2 y}{\tan x + \tan y + \tan z}$$

$$\frac{\partial u}{\partial z} = \frac{\sec^2 z}{\tan x + \tan y + \tan z}$$



Sinhgad Institutes

Now

$$\begin{aligned} & \sin 2x \frac{\partial u}{\partial x} + \sin 2y \frac{\partial u}{\partial y} + \sin 2z \frac{\partial u}{\partial z} \\ &= \frac{\sin 2x \sec^2 x + \sin 2y \sec^2 y + \sin 2z \sec^2 z}{\tan x + \tan y + \tan z} \\ &= \frac{\frac{2 \sin x \cos x}{\cos^2 x} + \frac{2 \sin y \cos y}{\cos^2 y} + \frac{2 \sin z \cos z}{\cos^2 z}}{\tan x + \tan y + \tan z} \\ &= \frac{2 \tan x + 2 \tan y + 2 \tan z}{\tan x + \tan y + \tan z} = 2 \end{aligned}$$



Sinhgad Institutes

4) If  $u = \phi(x + ay) + \varphi(x - ay)$  then show that  $\frac{\partial^2 u}{\partial y^2} = a^2 \frac{\partial^2 u}{\partial x^2}$

Solution:  $\frac{\partial u}{\partial y} = \phi'(x + ay) \cdot a + \varphi'(x - ay) \cdot (-a)$

Again differentiating w.r.t.  $y$

$$\begin{aligned} \frac{\partial^2 u}{\partial y^2} &= \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial y} \right) \\ &= \frac{\partial}{\partial y} (\phi'(x + ay) \cdot a + \varphi'(x - ay) \cdot (-a)) \end{aligned}$$





Sinhgad Institutes

$$= a. \phi''(x + ay).a + (-a)\phi''(x - ay)(-a)$$

$$= a^2(\phi''(x + ay) + \phi''(x - ay))$$

$$\text{Now } \frac{\partial u}{\partial x} = \phi'(x + ay) + \phi'(x - ay)$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial u}{\partial x} \right) = \phi''(x + ay). + \phi''(x - ay)$$

$$\therefore \frac{\partial^2 u}{\partial y^2} = a^2 \frac{\partial^2 u}{\partial x^2}$$



Sinhgad Institutes

## Partial derivative of higher order:-

Let  $z = f(x, y)$ , then  $\frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial x} \right) = z_{xx}$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial z}{\partial y} \right) = z_{xy} \quad \frac{\partial^2 z}{\partial y \partial x} = \frac{\partial}{\partial y} \left( \frac{\partial z}{\partial x} \right) = z_{yx}$$

In general  $\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$

Order of differentiation is commutative where  $z$  is a continuous function

Therefore further differentiation of any of these w .r. t either  $x$  or  $y$  or both possible.



Sinhgad Institutes

## To find the value of Parameter 'n'

Ex.1) Find the value of 'n' for which  $u = A e^{-gx} \sin(nt - gx)$  satisfies the partial differential equation  $\frac{\partial u}{\partial t} = m \frac{\partial^2 u}{\partial x^2}$  where A, g, m are constant.

Solution : Given  $u = A e^{-gx} \sin(nt - gx)$

Differentiating w.r.t. 't' keeping x constant

$$\frac{\partial u}{\partial t} = A e^{-gx} \cos(nt - gx)(n)$$

Now differentiating w.r.t. 'x' keeping t constant



Sinhgad Institutes

Now differentiating.w.r.t. 'x' keeping t constant

$$\frac{\partial u}{\partial x} = Ae^{-gx}(-g) \sin(nt - gx) + Ae^{-gx} \cdot \cos(nt - gx)(-g)$$

Again differentiating w.r.t. x

$$\begin{aligned} \frac{\partial^2 u}{\partial x^2} = & A(-g)e^{-gx}(-g) \cdot \sin(nt - gx) + \\ & Ae^{-gx}(-g) \cdot \cos(nt - gx)(-g) + \\ & Ae^{-gx}(-g) \cos(nt - gx)(-g) + Ae^{-gx}(-\sin(nt - \\ & gx))(-g)(-g) \end{aligned}$$



Sinhgad Institutes

$$= A g^2 e^{-gx} [\sin(nt - gx) + \cos(nt - gx) + \cos(nt - gx) - \sin(nt - gx)]$$

$$\frac{\partial^2 u}{\partial x^2} = A g^2 e^{-gx} 2 \cos(nt - gx)$$

Now as  $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$

$$A e^{-gx} \cos(nt - gx)(n) = m A g^2 e^{-gx} 2 \cos(nt - gx)$$

$$\text{Thus, } n = 2mg^2$$



Sinhgad Institutes

Q2 Find 'n' such that  $v = r^n(3 \cos^2 \theta - 1)$

satisfies the equation  $\frac{\partial}{\partial r} \left( r^2 \frac{\partial v}{\partial r} \right) +$

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial v}{\partial \theta} \right) = 0$$

*solution:*

$$v = r^n(3 \cos^2 \theta - 1) \therefore \frac{\partial v}{\partial r} = nr^{n-1}(3 \cos^2 \theta - 1).$$

$$\therefore \frac{\partial}{\partial r} \left( r^2 \frac{\partial v}{\partial r} \right) = n(n+1)v$$

$$\text{Also } \frac{\partial v}{\partial \theta} = r^n[3 \cdot 2 \cos^1 \theta (-\sin \theta)].$$

$$\therefore \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial v}{\partial \theta} \right) = -6r^n[-\sin \theta \sin^2 \theta + \cos \theta \cdot 2 \sin \theta \cos \theta]$$



Sinhgad Institutes

$$\therefore \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left( \sin\theta \frac{\partial v}{\partial\theta} \right) = -6r^n [2 \cos^2 \theta - \sin^2 \theta]$$

$$= -6r^n [3 \cos^2 \theta - 1] = -6v$$

$$\therefore \frac{\partial}{\partial r} \left( r^2 \frac{\partial v}{\partial r} \right) + \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left( \sin\theta \frac{\partial v}{\partial\theta} \right) = 0$$

$$\Rightarrow n(n+1)v - 6v = 0$$

$$\Rightarrow n^2 + n - 6 = 0, v \neq 0$$

$$\Rightarrow n = -3, 2.$$

## Example on verification of $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$

Ex.1) If  $u = x^y + y^x$  then prove that  $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$

Solution :  $u = x^y + y^x$ -----(1)

Differentiating equation (1) partially w.r.t.  $y$  keeping 'x' as constant

$$\frac{\partial u}{\partial y} = x^y \log x + xy^{x-1}$$

Differentiating above partially w.r.t.  $x$  keeping  $y$  as constant





Sinhgad Institutes

$$\begin{aligned}\frac{\partial^2 u}{\partial x \partial y} &= \frac{\partial}{\partial x} \left( \frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial x} [x^y \log x + xy^{x-1}] \\ &= x^y \frac{1}{x} + \log x \cdot yx^{y-1} + xy^{x-1} \log y + y^{x-1} \\ \frac{\partial^2 u}{\partial x \partial y} &= x^{y-1} + yx^{y-1} \log x + xy^{x-1} \log y + y^{x-1}\end{aligned}$$

Now diff.eq(1) partially w.r.t. x keeping 'y' as constant

$$\frac{\partial u}{\partial x} = yx^{y-1} + y^x \log y$$

Again diff. above w.r.t.y keeping x as constant



Sinhgad Institutes

$$\begin{aligned}\frac{\partial^2 u}{\partial y \cdot \partial x} &= \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial x} \right) = \frac{\partial}{\partial y} [y x^{y-1} + y^x \log y] \\ &= y x^{y-1} \log x + x^{y-1} + y^x \frac{1}{y} + \log y \cdot x \cdot y^{x-1} \\ &= y x^{y-1} \log x + x^{y-1} + y^{x-1} + x y^{x-1} \log y \text{ -----(B)}\end{aligned}$$

From (A) and (B)

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}$$



Sinhgad Institutes

# Application Of P D

Partial Differentiation is used in the problems of Jacobian, vibration of string, Theory of approximation, Maxima & Minima .

Boundary value problems, Laplace Equation, Vectors, Heat ,Wave equations & solving some problems in Electrical engineering.

PDEs can be used to describe a wide variety of phenomena such as Sound ,Heat, Electrostatics, Electrodynamics , Fluid flow, Elasticity or Quantum Mechanics



Sinhgad Institutes

## University Questions

1) If  $u = \log(x^3 + y^3 - x^2y - y^2x)$  then Show that

$$\frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial^2 u}{\partial y^2} = -\frac{4}{(x+y)^2}$$

2) If  $u = \log(x^2 + y^2)$ , verify  $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial x \partial y}$

3) If  $\theta = t^n \cdot e^{\frac{-r^2}{4t}}$  Find  $n$  such that  $\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \theta}{\partial r} \right) = \frac{\partial \theta}{\partial t}$

4) Prove that at a point of surface  $x^x y^y z^z = c$  where  $x = y = z$ ,  $\frac{\partial^2 z}{\partial x \partial y} =$


$$-(x \log x)^{-1}$$




Sinhgad Institutes

# Partial Derivative of Composite Function

- If  $u$  is a function of  $r$  i.e.  $u=f(r)$  and  $r$  again is a function of two independent variables  $x, y$  i.e.  $r=g(x, y)$ . Thus  $u$  becomes a composite function of  $x$  and  $y$ .

- $$u \rightarrow r \rightarrow (x, y) \therefore \frac{\partial u}{\partial x} = \frac{du}{dr} \cdot \frac{\partial r}{\partial x} \quad , \quad \frac{\partial u}{\partial y} = \frac{du}{dr} \cdot \frac{\partial r}{\partial y}$$


- $$u \rightarrow r \rightarrow (x, y, z) \therefore \frac{\partial u}{\partial x} = \frac{du}{dr} \cdot \frac{\partial r}{\partial x} \quad \frac{\partial u}{\partial y} = \frac{du}{dr} \cdot \frac{\partial r}{\partial y}$$

$$\frac{\partial u}{\partial z} = \frac{du}{dr} \cdot \frac{\partial r}{\partial z}$$




Sinhgad Institutes

$$z \rightarrow x, y \rightarrow u, v$$

then  $z$  is a composite function of two variables  $u$  &  $v$

$$\therefore \frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial u} \quad \frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v}$$

- $l \rightarrow x, y, z \rightarrow u, v, w$

$$\therefore \frac{\partial l}{\partial u} = \frac{\partial l}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial l}{\partial y} \frac{\partial y}{\partial u} + \frac{\partial l}{\partial z} \frac{\partial z}{\partial u} \text{ \&so on..}$$

- Note: For a function of single variable ,derivative is total whereas for a function of more than one variable derivative is Partial.



Sinhgad Institutes

## Example on Composite Function

$$1) \text{ If } z = f(u, v) \text{ and } u = x \cos t - y \sin t;$$
$$v = x \sin t + y \cos t$$

where  $t$  is a constant Then show that

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = u \frac{\partial z}{\partial u} + v \frac{\partial z}{\partial v}$$

Solution :  $z = f(u, v)$  and

$$u = x \cos t - y \sin t \text{-----}(1)$$

$$v = x \sin t + y \cos t \text{-----}(2)$$

## Differentiating $z$ w.r.t. $x$ keeping $y$ constant

$$\left(\frac{\partial z}{\partial x}\right) = \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial x}$$

&

$$\left(\frac{\partial z}{\partial y}\right) = \frac{\partial z}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial y}$$

Differentiating equation (1) w.r.t. ' $x$ ' keeping  $t$  as constant

$$\left(\frac{\partial u}{\partial x}\right) = \frac{\partial}{\partial x} (x \cos t - y \sin t) = \cos t$$

$$\left(\frac{\partial u}{\partial y}\right) = \frac{\partial}{\partial y} (x \cos t - y \sin t) = -\sin t$$

$$\text{Now } \left(\frac{\partial v}{\partial x}\right) = \sin t ; \left(\frac{\partial v}{\partial y}\right) = \cos t$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial u} (\cos t) + \frac{\partial z}{\partial v} (\sin t) \text{-----(A)}$$

$$\left(\frac{\partial z}{\partial y}\right) = \frac{\partial z}{\partial u} (-\sin t) + \frac{\partial z}{\partial v} (\cos t) \text{-----(B)}$$

- Multiplying (A) by x & (B) by y

$$\begin{aligned}
 x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} &= x \cos t \frac{\partial z}{\partial u} + x \sin t \frac{\partial z}{\partial v} - y \sin t \frac{\partial z}{\partial u} \\
 &+ y \cos t \frac{\partial z}{\partial v} \\
 &= \frac{\partial z}{\partial u} (x \cos t - y \sin t) + \frac{\partial z}{\partial v} (x \sin t + y \cos t)
 \end{aligned}$$

$$\text{Thus } x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = u \frac{\partial z}{\partial u} + v \frac{\partial z}{\partial v}$$

2) If  $u = f(x - y, y - z, z - x)$  then prove that  $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$

- Solution : Let  $x - y = l ; y - z = m ; z - x = n$

Differentiating by chain rule

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l} \frac{\partial l}{\partial x} + \frac{\partial u}{\partial m} \frac{\partial m}{\partial x} + \frac{\partial u}{\partial n} \frac{\partial n}{\partial x} \text{----- (A)}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l} \frac{\partial l}{\partial y} + \frac{\partial u}{\partial m} \frac{\partial m}{\partial y} + \frac{\partial u}{\partial n} \frac{\partial n}{\partial y} \text{----- (B)}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l} \frac{\partial l}{\partial z} + \frac{\partial u}{\partial m} \frac{\partial m}{\partial z} + \frac{\partial u}{\partial n} \frac{\partial n}{\partial z} \text{----- (C)}$$

Since  $l = x - y ; m = y - z ; n = z - x$



Sinhgad Institutes

$$\begin{aligned} \frac{\partial l}{\partial x} &= 1 ; \frac{\partial m}{\partial x} = 0 ; \frac{\partial n}{\partial x} = -1 \\ \frac{\partial l}{\partial y} &= -1 ; \frac{\partial m}{\partial y} = 1 ; \frac{\partial n}{\partial y} = 0 \\ \frac{\partial l}{\partial z} &= 0 ; \frac{\partial m}{\partial z} = -1 ; \frac{\partial n}{\partial z} = 1 \end{aligned}$$

Equation (A), (B),(C) becomes

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l}(1) + \frac{\partial u}{\partial m}(0) + \frac{\partial u}{\partial n}(-1) \text{ ----- (1)}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l}(-1) + \frac{\partial u}{\partial m}(1) + \frac{\partial u}{\partial n}(0) \text{ -----(2)}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial l}(0) + \frac{\partial u}{\partial m}(-1) + \frac{\partial u}{\partial n}(1) \text{ -----(3)}$$

• Adding (1),(2) ,(3) we get

$$\bullet \quad \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$$



Sinhgad Institutes

## Example on Composite Function

Q3) If  $u = f(x^2 - y^2, y^2 - z^2, z^2 - x^2)$ , Prove that

$$\frac{1}{x} \frac{\partial u}{\partial x} + \frac{1}{y} \frac{\partial u}{\partial y} + \frac{1}{z} \frac{\partial u}{\partial z} = 0$$

Solution : Let  $x^2 - y^2 = l$ ,  $y^2 - z^2 = m$ ,  $z^2 - x^2 = n$ ,

$$\therefore u = f(l, m, n)$$

$$\therefore \frac{\partial u}{\partial x} = \frac{\partial u}{\partial l} \cdot \frac{\partial l}{\partial x} + \frac{\partial u}{\partial m} \cdot \frac{\partial m}{\partial x} + \frac{\partial u}{\partial n} \cdot \frac{\partial n}{\partial x}$$

$$\therefore \frac{\partial u}{\partial x} = \frac{\partial u}{\partial l} \cdot (2x) + \frac{\partial u}{\partial m} \cdot (0) + \frac{\partial u}{\partial n} \cdot (-2x)$$

$$\text{similarly } \frac{\partial u}{\partial y} = \frac{\partial u}{\partial l} \cdot (-2y) + \frac{\partial u}{\partial m} \cdot (2y) + \frac{\partial u}{\partial n} \cdot (0)$$



Sinhgad Institutes

similarly

$$\frac{\partial u}{\partial y} \cdot (-2y) + \frac{\partial u}{\partial m} \cdot (2y) + \frac{\partial u}{\partial n} \cdot (0)$$

$$\& \frac{\partial u}{\partial z} = \frac{\partial u}{\partial l} \cdot (0) + \frac{\partial u}{\partial m} \cdot (-2z) + \frac{\partial u}{\partial n} \cdot (2z)$$

$$\therefore \frac{1}{x} \frac{\partial u}{\partial x} + \frac{1}{y} \frac{\partial u}{\partial y} + \frac{1}{z} \frac{\partial u}{\partial z} = 0$$

Hence Proved



Sinhgad Institutes

# Variable to be Treated As Constant

Consider the

Notation  $\left(\frac{\partial u}{\partial x}\right)_y$ ,  $\left(\frac{\partial r}{\partial x}\right)_\theta$ ,  $\left(\frac{\partial v}{\partial y}\right)_{x,u}$  all of them have definite meaning.

- These notations arise when there are more than one functional relations given.
- $\left(\frac{\partial u}{\partial x}\right)_y$  means express  $u$  in terms of  $x$  &  $y$  and differentiate  $u$  partially w . r . t .  $x$  keeping  $y$  as constant using elimination method.



Sinhgad Institutes

$\left(\frac{\partial r}{\partial x}\right)_{\theta}$  means express  $r$  in terms of  $x$  &  $\theta$  and differentiate

$r$  partially w . r . t .  $x$  keeping  $\theta$  as constant using elimination method.

$\left(\frac{\partial v}{\partial y}\right)_{x,u}$  means express  $v$  in terms of  $x$  ,  $y$  &  $u$  and

differentiate  $v$  partially w . r . t .  $y$  keeping  $x$  ,  $u$  as constant using elimination method.

## Note



Sinhgad Institutes

In case the elimination of certain variables is not possible or difficult then we interpret these notations as follows :

1) For a given notation variables in denominator are to be treated as independent variables & remaining variables as dependent .For e.g.  $(\frac{\partial u}{\partial z})_{x,v}$   $x, z, v$  are independent variables &  $u, z, y$  are dependent variables if there are two equations in 5 variables  $u, v, x, y, z$  .

2 In these cases we differentiate all the given equations w.r.t. one of the independent variables keeping other independent variables as constants. Therefore we get simultaneous equations in terms of number of derivatives.

3 We eliminate other derivatives to get the required derivative





Sinhgad Institutes

## Examples on variable to be treated as constant

Q1 If  $x^2 = au + bv$ ,  $y^2 = au - bv$  then show that

$$\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v = \frac{1}{2} = \left(\frac{\partial v}{\partial y}\right)_x \left(\frac{\partial y}{\partial v}\right)_u$$

solution:  $x^2 = au + bv$ ,  $y^2 = au - bv$

$$\left(\frac{\partial x}{\partial u}\right)_v = \frac{a}{2x}$$

$$\left(\frac{\partial y}{\partial v}\right)_u = -\frac{b}{2y}$$

Since  $x^2 + y^2 = 2au$        $x^2 - y^2 = 2bv$

$$u = \frac{x^2 + y^2}{2a} \left(\frac{\partial u}{\partial x}\right)_y = \frac{x}{a}$$

$$v = \frac{x^2 - y^2}{2b} \left(\frac{\partial v}{\partial y}\right)_x = -\frac{y}{b}$$



Sinhgad Institutes

- $\therefore \left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v = \frac{x}{a} \frac{a}{2x} = \frac{1}{2}$
- $\left(\frac{\partial v}{\partial y}\right)_x \left(\frac{\partial y}{\partial v}\right)_u = \left(-\frac{y}{b}\right) \left(-\frac{b}{2y}\right) = \frac{1}{2}$



Sinhgad Institutes

2) If  $x = \frac{r}{2}(e^{\theta} + e^{-\theta})$ ;  $y = \frac{r}{2}(e^{\theta} - e^{-\theta})$

then prove that  $\left(\frac{\partial x}{\partial r}\right)_{\theta} = \left(\frac{\partial r}{\partial x}\right)_y$

Solution : Given  $x = \frac{r}{2}(e^{\theta} + e^{-\theta})$ ;  $y = \frac{r}{2}(e^{\theta} - e^{-\theta})$

$$\therefore x = r \cosh \theta ; y = r \sinh \theta$$

Differentiating w.r.t.  $r$  keeping  $\theta$  as constant

$$\therefore \left(\frac{\partial x}{\partial r}\right)_{\theta} = \cosh \theta \text{ ----(1)}$$

Now we express  $r$  as a function of  $x$  and  $y$

$$x = r \cosh \theta ; y = r \sinh \theta$$

$$\therefore x^2 - y^2 = r^2 \cosh^2 \theta - r^2 \sinh^2 \theta = r^2 (1) = r^2$$



Sinhgad Institutes

$$\therefore r^2 = x^2 - y^2$$

Differentiating  $r$  with respect to  $x$

keeping  $y$  constant

$$2r \left( \frac{\partial r}{\partial x} \right)_y = 2x \therefore \left( \frac{\partial r}{\partial x} \right)_y = \frac{x}{r} = \frac{r \cosh \theta}{r} = \cosh \theta$$

Thus from (1) and (2)

$$\left( \frac{\partial x}{\partial r} \right)_\theta = \left( \frac{\partial r}{\partial x} \right)_y$$



Sinhgad Institutes

3)  $x = u \tan v$ ;  $y = u \sec v$ , then prove that:

$$\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial v}{\partial x}\right)_y = \left(\frac{\partial u}{\partial y}\right)_x \left(\frac{\partial v}{\partial y}\right)_x$$

• Solution : Given  $x = u \tan v$ ;  $y = u \sec v$

We have to find  $u, v$  are functions of  $x, y$

i.e. we have to find  $u = f(x, y)$ ;  $v = f(x, y)$

squaring  $x, y$  and subtracting

- $y^2 - x^2 = u^2 \sec^2 v - u^2 \tan^2 v$
- $= u^2 [\sec^2 v - \tan^2 v] = u^2$
- $y^2 - x^2 = u^2$



Sinhgad Institutes

*differentiate*  $u^2$  w.r.t.  $x$  keeping  $y$  as constant

$$u^2 = y^2 - x^2$$

$$2u \frac{\partial u}{\partial x} = \frac{\partial}{\partial x} (y^2 - x^2)$$

$$2u \frac{\partial u}{\partial x} = -2x$$

$$\left( \frac{\partial u}{\partial x} \right)_y = \frac{-2x}{2u} = \frac{x}{u} = - \left[ \frac{u \tan v}{u} \right]$$

$$\left( \frac{\partial u}{\partial x} \right)_y = -\tan v \text{ -----(1)}$$



Sinhgad Institutes

differentiate  $u^2$  w.r.t.  $y$ , keeping  $x$  as constant

$$2u \left( \frac{\partial u}{\partial y} \right)_x = \frac{\partial}{\partial y} (y^2 - x^2) = 2y$$

$$\left( \frac{\partial u}{\partial y} \right)_x = \frac{y}{u} = \frac{u \sec v}{u} = \sec v \text{-----}(2)$$

Now to find  $v$  as function of  $x, y$

$$\frac{x}{y} = \frac{u \tan v}{u \sec v} = \frac{\sin v}{\cos v \left( \frac{1}{\cos v} \right)} = \sin v$$

$$\frac{x}{y} = \sin v \text{-----}(1)$$



Sinhgad Institutes

Differentiate eq(1) w.r.t.x, keeping y as constant

$$\frac{1}{y} \frac{\partial}{\partial x} (x) = \frac{\partial}{\partial x} (\sin v)$$

$$\frac{1}{y} = \cos v \left( \frac{\partial v}{\partial x} \right)_y$$

$$\left( \frac{\partial v}{\partial x} \right)_y = \frac{1}{y \cos v} \text{-----(3)}$$

Differentiate (I) w.r.t.y keeping x as constant

$$x \cdot \frac{\partial}{\partial y} \left( \frac{1}{y} \right) = \frac{\partial}{\partial y} (\sin v)$$





Sinhgad Institutes

$$\frac{-x}{y^2} = \cos v \cdot \left( \frac{\partial v}{\partial y} \right)_x$$

$$\left( \frac{\partial v}{\partial y} \right)_x = \frac{-x}{y^2 \cos v} \text{-----(4)}$$

From eq(1) and (3) we get,

$$\left( \frac{\partial u}{\partial x} \right)_y \left( \frac{\partial v}{\partial x} \right)_y = -\tan v \frac{1}{y \cos v} = \frac{-\sin v}{y \cos^2 v}$$

And from eq(2) and (4), we get

$$\left( \frac{\partial u}{\partial y} \right)_x \left( \frac{\partial v}{\partial y} \right)_x = \sec v \cdot \frac{-x}{y^2 \cos v} = \frac{-x}{y^2 \cos^2 v}$$



Sinhgad Institutes

This shows that  $\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial v}{\partial x}\right)_y = \left(\frac{\partial u}{\partial y}\right)_x \left(\frac{\partial v}{\partial y}\right)_x$

4) If  $u = 2x + 3y$ ;  $v = 3x - 2y$  then find the value of

i)  $\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v \left(\frac{\partial y}{\partial v}\right)_x \left(\frac{\partial v}{\partial y}\right)_u$

ii)  $\left(\frac{\partial y}{\partial v}\right)_x \left(\frac{\partial v}{\partial y}\right)_x = \frac{13}{4}$

iii)  $\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v = \frac{4}{13}$



Sinhgad Institutes

Solution : given  $u = 2x + 3y$ ---(1)

$$v = 3x - 2y$$
---(2)

We have to find

$$y \rightarrow f(x, v); v \rightarrow f(y, u)$$

$$u \rightarrow f(x, y); x \rightarrow f(u, v)$$

Differentiate eq(1) w.r.t.x keeping y as constant

$$\left(\frac{\partial u}{\partial x}\right)_y = 2$$
---(1)

To find  $y \rightarrow f(x, v)$  From eq(2)

$$y = \frac{nx-v}{m}$$



Sinhgad Institutes

$$\therefore \left( \frac{\partial y}{\partial v} \right)_x = \left( \frac{-1}{2} \right) \text{----(II)}$$

To find  $v \rightarrow f(y, u)$  from eq(1)

$$x = \frac{u-3y}{2} \text{ put in eq(2)}$$

$$v = n \left[ \frac{u-3y}{2} \right] - 2y$$

$$\left( \frac{\partial v}{\partial y} \right)_u = \frac{3}{2} (-3) - 2 = \frac{-3^2}{2} - 2$$

$$\left( \frac{\partial v}{\partial y} \right)_u = - \left[ \frac{(2^2 + 3^2)}{2} \right] = \left( \frac{-13}{2} \right) \text{----(III)}$$



Sinhgad Institutes

To find  $x \rightarrow f(u, v)$

We do,  $m \times eq(1) + n \times eq(2)$

$$2u = 2^2x + 6y$$

$$3v = 3^2x - 6y$$

$$2u + 3v = 13x$$

$$x = \frac{2u+3v}{13}$$

$$\left(\frac{\partial x}{\partial u}\right)_v = \frac{2}{13} \text{-----(IV)}$$



Sinhgad Institutes

From (I),(II),(III),(IV)

$$\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v \left(\frac{\partial y}{\partial v}\right)_x \left(\frac{\partial v}{\partial y}\right)_u = 2 \left(\frac{-1}{2}\right) \left(\frac{-13}{2}\right) \frac{2}{13} = 1$$

from(II)and(III)

$$\text{ii) } \left(\frac{\partial y}{\partial v}\right)_x \left(\frac{\partial v}{\partial y}\right)_x = \frac{13}{4}$$

from(I) and(IV)

$$\left(\frac{\partial u}{\partial x}\right)_y \left(\frac{\partial x}{\partial u}\right)_v = \frac{4}{13}$$



Sinhgad Institutes

## University Questions

1) If  $z = f(u, v)$  ;  $u = x^2 - 2xy - y^2$  and  $v = y$

Show that  $(x + y) \frac{\partial z}{\partial x} + (x - y) \frac{\partial z}{\partial y} = (x - y) \frac{\partial z}{\partial v}$

2 ) If  $x = e^u \tan v$  and  $y = e^u \sec v$  find the value of

$$\left( x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) \left( x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} \right)$$

3) If  $z = e^{ax+by} f(ax - by)$  , prove that  $b \frac{\partial z}{\partial x} +$

$$a \frac{\partial z}{\partial y} = 2abz$$



Sinhgad Institutes

4) If  $x = \frac{r}{2}(e^\theta + e^{-\theta})$  &  $y = \frac{r}{2}(e^\theta - e^{-\theta})$

*Prove that*

$$(\partial x / \partial r)_\theta = (\partial r / \partial x)_y$$

5) If  $ux + vy = 0$  &  $\frac{u}{x} + \frac{v}{y} = 1$  *prove that*

$$\left(\frac{\partial u}{\partial x}\right)_y - \left(\frac{\partial v}{\partial y}\right)_x = \frac{x^2 + y^2}{y^2 - x^2}$$





Sinhgad Institutes

# UNIT III: Partial Differentiation

## Topic: Euler's Theorem



Sinhgad Institutes

# HOMOGENEOUS FUNCTIONS

A function  $z$  of two variables  $x$  and  $y$  is said to be *homogeneous function of degree 'n'* if it is possible to express  $z$  in the form

$$z = x^n \cdot f\left(\frac{y}{x}\right) \quad \text{OR} \quad z = y^n \cdot f\left(\frac{x}{y}\right)$$

**Euler's Theorem** : If  $z$  is a *homogeneous* expression of two variables  $x$  and  $y$  of degree  $n$  then ,

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$$

$$z = x^n f\left(\frac{y}{x}\right) \Rightarrow \frac{\partial z}{\partial x} = x^n f' \left(\frac{y}{x}\right) \left(-\frac{y}{x^2}\right) + nx^{n-1} f\left(\frac{y}{x}\right)$$



Sinhgad Institutes

$$\therefore x \frac{\partial z}{\partial x} = -yx^{n-1} f' \left( \frac{y}{x} \right) + nx^n f \left( \frac{y}{x} \right) - - - (1)$$

$$\frac{\partial z}{\partial y} = x^n f' \left( \frac{y}{x} \right) \left( \frac{1}{x} \right)$$

$$y \frac{\partial z}{\partial y} = yx^{n-1} f' \left( \frac{y}{x} \right) - - - - (2)$$

Adding (1) and (2)

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = -yx^{n-1} f' \left( \frac{y}{x} \right) + nx^n f \left( \frac{y}{x} \right) + yx^{n-1} f' \left( \frac{y}{x} \right)$$

$$\text{Hence } x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = nx^n f \left( \frac{y}{x} \right) = nz$$



Sinhgad Institutes

## Example On Euler's Theorem

1) If  $T = \sin \left[ \frac{xy}{x^2+y^2} \right] + \sqrt{x^2 + y^2} + \frac{x^2y}{x+y}$  then find the value of  $x \frac{\partial T}{\partial x} + y \frac{\partial T}{\partial y}$  at (3,4)

Solution :  $T = \sin \left[ \frac{xy}{x^2+y^2} \right] + \sqrt{x^2 + y^2} + \frac{x^2y}{x+y}$

Let  $T = U + V + W$

Where  $U = \sin \left[ \frac{xy}{x^2+y^2} \right]$  ;  $V = \sqrt{x^2 + y^2}$  ;  $W = \frac{x^2y}{x+y}$



Sinhgad Institutes

Put  $x = xt$ ;  $y = yt$ ; in  $U, V, W$  we get

$$U = \sin \left[ \frac{xt \cdot yt}{(xt)^2 + (yt)^2} \right] = t^0 \sin \left[ \frac{xy}{x^2 + y^2} \right]$$

$$V = \sqrt{(xt)^2 + (yt)^2} = t^1 \sqrt{x^2 + y^2} = t V$$

$$W = \frac{xt^2 \cdot yt}{xt + yt} = t^2 \frac{x^2 y}{x + y} = t^2 W$$

Thus  $U, V, W$  are homogeneous functions of degree 0, 1, 2 respectively

Thus by Euler's theorem

$$x \frac{\partial U}{\partial x} + y \frac{\partial U}{\partial y} = nU = 0. U = 0 \text{-----(1)}$$

$$x \frac{\partial V}{\partial x} + y \frac{\partial V}{\partial y} = nV = 1. V = V \text{-----(2)}$$

$$x \frac{\partial W}{\partial x} + y \frac{\partial W}{\partial y} = nW = 2. w = 2W \text{-----(3)}$$

Adding, We get

$$x \left( \frac{\partial U}{\partial x} + \frac{\partial V}{\partial x} + \frac{\partial W}{\partial x} \right) + y \left( \frac{\partial U}{\partial y} + \frac{\partial V}{\partial y} + \frac{\partial W}{\partial y} \right) = 0 + 1V + 2W \text{-----(4)}$$

But as  $T = U + V + W$



Sinhgad Institutes

Differentiating T w.r.t. x & y

$$\therefore \frac{\partial T}{\partial x} = \frac{\partial U}{\partial x} + \frac{\partial V}{\partial x} + \frac{\partial W}{\partial x}$$

$$\frac{\partial T}{\partial y} = \frac{\partial U}{\partial y} + \frac{\partial V}{\partial y} + \frac{\partial W}{\partial y}$$

Equation (4) becomes

$$x \cdot \frac{\partial T}{\partial x} + y \cdot \frac{\partial T}{\partial y} = V + 2W$$

$$\therefore x \cdot \frac{\partial T}{\partial x} + y \cdot \frac{\partial T}{\partial y} = \sqrt{x^2 + y^2} + 2 \frac{x^2 y}{x+y}$$

At (x , y ) = (3,4)



Sinhgad Institutes

$$x \cdot \frac{\partial T}{\partial x} + y \cdot \frac{\partial T}{\partial y} = (5) + 2 \left( \frac{36}{7} \right) = \frac{107}{7}$$





Sinhgad Institutes

## Deductions From Euler's Theorem :

(1) If  $z$  is a *homogeneous* expression of two variables  $x$  and  $y$  of degree  $n$  then ,

$$x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = n(n-1) z$$

(2) If  $z$  is a *homogeneous* function of two variables  $x$  and  $y$  of degree  $n$  and  $z = f(u)$  then ,

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)}$$

$$(3) \quad x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = g(u)[g'(u) - 1]$$

$$\text{Where } g(u) = n \frac{f(u)}{f'(u)}$$



Sinhgad Institutes

## HOMOGENEOUS FUNCTIONS OF THREE VARIABLES.

A function  $u = f(x, y, z)$  is said to be homogeneous function of three variables  $x, y, z$  of degree 'n' if it is possible to express  $u$  in the form

$$u = x^n \cdot f\left(\frac{y}{x}, \frac{z}{x}\right) \quad \text{OR} \quad u = y^n \cdot f\left(\frac{x}{y}, \frac{z}{y}\right)$$



Sinhgad Institutes

## Example On Deduction (1)

1) If  $z = x^n f\left(\frac{y}{x}\right) + y^{-n} \phi\left(\frac{x}{y}\right)$  then prove that  
 $x^2 z_{xx} + 2xy z_{yy} + x \cdot z_x + y \cdot z_y = n^2 z$

Solution: Given  $z = x^n f\left(\frac{y}{x}\right) + y^{-n} \phi\left(\frac{x}{y}\right)$

$$Z = U + V \text{ -----(1)}$$

Where  $U = x^n f\left(\frac{y}{x}\right)$  ;  $V = y^{-n} \cdot \phi\left(\frac{x}{y}\right)$

$\therefore$  U and V are homogeneous functions  
in x & y of degree n and  $-n$  respectively

$\therefore$  By Euler's theorem

$$x U_x + y U_y = nU \text{ -----(2)}$$

$$x V_x + y V_y = -nV \text{ -----(3)}$$

And

$$x^2 U_{xx} + 2xy U_{xy} + y^2 U_{yy} = n(n-1)U \text{ -----(4)}$$

$$\begin{aligned} x^2 V_{xx} + 2xy V_{xy} + y^2 V_{yy} &= -n(-n-1)V \\ &= n(n+1) \text{ -----(5)} \end{aligned}$$



Sinhgad Institutes

Adding (2) ,(3) ,(4) and (5) we get

$$\begin{aligned} & x^2(U_{xx} + V_{xx}) + 2xy(U_{xy} + V_{xy}) \\ & + y^2(U_{yy} + V_{yy}) + x(U_x + V_x) + y(U_y + V_y) \\ & = n(n-1)U + n(n+1)V + nU - nV \\ & = n^2U - nU + n^2V + nV + nU - nV \\ & = n^2(U + V) \text{-----(6)} \end{aligned}$$

Now from equation (1)

$$Z = U + V$$

Differentiating Z w.r.t. x & y

$$Z_x = U_x + V_x \text{ and } Z_y = U_y + V_y$$



Sinhgad Institutes

$$\therefore Z_{xy} = U_{xy} + V_{xy}; Z_{xx} = U_{xx} + V_{xx};$$
$$Z_{xy} = U_{yy} + V_{yy};$$

$$x^2 Z_{xx} + 2xy Z_{xy} + y^2 Z_{yy} + x Z_x + y Z_y = n^2 Z$$

Equation (6) becomes



Sinhgad Institutes

## EXAMPLES On Deduction (2)

**EX 1)** : If  $x = e^u \tan v$  and  $y = e^u \sec v$  find the value of

$$\left( x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) \left( x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} \right) = 0$$

Eliminating  $v$  between the given relation

$$y^2 - x^2 = e^{2u} = z \text{ (say)}$$

Then  $z = x^2 \left( \frac{y^2}{x^2} - 1 \right) = x^2 f \left( \frac{y}{x} \right)$  and also  $z = f(u) = e^{2u}$

$\Rightarrow z$  is a Homogeneous function of  $x$  and  $y$  of degree 2.

$\therefore$  from deduction of Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} = 2 \frac{e^{2u}}{2e^{2u}} = 1 \text{ --- (1)}$$



**Sinhgad Institutes**

Now eliminating  $u$  between the given relation

$$\frac{y}{x} = \frac{\sec v}{\tan v} = \operatorname{cosec} v = z \text{ (say), Here } z = x^0 \left( \frac{y}{x} \right), \text{ also } z = g(v) = \operatorname{cosec} v$$

$\Rightarrow z$  is a Homogeneous function of  $x$  and  $y$  of degree 0.

$\therefore$  from deduction of Euler's theorem

$$x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} = n \frac{g(v)}{g'(v)} = 0 \times \frac{\operatorname{cosec} v}{-\cot^2 v} = 0 \text{ --- (2)}$$

$\therefore$  from (1) and (2) we get the required result.

$$\left( x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} \right) \left( x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} \right) = 1 \times 0 = 0$$





Sinhgad Institutes

2) If  $u = \sin^{-1} \left( \frac{x+y}{\sqrt{x} + \sqrt{y}} \right)$  then prove that

$$2x \frac{\partial u}{\partial x} + 2y \frac{\partial u}{\partial y} = \tan u$$

Solution : Given  $u = \sin^{-1} \left( \frac{x+y}{\sqrt{x} + \sqrt{y}} \right)$  -----(1)

Put  $x = xt : y = yt$  in equation (1) we get

$$u = \sin^{-1} \left( \frac{xt + yt}{\sqrt{xt} + \sqrt{yt}} \right) = \sin^{-1} \left[ t^{1/2} \left( \frac{x + y}{\sqrt{x} + \sqrt{y}} \right) \right]$$



Sinhgad Institutes

Is not homogeneous function

But  $f(u) = \sin u = t^{1/2} \left( \frac{x+y}{\sqrt{x} + \sqrt{y}} \right) = t^{1/2} u$  is homogeneous function in  $x$  &  $y$  of degree  $\frac{1}{2}$

Thus, by Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} = \frac{1}{2} \frac{\sin u}{\cos u}$$

$$\therefore 2x \frac{\partial u}{\partial x} + 2y \frac{\partial u}{\partial y} = \tan u$$



Sinhgad Institutes

**EX :** If  $u = \tan^{-1} \left( \frac{x^3+y^3}{x-y} \right)$ , Show that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = (1 - 4\sec^2 u) \sin 2u$$

$$\text{Given } \tan u = \frac{x^3+y^3}{x-y} = x^2 \left[ \frac{1+y^3/x^3}{1-y/x} \right] = x^2 f\left(\frac{y}{x}\right) = z(\text{say})$$

$$\text{Also } z = f(u) = \tan u$$

Homogeneous function of  $x$  and  $y$  of degree 2.

By Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} = 2 \frac{\tan u}{\sec^2 u}$$

$$= 2 \sin u \cos u = g(u) \text{ say } \dots (1)$$

Other Subjects: <https://www.studymedia.in/fe/notes>



Sinhgad Institutes

$$\begin{aligned}x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} &= g(u)[g'(u) - 1], \text{ Where } g(u) = n \frac{f(u)}{f'(u)} \\&= \sin 2u [2\cos 2u - 1] \\&= \sin 2u [2(1 - 2\sin^2 u) - 1] \\&= \sin 2u [1 - 4\sin^2 u]\end{aligned}$$



Sinhgad Institutes

**EX :** If  $T = \sin \left( \frac{xy}{x^2+y^2} \right) + \sqrt{x^2+y^2} + \frac{x^2y}{x+y}$  then

find  $x \frac{\partial T}{\partial x} + y \frac{\partial T}{\partial y}$

Let  $T = u + v + w$  Then

$$x \frac{\partial T}{\partial x} + y \frac{\partial T}{\partial y} = \left( x \frac{\partial U}{\partial x} + y \frac{\partial U}{\partial y} \right) + \left( x \frac{\partial V}{\partial x} + y \frac{\partial V}{\partial y} \right) + \left( x \frac{\partial W}{\partial x} + y \frac{\partial W}{\partial y} \right) \text{ --- (1)}$$

$$\begin{aligned} u = \sin \left( \frac{xy}{x^2+y^2} \right) &\Rightarrow \sin^{-1} u = \frac{xy}{x^2+y^2} \\ &= \frac{x^2(y/x)}{x^2 \left( 1 + \frac{y^2}{x^2} \right)} = x^0 f \left( \frac{y}{x} \right) = z \end{aligned}$$

and also  $\sin^{-1} u = z$

$u$  is homogeneous function of  $x, y$  order 0.

By Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} = 0 \text{ --- (1)}$$

Other Subjects: <https://www.studymedia.in/fe/notes>



Sinhgad Institutes

$$v = \sqrt{x^2 + y^2} = x \sqrt{1 + \frac{y^2}{x^2}} = x f\left(\frac{y}{x}\right)$$

It is homogeneous function of x, y, of order 1

By Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n v = 1 \cdot \sqrt{x^2 + y^2} = \sqrt{x^2 + y^2} \quad \text{--- (2)}$$

$$w = \frac{x^2 y}{x+y} = \frac{x^3 (y/x)}{x(1+y/x)} = x^2 f\left(\frac{y}{x}\right)$$

So w is homogeneous function of x, y of order 2.

By Euler's theorem

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n v = 2 \cdot \frac{x^2 y}{x+y} \quad \text{--- (3)}$$

Putting the values from (1), (2) and (3)

$$x \frac{\partial T}{\partial x} + y \frac{\partial T}{\partial y} = 0 + \sqrt{x^2 + y^2} + \frac{2x^2 y}{x+y} = \sqrt{x^2 + y^2} + \frac{2x^2 y}{x+y}$$



Sinhgad Institutes

## Example On Deduction (3)

- If  $u = \operatorname{cosec}^{-1} \sqrt{\frac{x^{1/2} + y^{1/2}}{x^{1/3} + y^{1/3}}}$

Put  $x = xt$ ;  $y = yt$  in  $u$ , we get

$$u = \operatorname{cosec}^{-1} \sqrt{\frac{(xt)^{1/2} + (yt)^{1/2}}{(xt)^{1/3} + (yt)^{1/3}}}$$

$$u = \operatorname{cosec}^{-1} \sqrt{\frac{t^{1/2}(x^{1/2} + y^{1/2})}{t^{1/3}(x^{1/3} + y^{1/3})}}$$



Sinhgad Institutes

$$u = \operatorname{cosec}^{-1} \sqrt{\frac{t^{1/4} \sqrt{x^{1/2} + y^{1/2}}}{t^{1/6} \sqrt{x^{1/3} + y^{1/3}}}}$$

$$u = \operatorname{cosec}^{-1} \left[ t^{1/2} \sqrt{\frac{x^{1/2} + y^{1/2}}{x^{1/3} + y^{1/3}}} \right]$$

U is not homogeneous function

$$\text{But } f(u) = \operatorname{cosec} u = t^{1/2} \sqrt{\frac{x^{1/2} + y^{1/2}}{x^{1/3} + y^{1/3}}}$$





Sinhgad Institutes

is homogeneous function in  $x$  &  $y$  of degree  $\frac{1}{2}$

Thus, by Euler's theorem on homogeneous function, we get

$$x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = G(u)[G'(u) - 1] \text{ ---- (1)}$$

$$\text{Where } G(u) = \frac{n f(u)}{f'(u)} = \frac{1}{12} \cdot \frac{\operatorname{cosec} u}{-\operatorname{cosec} u \cdot \cot u} = \frac{-1}{12} \sec^2 u$$

Equation (1) becomes

$$x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = \frac{-1}{12} \tan u \left[ \frac{-1}{12} \sec^2 u - 1 \right]$$



Sinhgad Institutes

$$\begin{aligned} &= \frac{1}{144} \tan u [\sec^2 u + 12] \\ &= \frac{1}{144} \tan u [\tan^2 u + 1 + 12] \\ &= \frac{1}{144} \tan u [\tan^2 u + 13] \end{aligned}$$



Sinhgad Institutes

# TOTAL DERIVATIVE

If  $u = f(x, y)$ , where  $x = \phi(t)$  and  $y = \psi(t)$  then

$u$  can be expressed as function of single variable  $t$ .

and derivative of  $u$  w.r.t.  $t$  is ordinary differential coefficient  $\frac{du}{dt}$

This  $\frac{du}{dt}$  is called as Total Derivative.

If we put  $t = x$  in (1) then

$$\frac{du}{dx} = \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} \frac{dy}{dx}$$

Similarly if we put  $t = y$  in (1) then

$$\frac{du}{dy} = \frac{\partial u}{\partial x} \frac{dx}{dy} + \frac{\partial u}{\partial y}$$



Sinhgad Institutes

## DIFFERENTIATION OF IMPLICIT FUNCTION

If  $z = f(x, y) = 0$ ,  $y$  is function of  $x$  then

$$\frac{dz}{dx} = \frac{\partial z}{\partial x} \frac{dx}{dx} + \frac{\partial z}{\partial y} \frac{dy}{dx} \quad \text{i. e.} \quad \frac{dz}{dx} = \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} \frac{dy}{dx} \quad \text{--- (1)}$$

$$\text{Since } z = f(x, y) = 0 \Rightarrow \frac{dz}{dx} = 0$$

$$\therefore \frac{\partial z}{\partial x} + \frac{\partial z}{\partial y} \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = - \frac{\frac{\partial z}{\partial x}}{\frac{\partial z}{\partial y}} = - \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} = - \frac{p}{q}$$

$$\text{as } p = \frac{\partial z}{\partial x}, \quad q = \frac{\partial z}{\partial y}, \quad r = \frac{\partial^2 z}{\partial x \partial y}, \quad s = \frac{\partial^2 z}{\partial x^2}, \quad t = \frac{\partial^2 z}{\partial y^2}$$

$$\text{or } p = \frac{\partial f}{\partial x}, \quad q = \frac{\partial f}{\partial y}, \quad r = \frac{\partial^2 f}{\partial x \partial y}, \quad s = \frac{\partial^2 f}{\partial x^2}, \quad t = \frac{\partial^2 f}{\partial y^2}$$



Sinhgad Institutes

Differentiating once more we get

$$\therefore \frac{d^2y}{dx^2} = - \left[ \frac{q^2r - 2pqs + p^2t}{q^3} \right]$$

This can also be remembered as

$$\therefore \frac{d^2y}{dx^2} = \frac{\det \begin{bmatrix} r & s & p \\ s & t & q \\ p & q & 0 \end{bmatrix}}{q^3}$$

~~~~~

## EXAMPLES.

**EX :** If  $u = x \log xy$  and  $x^3 + y^3 + 3xy = 0$  then find  $\frac{du}{dx}$

Given  $u = x \log xy$  and  $x^3 + y^3 + 3xy = 0$

$u$  is composite function of  $x$

$$\therefore \frac{du}{dx} = \frac{\partial u}{\partial x} \frac{dx}{dx} + \frac{\partial u}{\partial y} \frac{dy}{dx} \text{ --- (1)}$$

$u = x \log xy$  and  $f(x, y) \equiv x^3 + y^3 + 3xy = 0$

$$\frac{\partial u}{\partial x} = x \left( \frac{1}{xy} \right) y + \log xy = 1 + \log xy \text{ --- (2)}$$

$$\frac{\partial u}{\partial y} = x \left( \frac{1}{xy} \right) x = \frac{x}{y} \text{ --- (3)}$$

$$\frac{dy}{dx} = -\frac{p}{q} = -\frac{\partial f / \partial x}{\partial f / \partial y} = -\frac{3x^2 + 3y}{3y^2 + 3x} \text{ --- (4)}$$



Sinhgad Institutes

Putting the values from (2) , (3) and (4) in (1)

$$\frac{du}{dx} = 1 + \log xy + \frac{x}{y} \left( -\frac{3x^2 + 3y}{3y^2 + 3x} \right)$$

$$= 1 + \log xy - \left( \frac{x^3 + xy}{y^3 + xy} \right)$$

$$= \log xy + \frac{y^3 + xy - x^3 - xy}{y^3 + xy}$$

$$\therefore \frac{du}{dx} = \log xy + \frac{y^3 - x^3}{y^3 + xy}$$

EX: If  $(\cos x)^y = (\sin y)^x$  then find  $\frac{dy}{dx}$



## EXAMPLES ON TOTAL DERIVATIVES

Sinhgad Institutes **Ex :** If  $x^2y - e^z + x \sin z = 0$  and  $x^2 + y^2 + z^2 = a^2$

Then evaluate  $\frac{dy}{dx}$  and  $\frac{dx}{dz}$

$$\text{Let } f(x, y, z) = x^2y - e^z + x \sin z = 0$$

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz$$

$$0 = (2xy + \sin z)dx + (x^2)dy + (-e^z + x \cos z)dz \quad \text{--- (1)}$$

$$\text{Let } \phi(x, y, z) = x^2 + y^2 + z^2 - a^2 = 0$$

$$d\phi = \frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz$$

$$0 = x dx + y dy + z dz \quad \text{--- (2)}$$





Sinhgad Institutes

From (1) and (2) by Cramer's rule for  $dx$ ,  $dy$ ,  $dz$

$$\frac{dx}{y(-e^z + x \cos z) - zx^2} = \frac{dy}{z(2xy + \sin z) - x(-e^z + x \cos z)}$$
$$= \frac{dz}{x^3 - y(2xy + \sin z)}$$

$$\frac{dy}{dx} = \frac{z(2xy + \sin z) - x(-e^z + x \cos z)}{y(-e^z + x \cos z) - zx^2}$$

$$\frac{dx}{dz} = \frac{y(-e^z + x \cos z) - zx^2}{x^3 - y(2xy + \sin z)}$$



Sinhgad Institutes

2) If  $u = x \log(xy)$  and  $x^3 + y^3 + 3xy = 0$  then find  $\frac{du}{dx}$  at (1,1)

Solution: Given  $u = x \log(xy) = x[\log x + \log y]$

&

$$x^3 + y^3 + 3xy = 0$$

By differentiating  $u$  w.r.t.  $x$  by chain rule

$$\frac{du}{dx} = \frac{\partial u}{\partial x} \cdot \frac{dx}{dx} + \frac{\partial u}{\partial y} \cdot \frac{dy}{dx} \text{ -----(1)}$$

$$\frac{\partial u}{\partial x} = \frac{\partial}{\partial x} x[\log x + \log y] = 1 + \log x + \log y$$



Sinhgad Institutes

And

$$\frac{\partial u}{\partial y} = \frac{x}{y}$$

Eq(1) becomes

$$\frac{du}{dx} = [1 + \log x + \log y] + \frac{x}{y} \cdot \frac{dy}{dx} \text{ -----(2)}$$

$$\text{As } f = x^3 + y^3 + 3xy - 1$$



Sinhgad Institutes

$$\begin{aligned}\text{We have } \frac{dy}{dx} &= - \left[ \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} \right] = - \left[ \frac{3x^2 + 3y}{3y^2 + 3x} \right] = \\ &= - \left[ \frac{x^2 + y}{y^2 + x} \right]\end{aligned}$$

eq(2) becomes

$$\frac{du}{dx} = [1 + \log x + \log y] + \frac{x}{y}$$



Sinhgad Institutes

$$= [1 + \log x + \log y] - \frac{x}{y} \cdot \left[ \frac{x^2 + y}{y^2 + x} \right]$$

$$\left( \frac{du}{dx} \right)_{(1,1)} = (1 + 0 + 0) - \frac{1}{1} \left[ \frac{1+1}{1+1} \right]$$

$$\left( \frac{du}{dx} \right)_{(1,1)} = 1 - 1$$

$$\left( \frac{du}{dx} \right)_{(1,1)} = 0$$



Sinhgad Institutes

3) If  $f(x, y) = 0$ ;  $\phi(x, z) = 0$  then prove that

$$\frac{\partial \phi}{\partial x} \frac{\partial f}{\partial y} \frac{dy}{dz} = \frac{\partial f}{\partial x} \frac{\partial \phi}{\partial z}$$

Solution : Given

$$f(x, y) = 0; \phi(x, z) = 0$$

$$f \rightarrow x, y \rightarrow x \text{ \& } \phi \rightarrow x, z \rightarrow x$$

$$\text{Means } f \rightarrow x \text{ \& } y \rightarrow x$$



Sinhgad Institutes

$$\emptyset \rightarrow x \text{ \& } z \rightarrow x$$

Thus by chain rule

$$df = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx}$$

$$df = \frac{\partial \emptyset}{\partial x} \frac{dx}{dx} + \frac{\partial \emptyset}{\partial y} \frac{dy}{dx}$$

$$f = 0 \text{ and } \emptyset = 0$$

$$\text{Thus } df = 0 \text{ and } d\emptyset = 0$$



Sinhgad Institutes

$$\therefore \frac{dy}{dx} = - \left[ \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} \right] \text{ And } \frac{dz}{dx} = - \left[ \frac{\frac{\partial \phi}{\partial x}}{\frac{\partial \phi}{\partial z}} \right]$$

$$\frac{\left( \frac{dy}{dx} \right)}{\left( \frac{dz}{dx} \right)} = \frac{- \left[ \frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} \right]}{- \left[ \frac{\frac{\partial \phi}{\partial x}}{\frac{\partial \phi}{\partial z}} \right]}$$





Sinhgad Institutes

$$\text{Therefore } \frac{dy}{dz} = \frac{\frac{\partial f}{\partial x} \frac{\partial \phi}{\partial z}}{\frac{\partial \phi}{\partial x} \frac{\partial f}{\partial y}}$$

$$\text{And } \frac{\partial \phi}{\partial x} \frac{\partial f}{\partial y} \frac{dy}{dz} = \frac{\partial f}{\partial x} \frac{\partial \phi}{\partial z}$$

3) Find  $\frac{dz}{dx}$  if  $z = x^2y$  &  $x^2 + xy + y^2 = 1$

Solution :  $z = x^2y$  &  $x^2 + xy + y^2 = 1$



Sinhgad Institutes

Differentiate  $z$  w.r.t  $x$ , keeping  $y$  as constant and then differentiate  $z$  w.r.t  $y$ , keeping  $x$  as constant

$$\frac{\partial z}{\partial x} = 2xy \text{ \& } \frac{\partial z}{\partial y} = x^2$$

$$\text{As } f(x, y) = x^2 + xy + y^2 - 1$$

$$\frac{dy}{dx} = - \left[ \frac{2x+y}{x+2y} \right]$$

By chain rule  $z \rightarrow x, y \rightarrow x$



Sinhgad Institutes

$$\frac{dz}{dx} = \frac{\partial z}{\partial x} \frac{dx}{dx} + \frac{\partial z}{\partial y} \frac{dy}{dx} = (2xy)(1) + (x^2)\left(-\left[\frac{2x+y}{x+2y}\right]\right)$$

$$\text{Thus } \frac{dz}{dx} = 2xy - x^2 \left[\frac{2x+y}{x+2y}\right]$$



Sinhgad Institutes

**Ex :**  $lax^2 + by^2 + cz^2 = 1$  and  $lx + my + nz = 0$

Then prove that

$$\frac{dx}{bny - cmz} = \frac{dy}{clz - anx} = \frac{dz}{amx - bly}$$

---

---

---



Sinhgad Institutes

## References:-

- 1) Higher Engineering Mathematics –  
B. V. Raman
- 2) Advanced Engineering Mathematics -  
H. K. Dass
- 3) Advanced Engineering Mathematics --