

JOIN



Telegram
@PuneEngineers

For more Subjects

<https://www.studymedia.in/fe/notes>



 SCAN ME



Total No. of Questions : 8]

SEAT No. :

PA-4295

[Total No. of Pages : 3

[5924]-4

F.E.

ENGINEERING PHYSICS

(2019 Pattern) (Semester - I/II) (107002)

Time : 2½ Hours]

[Max. Marks :70

Instructions to the candidates:

- 1) Answer Q1 or Q2, Q3 or Q4, Q5 or Q6, Q7 or Q8.
- 2) Neat diagrams must be drawn wherever necessary.
- 3) Figures to the right side indicate full marks.
- 4) Use of logarithmic tables slide rule, Mollier charts, electronic pocket calculator and steam tables is allowed.
- 5) Assume suitable data, if necessary.

Physical Constants :

- 1) Mass of Electron (M_e) = 9.1×10^{-31} kg
- 2) Charge on Electron (e) = 1.6×10^{-19} C
- 3) Mass of Proton (M_p) = 1.673×10^{-27} kg
- 4) Mass of neutron (M_n) = 1.673×10^{-27} kg
- 5) Velocity of light (c) = 3×10^8 m/s
- 6) Plank's constant (h) = 6.63×10^{-34} J.s

- Q1)** a) What is the significance of Schrodinger's equation? Derive schrodinger's time independent equation. [6]
- b) Derive the expression for de Broglie wavelength for a particle when it is moving with kinetic energy 'E'. [4]
- c) With a schematic diagram, explain in brief the principle of scanning tunneling microscope. [4]
- d) If a proton is moving with velocity 2% of the velocity of light, calculate de Broglie wavelength. [4]

OR

- Q2)** a) For a particle trapped in one-dimensional rigid box (infinite potential well), derive the equation for its energy. Explain in brief why this energy is quantized. [6]

P.T.O.

- b) For an electron accelerated by potential difference V , derive the expression for its de Broglie wavelength. [4]
- c) What is de Broglie hypothesis of matter waves? Explain in brief why matter waves are neither electromagnetic nor mechanical waves. [4]
- d) An electron is trapped in an infinite potential well of width 1.75\AA . Calculate energy difference between ground and first energy level. [4]

- Q3)**
- a) Explain Hall effect with suitable diagram. Derive the expression for Hall voltage and Hall coefficient. [6]
 - b) Draw a neat and labelled diagram showing I-V characteristics of a solar cell. Write the equations for fill factor and efficiency of solar cell. [4]
 - c) Define Fermi level in metals. Write Fermi-Dirac probability distribution function and explain the meaning of each term. [4]
 - d) Calculate number of donor atoms which must be added to an intrinsic semiconductor to obtain a resistivity of 12 ohm-cm (Mobility of electrons, $\mu_e = 500\text{ cm}^2/\text{V.S.}$). [3]

OR

- Q4)**
- a) Explain classification of solids into conductors, semiconductors and insulators on the basis of band theory. [6]
 - b) Define Fermi level in semiconductor. Draw neat & labelled diagram showing position of Fermi level in intrinsic, N-type and P-type semiconductors. [4]
 - c) State the advantages (any two) and applications (any two) of solar cell. [4]
 - d) An N type semiconductor has a thickness 0.12 mm and a current of 1mA is flowing along its length. When a magnetic field of 1.5 T is applied along its thickness, calculate voltage developed across its width. [Hall coefficient, $R_H = 3.68 \times 10^{-4}\text{ m}^3/\text{C}$] [3]
- Q5)**
- a) On the basis of orbital and spin motion of electron and spin motion of nucleus, explain origin of magnetism. [6]
 - b) Define (i) magnetic field strength (ii) magnetic induction (iii) magnetic permeability. Write relation between them [4]

- c) What is super conductivity? Differentiate between type I & type II super conductors (any three). [4]
- d) The transition temperature T_c for lead is 7.26 K. The critical magnetic field at 0K i.e. $H_c(0)$ is 8×10^5 A/m. It is to be used as a super conductor subjected to a magnetic field $H_c(T)$ of 4×10^4 A/m. At what maximum temperature T, it can be operated without losing its superconductivity. [4]

OR

- Q6)**
- a) Explain Meissner effect in brief. What is the cause of Meissner effect? Show that super conductors exhibit perfect diamagnetism. [6]
 - b) What is magnetic permeability (μ) & magnetic susceptibility (χ). Obtain relation between them. [4]
 - c) Differentiate between diamagnetism and ferromagnetism (Any two points) [4]
 - d) For Niobium, if critical temperature T_c is 11.3 K and critical magnetic field at 0K is. $H_c(0)$ is 2×10^5 A/m, Calculate critical magnetic field $H_c(T)$ at $T = 4.2K$. [4]
- Q7)**
- a) Explain electrical and mechanical properties of nanoparticles. [6]
 - b) Write applications of nanotechnology in the field of electronics. Explain any one application in brief. [4]
 - c) What are ultrasonic waves? An ultrasonic wave of velocity 6400 m/s is sent through the top of an aluminium block of thickness 4.5 cm. It is reflected back from the internal flaw. After what time echo would be obtained. [4]
 - d) Explain in brief, the principle of radiography testing technique. [3]

OR

- Q8)**
- a) What is Non-Destructive Testing (NDT)? State objectives of NDT (any two). State advantages of NDT over DT (any two). [6]
 - b) Write applications of nanotechnology in the field of automobiles. Explain any one application in brief. [4]
 - c) What are ultrasonic waves? An ultrasonic wave is sent through a block of steel and its echo is received after 1.5 μs . if velocity of ultrasonic waves is 5890 m/s, determine location of the flaw. [4]
 - d) Explain optical properties of nanoparticles. [3]

