

Exercise 1.

Let $X = \{a\}$ and $Y = \{b, c\}$.

- List all possible functions from X to Y (you may draw a diagram).
- List all functions from Y to X .
- Compose a function from the first list with a function from the second list (in either order). What are all the different functions that arise in this way?

- Solution.*
- To specify a function $f: X \rightarrow Y$, we need to describe the image of each element of X . Since X has only one element a , any function $f: X \rightarrow Y$ is uniquely determined by $f(a) \in Y = \{b, c\}$. That is, there are exactly two functions f_1 and f_2 in Y^X , which are defined by $f_1(a) = b$ and $f_2(a) = c$.
 - To define a function $g: Y \rightarrow X$ is equivalent to choose the value of $g(y) \in X$ for all $y \in Y$. Since $X = \{a\}$ has only one element, we are forced to choose $g(y) = a$ for all $y \in Y$. So there exists a unique function $g: Y \rightarrow X$, defined by $g(b) = g(c) = a$.
 - Let us give the details for $f_1 \circ g$:

$$\begin{aligned} f_1 \circ g: Y &\xrightarrow{g} X \xrightarrow{f_1} Y \\ b &\mapsto a \mapsto b \\ c &\mapsto a \mapsto b. \end{aligned}$$

So $f_1 \circ g$ is the constant function b . That is: $(f_1 \circ g)(y) = b$ for all $y \in Y$. Similarly, $f_2 \circ g$ is the constant function c . Composing in the other direction we obtain that both $g \circ f_1$ and $g \circ f_2$ are the constant function a .

Observe that we have $g \circ f_1 = g \circ f_2 = \text{Id}_X$ with $f_1 \neq f_2$, but $f_1 \circ g \neq f_2 \circ g$ and none of them is equal to Id_Y . ■

Exercise 2 (Proposition 1.2.17).

Let X, Y, Z be sets and suppose that $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are functions.

- Show that if f and g are injective then $g \circ f$ is injective.
- Show that if f and g are surjective then $g \circ f$ is surjective.
- If $g \circ f$ is injective then what can you say about f or g ? What if $g \circ f$ is surjective?

- Solution.*
- Suppose that both f and g are injective and let $a, b \in X$ be two elements of X such that $(g \circ f)(a) = (g \circ f)(b)$. We need to prove that in this case one

necessary have $a = b$. From $(g \circ f)(a) = (g \circ f)(b)$ one obtain $g(f(a)) = g(f(b))$ and so $f(a) = f(b)$ by injectivity of g and finally $a = b$ by injectivity of f .

- b. Suppose that both f and g are surjective and let $z \in Z$ be any element of Z . By surjectivity of g , there exists $y \in Y$ such that $g(y) = z$. By surjectivity of f , there exists $x \in X$ such that $f(x) = y$. But then $(g \circ f)(x) = g(y) = z$, proving surjectivity of $g \circ f$.

- c. We will show that $g \circ f$ injective implies f injective, but g might not be injective; and that $g \circ f$ surjective implies g surjective, but f might not be surjective.

We start by providing a counterexample to the two incorrect implications. Let f_1 and g be the functions from the solution to Exercise 1.a. Then $g \circ f_1 = \text{Id}_X$ is both injective and surjective. However, g is not injective as $g(b) = g(c)$ even if $b \neq c$, and f_1 is not surjective as $c \notin \text{Im}(f_1)$.

Suppose now that $g \circ f$ is injective. Let $a, b \in X$ be any two elements with $f(a) = f(b)$. By applying g on both sides we have $(g \circ f)(a) = (g \circ f)(b)$ and hence $a = b$ by injectivity of $g \circ f$. We have just proved injectivity of f .

Finally, suppose that $g \circ f$ is surjective. Let $z \in Z$ be any element. Then, by surjectivity of $g \circ f$, there exists $x \in X$ such that $z = (g \circ f)(x) = g(f(x))$. Since $y := f(x)$ is in Y , we have $z \in \text{Im}(g)$, proving surjectivity of g .

■

Recall that the **identity map** on a set X is the function $\text{Id}_X: X \rightarrow X$ given by $\text{Id}_X(x) = x$ for all $x \in X$.

Exercise 3.

Let $f: X \rightarrow Y$ and $g: Y \rightarrow X$ be functions. Suppose that $g \circ f = \text{Id}_X$. Does it follow that $f \circ g = \text{Id}_Y$?

Solution. No. See the solution of Question 1 for a counterexample. ■

Exercise 4 (See Proposition 2.1.3).

Recall that $\mathbf{F}$ stands for $\mathbf{R}$ or $\mathbf{C}$. Let m be an integer. Prove that $\mathbf{F}^m$ satisfies the *distributive properties*, that is, for all $a, b \in \mathbf{F}$ and $u, v \in \mathbf{F}^m$ we have:

1. $a(u + v) = au + av$;
2. $(a + b)v = av + bv$.

Solution. Let $u = [u_1, \dots, u_m]^\top$ and $v = [v_1, \dots, v_m]^\top$. Then

$$a(u + v) = a \left(\begin{bmatrix} u_1 \\ \vdots \\ u_m \end{bmatrix} + \begin{bmatrix} v_1 \\ \vdots \\ v_m \end{bmatrix} \right) = a \begin{bmatrix} u_1 + v_1 \\ \vdots \\ u_m + v_m \end{bmatrix}$$

$$\begin{aligned}
&= \begin{bmatrix} a(u_1 + v_1) \\ \vdots \\ a(u_m + v_m) \end{bmatrix} = \begin{bmatrix} au_1 + av_1 \\ \vdots \\ au_m + av_m \end{bmatrix} \\
&= \begin{bmatrix} au_1 \\ \vdots \\ au_m \end{bmatrix} + \begin{bmatrix} av_1 \\ \vdots \\ av_m \end{bmatrix} = au + av,
\end{aligned}$$

where we used the distributivity property $a(u_1 + v_1) = au_1 + av_1$ in $\mathbf{F}$. ■

Further Questions

Exercise 5 (Proposition 1.2.23).

Let X and Y be finite sets and $f: X \rightarrow Y$ be a function.

- a. Assume that f is injective. Find a function $g: Y \rightarrow X$ such that $g \circ f = \text{Id}_X$.
- b. Assume that f is surjective. Find a function $g: Y \rightarrow X$ such that $f \circ g = \text{Id}_Y$.

Solution. a. Suppose X non-empty and let x_0 be any element of X . Define $g: Y \rightarrow X$ by:

$$g(y) := \begin{cases} \text{the unique } x \in X \text{ such that } f(x) = y & \text{if } y \in \text{Im}(f) \\ x_0 & \text{otherwise.} \end{cases}$$

It is then straightforward that $g \circ f = \text{Id}_X$.

To go further

If $X = \emptyset$ there is a unique function from X to Y : the empty function. So if $X = \emptyset = Y$, the empty function is its own inverse. If Y is not empty, then there is no function from Y to $\emptyset$.

Observe that we did not use that X and Y are finite. That is, we actually proved a bit more than what was stated. Moreover, one can check that the g we constructed is surjective. So we have proved: if $f: X \rightarrow Y$ is an injective function between arbitrary sets, there exists a surjective function $g: Y \rightarrow X$ with $g \circ f = \text{Id}_X$.

- b. If f is surjective, then $Y = \text{Im}(f)$. One can define $g: Y \rightarrow X$ by for each $y \in Y = \text{Im}(f)$ choosing a $x \in X$ such that $f(x) = y$. One can check that such a g is injective and $f \circ g = \text{Id}_Y$. So we have proved: if $f: X \rightarrow Y$ is an surjective function with Y finite, there exists an injective function $g: Y \rightarrow X$ with $f \circ g = \text{Id}_Y$.

To go further

If Y is infinite, we need the axiom of choice (AC) to be able to make infinitely many simultaneous choices. So, if we assume (AC) we have: if $f: X \rightarrow Y$ is a surjective function between arbitrary sets, there exists an injective function $g: Y \rightarrow X$ with $f \circ g = \text{Id}_Y$. Actually, this statement is equivalent to (AC). This is one of the reasons to assume (AC): it implies that injective and surjective functions behave “dually”.

■

The following questions are intended as revision for Year 1 Linear Algebra.

Exercise 6.

Solve the system of linear equations

$$\begin{aligned}x_1 + 3x_2 - 5x_3 &= 4 \\x_1 + 4x_2 - 8x_3 &= 7 \\-3x_1 - 7x_2 + 9x_3 &= -6.\end{aligned}$$

Solution. By doing the row transformations $r_2 \mapsto r_2 - r_1$ and $r_3 \mapsto r_3 + 3r_1$ we obtain the equivalent system

$$\begin{aligned}x_1 + 3x_2 - 5x_3 &= 4 \\x_2 - 3x_3 &= 3 \\2x_2 - 6x_3 &= 6.\end{aligned}$$

Doing one last row transformation $r_3 \mapsto r_3 - 2r_2$ one has the, also equivalent, system

$$\begin{aligned}x_1 + 3x_2 - 5x_3 &= 4 \\x_2 - 3x_3 &= 3 \\0 &= 0.\end{aligned}$$

We conclude that x_3 is free, $x_2 = 3 + 3x_3$ and $x_1 = 4 - 3x_2 + 5x_3 = -5 - 4x_3$. ■

Exercise 7.

Let $A = \begin{bmatrix} 2 & -2 & 1 \\ 3 & 5 & 0 \\ -1 & 3 & 4 \end{bmatrix}$. Compute $\det(A)$ and A^{-1} .

Solution. There are many ways to compute the determinant. We will compute it by

developing the determinant according to the last column, giving:

$$\begin{aligned}\det(A) &= 1 \cdot \det \begin{bmatrix} 3 & 5 \\ -1 & 3 \end{bmatrix} + 0 + 4 \cdot \det \begin{bmatrix} 2 & -2 \\ 3 & 5 \end{bmatrix} \\ &= (9 + 5) + 4(10 + 6) = 78.\end{aligned}$$

There are also many methods to compute the inverse of A . We will show some of them. Firstly, we will use the formula

$$A^{-1} = \frac{1}{\det(A)} \text{adj}(A),$$

where $\text{adj}(A) = ((-1)^{i+j} \det(\text{cofact}_{ij}))_{ij}^T$. Recall that cofact_{ij} is the matrix obtained from A by erasing the i -row and j -column. Let us unravel this for our specific example. So we have

$$\text{adj}(A)^T = ((-1)^{i+j} \det(\text{cofact}_{ij}))_{ij} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix},$$

where we need to compute the coefficients a to i . For example,

$$a = 1 \cdot \det \begin{bmatrix} 5 & 0 \\ 3 & 4 \end{bmatrix} = 20 \quad b = -1 \cdot \det \begin{bmatrix} 3 & 0 \\ -1 & 4 \end{bmatrix} = -12.$$

By computing all the coefficients, one obtains

$$\text{adj}(A)^T = \begin{bmatrix} 20 & -12 & 14 \\ 11 & 9 & -4 \\ -5 & 3 & 16 \end{bmatrix},$$

And finally

$$A^{-1} = \frac{1}{78} \begin{bmatrix} 20 & 11 & -5 \\ -12 & 9 & 3 \\ 14 & -4 & 16 \end{bmatrix}.$$

While the above method always works, it is lengthy. Another way to compute the inverse is to use operations on rows. The idea is to start with $[A \mid \text{Id}]$ and to do row transformations to obtain $[\text{Id} \mid B]$. The matrix B obtained with this process is A^{-1} . If we apply that to our concrete example, we obtain

$$A^{-1} = \begin{bmatrix} \frac{10}{39} & \frac{11}{78} & -\frac{5}{78} \\ -\frac{2}{13} & \frac{3}{26} & \frac{1}{26} \\ \frac{7}{39} & -\frac{2}{39} & \frac{8}{39} \end{bmatrix} = \frac{1}{78} \begin{bmatrix} 20 & 11 & -5 \\ -12 & 9 & 3 \\ 14 & -4 & 16 \end{bmatrix},$$

which is identical to the answer obtained via the cofactors method. The details are shown on the next page.

$$\begin{array}{c}
 r_1 \mapsto \frac{1}{2} r_1 \\
 r_2 \mapsto r_2 - \frac{3}{2} r_1 \\
 r_3 \mapsto r_3 + \frac{1}{2} r_1 \\
 \left[\begin{array}{ccc|ccc} 2 & -2 & 1 & 1 & 0 & 0 \\ 3 & 5 & 0 & 0 & 1 & 0 \\ -1 & 3 & 4 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\substack{r_1 \mapsto \frac{1}{2} r_1 \\ r_2 \mapsto r_2 - \frac{3}{2} r_1 \\ r_3 \mapsto r_3 + \frac{1}{2} r_1}} \left[\begin{array}{ccc|ccc} 1 & -1 & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & 8 & -\frac{3}{2} & -\frac{3}{2} & 1 & 0 \\ 0 & 2 & \frac{9}{2} & \frac{1}{2} & 0 & 1 \end{array} \right] \\
 \begin{array}{l} r_2 \mapsto \frac{1}{8} r_2 \\ r_3 \mapsto r_3 - \frac{1}{4} r_2 \end{array} \swarrow \\
 \left[\begin{array}{ccc|ccc} 1 & -1 & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & 1 & -\frac{3}{16} & -\frac{3}{16} & \frac{1}{8} & 0 \\ 0 & 0 & \frac{39}{8} & \frac{7}{8} & -\frac{1}{4} & 1 \end{array} \right] \xrightarrow{r_3 \mapsto \frac{8}{39} r_3} \left[\begin{array}{ccc|ccc} 1 & -1 & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & 1 & -\frac{3}{16} & -\frac{3}{16} & \frac{1}{8} & 0 \\ 0 & 0 & 1 & \frac{7}{39} & -\frac{2}{39} & \frac{8}{39} \end{array} \right] \\
 \begin{array}{l} r_1 \mapsto r_1 - \frac{1}{2} r_3 \\ r_2 \mapsto r_2 + \frac{3}{16} r_3 \end{array} \swarrow \\
 \left[\begin{array}{ccc|ccc} 1 & -1 & 0 & \frac{16}{39} & \frac{1}{39} & -\frac{4}{39} \\ 0 & 1 & 0 & -\frac{2}{13} & \frac{3}{26} & \frac{1}{26} \\ 0 & 0 & 1 & \frac{7}{39} & -\frac{2}{39} & \frac{8}{39} \end{array} \right] \xrightarrow{r_1 \mapsto r_1 + r_2} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{10}{39} & \frac{11}{78} & -\frac{5}{78} \\ 0 & 1 & 0 & -\frac{2}{13} & \frac{3}{26} & \frac{1}{26} \\ 0 & 0 & 1 & \frac{7}{39} & -\frac{2}{39} & \frac{8}{39} \end{array} \right].
 \end{array}$$

■

Exercise 1 (Lemma 2.2.19).

Let V be a vector space. Prove that $(-1)v = -v$ for all $v \in V$. (Before starting, think about what the minus sign on each side means.)

Solution. The left hand side of the equation is $-1_{\mathbf{F}} \cdot_{\mathbf{F}} v$. That is: the scalar multiplication of $-1 \in \mathbf{F}$ and $v \in V$, where -1 is the additive inverse of 1 in $\mathbf{F}$. The right hand side of the equation is $-_V v$, that is the additive inverse of v .

We have

$$v + (-1) \cdot v = 1 \cdot v + (-1) \cdot v = (1 + (-1))v = 0 \cdot v = 0.$$

We conclude that $(-1)v$ is an additive inverse of v and hence $(-1)v = -v$ by unicity of the additive inverse. ■

Exercise 2.

Determine if the following subsets of $\mathbf{R}^3$ are subspaces. Give reasons.

$$\text{a. } U = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \mathbf{R}^3 \mid 2x - 3y + z = 0 \right\}.$$

$$\text{b. } W = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \mathbf{R}^3 \mid xyz = 0 \right\}.$$

Solution. a. U is a subspace of $\mathbf{R}^3$. Firstly, $2 \cdot 0 - 3 \cdot 0 + 0 = 0$ so $[0, 0, 0]^T$ is in U . Secondly, let $u_1 = [x_1, y_1, z_1]^T$ and $u_2 = [x_2, y_2, z_2]^T$ be two elements of U and let $\lambda \in \mathbf{R}$ be a scalar. By assumption, we have

$$2x_1 - 3y_1 + z_1 = 0 \quad \text{and} \quad 2x_2 - 3y_2 + z_2 = 0.$$

By multiplying the first equation by λ and adding the result to the second equation, we obtain

$$\begin{aligned} 0 &= \lambda(2x_1 - 3y_1 + z_1) + (2x_2 - 3y_2 + z_2) \\ &= 2(\lambda x_1 + x_2) - 3(\lambda y_1 + y_2) + (\lambda z_1 + z_2). \end{aligned}$$

Therefore, $\lambda u_1 + u_2$ is in U , finishing the proof that U is a subspace.

One can show that U is in a plane containing the origin in $\mathbf{R}^3$.

b. W is not a subspace of $\mathbf{R}^3$. Indeed, both $[1, 1, 0]^T$ and $[0, 0, 1]^T$ belong to W , but their sum $[1, 1, 0]^T + [0, 0, 1]^T = [1, 1, 1]^T$ is not in W .

Geometrically, W is the union of the three coordinate planes (the xy -plane, the xz -plane and the yz plane), see Figure 1.

Observe that the fact that U is a subspace of $\mathbf{F}^3$ but W is not remains true for an arbitrary field $\mathbf{F}$. ■




Figure 1: The union of the three coordinates planes.

Exercise 3.

A sequence $(x_0, x_1, x_2, \dots) \in \mathbf{R}^{\mathbf{N}}$ is *increasing* if $x_m \leq x_{m+1}$ for all $m \in \mathbf{N}$. Does the set X of all increasing sequences form a subspace of $\mathbf{R}^{\mathbf{N}}$? Give reasons.

Solution. No, because X is not closed under scalar multiplication. For example $(0, 1, 1, \dots)$ is in X , but $-1 \cdot (0, 1, 1, \dots) = (0, -1, -1, \dots)$ is not. ■

Exercise 4.

Let V be a vector space and $U \subseteq V$ be a subspace. Suppose that $u \in U$ and $v \in V$. Prove that $v \in U$ if and only if $u + v \in U$.

Solution. If v is in U , then $u + v$ is also in U because U is a subspace.

If $u + v$ is in U , then $v = -1 \cdot u + (u + v)$ is also in U because U is a subspace. ■

Exercise 5.

Let U be the xy -plane in $\mathbf{R}^3$. Find distinct subspaces W_1, W_2 of $\mathbf{R}^3$ such that $U \oplus W_1 = U \oplus W_2 = \mathbf{R}^3$.

Solution. Let $W_1 := \{[0, 0, z]^T \mid z \in \mathbf{R}\}$ and let $W_2 := \{[0, z, z]^T \mid z \in \mathbf{R}\}$. Then $U \oplus W_1 = U \oplus W_2 = \mathbf{R}^3$.

For each W_i , we need to prove three things: 1. that it is a subspace of $\mathbf{R}^3$, 2. that $U + W_i = \mathbf{R}^3$, 3. and finally that $U + W_i$ is a direct sum. We will only check the last two properties, and only do it for W_2 . The proof for W_1 is similar (and even simpler) and the proof that the W_i are subspaces of $\mathbf{R}^3$ is elementary.

Let $[x, y, z]^T$ be any vector of $\mathbf{R}^3$. Then $[x, y, z]^T = [x, y - z, 0]^T + [0, z, z]^T$ is in $U + W_2$ and so $U + W_2 = \mathbf{R}^3$. Finally, suppose that $0 = u + w$ with $u \in U$ and $w \in W_2$. So

$[0, 0, 0]^T = [x, y, 0]^T + [0, z, z]^T$ for some $x, y, z \in \mathbf{R}$. But this implies

$$\begin{cases} 0 = x \\ 0 = y + z \\ 0 = z \end{cases}$$

and we hence conclude that both u and w are the 0 vector. This shows that the sum $U + W_2$ is direct.

Observe that a similar proof shows that $U \oplus W = \mathbf{R}^3$ if W is any line containing the origin and not included in U .

To go further

The above statement remains true if we replace $\mathbf{R}$ by $\mathbf{F}$, in which case a *line* simply means a subspace of dimension 1. That is, a *line containing the origin* is any subspace of the form $\{[\lambda_1 z, \lambda_2 z, \lambda_3 z]^T \mid z \in \mathbf{F}\}$ for some $\lambda_1, \lambda_2, \lambda_3$ with at least one $\lambda_i \neq 0$.

■

Further Questions

Exercise 6 (See Lemma 2.3.28).

Is addition of subspaces commutative? That is, for subspaces U, W of V , is it true that $U + W = W + U$? Explain.

Solution. Addition of subspaces is commutative, because addition of vectors is. Indeed, a vector $v \in V$ belongs to $U + W$ if and only if there exists $u \in U$ and $w \in W$ such that $v = u + w$. By commutativity of *vectors addition*: $u + w = w + u$. So $v \in V$ belongs to $U + W$ if and only if there exists $u \in U$ and $w \in W$ such that $v = w + u$, that is if and only if v belongs to $W + U$.

Observe that we did not use the fact that U and W are subspaces. Therefore, addition of subsets (of a vector space) is commutative. ■

Exercise 7.

Does every subspace of V have an *additive inverse*? That is, for each subspace U of V , does there exist a subspace W such that $U + W = \{0\}$?

Solution. No. This is the case if and only if $U = \{0\}$, in which case $W = \{0\}$. Indeed, we always have $U \subseteq U + W$. So if $U + W = \{0\}$ for some W , then $U = \{0\}$. ■

Exercise 8.

Recall that a function $f: \mathbf{R} \rightarrow \mathbf{R}$ is *even* if $f(-x) = f(x)$ for all $x \in \mathbf{R}$; and *odd* if $f(-x) = -f(x)$ for all $x \in \mathbf{R}$. Let U_e and U_o respectively denote the set of even and odd functions in $\mathbf{R}^{\mathbf{R}}$. Prove that $\mathbf{R}^{\mathbf{R}} = U_e \oplus U_o$.

Solution. Once again, we need to prove 3 things: 1. that both U_e and U_o are subspaces of $\mathbf{R}^{\mathbf{R}}$, 2. that $U_e + U_o = \mathbf{R}^{\mathbf{R}}$, 3. and finally that $U_e + U_o$ is a direct sum.

Firstly, the constant function 0 is both even and odd. Moreover, if f and g are two even functions (respectively odd functions) and $\lambda \in \mathbf{R}$ a scalar, one easily verify that $\lambda f + g$ is still even (respectively odd). Indeed, this directly follows from $(\lambda f + g)(x) = \lambda f(x) + g(x)$. We conclude that both U_e and U_o are subspaces of $\mathbf{R}^{\mathbf{R}}$.

Then, let f be any function of $\mathbf{R}^{\mathbf{R}}$. Define two functions f_e and f_o by:

$$f_e(x) := \frac{f(x) + f(-x)}{2}, \quad f_o(x) := \frac{f(x) - f(-x)}{2}.$$

One directly have $(f_e + f_o)(x) = f(x)$ for all $x \in \mathbf{R}$ and so $f_e + f_o = f$. We also have

$$f_e(-x) = \frac{f(-x) + f(x)}{2} = \frac{f(x) + f(-x)}{2} = f_e(x)$$

and

$$f_o(-x) = \frac{f(-x) - f(x)}{2} = -\frac{f(x) - f(-x)}{2} = -f_o(x).$$

So f_e is even and f_o is odd, showing that f is in $U_e + U_o$. This finishes the proof that $U_e + U_o = \mathbf{R}^{\mathbf{R}}$.

Finally, let f be in $U_e \cap U_o$. Then for every $x \in \mathbf{R}$, let $y = -x$. We have:

$$f(-y) = f(-y) = -f(-y).$$

But this implies that $f(x) = f(-y) = -f(-y) = -f(x)$ and so $f(x) = 0$. Therefore, f is the zero function, $U_e \cap U_o = \{0\}$ and the sum is direct. ■

Exercise 1.

Let (u_1, u_2, u_3) be a list of vectors of a vector space V .

- Show that if (u_1, u_2, u_3) span V then so does the list $(u_1 - u_2, u_2 - u_3, u_3)$.
- Show that if (u_1, u_2, u_3) is linearly independent then so is the list $(u_1 - u_2, u_2 - u_3, u_3)$.

Solution. a. We will show that $\text{span}(u_1, u_2, u_3) = \text{span}(u_1 - u_2, u_2 - u_3, u_3)$, even though they are not necessarily equal to V . The $\supseteq$ inclusion is clear, so let us prove $\text{span}(u_1, u_2, u_3) \subseteq \text{span}(u_1 - u_2, u_2 - u_3, u_3)$. Let $v \in \text{span}(u_1, u_2, u_3)$. There exists scalars λ_1, λ_2 and λ_3 such that

$$\begin{aligned} v &= \lambda_1 u_1 + \lambda_2 u_2 + \lambda_3 u_3 \\ &= \lambda_1(u_1 - u_2) + (\lambda_2 + \lambda_1)(u_2 - u_3) + (\lambda_3 + \lambda_2)u_3. \end{aligned}$$

Since all of $\lambda_1, \lambda_2 + \lambda_1$ and $\lambda_3 + \lambda_2$ are in $\mathbf{F}$, we have just shown that v belongs to $\text{span}(u_1 - u_2, u_2 - u_3, u_3)$, which finishes the proof.

- Suppose that (u_1, u_2, u_3) is linearly independent and let λ_1, λ_2 and λ_3 be scalars such that

$$0 = \lambda_1(u_1 - u_2) + \lambda_2(u_2 - u_3) + \lambda_3 u_3.$$

We have to show that this implies that all the λ_i are zeros. We have

$$0 = \lambda_1 u_1 + (\lambda_2 - \lambda_1)u_2 + (\lambda_3 - \lambda_2)u_3.$$

By linear independence of (u_1, u_2, u_3) , all the coefficients $\lambda_1, \lambda_2 - \lambda_1$ and $\lambda_3 - \lambda_2$ are zeros. This directly implies that all the λ_i are 0, as required. ■

Exercise 2.

Consider the subspace $U = \{p \in \mathcal{P}(\mathbf{R})_2 \mid p(1) = 0\}$ of $\mathcal{P}(\mathbf{R})_2$.

- Find a basis for U . (Hint: if a polynomial p satisfies $p(a) = 0$ then $(x - a)$ is a factor of $p(x)$.)
- Extend the basis you found to a basis of $\mathcal{P}(\mathbf{R})_2$.
- Find a subspace W of $\mathcal{P}(\mathbf{R})_2$ such that $U \oplus W = \mathcal{P}(\mathbf{R})_2$.

Solution. We will show the result for a general field $\mathbf{F}$.

- On one hand, we know that $\mathcal{P}(\mathbf{F})_2$ is of dimension 3 (with basis $(1, x, x^2)$). On the other hand, $U \neq \mathcal{P}(\mathbf{F})_2$ (for example, because $x \notin U$). By [Question 3](#), this implies that U is of dimension at most 2. So if we find a linearly independent family of

length 2 we are done. For example, one can take the family $(p_1 = x - 1, p_2 = x^2 - 1)$. We claim that this family is linearly independent. Indeed, if $0 = \lambda p_1 + \mu p_2$, then

$$0 = \mu x^2 + \lambda x + (-\lambda - \mu)$$

and therefore $\mu = 0 = \lambda$, which proves the claim.

- b. We already have a length 2 linearly independent family. So for any $q \in \mathcal{P}(\mathbf{F})_2$ not in $\text{span}(p_1, p_2) = U$, the family (p_1, p_2, q) is linearly independent by the linear dependence lemma. Since it is a length 3 linear independent family in a dimension 3 vector space, it is linearly independent.

For a concrete example, one can take $q = 1$ the constant polynomial 1. But $q = x$ or $q = x^2$ also work.

- c. Let $q \notin U$ be any polynomial in $\mathcal{P}(\mathbf{F})_2$ but not in U . We claim that for $W := \text{span}(q)$ we have $U \oplus W = \mathcal{P}(\mathbf{F})_2$. It is clear from above that $U + W = \mathcal{P}(\mathbf{F})_2$. It remains to prove that the sum is direct. Let p be in $U \cap W$. We need to show that $p = 0$. We have $p = \lambda q \in U$. If $p \neq 0$, then $q = \lambda^{-1}p \in U$; a contradiction.

■

Exercise 3 (Corollary 2.4.45).

Let V be a finite dimensional vector space and suppose that $U \subseteq V$ is a subspace. Prove that if $\dim(U) = \dim(V)$ then $U = V$.

Solution. By assumption, $\dim(U) = \dim(V)$. So there exists a finite basis $\mathcal{B} = (u_1, \dots, u_n)$ of U , where $n = \dim(V)$. Then $\mathcal{B}$ is a linearly independent family of vectors of U , hence also of vectors of V . Since it has $\dim(V)$ elements it is spanning in V . So $U = \text{span}(\mathcal{B}) = V$.

■

Exercise 4.

Let U and W be 3-dimensional subspaces of $\mathbf{R}^5$. What are the possible values of $\dim(U \cap W)$? Can $U + W$ be a direct sum?

Solution. Once again, we will prove the statement for a general $\mathbf{F}$.

We have

$$\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W) \leq \dim(\mathbf{R}^5) = 5.$$

In other words, $\dim(U \cap W) = \dim(U) + \dim(W) - \dim(U + W) \geq 3 + 3 - 5 = 1$. But we also have $\dim(U + W) \leq \dim(U) = 3$. So, $\dim(U + W)$ belongs to $\{1, 2, 3\}$. We now show that all this value can happens.

If $U = W_1$, for example $U = W_1 = \{[x_1, x_2, x_3, 0, 0]^T \mid x_i \in \mathbf{F}\}$, then $\dim(U \cap W_1) = 3$.

For U as above and $W_2 = \{[0, x_2, x_3, x_4, 0]^T \mid x_i \in \mathbf{F}\}$ the subspace $U \cap W_2 = \{[0, x_2, x_3, 0, 0]^T \mid x_i \in \mathbf{F}\}$ is of dimension 2.

Finally, for U as above and $W_3 = \{[0, 0, x_3, x_4, x_5]^T \mid x_i \in \mathbf{F}\}$ the subspace $U \cap W_3 = \{[0, 0, x_3, 0, 0]^T \mid x_i \in \mathbf{F}\}$ is of dimension 1.

Since $\dim(U \cap W) \geq 1$, the sum $U + W$ is never direct. Indeed, the sum $U + W$ is direct if and only if $U \cap W = \{0\}$, if and only if $\dim(U \cap W) = 0$. ■

Exercise 5.

Let V be a vector space. Suppose that for each integer $m > 0$, there exists a list of m linearly independent vectors in V . Prove that V is infinite dimensional.

Solution. By Grassman's exchange lemma, if V has a linearly independent family of length m , then any spanning family has length at least m . In particular, any base of V has at least m elements, for all $m \in \mathbf{N}$. We conclude that a basis cannot be finite. ■

Further Questions

Exercise 6.

Prove that $\mathbf{R}^{\mathbf{N}}$ is infinite dimensional. (More generally, if S is an infinite set then $\mathbf{F}^{(S)}$ and $\mathbf{F}^S$ are infinite dimensional.)

Solution. By Question 5, it is enough to exhibit arbitrary long linearly independent lists. For any $i \in \mathbf{N}_{\geq 1}$, define $e_i = (0, \dots, 0, 1, 0, \dots)$ to be list with a 1 in coordinate i and 0 everywhere else. We claim that for any m , the family $(e_1, \dots, e_m)$ is linearly independent. Indeed, let λ_i be scalars such that

$$0 = \sum_{i=1}^m \lambda_i e_i.$$

Then

$$(0, 0, \dots) = (\lambda_1, \dots, \lambda_m, 0, \dots)$$

forcing all the λ_i to be 0.

A similar proof shows that $\mathbf{F}^{(S)}$ (and thus also $\mathbf{F}^S$) are infinite dimensional as soon as S is infinite. Indeed, for $s \in S$ one can define a map $\chi_s: S \rightarrow \mathbf{F}$ by

$$\chi_s(t) = \begin{cases} 1 & \text{if } t = s \\ 0 & \text{otherwise.} \end{cases}$$

We now claim that for any finite subset $A \subseteq S$, the family $(\chi_a)_{a \in A}$ is linearly independent. Indeed, let $(\lambda_a)_{a \in A}$ be a family of scalars such that

$$0 = \sum_{a \in A} \lambda_a \chi_a.$$

Then, for $a \in A$ one can evaluate both sides of the above equality on a to obtain $0 = \lambda_a \cdot 1$, and thus $\lambda_a = 0$.

To go further

It is in fact possible to show that for any S , finite or infinite, we have $\dim(\mathbf{F}^{(S)}) = \#S$. Recall that for finites S , one has $\mathbf{F}^{(S)} = \mathbf{F}^S$.

■

Exercise 7.

Explain why all the subspaces of $\mathbf{R}^3$ are precisely $\{0\}$, all straight lines through the origin, all planes through the origin, and $\mathbf{R}^3$.

Solution. First of all, let us remind that a line in $\mathbf{R}^3$ is a subset of the form

$$L_{u,v} = \{\lambda u + v \mid \lambda \in \mathbf{R}\}$$

for some vectors v and $u \neq 0$. A line $L_{u,v}$ goes through the origin if and only if $v = 0$. Let us fix $u \in \mathbf{R}^3$. One easily check that $0 \in L_{u,0}$ and that $L_{u,0}$ is closed under addition and scalar multiplication. In consequence $L_{u,0}$ is a subspace of $\mathbf{R}^3$, of dimension 1 ((u) is a basis).

Similarly, a plane in $\mathbf{R}^3$ is a subset of the form

$$P_{u_1,u_2,v} = \{\lambda_1 u_1 + \lambda_2 u_2 + v \mid \lambda_1, \lambda_2 \in \mathbf{R}\}$$

for some vectors v and non-collinear (that is linearly independent) u_1 and u_2 . Once again, a plane $P_{u_1,u_2,v}$ goes through the origin if and only if $v = 0$ and $P_{u_1,u_2,0}$ is a subspace of $\mathbf{R}^3$, of dimension 2 ((u_1, u_2) is a basis).

Finally, it is clear that both $\{0\}$ and $\mathbf{R}^3$ are subspaces of $\mathbf{R}^3$ (of respective dimension 0 and 3). We will now prove that any subspace U of $\mathbf{R}^3$ is of the form $\{0\}$, $L_{u,0}$ (for some non-zero u), $P_{u_1,u_2,0}$ (for some linearly independent u_1 and u_2) or $\mathbf{R}^3$. To do this, we will do a case-by-case analysis depending on $\dim(U) \in \{0, \dots, 3\}$.

If $\dim(U) = 0$, then $U = \{0\}$.

If $\dim(U) = 3 = \dim(\mathbf{R}^3)$, then $U = \mathbf{R}^3$ by [Question 3](#).

Suppose now that $\dim(U) = 1$ and take any $u \neq 0$ in U . Then $\dim(\text{span}(u)) = 1 = \dim(U)$ and thus $U = \text{span}(u) = \{\lambda u \mid \lambda \in \mathbf{R}\} = L_{u,0}$ is a line through the origin. We also have the converse: any line through the origin is a subspace of dimension 1.

Suppose finally that $\dim(U) = 2$. Take any $u_1 \neq 0$ in U . Since $\dim(\text{span}(u_1)) = 1 < 2 = \dim(U)$, there exists $u_2 \in U \setminus \text{span}(u_1)$. By the linear dependence lemma, the family (u_1, u_2) is linearly independent and hence a basis of U . So $U = \text{span}(u_1, u_2) = \{\lambda_1 u_1 + \lambda_2 u_2 \mid \lambda_i \in \mathbf{R}\} = P_{u_1,u_2,0}$ is a plane containing the origin. We also have the converse: any plane containing the origin is a subspace of dimension 2. ■




Figure 2: Subspaces of dimension 0 (the origin), 1 (line through the origin) and 2 (plane through the origin) of $\mathbb{R}^3$. The last kind of subspace is $\mathbb{R}^3$ itself.

Exercise 1.

Determine whether the following functions are linear maps.

- $T_1: \mathbf{R}^2 \rightarrow \mathbf{R}$ given by $T_1\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = xy$.
- $T_2: \mathbf{C} \rightarrow \mathbf{C}$ given by $T_2(z) = \bar{z}$, where $\mathbf{C}$ is regarded as a real vector space ($a + bi = a - bi$ is the complex conjugation).
- $T_2: \mathbf{C} \rightarrow \mathbf{C}$ given by $T_2(z) = \bar{z}$, where $\mathbf{C}$ is regarded as a complex vector space.
- $T_3: \mathcal{P}(\mathbf{R}) \rightarrow \mathbf{R}$ given by $T_3(p) = p(2)$.

Solution. a. T_1 is not linear. Indeed, $T_1\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = T_1\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}\right) = 1 \cdot 1 = 1$ while $T_1\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + T_1\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = 1 \cdot 0 + 0 \cdot 1 = 0 \neq 1$.

To go further

We have actually showed for any $\mathbf{F}$ the map $T_1: \mathbf{F}^2 \rightarrow \mathbf{F}, \begin{bmatrix} x \\ y \end{bmatrix} \mapsto xy$ is not linear.

- T_2 is $\mathbf{R}$ -linear. Indeed, for $z, w \in \mathbf{C}$ and $\lambda \in \mathbf{R}$ we have $T_2(\lambda z + w) = \overline{\lambda z + w} = \overline{\lambda z} + \bar{w} = \lambda \bar{z} + \bar{w} = \lambda T_2(z) + T_2(w)$. The above equalities can be verified explicitly by writing $z = a + bi$ and $w = c + di$.
- T_2 is not $\mathbf{C}$ -linear. Indeed, $iT_2(1) = i\bar{1} = i$, but $T_2(i1) = \bar{i} = -i \neq i$.
- T_3 is linear. Indeed, let p and q be two polynomials in $\mathcal{P}(\mathbf{R})$ and $\lambda \in \mathbf{R}$. Then $T_3(\lambda p + q) = (\lambda p + q)(2) = \lambda p(2) + q(2) = \lambda T_3(p) + T_3(q)$.

To go further

The same proof shows that for any field $\mathbf{F}$, and any element $a \in \mathbf{F}$, the map $\mathcal{P}(\mathbf{F}) \rightarrow \mathbf{F}, p \mapsto p(a)$ is linear. Even more generally, if S is any set then the map $\mathbf{F}^S \rightarrow \mathbf{F}, f \mapsto f(a)$ is linear. This map is called the **evaluation at a** .

■

For a linear map $T \in \mathcal{L}(U, V)$, we write $\ker(T) = \text{null}(T)$ for its kernel (also called nullspace): $\ker(T) = \{u \in U \mid T(u) = 0\}$. We write $\text{Im}(T) = \text{range}(T)$ for the image of T (also called range): $\text{Im}(T) = \{v \in V \mid \exists u \in U : T(u) = v\}$.

Exercise 2.

Let U, V , and W be vector spaces, and suppose $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$ are linear maps.

- Prove that $\ker(T)$ is a subspace of $\ker(ST)$.

b. Prove that $\text{Im}(ST)$ is a subspace of $\text{Im}(S)$.

Solution. We have

$$\begin{array}{ccccc} U & \xrightarrow{T} & V & \xrightarrow{S} & W. \\ & \searrow & \nearrow & & \\ & & ST & & \end{array}$$

- a. Both T and ST have domain U , so both $\ker(T)$ and $\ker(ST)$ are subspaces of U . It is thus enough to prove that $\ker(T)$ is a *subset* of $\ker(ST)$. Let u be any element of $\ker(T)$, so $T(u) = 0$. Then $(ST)(u) = S(T(u)) = S(0) = 0$ and hence u is in $\ker(ST)$ as desired.
- b. Both S and ST have codomain W , so both $\text{Im}(S)$ and $\text{Im}(ST)$ are subspaces of W . It is thus enough to prove that $\text{Im}(ST)$ is a *subset* of $\text{Im}(S)$. Let w be any element in $\text{Im}(ST)$. So there exists $u \in U$ with $(ST)u = w$. In particular, there exists $v = T(u)$ in V (the domain of S) such that $S(v) = S(T(u)) = (ST)(u) = w$ and hence w is in $\text{Im}(S)$ as desired.

Remark 0.1.

The inclusion $\ker(T) \subseteq \ker(ST)$ might be interpreted as *ST is less injective than T* . We recover the particular case: if ST is injective then so is T .

The inclusion $\text{Im}(ST) \subseteq \text{Im}(S)$ might be interpreted as *ST is less surjective than S* . We recover the particular case: if ST is surjective then so is S .

■

Exercise 3.

Let $T \in \mathcal{L}(V, W)$ be an injective linear map. Suppose that $(v_1, \dots, v_k)$ is a linearly independent list of vectors in V . Prove that $(Tv_1, \dots, Tv_k)$ is a linearly independent list in W .

Solution. Let $\lambda_1, \dots, \lambda_k$ be scalars such that

$$0 = \sum_{j=1}^k \lambda_j Tv_j.$$

We need to prove that all the λ_j are 0. By linearity of T we have $0 = T\left(\sum_{j=1}^k \lambda_j v_j\right)$. By injectivity of T one obtains

$$0 = \sum_{j=1}^k \lambda_j v_j.$$

By linear independence of the $(v_1, \dots, v_k)$ we conclude that all the λ_j are 0. ■

Exercise 4.

Let V, W be vector spaces such that W is finite-dimensional. Let $T \in \mathcal{L}(V, W)$ be a linear map.

- Prove that T is surjective if and only if there exists a linear map $S \in \mathcal{L}(W, V)$ such that $TS = \text{Id}_W$. (Hint: Define S on any basis of W then extend linearly.)
- Prove that T is injective if and only if there exists a linear map $S \in \mathcal{L}(W, V)$ such that $ST = \text{Id}_V$. (Hint: Similar to the above, but choose a basis of W carefully.)

Solution. First of all, we know that for functions $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ we have:

- if $g \circ f$ is injective then f is injective;
- if $g \circ f$ is surjective then g is surjective.

Since Id_X is bijective, this directly shows:

- if $ST = \text{Id}_V$ then T is injective;
- if $TS = \text{Id}_W$ then T is surjective.

Observe that we don't need to suppose that W is finite dimensional, or that S is a linear map (S a function is enough).

For the other direction, we will use bases to translate our problem for linear maps on vector spaces to a problem about functions on sets. We can do this, because a linear map $T: V \rightarrow W$ is uniquely determined by what it does on a given basis $\mathcal{B}$ of V .

- Let $n = \dim W$. Suppose that T is surjective and let $\mathcal{C} = (w_1, \dots, w_n)$ be any basis of W . For any $w_j \in \mathcal{C}$, choose a preimage $v_j \in V$ so $T(v_j) = w_j$. There exists at least one such v_j by surjectivity. We define $S: \mathcal{C} \rightarrow V$ by $S(w_j) = v_j$ and extend it to a linear map $S \in \mathcal{L}(W, V)$. Then for any w_j we have $TSw_j = w_j$. Since TS is the identity on the basis $\mathcal{C}$, it is the identity everywhere: $TS = \text{Id}_W$.
- Suppose that T is injective and let $\mathcal{D} = (w_1, \dots, w_k)$ be a basis of $\text{Im}(T)$. Since T is injective, for every $j \in \{1, \dots, k\}$ there exists a unique $v_j \in V$ with $Tv_j = w_j$. We claim that $\mathcal{A} = (v_1, \dots, v_k)$ is a basis of V .

First, we prove that $\mathcal{A}$ is a spanning family for V . Let v be any vector in V . Then $Tv \in \text{Im}(T) = \text{span}(\mathcal{D})$ so there exists $\lambda_j, j \in \{1, \dots, k\}$ such that

$$T(v) = \sum_{j=1}^k \lambda_j w_j = \sum_{j=1}^k \lambda_j T v_j = T\left(\sum_{j=1}^k \lambda_j v_j\right).$$

By injectivity, $v = \sum_{j=1}^k \lambda_j v_j$ is in $\text{span}(\mathcal{A})$.

We now prove that $\mathcal{A}$ is linearly independent. Suppose that

$$0 = \sum_{j=1}^k \lambda_j v_j$$

for some λ_j . Applying T on both sides, one obtain $0 = \sum_{j=1}^k \lambda_j w_j$ and conclude that all the λ_j are 0, proving linear independence of $\mathcal{A}$.

Now, since $\mathcal{D}$ is a basis of $\text{Im}(T) \subseteq W$, we can extend it to a basis $\mathcal{C} = (w_1, \dots, w_k, w_{k+1}, \dots, w_m)$ of W . We define S on $\mathcal{C}$ by:

$$S(w_j) = \begin{cases} v_j & \text{if } j \in \{1, \dots, k\} \\ 0 & \text{otherwise} \end{cases}$$

and then extend it to a linear map $S \in \mathcal{L}(W, V)$. One have $ST(v_j) = v_j$ for all $j \in \{1, \dots, k\}$ and so $ST = \text{Id}_V$.

Infinite dimensional vector spaces

The statements remain true for a general, not necessarily finite dimensional, vector space W if we assume (AC). We use (AC) in two different places. Firstly, for the existence of a basis of W , and secondly when T is injective and we *choose* a preimage of w_j .

■

Exercise 5.

Rephrase the following problems in terms of linear maps between vector spaces. (Do not solve the problems.)

- Find all smooth functions $f: \mathbf{R} \rightarrow \mathbf{R}$ that satisfy $f'' - 3f' + 2f = 0$.
- Find all real sequences $(x_0, x_1, \dots)$ that satisfy $x_{n+2} = x_{n+1} + x_n + 1$ for all $n \in \mathbf{N}$.

Solution. a. The space of smooth functions is $\mathcal{C}^\infty(\mathbf{R}) = \{f: \mathbf{R} \rightarrow \mathbf{R} \mid f \text{ smooth}\} = \{f: \mathbf{R} \rightarrow \mathbf{R} \mid \forall n: f^{(n)} \text{ exists}\}$. Let $D: \mathcal{C}^\infty(\mathbf{R}) \rightarrow \mathcal{C}^\infty(\mathbf{R}), f \mapsto f'$ be the differentiation operator. This is a linear map. Define $T := D^2 - 3D + 2\text{Id}$, where $D^2 = D \circ D$. This is also a linear map from $\mathcal{C}^\infty(\mathbf{R})$ to itself and $T(f) = f'' - 3f' + 2f$. We have $f'' - 3f' + 2f = 0$ if and only if $f \in \ker(T)$.

- The space of real sequences is $\mathbf{R}^{\mathbf{N}} = \{(x_0, x_1, x_2, \dots) \mid \forall j: x_j \in \mathbf{R}\}$. Let $S: \mathbf{R}^{\mathbf{N}} \rightarrow \mathbf{R}^{\mathbf{N}}$ be the backward shift: $S(x_0, x_1, \dots) = (x_1, x_2, \dots)$. Finally, let $c = (1, 1, \dots)$ be the constant sequence 1. We want to find all sequences $x \in \mathbf{R}^{\mathbf{N}}$ such that $S^2(x) = S(x) + x + c$. So if we define $T = S^2 - S - \text{Id}$, a sequence x satisfies the questions if and only if $T(x) = c$.

■

Further Questions**Exercise 6.**

Let U , V , and W be vector spaces, and suppose $S_1, S_2 \in \mathcal{L}(U, V)$ and $T \in \mathcal{L}(V, W)$ are linear maps. Prove that the following distributive property holds: $T(S_1 + S_2) = TS_1 + TS_2$.

Solution. This is similar to the proof of $(T_1 + T_2)S = T_1S + T_2S$ viewed in class. Let u be any vector in U . Then

$$\begin{aligned} (T(S_1 + S_2))u &= T((S_1 + S_2)u) \\ &= T(S_1u + S_2u) \\ &= T(S_1u) + T(S_2u) \\ &= (TS_1)u + (TS_2)u = (TS_1 + TS_2)u \end{aligned}$$

and thus $T(S_1 + S_2) = TS_1 + TS_2$. ■

Exercise 7.

Let $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$ be linear maps. Show that $ST = 0 \in \mathcal{L}(U, W)$ (the zero map) if and only if $\text{Im}(T)$ is a subspace of $\ker S$.

Solution. “ $\Rightarrow$ ” Suppose $ST = 0$. Let v be any element in $\text{Im}(T)$. By assumption, there exists $u \in U$ such that $Tu = v$. But then, $Sv = S(Tu) = (ST)u = 0u = 0$ and thus v is in $\ker(S)$ as desired.

“ $\Leftarrow$ ” Suppose that $\text{Im}(T) \subseteq \ker(S)$. Then for any $u \in U$ we have $(ST)(u) = S(Tu) = 0$, showing that ST is the 0 map. ■

Exercise 1.

Let $D: \mathcal{P}(\mathbf{R}) \rightarrow \mathcal{P}(\mathbf{R})$ be the differentiation operator. Is D invertible?

Solution. The operator D is not injective and hence not invertible. Indeed, both the constant polynomial 0 and the constant polynomial 1 are sent onto 0. ■

Exercise 2.

Let U be the subspace of $\mathcal{C}^\infty(\mathbf{R})$ spanned by the basis $\mathcal{B}$ given by (e^x, xe^x, x^2e^x) . Let $D \in \mathcal{L}(U)$ be the differentiation operator. Find the matrix $[D]_{\mathcal{B}}^{\mathcal{B}}$.

Solution. We compute each column of $[D]_{\mathcal{B}}^{\mathcal{B}}$.

Firstly, $D(e^x) = e^x = 1 \cdot e^x + 0 \cdot xe^x + 0 \cdot x^2e^x$ so $[D(e^x)]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Then, $D(xe^x) = e^x + xe^x$ and therefore $[D(xe^x)]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$. Finally, $D(x^2e^x) = 2xe^x + x^2e^x$ and therefore $[D(x^2e^x)]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$.

Altogether, we have

$$[D]_{\mathcal{B}}^{\mathcal{B}} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

■

Exercise 3.

Let $T \in \mathcal{L}(V, W)$ be a linear map and suppose $S_1, S_2 \in \mathcal{L}(W, V)$ satisfy $S_1T = \text{Id}_V$ and $TS_2 = \text{Id}_W$. Prove that T is invertible. What is the inverse of T ?

Solution. We have

$$S_1 = S_1 \text{Id}_W = S_1TS_2 = \text{Id}_V S_2 = S_2.$$

Therefore $S_1 = S_2$ is the inverse of T , which is therefore invertible. ■

Exercise 4.

Let $T \in \mathcal{L}(V, W)$ be a surjective linear map. Suppose that U is a subspace of V such that $\ker(T) \oplus U = V$. Prove that the restriction $T|_U: U \rightarrow W$ is an isomorphism (recall that the restriction is defined by $T|_U(u) = Tu$ for all $u \in U$).

Solution. The map $T|_U$ is the restriction of a linear map, and thus linear. It remains to show that it is injective and surjective.

We first prove injectivity. We both have $\ker(T|_U) \subseteq U$ and $\ker(T|_U) \subseteq \ker(T)$. We conclude $\ker(T|_U) \subseteq U \cap \ker(T)$. But since the sum $U \oplus \ker(T)$ is direct, we have $U \cap \ker(T) = \{0\}$, proving that $T|_U$ is injective.

We now prove surjectivity. Let $w \in W$ be any vector. By surjectivity of T , there exists $v \in V$ such that $T(v) = w$. Since $V = \ker(T) \oplus U$, there exist (unique) $u_1 \in \ker(T)$ and $u_2 \in U$ such that $v = u_1 + u_2$. So $w = T(v) = T(u_1 + u_2) = T(u_1) + T(u_2) = 0 + T(u_2) = T|_U(u_2)$, proving surjectivity of $T|_U$. ■

Exercise 5.

Let $x_1, \dots, x_{n+1} \in \mathbf{R}$ be distinct. Define a linear map $T: \mathcal{P}(\mathbf{R})_n \rightarrow \mathbf{R}^{n+1}$ by $T(p) := [p(x_1), \dots, p(x_{n+1})]^\top$.

- Prove that T is injective. (Hint: By the division algorithm, any non-zero degree n polynomial has at most n distinct roots.)
- Show that for any $y_1, \dots, y_{n+1} \in \mathbf{R}$, there exists a unique polynomial p of degree at most n such that $p(x_i) = y_i$ for all $1 \leq i \leq n+1$. (Hint: Show that T is an isomorphism.)

Solution. a. Let p be any polynomial in $\ker(T)$. So $p(x_1) = \dots = p(x_{n+1}) = 0$. That is, p is a polynomial of degree at most n with (at least) $n+1$ distinct roots. We conclude that p is the zero polynomial and so that $\ker(T) = \{0\}$.

- Both $\mathcal{P}(\mathbf{R})_n$ and $\mathbf{R}^{n+1}$ have dimension $n+1$. Therefore T being injective is also surjective, and hence bijective. In particular, for any $y_1, \dots, y_{n+1} \in \mathbf{R}$ there exists a unique polynomial $p \in \mathcal{P}(\mathbf{R})_n$ such that $T(p) = [y_1, \dots, y_{n+1}]^\top$. ■

Further Questions

Exercise 6 (Continuing from Question 5, if you want to solve it explicitly).

Let $(e_1, \dots, e_{n+1})$ be the standard basis for $\mathbf{R}^{n+1}$ and $p_i = T^{-1}(e_i) \in \mathcal{P}(\mathbf{R})_n$. In other words, p_i is the unique polynomial of degree at most n satisfying $p_i(x_i) = 1$ and $p_i(x_j) = 0$ for $i \neq j$.

- Find an explicit formula for $p_i(x)$. (Hint: a non-zero polynomial p satisfies $p(a) = 0$ if and only if $(x - a)$ is a factor of p .)
- Write down $T^{-1}[y_1, \dots, y_{n+1}]^\top$ in terms of the p_i 's and y_i 's. (This is known as Lagrange's Interpolation Formula.)

Solution. a. We first construct polynomials that satisfy $q_i(x_j) = 0$ if $i \neq j$, but might not have the correct value for $q_i(x_i) = 0$. This is quite easy.

Let $q_1(x) := (x - x_2) \cdots (x - x_{n+1})$ and define similarly $q_2 := (x - x_1)(x - x_3) \cdots (x - x_{n+1})$ and $q_3, \dots, q_n$ up to $q_{n+1}(x) := (x - x_1)(x - x_2) \cdots (x - x_{n+1} + 1)$. In other

words,

$$q_i = \prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} (x - x_k).$$

Then for any $1 \leq j \leq n+1$ one have

$$q_i(x_j) = \begin{cases} 0 & \text{if } i \neq j, \\ \prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} (x_i - x_k) \neq 0 & \text{if } i = j. \end{cases}$$

It is now easy to construct the p_i , by normalising their value at x_i :

$$p_i = \frac{\prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} (x - x_k)}{\prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} (x_i - x_k)} = \prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} \frac{(x - x_k)}{(x_i - x_k)}.$$

b. We have $[y_1, \dots, y_{n+1}]^T = y_1 e_1 + \dots y_{n+1} e_{n+1}$. By linearity

$$T^{-1}[y_1, \dots, y_{n+1}]^T = \sum_{i=1}^{n+1} y_i T^{-1}(e_i) = \sum_{i=1}^{n+1} y_i p_i = \sum_{i=1}^{n+1} y_i \prod_{\substack{1 \leq k \leq n+1 \\ k \neq i}} \frac{(x - x_k)}{(x_i - x_k)}.$$

■

Exercise 7.

Prove that V is isomorphic to $\mathcal{L}(\mathbf{F}, V)$. (Hint: consider the function $F: \mathcal{L}(\mathbf{F}, V) \rightarrow V$ defined by $F(T) = T(1)$. This function is called evaluation at 1.)

Solution. We need to prove that F is a bijective linear map.

We first prove injectivity. Let T be an element of the kernel of F . That is: $T: \mathbf{F} \rightarrow V$ is a linear map such that $T(1) = 0$. But then, for every $\lambda \in \mathbf{F}$ we have $T(\lambda) = T(\lambda \cdot 1) = \lambda T(1) = \lambda \cdot 0 = 0$. This proves that $\ker(F) = \{0\}$ and so F is injective.

For surjectivity, let v be any vector in V . Since 1 is a basis of $\mathbf{F}$, there exists a unique linear map $T: \mathbf{F} \rightarrow V$ satisfying $T(1) = v$. We have $F(T) = v$ as desired.


Alternatively, one can prove bijectivity of F by finding an inverse. Let $G: V \rightarrow \mathcal{L}(\mathbf{F}, V)$ be the map defined by $G(v): \mathbf{F} \rightarrow V, \lambda \mapsto \lambda \cdot v$. That is, each v is sent by G onto the map $G(v) = T$ from the surjectivity proof. One easily verify that $(G \circ F)(T) = T$ and $(F \circ G)(v) = v$, proving that G is the inverse of F .

Finally, we show that F is linear. Let $S, T \in \mathcal{L}(\mathbf{F}, V)$ be two linear maps and let $\lambda \in \mathbf{F}$ be a scalar. Then $F(\lambda S + T) = (\lambda S + T)(1) = \lambda \cdot S(1) + T(1) = \lambda F(S) + F(T)$ as desired. ■

Exercise 1.

Let $R_\theta: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be a rotation about the origin through an angle of θ . Find the matrix $[R_\theta]_{\mathcal{E}}$, where $\mathcal{E}$ is the standard basis for $\mathbf{R}^2$. Use this to prove that $R_\theta \circ R_\phi = R_{\theta+\phi}$. (Hint: use the trigonometric formulas $\cos(a+b) = \cos(a)\cos(b) - \sin(a)\sin(b)$ and $\sin(a+b) = \sin(a)\cos(b) + \cos(a)\sin(b)$.)

Solution. We have $\mathcal{E} = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right)$. Simple trigonometry allows us to compute the coordinates of $R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.



So we have

$$R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \quad \text{and} \quad R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix},$$

which implies

$$[R_\theta]_{\mathcal{E}} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

We then compute

$$\begin{aligned} [R_\theta \circ R_\phi]_{\mathcal{E}} &= [R_\theta]_{\mathcal{E}} [R_\phi]_{\mathcal{E}} \\ &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta \cos \phi - \sin \theta \sin \phi & -\cos \theta \sin \phi - \sin \theta \cos \phi \\ \sin \theta \cos \phi + \cos \theta \sin \phi & -\sin \theta \sin \phi + \cos \theta \cos \phi \end{bmatrix} \\ &= \begin{bmatrix} \cos(\theta + \phi) & -\sin(\theta + \phi) \\ \sin(\theta + \phi) & \cos(\theta + \phi) \end{bmatrix} = [R_{\theta+\phi}]_{\mathcal{E}}. \end{aligned}$$


We conclude $R_\theta \circ R_\phi = R_{\theta+\phi}$ as desired. ■

Exercise 2.

Let $S_\theta: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ be a reflection across the line through the origin which makes an angle of θ with the positive x -axis. Let $\mathcal{B}$ be the basis of $\mathbf{R}^2$ given by $v_1 = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$

and $v_2 = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$. Find $[S_\theta]_{\mathcal{B}}$. Then find $[S_\theta]_{\mathcal{E}}$ by applying the change of basis formula, where $\mathcal{E}$ is the standard basis of $\mathbf{R}^2$.

Solution.



Since S_θ is a reflection along $l = \text{span}(v_1)$ and $v_2 \perp v_1$ we have $S_\theta(v_1) = v_1$ and $S_\theta(v_2) = -v_2$. Therefore, $[S_\theta(v_1)]_{\mathcal{B}} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $[S_\theta(v_2)]_{\mathcal{B}} = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$. Altogether we obtain

$$[S_\theta]_{\mathcal{B}} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

The change of basis formula gives us:

$$[S_\theta]_{\mathcal{E}} = [\text{Id}]_{\mathcal{B}}^{\mathcal{E}} [R_\theta]_{\mathcal{B}} [\text{Id}]_{\mathcal{E}}^{\mathcal{B}}.$$

So we need to compute $[\text{Id}]_{\mathcal{B}}^{\mathcal{E}}$ and $[\text{Id}]_{\mathcal{E}}^{\mathcal{B}} = ([\text{Id}]_{\mathcal{B}}^{\mathcal{E}})^{-1}$. We have $\text{Id } v_1 = \cos \theta e_1 + \sin \theta e_2$ and $\text{Id } v_2 = -\sin \theta e_1 + \cos \theta e_2$. This gives us

$$[\text{Id}]_{\mathcal{B}}^{\mathcal{E}} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad \text{and} \quad [\text{Id}]_{\mathcal{E}}^{\mathcal{B}} = ([\text{Id}]_{\mathcal{B}}^{\mathcal{E}})^{-1} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

Finally, one have

$$\begin{aligned} [S_\theta]_{\mathcal{E}} &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos^2 \theta - \sin^2 \theta & 2 \cos \theta \sin \theta \\ 2 \cos \theta \sin \theta & \sin^2 \theta - \cos^2 \theta \end{bmatrix} = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}. \end{aligned}$$

■

Exercise 3.

Consider a basis $\mathcal{B}$ of $\mathbf{R}^2$ given by $v_1 = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$ and $v_2 = \begin{bmatrix} -4 \\ 3 \end{bmatrix}$. Let T be a transformation that stretches in the v_1 -direction by a factor of 2, and in the v_2 -direction by a factor of 3. Compute $[T]_{\mathcal{B}}^{\mathcal{B}}$.

Solution. We have $Tv_1 = 2v_1$ and $Tv_2 = 3v_2$, which gives $[T]_{\mathcal{B}}^{\mathcal{B}} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$. The change of basis matrices are given by

$$[\text{Id}]_{\mathcal{B}}^{\mathcal{E}} = \begin{bmatrix} 3 & -4 \\ 4 & 3 \end{bmatrix} \quad \text{and} \quad [\text{Id}]_{\mathcal{E}}^{\mathcal{B}} = ([\text{Id}]_{\mathcal{B}}^{\mathcal{E}})^{-1} = \frac{1}{25} \begin{bmatrix} 3 & 4 \\ -4 & 3 \end{bmatrix}.$$

Finally,

$$[T]_{\mathcal{E}}^{\mathcal{E}} = \begin{bmatrix} 3 & -4 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \frac{1}{25} \begin{bmatrix} 3 & 4 \\ -4 & 3 \end{bmatrix} = \frac{1}{25} \begin{bmatrix} 3 & -4 \\ 4 & 3 \end{bmatrix} \begin{bmatrix} 6 & 8 \\ -12 & 9 \end{bmatrix} = \frac{1}{25} \begin{bmatrix} 66 & -12 \\ -12 & 59 \end{bmatrix}.$$

■

Exercise 4.

Let U be the subspace of $\mathcal{C}^\infty(\mathbf{R})$ spanned by the basis $\mathcal{B}$ given by (e^x, xe^x, x^2e^x) . Let $D \in \mathcal{L}(U)$ be the differentiation operator. Recall from Tutorial 5 that

$$[D]_{\mathcal{B}}^{\mathcal{B}} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

Is D invertible? If so, compute $[D^{-1}]_{\mathcal{B}}^{\mathcal{B}}$ and use this to find the unique function $f \in U$ satisfying $f' = x^2e^x$.

Solution. Since $\det([D]_{\mathcal{B}}^{\mathcal{B}}) = 1$, the matrix $[D]_{\mathcal{B}}^{\mathcal{B}}$ is invertible and so D is invertible. We compute

$$\underbrace{\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}}_{[D]_{\mathcal{B}}^{\mathcal{B}}} \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}}_{\text{Id}} \xrightarrow[r_2 \mapsto r_2 - 2r_3]{r_1 \mapsto r_1 - r_2 + 2r_3} \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}}_{\text{Id}} \underbrace{\begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}}_{([D]_{\mathcal{B}}^{\mathcal{B}})^{-1}}.$$

So

$$([D]_{\mathcal{B}}^{\mathcal{B}})^{-1} = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

and f is given by

$$[f]_{\mathcal{B}} = ([D]_{\mathcal{B}}^{\mathcal{B}})^{-1} [x^2e^x]_{\mathcal{B}} = \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}.$$

In other words, $f = 2e^x - 2xe^x + x^2e^x$. ■

Further Questions

Exercise 5 (Continuing from Question 2).

Compute $[S_\theta \circ S_\phi]_{\mathcal{E}}^{\mathcal{E}}$. Explain what the transformation $S_\theta \circ S_\phi$ does geometrically.

Solution.

$$\begin{aligned} [S_\theta \circ S_\phi]_{\mathcal{E}}^{\mathcal{E}} &= [S_\theta]_{\mathcal{E}}^{\mathcal{E}} [S_\phi]_{\mathcal{E}}^{\mathcal{E}} \\ &= \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix} \begin{bmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{bmatrix} \\ &= \begin{bmatrix} \cos 2\theta \cos 2\phi + \sin 2\theta \sin 2\phi & \cos 2\theta \sin 2\phi - \sin 2\theta \cos 2\phi \\ \sin 2\theta \cos 2\phi - \cos 2\theta \sin 2\phi & \sin 2\theta \sin 2\phi + \cos 2\theta \cos 2\phi \end{bmatrix} \\ &= \begin{bmatrix} \cos(2\theta - 2\phi) & -\sin(2\theta - 2\phi) \\ \sin(2\theta - 2\phi) & \cos(2\theta - 2\phi) \end{bmatrix} = [R_{2\theta-2\phi}]_{\mathcal{E}}^{\mathcal{E}}. \end{aligned}$$

We conclude that the linear map $S_\theta \circ S_\phi$ is the rotation of angle $2\theta - 2\phi$. ■

Exercise 6.

Let B and D be invertible $n \times n$ matrices over $\mathbf{F}$. Let $\mathcal{B} = (b_1, \dots, b_n)$ and $\mathcal{D} = (d_1, \dots, d_n)$ be bases of $\mathbf{F}^n$, where b_j and d_j are respectively the j^{th} column of B and D . Explain how you can compute $[\text{Id}_{\mathbf{F}^n}]_{\mathcal{D}}^{\mathcal{D}}$ from B and D .

Solution. Let $\mathcal{E}$ be the standard basis of $\mathbf{F}^n$. Since the b_j s and d_j s are column vectors in $\mathbf{F}^n$ we have $b_j = [b_j]_{\mathcal{E}}$ and $d_j = [d_j]_{\mathcal{E}}$. So

$$B = [b_1 \ b_2 \ \dots \ b_n] = [[b_1]_{\mathcal{E}} \ [b_2]_{\mathcal{E}} \ \dots \ [b_n]_{\mathcal{E}}] = [\text{Id}]_{\mathcal{B}}^{\mathcal{E}} \quad \text{and} \quad D = [\text{Id}]_{\mathcal{D}}^{\mathcal{E}}.$$

Since D is invertible, we obtain

$$[\text{Id}_{\mathbf{F}^n}]_{\mathcal{D}}^{\mathcal{D}} = [\text{Id}]_{\mathcal{E}}^{\mathcal{D}} [\text{Id}]_{\mathcal{B}}^{\mathcal{E}} = D^{-1}B.$$

■

Exercise 1.

Let $P \in \mathcal{L}(V)$ be a projection, i.e. $P^2 = P$. Let $b \in V$. Show that there exists a solution $x \in V$ to $Px = b$ if and only if $Pb = b$.

Solution. If $Pb = b$, then $x = b$ is a solution to $Px = b$.

Suppose that x is a solution to $b = Px$. Then applying P to both sides we have $Pb = P^2x = Px = b$. ■

Exercise 2.

Suppose that $P \in \mathcal{L}(V)$ is a projection. Prove that $\text{Id} - P \in \mathcal{L}(V)$ is also a projection (it is sometimes called the *complementary projection* to P). What is the relationship between the image and kernel of P and $\text{Id} - P$?

Solution. The identity function $\text{Id} = \text{Id}_V$ is in $\mathcal{L}(V)$. So P is in $\mathcal{L}(V)$ if and only if $\text{Id} - P$ is in $\mathcal{L}(V)$, which proves that $\text{Id} - P$ is linear. We have

$$(\text{Id} - P)^2 = \text{Id}^2 - \text{Id}P - P\text{Id} + P^2 = \text{Id} - 2P + P = \text{Id} - P,$$

showing that $\text{Id} - P$ is a projection.

A vector v is in $\ker(P)$ if and only if $P(v) = 0$, if and only if $(\text{Id} - P)v = v$, if and only if (by Question 1) v is in the image of $\text{Id} - P$. We conclude that $\ker(P) = \text{Im}(\text{Id} - P)$.

Since $\text{Id} - P$ is a projection, we have $\ker(\text{Id} - P) = \text{Im}(\text{Id} - (\text{Id} - P)) = \text{Im}(P)$. ■

Exercise 3.

Show that the matrix

$$A = \frac{1}{3} \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

represents a projection map on $\mathbf{F}^3$. Use this to determine whether the equations $Ax = b$ and $Ax = c$ are consistent, where $b = [1, 1, 0]^\top$ and $c = [1, -1, -1]^\top$. Write b and c in the form $u + u'$ where $u \in \text{col } A$ and $u' \in \text{null } A$.

Solution. Recall that a $m \times n$ matrix A , represents the linear map $L_A \in \mathcal{L}(\mathbf{F}^n, \mathbf{F}^m)$ defined by $L_A(v) = Av$. The linear map L_A is a projection if and only if $A^2 = A$.

We need to show that $A^2 = A$. A simple computation gives us

$$A^2 = \frac{1}{9} \begin{bmatrix} 6 & 3 & 3 \\ 3 & 6 & -3 \\ 3 & -3 & 6 \end{bmatrix} = A.$$

We verify

$$Ab = \frac{1}{3} \begin{bmatrix} 3 \\ 3 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = b \quad \text{and} \quad Ac = \frac{1}{3} \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \neq c.$$

By Question 1, $Ax = b$ is consistent, but $Ax = c$ is not.

We have $\text{col}(A) = \text{Im}(L_A)$ and $\text{null } A = \ker(L_A)$. It immediately follows that $b = b + 0$ and $c = 0 + c$ as b belongs to $\text{Im}(L_A)$ and c belongs to $\ker(L_A)$. ■

Exercise 4.

Let $\mathbf{F}$ be either $\mathbf{R}$ or $\mathbf{C}$ and let V be an $\mathbf{F}$ -vector space. Suppose that $P \in \mathcal{L}(V)$ is an operator. Let $T = 2P - \text{Id} \in \mathcal{L}(V)$.

- Prove that P is a projection if and only if $T^2 = \text{Id}$.
- Suppose that P is an orthogonal projection to a line L through the origin in $\mathbf{R}^2$. Give a geometric description of the operator T .


Solution. a. We have $T^2 = (2P - \text{Id})^2 = 4P^2 - 4P + \text{Id} = \text{Id} + 4(P^2 - P)$. Therefore, $T^2 = \text{Id}$ if and only if $P^2 = P$.

To go further

In any field $\mathbf{F}$, if P is a projection, then $(2P - \text{Id})^2 = \text{Id}$. The converse is not true in general. Indeed, it might happen that $4 = 0$ in $\mathbf{F}$. This is the case in $\mathbf{F}_2 = \{0, 1\}$ for example. If $4 = 0$ in $\mathbf{F}$ then one can show that $2 = 0$ and thus $-1 = 1$. This implies $T = \text{Id}$ is independent of P .

For a concrete example, let $V = (\mathbf{F}_2)^2$. This is a $\mathbf{F}_2$ -vector space with 4 elements: $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Then the map $S \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ is a linear map but not a projection as $S^2 = \text{Id} \neq S$. However, $(2S - \text{Id}) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2y - x \\ 2x - y \end{bmatrix} = \begin{bmatrix} -x \\ -y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}$.

- T is the orthogonal reflection across L as demonstrated in the following pictures.



Exercise 5.

Let $T \in \mathcal{L}(V, W)$ and $S \in \mathcal{L}(W, V)$. Given $b \in W$, consider the linear problem $Tx = b$. Prove the following:

- If $TS = \text{Id}_W$ then $x = Sb$ is a solution to $Tx = b$ (in particular, the problem is consistent for all $b \in W$).
- If $ST = \text{Id}_V$ then $Tx = b$ is consistent if and only if $TSb = b$; in which case there is a unique solution $x = Sb$.

Solution. a. We have $T(Sb) = (TS)b = \text{Id}b = b$.

b. If $b = TSb = T(Sb)$, the system is consistent.

If the system is consistent, then $Tx = b$ for some x . Applying S on both sides we obtain $Sb = STx = x$. This shows both that Sb is a solution and also that it is the unique solution. ■

Further Questions**Exercise 6.**

Let $D \in \mathcal{L}(\mathcal{P}(\mathbf{R}))$ be the differentiation operator. Find a right-inverse for D . Does D have a left-inverse?

Solution. Intuitively, we want the right inverse to be the integration map $\int p(x) dx$. Integration is defined up to an additive constant only. Since we want our map to be linear we probably want to impose $T0 = 0$ and more generally that the constant coefficient of Tp is 0 for all polynomial.

Formally, let $T: \mathcal{P}(\mathbf{R}) \rightarrow \mathcal{P}(\mathbf{R})$ be the unique linear map such that $T(x^n) = \frac{1}{n+1}x^{n+1}$. We have $T(a_0 + \dots + a_n x^n) = a_0 x + \dots + \frac{1}{n+1}x^{n+1}$. One easily verify that we have $DT(p) = p$ for any polynomial.

The map D has no left-inverse, as the equality $SD = \text{Id}$ would implies that D is injective, which is absurd. ■

Exercise 7.

Suppose $T \in \mathcal{L}(V, W)$ and $S \in \mathcal{L}(W, V)$ satisfy $TST = T$ and $STS = S$. Show that TS and ST are both projections. What are their images/kernel? (Hint: The answers can be chosen from $\ker(S)$, $\text{Im}(S)$, $\ker(T)$, $\text{Im}(T)$.)

Solution. We always have $\ker(T) \subseteq \ker(ST)$ and $\text{Im}(ST) \subseteq \text{Im}(S)$ (Tutorial 4, Question 2).

Suppose that $TST = T$. Then $(ST)^2 = STST = S(TST) = ST$ and so ST is a projection. A vector v is in $\ker(ST)$ if and only if $STv = 0$. This implies $0 = TSTv = Tv$. We conclude that $\ker(ST) \subseteq \ker(T)$ as soon as $TST = T$.

Suppose now that $STS = S$. Then $(ST)^2 = STST = (STS)T = ST$ is a projection. Finally, let $v \in \operatorname{Im}(S)$. Then there exists w such that $v = Sw$. But then $STv = STSw = Sw = v$ and therefore $\operatorname{Im}(S) = \operatorname{Im}(ST)$.

The proofs for TS are similar. ■

Exercise 1.

Let V be an inner product space. Suppose that $u, v \in V$ are vectors satisfying $\langle u, u \rangle = 4$, $\langle v, v \rangle = 6$ and $\langle u, v \rangle = -2$. What is $\|3u - 2v\|$?

Solution. We have

$$\begin{aligned}\|3u - 2v\|^2 &= \langle 3u - 2v, 3u - 2v \rangle = 9\langle u, u \rangle - 6\langle u, v \rangle - 6\langle v, u \rangle + 4\langle v, v \rangle \\ &= 9 \cdot 4 - 6 \cdot (-2) - 6 \cdot \overline{-2} + 4 \cdot 6 = 84.\end{aligned}$$

So $\|3u - 2v\| = \sqrt{84} = 2\sqrt{21}$. ■

Exercise 2.

Consider the vector space $\mathcal{P}(\mathbf{R})$ equipped with the inner product $\langle p, q \rangle = \int_0^1 p(x)q(x) dx$. Write the polynomial x as a sum $p + q$, where p is a scalar multiple of 1 and q is orthogonal to 1.

Solution. Recall that if V is an inner product space and u and v are two vectors with $v \neq 0$ we have $u = \frac{\langle u, v \rangle}{\langle v, v \rangle}v + w$ with w orthogonal to v . In our case, we have $p = \frac{\langle x, 1 \rangle}{\langle 1, 1 \rangle}1$ and $q = x - p$. To find p we compute

$$\langle x, 1 \rangle = \int_0^1 x dx = \left[\frac{1}{2}x^2 \right]_0^1 = \frac{1}{2} \quad \text{and} \quad \langle 1, 1 \rangle = \int_0^1 1 dx = [x]_0^1 = 1.$$

So $p = \frac{1}{2}$ and $q = x - \frac{1}{2}$. ■

Exercise 3.

Prove the converse of the Pythagorean Theorem:

If u and v are vectors in a real inner product space V satisfying $\|u\|^2 + \|v\|^2 = \|u + v\|^2$ then u and v must be orthogonal.

Explain why the assumption that V is a real inner product space is necessary.

Solution. We have

$$\begin{aligned}\|u + v\|^2 &= \langle u + v, u + v \rangle = \langle u, u \rangle + \langle u, v \rangle + \langle v, u \rangle + \langle v, v \rangle \\ &= \|u\|^2 + \|v\|^2 + \langle u, v \rangle + \langle v, u \rangle.\end{aligned}$$

Therefore, the equality $\|u\|^2 + \|v\|^2 = \|u + v\|^2$ implies (in fact is equivalent to) $0 = \langle u, v \rangle + \langle v, u \rangle = \langle u, v \rangle + \langle u, v \rangle$. If V is a real inner product space we conclude $0 = \langle u, v \rangle$. In other words, u and v are orthogonal.

The above proof does not work for complex inner product spaces. But what about the conclusion? It also fails for any complex inner product space $V \neq \{0\}$. Indeed,

let $v \neq 0$ be a non-zero vector and let $u = iv$. Then u and v are not orthogonal since $\langle u, v \rangle = i \cdot \langle v, v \rangle \neq 0$. However, we have $\langle u, v \rangle + \langle v, u \rangle = i\langle v, v \rangle + \bar{i}\langle v, v \rangle = 0$ and thus $\|u\|^2 + \|v\|^2 = \|u + v\|^2$. ■

Exercise 4.

Recall that the **mean** of n real numbers $x_1, \dots, x_n$ is equal to $\frac{1}{n} \sum_{j=1}^n x_j$. Prove that the square of the mean of $x_1, \dots, x_n$ is less than or equal to the mean of $x_1^2, \dots, x_n^2$.

Solution. We want to prove that

$$\left(\frac{1}{n} \sum_{j=1}^n x_j \right)^2 \leq \frac{1}{n} \sum_{j=1}^n x_j^2,$$

or equivalently that

$$\left(\sum_{j=1}^n x_j \right)^2 \leq n \sum_{j=1}^n x_j^2.$$

In order to do that, we will use the Cauchy-Schwarz inequality: $|\langle u, v \rangle|^2 \leq \|u\|^2 \|v\|^2$ for $u = [x_1, \dots, x_n]^\top$ and $v = [1, \dots, 1]^\top$ in $\mathbf{R}^n$ and the standard dot product and the corresponding norm $\|v\| = \sqrt{v \bullet v}$. One easily check

$$\begin{aligned} (u \bullet v)^2 &= (x_1 + \dots + x_n)^2, \\ \|u\|^2 &= x_1^2 + \dots + x_n^2, \\ \|v\|^2 &= n \end{aligned}$$

and thus the desired formula follows. ■

Exercise 5.

[From Calculus] Let $\gamma: \mathbf{R} \rightarrow \mathbf{R}^n$ be a **smooth path**, i.e., $\gamma(t) = [\gamma_1(t), \dots, \gamma_n(t)]^\top$ where each $\gamma_i: \mathbf{R} \rightarrow \mathbf{R}$ is a smooth function. Its derivative is given by $\gamma'(t) = [\gamma'_1(t), \dots, \gamma'_n(t)]^\top$. Prove that if γ and δ are smooth paths in $\mathbf{R}^n$, then for any t we have $(\gamma(t) \bullet \delta(t))' = (\gamma'(t) \bullet \delta(t)) + (\gamma(t) \bullet \delta'(t))$ for the standard dot product on $\mathbf{R}^n$.

Solution. For a fixed t , the dot product is $\gamma(t) \bullet \delta(t) = \sum_{i=1}^n \gamma_i(t) \delta_i(t)$. So the function $t \mapsto (\gamma(t) \bullet \delta(t))$ is a smooth function from $\mathbf{R}$ to $\mathbf{R}$ and

$$\begin{aligned} (\gamma(t) \bullet \delta(t))' &= \left(\sum_{i=1}^n \gamma_i(t) \delta_i(t) \right)' = \sum_{i=1}^n (\gamma_i(t) \delta_i(t))' \\ &= \sum_{i=1}^n (\gamma'_i(t) \delta_i(t) + \gamma_i(t) \delta'_i(t)) = \sum_{i=1}^n \gamma'_i(t) \delta_i(t) + \sum_{i=1}^n \gamma_i(t) \delta'_i(t) \\ &= (\gamma'(t) \bullet \delta(t)) + (\gamma(t) \bullet \delta'(t)). \end{aligned}$$

■

Further Questions

Exercise 6 (Continuing from Question 5).

Suppose that $\gamma: \mathbf{R} \rightarrow \mathbf{R}^n$ is a smooth path of constant speed, i.e. $\|\gamma'(t)\|$ is constant. Prove that $\gamma'(t)$ and $\gamma''(t)$ are orthogonal for all t . (This shows that the velocity and acceleration along a constant speed path are always orthogonal.)

Solution. Since $\|\gamma'(t)\|^2 = \gamma'(t) \bullet \gamma'(t)$ is constant, we have

$$0 = (\gamma'(t) \bullet \gamma'(t))' = (\gamma''(t) \bullet \gamma'(t)) + (\gamma'(t) \bullet \gamma''(t)) = 2\gamma''(t) \bullet \gamma'(t)$$

since $\gamma''(t) \bullet \gamma'(t)$ is a real number. Therefore $0 = \gamma''(t) \bullet \gamma'(t)$ for every $t \in \mathbf{R}$. ■

Exercise 7 (From Statistics).

Suppose that we have some data given as n pairs of real numbers $(x_1, y_1), \dots, (x_n, y_n)$. The **sample correlation coefficient** is given by the following formula:

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}},$$

where $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ and $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$. Prove that $-1 \leq r \leq 1$. What can you say about the value of r if:

- A fixed constant is added to all x values (or all y values)?
- We scale each x value (or each y value) by a fixed positive constant?

Solution. Let $x = [x_1, \dots, x_n]^T$, $y = [y_1, \dots, y_n]^T$ and $u = [1, \dots, 1]^T$ be vectors in $\mathbf{R}^n$. Then

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i = \frac{x \bullet u}{u \bullet u} \quad \text{and} \quad \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i = \frac{y \bullet u}{u \bullet u}.$$

Now, let

$$\tilde{x} = x - \frac{x \bullet u}{u \bullet u} u = [x_1 - \bar{x}, \dots, x_n - \bar{x}]^T \quad \text{and} \quad \tilde{y} = y - \frac{y \bullet u}{u \bullet u} u = [y_1 - \bar{y}, \dots, y_n - \bar{y}]^T.$$

We therefore have

$$r = \frac{\tilde{x} \bullet \tilde{y}}{\|\tilde{x}\| \|\tilde{y}\|}.$$

By the Cauchy-Schwarz inequality we have $|\tilde{x} \bullet \tilde{y}| \leq \|\tilde{x}\| \|\tilde{y}\|$, that is: $-1 \leq r \leq 1$.

- Adding a fixed constant λ to all x_i is the same as replacing x by $x + \lambda u$. But then $\frac{x + \lambda u}{u \bullet u} = \bar{x} + \lambda$ and $\tilde{x}$ is replaced by

$$\widetilde{x + \lambda u} = [(x_1 + \lambda) - (\bar{x} + \lambda), \dots, (x_n + \lambda) - (\bar{x} + \lambda)]^T = \tilde{x}.$$

So r remains the same.

- b. Scaling each value by a positive λ is the same as replacing x by $\lambda x = [\lambda x_1, \dots, \lambda x_n]^\top$. We have $\overline{\lambda x} = \lambda \bar{x}$ and thus $\widetilde{\lambda x} = \lambda \tilde{x}$. Therefore r becomes

$$r = \frac{\lambda \tilde{x} \bullet \tilde{y}}{\|\lambda \tilde{x}\| \|\tilde{y}\|} = \frac{\lambda \tilde{x} \bullet \tilde{y}}{|\lambda| \|\tilde{x}\| \|\tilde{y}\|} = \frac{\tilde{x} \bullet \tilde{y}}{\|\tilde{x}\| \|\tilde{y}\|}.$$

So r remains the same (observe that it is crucial that λ is positive).

■

For this tutorial, the field $\mathbf{F}$ is either $\mathbf{R}$ or $\mathbf{C}$.

Exercise 1.

Consider the vector space $\mathcal{P}(\mathbf{R})_2$ with the inner product given by $\langle p, q \rangle = \int_0^1 p(x)q(x) dx$. Apply the Gram-Schmidt procedure to $(x, 1, x^2)$ to obtain an orthonormal basis for $\mathcal{P}(\mathbf{R})$. (Note that the order matters.)

Solution. Let $v_1 = x$, $v_2 = 1$ and $v_3 = x^2$. Then $u_1 = v_1 = x$. We compute

$$\langle u_1, u_1 \rangle = \int_0^1 x^2 dx = \frac{1}{3} \quad \text{and} \quad \langle v_2, u_1 \rangle = \int_0^1 x dx = \frac{1}{2}.$$

This gives us

$$u_2 = v_2 - \frac{\langle v_2, u_1 \rangle}{\langle u_1, u_1 \rangle} u_1 = 1 - \frac{\frac{1}{2}}{\frac{1}{3}} x = 1 - \frac{3}{2} x.$$

We now compute

$$\begin{aligned} \langle v_3, u_1 \rangle &= \int_0^1 x^3 dx = \frac{1}{4}, & \langle v_3, u_2 \rangle &= \int_0^1 x^2 - \frac{3}{2} x^3 dx = \frac{1}{3} - \frac{3}{2} \frac{1}{4} = -\frac{1}{24}, \\ \langle u_2, u_2 \rangle &= \int_0^1 1 - 3x + \frac{9}{4} x^2 dx = 1 - 3 \frac{1}{2} + \frac{9}{4} \frac{1}{3} = \frac{1}{4}. \end{aligned}$$

We then have

$$u_3 = v_3 - \frac{\langle v_3, u_1 \rangle}{\langle u_1, u_1 \rangle} u_1 - \frac{\langle v_3, u_2 \rangle}{\langle u_2, u_2 \rangle} u_2 = x^2 - \frac{\frac{1}{4}}{\frac{1}{3}} x - \frac{-\frac{1}{24}}{\frac{1}{4}} \left(1 - \frac{3}{2} x\right) = x^2 - x + \frac{1}{6}.$$

We compute a last inner product

$$\langle u_3, u_3 \rangle = \int_0^1 x^4 - 2x^3 + \frac{4}{3} x^2 - \frac{1}{3} x + \frac{1}{36} dx = \frac{1}{5} - 2 \frac{1}{4} + \frac{4}{3} \frac{1}{3} - \frac{1}{3} \frac{1}{2} + \frac{1}{36} = \frac{1}{180}.$$

Finally, we normalize:

$$e_1 = \frac{u_1}{\|u_1\|_1} = \sqrt{3}x, \quad e_2 = \frac{u_2}{\|u_2\|_2} = 2 - 3x, \quad e_3 = \frac{u_3}{\|u_3\|_3} = 6\sqrt{5} \left(x^2 - x + \frac{1}{6} \right).$$

■

Exercise 2.

The conjugate transpose of an $m \times n$ matrix A is the $n \times m$ matrix $A^* = \overline{A}^T = \overline{A}^T$.

- Show that $u \bullet v = v^* u$ for vectors u and v in $\mathbf{F}^n$.
- Prove that $(\text{col } A)^\perp = \text{null } A^*$ for the standard dot product (where $\text{col } A$ is

the column space of A).

Solution. a. Let $u = [u_1, \dots, u_n]^T$ and $v = [v_1, \dots, v_n]^T$ be any two vectors in $\mathbf{F}^n$. We have

$$u \bullet v = \sum_{j=1}^n u_j \bar{v}_j = [\bar{v}_1 \ \dots \ \bar{v}_n] \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix} = v^* u.$$

b. Recall that $\text{col}(A) = \{Av \mid v \in \mathbf{F}^n\}$. Now, let u be any vector in $\mathbf{F}^m$. We have

$$\begin{aligned} u \in \text{col}(A)^\perp &\iff \forall v \in \mathbf{F}^n : u \bullet Av = 0 \\ &\iff \forall v \in \mathbf{F}^n : 0 = (Av)^* u = v^* A^* u = A^* u \bullet v \\ &\iff A^* u = 0 \\ &\iff u \in \text{null}(A^*). \end{aligned}$$

■

Exercise 3.

Let V be an inner product space and suppose that $P \in \mathcal{L}(V)$ is a projection onto a subspace U . Prove that P is an orthogonal projection if and only if $\|Pv\| \leq \|v\|$ for all $v \in V$. (*Hint: Show that $\ker P$ must be orthogonal to U .*)

Solution. “ $\Rightarrow$ ” Suppose that P is an orthogonal projection, so $P = P_{U, U^\perp}$. Then for any v in V the vectors $Pv \in U$ and $v - Pv \in U^\perp$ are orthogonal. By the Pythagorean theorem, $\|v\|^2 = \|Pv\|^2 + \|v - Pv\|^2$. Since $\|v - Pv\|^2$ is non-negative, we conclude $\|Pv\|^2 \leq \|v\|^2$ and hence $\|Pv\| \leq \|v\|$.




Figure 3: An orthogonal projection of v onto U and the decomposition $v = Pv + (v - Pv)$ with $v - Pv \in U^\perp$.

“ $\Leftarrow$ ” We will give a proof assuming that V is finite dimensional. For the general case, see the “Infinite dimensional vector spaces” box below.

We claim:

If $\|Pv\| \leq \|v\|$ for all $v \in V$, then $\ker(P)^\perp$ is contained in $\text{Im}(P)$.

Before proving this claim, let us see how it implies that P is an orthogonal projection (i.e. that $\text{Im}(P)^\perp = \ker(P)$ if V is finite dimensional. Since V is finite dimensional, so is $\ker(P)$. Therefore $V = \text{Im}(P) \oplus \ker(P) = \ker(P) \oplus \ker(P)^\perp$. Everything being finite dimensional,

we conclude $\dim(V) = \dim(\operatorname{Im} P) + \dim(\ker P) = \dim(\ker P) + \dim(\ker(P)^\perp)$. So $\dim(\ker(P)^\perp) = \dim(\operatorname{Im} P)$, which by the claim implies $\ker(P)^\perp = \operatorname{Im}(P)$.

Let us now prove the claim. Let v be any vector in $\ker(P)^\perp$. We have $Pv = v + (Pv - v)$ with $v \in \ker(P)^\perp$. One easily check $P(Pv - v) = P^2v - Pv = 0$ so $Pv - v$ is in $\ker(P)$ and hence orthogonal to v . By the Pythagorean Theorem we have $\|Pv\|^2 = \|v\|^2 + \|Pv - v\|^2$. By assumption, we also have $\|Pv\|^2 \leq \|v\|^2$. Altogether, this implies that $\|Pv - v\|^2 = 0$ and hence $v = Pv$ is in the image of P as desired.

Infinite dimensional vector spaces

The above proof works only for V finite dimensional. The result is still true for V infinite dimensional, but requires a different proof.

Since P is a projection onto U , we have $U = \operatorname{Im}(P)$ and $V = \operatorname{Im}(P) \oplus \ker(P)$. We claim:

If $\|Pv\| \leq \|v\|$ for all $v \in V$, then $\ker(P)$ is contained in $\operatorname{Im}(P)^\perp$ (and hence P is the orthogonal projection onto $\operatorname{Im}(P)$).

Let us prove this claim. Let $u \in \operatorname{Im}(P)$ and $w \in \ker(P)$. If $w = 0$, then it is orthogonal to u . We can hence suppose that $w \neq 0$. We want to show that u and w are orthogonal. The trick is to consider the orthogonal projection of u onto $\operatorname{span}(w)$. More precisely, we will evaluate the norm of $v := u - \frac{\langle u, w \rangle}{\langle w, w \rangle} w$. Since u is in the image of P and w in the kernel, we have $P(v) = u$. So by assumption, $\|u\| \leq \|v\|$. We hence have

$$\begin{aligned} 0 &\leq \|v\|^2 - \|u\|^2 = \left\| u - \frac{\langle u, w \rangle}{\|w\|^2} w \right\|^2 - \|u\|^2 \\ &= \left\| \frac{\langle u, w \rangle}{\|w\|^2} w \right\|^2 - \left\langle u, \frac{\langle u, w \rangle}{\|w\|^2} w \right\rangle - \left\langle \frac{\langle u, w \rangle}{\|w\|^2} w, u \right\rangle \\ &= \frac{|\langle u, w \rangle|^2}{\|w\|^2} - \frac{\overline{\langle u, w \rangle}}{\|w\|^2} \langle u, w \rangle - \frac{\langle u, w \rangle}{\|w\|^2} \langle w, u \rangle \\ &= -\frac{|\langle u, w \rangle|^2}{\|w\|^2}. \end{aligned}$$

We conclude that $\frac{|\langle u, w \rangle|^2}{\|w\|^2} \leq 0$ and thus $\langle u, w \rangle = 0$ as desired.

■

Exercise 4.

Prove that a $m \times m$ matrix A represents an orthogonal projection (with respect to the standard dot product) if and only if $A^2 = A = A^*$. (Hint: For the only if direction, show that A^* represents a projection and has the same image and kernel

as A .

Solution. “ $\Leftarrow$ ” Since $A = A^2$, it represents a projection L_A and $\mathbf{F}^n = \text{Im}(L_A) \oplus \ker(L_A) = \text{col}(A) \oplus \text{null}(A)$. By Question 2 $(\text{col } A)^\perp = \text{null}(A^*) = \text{null}(A)$. So A represents an orthogonal projection since $\text{col}(A)$ and $\text{null}(A)$ are orthogonal.

“ $\Rightarrow$ ” Suppose that L_A is an orthogonal projection, so $A^2 = A$ and $\mathbf{F}^n = \text{col}(A) \oplus \text{null}(A)$ with $(\text{col } A)^\perp = \text{null}(A)$. We have

$$(A^*)^2 = (\overline{A^\top})^2 = \overline{(A^\top)^2} = \overline{(A^2)^\top} = \overline{A^\top} = A^*,$$

so A^* represents a projection and $\mathbf{F}^n = \text{col}(A^*) \oplus \text{null}(A^*)$. By Question 2 $(\text{col } A)^\perp = \text{null}(A^*)$, and applying this to A^* and $(A^*)^* = A$ we also have $(\text{col } A^*)^\perp = \text{null}(A)$, or in other words $\text{col } A^* = (\text{null } A)^\perp$. Since A is an orthogonal projection, we have $\text{null}(A^*) = (\text{col } A)^\perp = \text{null}(A)$ and $\text{col}(A^*) = (\text{null } A)^\perp = \text{col}(A)$. That is, A^* represents the orthogonal projection onto $\text{col}(A)$ with direction parallel to $\text{null}(A)$. By unicity of such a projection, we conclude $A^* = A$. ■

Further Questions

Exercise 5.

Let A and B be matrices satisfying $ABA = A$, $BAB = B$, $(AB)^* = AB$ and $(BA)^* = BA$. Prove that AB and BA are both orthogonal projection matrices. What are their respective images? (*Hint: see Tutorial 7 Question 7.*)

Solution. By Tutorial 7 Question 7, AB represents a projection onto $\text{col}(AB)$ along direction $\text{null}(AB)$, $\text{col}(AB) = \text{col}(A)$ and $\text{null}(B) = \text{null}(AB)$. Altogether, AB represents a projection onto $\text{col}(A)$ along direction $\text{null}(B)$. Finally, $(AB)^* = AB$ implies (Question 4) that the projection represented by AB is orthogonal. ■

Exercise 6.

Let $T \in \mathcal{L}(V, W)$ be a linear map between finite-dimensional inner product spaces.

- Show that for each $w \in \text{Im}(T)$ there exists a unique $v \in \ker(T)^\perp$ such that $Tv = w$.
- Prove that there is a unique linear map $S \in \mathcal{L}(W, V)$ satisfying $TST = T$, $STS = S$, and TS , ST are orthogonal projections to $\text{Im}(T)$, $\ker(T)^\perp$ respectively. (This S is called the **pseudo-inverse** of T .)

Solution. a. Since w is in the image of T , there exists $u \in V$ such that $Tu = w$. Since V is finite dimensional, we have $V = \ker(T) \oplus \ker(T)^\perp$. Therefore, there exists (unique) $z \in \ker(T)$ and $v \in \ker(T)^\perp$ such that $u = z + v$. We have

$w = Tu = Tz + Tv = 0 + Tv = Tv$. This finishes the proof of existence. Now, suppose that v' also satisfies $Tv' = w$ and $v' \in \ker(T)^\perp$. Then $T(v - v') = 0$ implies that $v - v'$ belongs to $\ker(T) \cap \ker(T)^\perp = \{0\}$. In other words, we necessarily have $v' = v$, showing unicity.

- b. By finite dimension, we have $W = \text{Im}(T) \oplus \text{Im}(T)^\perp$. Use this to define a function

$$S: W = \text{Im}(T) \oplus \text{Im}(T)^\perp \longrightarrow V$$

$$w + z \mapsto \text{the unique } v \in \ker(T)^\perp \text{ such that } Tv = w.$$

The existence and unicity of v makes this a well-defined function. We now need to show it is linear. So let $w_1 + z_1$ and $w_2 + z_2$ be two vectors in $W = \text{Im}(T) \oplus \text{Im}(T)^\perp$ and let λ be a scalar. Let v_1 and v_2 be the unique $v_i \in \ker(T)^\perp$ such that $Tv_i = w_i$. Then $\lambda v_1 + v_2$ is in $\ker(T)^\perp$ and $T(\lambda v_1 + v_2) = \lambda w_1 + w_2$. It follows that $S(\lambda(w_1 + z_1) + (w_2 + z_2)) = \lambda v_1 + v_2 = \lambda S(w_1 + z_1) + S(w_2 + z_2)$. This shows the linearity of S .

For any $w + z$ in $W = \text{Im}(T) \oplus \text{Im}(T)^\perp$ we have $STS(w + z) = ST(v) = S(w) = S(w + z)$. For any $v \in V$, one has $TSTv = TS w = Tv$. It follows (Tutorial 7 Question 7) that TS is a projection onto $\text{Im}(T)$ along direction $\ker(S)$, while ST is a projection onto $\text{Im}(S)$ along direction $\ker(T)$.

For orthogonality, we need to show that $\ker(S)^\perp = \text{Im}(T)$ and $\ker(T)^\perp = \text{Im}(S)$. Let z be a vector in $\text{Im}(T)^\perp$, therefore we have $z = 0 + z \in \text{Im}(T) \oplus \text{Im}(T)^\perp$. Then $S(z)$ is the unique w in $\ker T^\perp$ such that $T(w) = 0$ and therefore $w \in \ker T \cap \ker T^\perp$ and $S(z) = w = 0$. This proves $\text{Im}(T)^\perp \subseteq \ker(S)$.

Now, let $w + z$ be any element in $\ker(S)$. So $0 = S(w + z) = S(w)$ satisfies $T0 = w$. So $w + z = 0 + z$ is in $\text{Im}(T)^\perp$. This proves the second inclusion $\ker(S) \subseteq \text{Im}(T)^\perp$.

We have $\text{Im}(S) \subseteq \ker(T)^\perp$ by definition of S . Now, for v in $\ker(T)^\perp$ we have Tv in $\text{Im}(T)$ and hence $S(Tv) = v$, so v belongs to $\text{Im}(S)$. This proves the inclusion $\ker(T)^\perp \subseteq \text{Im}(S)$.

We proved that there exists an S with the desired properties. To finish the proof, we need to show that such an S is unique. Suppose that Q also satisfies the same properties. Then both TS and TQ are orthogonal projections to $\text{Im}(T)$ along direction $\text{Im}(T)^\perp$. By unicity of orthogonal projections, $TS = TQ$. One prove similarly $ST = QT$. Finally, $S = STS = STQ = QTQ = Q$. ■

Exercise 1.

Consider the matrix $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$.

- Find the eigenvalues of A in $\mathbb{C}$.
- For what values of θ does A have real eigenvalues?

Solution. a. First we compute the characteristic polynomial:

$$\begin{aligned} \chi_A(t) &= \det \begin{bmatrix} t - \cos \theta & \sin \theta \\ -\sin \theta & t - \cos \theta \end{bmatrix} \\ &= (t - \cos \theta)^2 + \sin^2 \theta = t^2 - 2t \cos \theta + \cos^2 \theta + \sin^2 \theta \\ &= t^2 - 2t \cos \theta + 1 = (t - \lambda_1)(t - \lambda_2). \end{aligned}$$

Using Vieta's formula, we conclude:

$$\lambda_{\pm} = \frac{-(-2 \cos \theta) \pm \sqrt{(2 \cos \theta)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \cos \theta \pm \sin \theta i.$$

- b. A has real eigenvalues if and only if $\sin \theta = 0$ if and only if $\theta = k\pi$ for some $k \in \mathbb{Z}$. ■

Exercise 2.

Let V be a vector space and let $P = P_{U,W} \in \mathcal{L}(V)$ be a projection to a subspace U along a complementary subspace W .

- What are the possible eigenvalues of P ?
- What are the eigenspaces of P ?

Solution. a. By assumption $V = \text{Im}(P) \oplus \ker(P)$. Then $Pv = \lambda v$ and $P^2v = P\lambda v = \lambda Pv = \lambda^2 v$. Since P is a projection, $P^2 = P$ and so $Pv = \lambda^2 v$. We conclude that $\lambda^2 v = \lambda v$ and since $v \neq 0$ that $\lambda^2 = \lambda$. Therefore $\lambda \in \{0, 1\}$.

- b. The possible eigenvalues of P are 0 and 1. We have $E(0, P) = \ker(P - 0 \text{Id}) = \ker(P)$ and $E(1, P) = \ker(P - 1 \text{Id}) = \text{Im}(P)$. ■

Exercise 3.

Suppose that $T \in \mathcal{L}(\mathbb{F}^4)$ is an operator such that $\dim E(3, T) = 3$. Prove that at least one of $T - 2 \text{Id}$ or $T - 4 \text{Id}$ is invertible.

Solution. Since $\mathbb{F}^4$ is finite dimensional, the operator $T - \lambda \text{Id}$ is not invertible if and only if λ is an eigenvalue of T , if and only if $\dim E(\lambda, T) \geq 1$. We also have that

Proposition 5.1.11

$4 = \dim \mathbf{F}^4 \geq \sum_{j=1}^k E(\lambda_j, T)$ where the sum is taken over all the distinct eigenvalues. We already know that 3 is an eigenvalue, let us say $\lambda_1 = 3$, so

$$4 = \dim \mathbf{F}^4 \geq 3 + \sum_{j=2}^k E(\lambda_j, T).$$

We conclude that T has at most one other eigenvalue $\lambda \neq 3$ and thus $T - \lambda \text{Id}$ is *not* invertible for at most one $\lambda \neq 3$. In particular, at least one of $T - 2\text{Id}$ or $T - 4\text{Id}$ is invertible. ■

Exercise 4.

Define an operator $T \in \mathcal{L}(\mathcal{P}(\mathbf{R})_3)$ by $Tp(x) = xp'(x)$ for all $x \in \mathbf{R}$. Find all eigenvalues and eigenvectors of T .

Solution. Trying to solve $\lambda p(x) = xp'(x)$ to find an eigenvector $p \neq 0$ is a priori difficult. So we will instead compute the characteristic polynomial of T . Let $\mathcal{B} = (1, x, x^2, x^3)$ be the standard basis of $\mathcal{P}(\mathbf{R})_3$. We have $T(1) = 0$, $T(x) = x$, $Tx^2 = 2x^2$ and $Tx^3 = 3x^3$. So

$$[T]_{\mathcal{B}}^{\mathcal{B}} = \begin{bmatrix} 0 & & & \\ & 1 & & \\ & & 2 & \\ & & & 3 \end{bmatrix}.$$

Since this is a diagonal matrix, we conclude that the eigenvalues of T are 0, 1, 2 and 3.

Eigenvalue $\lambda = 0$ We have $\text{null}([T]_{\mathcal{B}}^{\mathcal{B}} - 0 \text{Id}) = \text{null}([T]_{\mathcal{B}}^{\mathcal{B}}) = \{[a, 0, 0, 0]^T \mid a \in \mathbf{R}\}$. So the $E(0, T) = \ker(T - 0 \text{Id}) = \{a \mid a \in \mathbf{R}\}$ (the constant polynomials). The eigenvectors are $E(0, T) \setminus \{0\}$.

Eigenvalue $\lambda = 1$ We have

$$\text{null}([T]_{\mathcal{B}}^{\mathcal{B}} - 1 \text{Id}) = \text{null} \begin{bmatrix} -1 & & & \\ & 0 & & \\ & & 1 & \\ & & & 2 \end{bmatrix} = \left\{ \begin{bmatrix} 0 \\ b \\ 0 \\ 0 \end{bmatrix} \mid b \in \mathbf{R} \right\}.$$

So the $E(1, T) = \ker(T - 1 \text{Id}) = \{bx \mid b \in \mathbf{R}\}$. The eigenvectors are $E(1, T) \setminus \{0\}$.

Eigenvalues $\lambda = 2$ and $\lambda = 3$ The same method as above gives $E(2, T) = \ker(T - 2 \text{Id}) = \{cx^2 \mid c \in \mathbf{R}\}$ and $E(3, T) = \ker(T - 3 \text{Id}) = \{dx^3 \mid d \in \mathbf{R}\}$. ■

Exercise 5.

Let $V = \mathcal{C}^\infty(\mathbf{R})$ be the space of smooth real functions and let $D \in \mathcal{L}(V)$ be the differentiation operator. Show that every $\lambda \in \mathbf{R}$ is an eigenvalue for D , and find a corresponding eigenvector.

Solution. For every $\lambda \in \mathbf{R}$ we have $De^{\lambda x} = \lambda e^{\lambda x}$. So $e^{\lambda x}$ is an eigenvector with eigenvalue λ . ■

Further Questions

Exercise 6.

Let $\lambda_1, \dots, \lambda_k$ be distinct real numbers. Show that the list $(e^{\lambda_1 x}, \dots, e^{\lambda_k x})$ is linearly independent in $\mathcal{C}^\infty(\mathbf{R})$. *Hint: Consider the differentiation operator restricted to the subspace $U = \text{span}(e^{\lambda_1 x}, \dots, e^{\lambda_k x})$ of $\mathcal{C}^\infty(\mathbf{R})$.*

Solution. Since $D e^{\lambda_j x} = \lambda_j e^{\lambda_j x}$, the subspace U is D invariant. Therefore, $D|_U$ is an operator on U . The λ_j for $j \in \{1, \dots, k\}$ are all eigenvalues of $D|_U$. This implies that U has dimension at least k . Therefore, U has dimension exactly k and $(e^{\lambda_1 x}, \dots, e^{\lambda_k x})$ is a basis for U (and hence linearly independent). ■

Exercise 7.

Let V be a finite dimensional vector space and suppose that $T \in \mathcal{L}(V)$ has $\dim(V)$ distinct eigenvalues. Let $S \in \mathcal{L}(V)$ be an operator with the same eigenvectors as T , but not necessarily the same eigenvalues. Prove that $TS = ST$.

Solution. Let $m := \dim(V)$ and let $\lambda_1, \dots, \lambda_m$ be the eigenvalues of T and $v_1, \dots, v_m$ be the corresponding eigenvectors. Since the v_j correspond to distinct eigenvalues, they are linearly independent and therefore $\mathcal{B} = (v_1, \dots, v_m)$ is a basis of V .

By assumption, there exists $\mu_1, \dots, \mu_m$ such that $S v_j = \mu_j v_j$. Now, $T S v_j = T \mu_j v_j = \lambda_j \mu_j v_j$ while $S T v_j = S \lambda_j v_j = \mu_j \lambda_j v_j = \lambda_j \mu_j v_j = T S v_j$. Since ST and TS agree on a basis, they are equal. ■

Exercise 1.

Consider the operator map $T \in \mathcal{L}(\mathbf{C}^3)$ given by $T(x) = Ax$ where

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

Find the eigenvalues and the generalised eigenspaces of T .

Solution. We first compute the characteristic polynomial. Since A is a triangular matrix, $\chi_T(t) = (t-2)^2(t-3)$ and the eigenvalues are $\lambda_1 = 2$ (of algebraic multiplicity 2) and $\lambda_2 = 3$ (of algebraic multiplicity 1).

For $\lambda_1 = 2$

$$2\text{Id} - A = \begin{bmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad \text{and} \quad (2\text{Id} - A)^2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We conclude

$$G(2, T) = \text{null}((2\text{Id} - A)^2) = \left\{ \begin{bmatrix} z_1 \\ z_2 \\ 0 \end{bmatrix} \mid z_1, z_2 \in \mathbf{C} \right\}.$$

For $\lambda_2 = 3$

$$3\text{Id} - A = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

We conclude

$$G(3, T) = \text{null}(3\text{Id} - A) = \left\{ \begin{bmatrix} 0 \\ 0 \\ z_3 \end{bmatrix} \mid z_3 \in \mathbf{C} \right\}.$$

■

Exercise 2.

Let T be an operator on $\mathbf{C}^5$ whose eigenvalues are 3, 4, and 5. Prove that $(T - 3\text{Id})^3(T - 4\text{Id})^3(T - 5\text{Id})^3 = 0$.

Solution. Since T is a complex operator with eigenvalues 3, 4 and 5, its characteristic polynomial is $\chi_T(t) = (t-3)^{m_{\text{alg}}(3, T)}(t-4)^{m_{\text{alg}}(4, T)}(t-5)^{m_{\text{alg}}(5, T)}$ with all the $m_{\text{alg}}(\lambda_j, T) \geq 1$ and $5 = m_{\text{alg}}(3, T) + m_{\text{alg}}(4, T) + m_{\text{alg}}(5, T)$. We conclude that for $\lambda \in \{3, 4, 5\}$ we have $m_{\text{alg}}(\lambda, T) \leq 3$.

The minimal polynomial $M_T(t)$ divides $\chi_T(t)$ which itself divides $(t-3)^3(t-4)^3(t-5)^3$ by the above. Since $M_T(T) = 0$ we conclude $(T - 3\text{Id})^3(T - 4\text{Id})^3(T - 5\text{Id})^3 = 0$. ■

Exercise 3.

Let V be an m -dimensional complex vector space. Suppose that P is a projection onto an l -dimensional subspace of V . Show that $\chi_P(t) = (t-1)^l t^{m-l}$.

Solution. We know (see Tutorial 10, Question 2) that the only possible eigenvalues of P are 0 and 1 and that $E(1, P) = \text{Im}(P)$ while $E(0, P) = \ker(P)$. Moreover, we have $V = G(1, P) \oplus G(0, P) = \text{Im } P \oplus \ker P = E(1, P) \oplus E(0, P)$ (we use that P is a projection for the second equality). Therefore, we must have $E(1, P) = G(1, P)$ and $E(0, P) = G(0, P)$.

We conclude that $m_{\text{alg}}(1, P) = \dim G(1, P) = \dim E(1, P) = l$, while $m_{\text{alg}}(0, P) = \dim \ker(P) = \dim V - \dim \text{Im } P = m - l$. This gives $\chi_P(t) = (t-1)^l t^{m-l}$ as desired. ■

Exercise 4.

Let $T \in \mathcal{L}(V)$ be an operator on a finite-dimensional complex vector space V . Suppose that the characteristic and minimal polynomials of T are respectively $\chi_T(z) = (z-2)(z+1)^4(z-3)^3$ and $M_T(z) = (z-2)(z+1)^3(z-3)$. Write down a Jordan form of T .

Solution. From the characteristic polynomial, the eigenvalues of T are 2, -1 and 3.

The sum of the sizes of the Jordan blocks with eigenvalue 2 is $1 = m_{\text{alg}}(2)$. So for $\lambda = 2$ we only have one block of size 1.

For $\lambda = -1$, the sum of the sizes of the Jordan blocks is $m_{\text{alg}}(-1) = 4$, while the size of the biggest block is 3 (the exponent of $(z+1)$ in the minimal polynomial). This means that we have one block of size 3 and one of size 1.

Finally, for $\lambda = 3$, the sum of the sizes of the Jordan blocks is $m_{\text{alg}}(3) = 3$, while the size of the biggest block is 1 (the exponent of $(z-3)$ in the minimal polynomial). This means that we have 3 blocks of size 1.

Altogether, this gives us

$$\begin{bmatrix} \boxed{2} & & & & & & \\ & \boxed{-1} & & & & & \\ & & \boxed{1} & & & & \\ & & & \boxed{-1} & & & \\ & & & & \boxed{1} & & \\ & & & & & \boxed{-1} & \\ & & & & & & \boxed{-1} \\ & & & & & & & \boxed{3} \\ & & & & & & & & \boxed{3} \\ & & & & & & & & & \boxed{3} \end{bmatrix}$$

■

Exercise 5.

Consider the matrix

$$A = \begin{bmatrix} 6 & 6 & -15 \\ 1 & 5 & -5 \\ 1 & 2 & -2 \end{bmatrix}.$$

- Find the characteristic polynomial and minimal polynomial of A
- What is the Jordan form for A ?
- (Optional) Find a Jordan basis for A .

Solution. a. We start by computing the characteristic polynomial:

$$\begin{aligned} \chi_A(t) &= \det \begin{bmatrix} t-6 & -6 & 15 \\ -1 & t-5 & 5 \\ -1 & -2 & t+2 \end{bmatrix} \\ &= (t-6) \det \begin{bmatrix} t-5 & 5 \\ -2 & t+2 \end{bmatrix} + 1 \det \begin{bmatrix} -6 & 15 \\ -2 & t+2 \end{bmatrix} - 1 \det \begin{bmatrix} -6 & 15 \\ t-5 & 5 \end{bmatrix} \\ &= (t-6)((t-5)(t+2) + 10) + (-6(t+2) + 30) - (-30 - 15(t-5)) \\ &= t^3 - 9t^2 + 27t - 27 = (t-3)^3. \end{aligned}$$

So A has only one eigenvalue, 3, of algebraic multiplicity 3. The geometric multiplicity of 3 is therefore 1, 2 or 3. We test

$$A - 3\text{Id} = \begin{bmatrix} 3 & 6 & -15 \\ 1 & 2 & -5 \\ 1 & 2 & -5 \end{bmatrix} \neq 0,$$

and

$$(A - 3\text{Id})^2 = \begin{bmatrix} 3 & 6 & -15 \\ 1 & 2 & -5 \\ 1 & 2 & -5 \end{bmatrix}^2 = 0.$$

We conclude that $m_{\text{geo}}(3, A) = 2$ and $M_A(t) = (t-3)^2$.

- b. The Jordan form for A has one Jordan block of size $m_{\text{geo}}(3, A) = 2$, and therefore one more block of size 1. That is, the Jordan form is

$$J = \begin{bmatrix} 3 & 1 & \\ & 3 & \\ & & 3 \end{bmatrix}.$$

- c. To start, we need to find a vector v_1 in $\text{null}(A - 3\text{Id})^2 = \mathbf{F}^3$ but not in $\text{null}(A - 3\text{Id})$.

We can for example take $v_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$. Then we compute

$$v_2 := (A - 3\text{Id})v_1 = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix} \in \text{null}(A - 3\text{Id}).$$

Finally, we need to complete v_2 to a basis of $\text{null}(A - 3\text{Id})$. That is, we need to find a vector v_3 in $\text{null}(A - 3\text{Id}) = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + 2y - 5z = 0 \right\}$ but not in $\text{span } v_2$. For example, we can take $v_3 = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}$. A Jordan basis is then given by (v_1, v_2, v_3) .

Remark 0.2.

Let P be the matrix $P = [v_1 \ v_2 \ v_3]$. We can easily verify that

$$AP = [Av_1 \ Av_2 \ Av_3] = [3v_1 \ v_1 + 3v_2 \ 3v_3] = [v_1 \ v_2 \ v_3] \begin{bmatrix} 3 & 1 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix} = PJ.$$

That is, $P^{-1}AP = J$.

■

Further Questions

Exercise 6.

Let $N \in \mathcal{L}(V)$ be a nilpotent operator.

- Prove that N can only have 0 as an eigenvalue.
- Show that $\text{Id}_V - N$ is invertible.

Solution. a. Let $v \neq 0$ be an eigenvector: $Nv = \lambda v$. Since N is nilpotent, there exists k with $N^k = 0$ and so $0 = N^k v = \lambda^k v$. Since $v \neq 0$ we necessarily have $\lambda^k = 0$, which implies that $\lambda = 0$.

- Let k be an integer such that $N^k = 0$, then the inverse of $\text{Id} - N$ is $\text{Id} + N + \dots + N^{k-1}$. This can easily be directly verified. The intuition to find it comes from the power series $\sum_{i=0}^{\infty} x^i = \frac{1}{1-x}$ for $|x| < 1$.

■

Exercise 7.

Let V be a finite-dimensional complex vector space. Suppose that $T \in \mathcal{L}(V)$ is an operator with only 0 as an eigenvalue.

- Show that T is nilpotent.
- What is the characteristic polynomial of T ?

Solution. a. The minimal polynomial of T is $M_T(t) = t^k$ for some $k \in \{1, \dots, \dim(V)\}$. Then $0_V = M_T(T) = T^k$ and so T is nilpotent.

- b. Since V is a complex vector space, its characteristic polynomial is

$$\chi_T(t) = (t - \lambda_1) \cdots (t - \lambda_{\dim(V)}) = t^{\dim(V)}.$$

■

Exercise 1.

Let $V \neq \{0\}$ be a finite dimensional complex vector space, and let $T \in \mathcal{L}(V)$ be an operator. Prove that there does not exist a direct sum decomposition of V into two nonzero subspaces invariant under T if and only if $M_T(z) = (z - \lambda)^{\dim V}$ for some $\lambda \in \mathbb{C}$.

Solution. Firstly, remind that the exponent of $z - \lambda$ in $M_T(z)$ is the size of the biggest block of eigenvalue λ in a Jordan normal form of T . Therefore, $M_T(z) = (z - \lambda)^{\dim V}$ if and only if the Jordan normal form of T has only one Jordan block (of eigenvalue λ).

Suppose that $V = V_1 \oplus V_2$ with V_1 and V_2 nonzero and T -invariant. For $j \in \{1, 2\}$, let $T_j := T|_{V_j}$ and let $\mathcal{B}_j$ be a Jordan basis for T_j (so $\mathcal{B}_j$ is a basis of V_j). Then $\mathcal{B} := \mathcal{B}_1 \cup \mathcal{B}_2$ is a basis of V for which

$$[T]_{\mathcal{B}} = \begin{bmatrix} [T_1]_{\mathcal{B}_1} & 0 \\ 0 & [T_2]_{\mathcal{B}_2} \end{bmatrix}.$$

So $[T]_{\mathcal{B}}$ is a Jordan normal form for T , with at least two blocks.

For the other direction, suppose that the Jordan normal form of T has at least two blocks and let $\mathcal{B}$ be a corresponding Jordan basis. Let $\mathcal{B}_1$ be the first vectors of $\mathcal{B}$ corresponding to the first block and let $\mathcal{B}_2 := \mathcal{B} \setminus \mathcal{B}_1$. So we have

$$[T]_{\mathcal{B}} = \begin{bmatrix} \boxed{J_1} & 0 \\ 0 & \boxed{J_2} \end{bmatrix}.$$

$\underbrace{\hspace{1.5cm}}_{\mathcal{B}_1} \quad \underbrace{\hspace{1.5cm}}_{\mathcal{B}_2}$

where J_1 is a Jordan block and J_2 is composed of one or more Jordan blocks. Let $V_i := \text{span}(\mathcal{B}_i)$. By construction, $V = V_1 \oplus V_2$. Moreover, the dimension of $V_i \neq \{0\}$ is equal to the size of J_i and hence at least 1. Finally, the fact that $[T]_{\mathcal{B}}$ is diagonal by blocks, with diagonal blocks J_1 and J_2 is equivalent to the fact that V_1 and V_2 are T invariant. ■