

Benchmarks for QDyn

All equations and calculations are in atomic units.

```
In[1]:= ħ = 1;  
me = 1;
```

Harmonic oscillator 1D

Harmonic oscillator with potential in form:

$$V(x) = \frac{1}{2} m \omega^2 x^2$$

Exact energies correspond to the following formula:

$$E_n = \hbar \omega \left(n + \frac{1}{2} \right), \quad n = 0, 1, 2, \dots$$

```
In[3]:= ω = 0.1;  
m = 1;  
V = 1 / 2 * m * ω ^ 2 * x ^ 2
```

```
Out[5]= 0.005 x2
```

Energies for benchmark

```
In[6]:= Table[{n, ħ * ω * (1 / 2 + n)}, {n, 0, 9}] // TableForm
```

```
Out[6]//TableForm=
```

0	0.05
1	0.15
2	0.25
3	0.35
4	0.45
5	0.55
6	0.65
7	0.75
8	0.85
9	0.95

```
In[7]:= Clear[ω, m, V]
```

Morse potential 1D

Harmonic oscillator with potential in form:

$$V(x) = D_e \left(1 - e^{-a(r-r_e)}\right)^2 - D_e$$

Exact energies correspond to the following formula:

$$E_n = h \nu_0 \left(n + \frac{1}{2}\right) - \frac{\left[h \nu_0 \left(n + \frac{1}{2}\right)\right]^2}{4 D_e}, \quad n = 0, 1, 2, \dots, \lambda - \frac{1}{2}$$


[Solution according to https://en.wikipedia.org/wiki/Morse_potential]

This Morse potential should correspond to one of the states of the N2 molecule.

```
In[8]:= h = ħ * 2 * Pi;
De = 0.364;
re = 2.041;
a = 1.51;
m = 1834 * 28;
ν0 = a / 2 / Pi * Sqrt[2 * De / m];
V[r_] = De * (1 - Exp[-a * (r - re)])^2 - De
```

```
Out[14]= -0.364 + 0.364 (1 - e-1.51 (-2.041+r))2
```

```
In[15]:= Plot[V[x], {x, 1, 4}]
```



Energies for benchmark

```
In[16]:= Table[{n, h * v0 * (1 / 2 + n) - (h * v0 * (1 / 2 + n)) ^ 2 / (4 * De) - De}, {n, 0, 9}] //
TableForm

Out[16]//TableForm=
  0    -0.361163
  1    -0.355522
  2    -0.349925
  3    -0.344373
  4    -0.338865
  5    -0.333402
  6    -0.327983
  7    -0.322608
  8    -0.317278
  9    -0.311992

In[17]:= Clear[v0, m, V, a, re, De, h]
```

Harmonic oscillator 2D - symmetric

Harmonic oscillator with potential in form:

$$V(x, y) = \frac{1}{2} m \omega_x^2 x^2 + \frac{1}{2} m \omega_y^2 y^2$$

Exact energies correspond to the following formula:

$$E_{n_x, n_y} = \hbar \omega (n_x + n_y + 1), \quad n_x, n_y = 0, 1, 2, \dots$$

```
In[18]:= wx = 0.1; wy = 0.1;
m = 1;
V = 1 / 2 * m * wx ^ 2 * x ^ 2 + 1 / 2 * m * wy ^ 2 * y ^ 2

Out[20]=
0.005 x^2 + 0.005 y^2
```

Energies for benchmark

```
In[21]:= data = Table[{nx, ny, h * (wx * (1 / 2 + nx) + wy * (1 / 2 + ny))}, {nx, 0, 4}, {ny, 0, 4}]

Out[21]=
{{{0, 0, 0.1}, {0, 1, 0.2}, {0, 2, 0.3}, {0, 3, 0.4}, {0, 4, 0.5}},
 {{1, 0, 0.2}, {1, 1, 0.3}, {1, 2, 0.4}, {1, 3, 0.5}, {1, 4, 0.6}},
 {{2, 0, 0.3}, {2, 1, 0.4}, {2, 2, 0.5}, {2, 3, 0.6}, {2, 4, 0.7}},
 {{3, 0, 0.4}, {3, 1, 0.5}, {3, 2, 0.6}, {3, 3, 0.7}, {3, 4, 0.8}},
 {{4, 0, 0.5}, {4, 1, 0.6}, {4, 2, 0.7}, {4, 3, 0.8}, {4, 4, 0.9}}}
```

```
In[22]:= Sort[ArrayReshape[data[[;;,;;,3],5*5]][[;;10]] // TableForm
Out[22]//TableForm=
0.1
0.2
0.2
0.3
0.3
0.3
0.4
0.4
0.4
0.4

In[23]:= Clear[m, ωx, ωy, V, data]
```

Harmonic oscillator 2D - asymmetric

Harmonic oscillator with potential in form:

$$V(x, y) = \frac{1}{2} m \omega_x^2 x^2 + \frac{1}{2} m \omega_y^2 y^2$$

Exact energies correspond to the following formula:

$$E_{n_x, n_y} = \hbar \left[\omega_x \left(n_x + \frac{1}{2} \right) + \omega_y \left(n_y + \frac{1}{2} \right) \right], \quad n_x, n_y = 0, 1, 2, \dots$$

```
In[24]:= ωx = 0.1; ωy = 0.15;
m = 1;
V = 1 / 2 * m * ωx^2 * x^2 + 1 / 2 * m * ωy^2 * y^2
Out[26]=
0.005 x^2 + 0.01125 y^2
```

Energies for benchmark

```
In[27]:= data = Table[{nx, ny, ħ * (ωx * (1 / 2 + nx) + ωy * (1 / 2 + ny))}, {nx, 0, 4}, {ny, 0, 4}]
Out[27]=
{{{0, 0, 0.125}, {0, 1, 0.275}, {0, 2, 0.425}, {0, 3, 0.575}, {0, 4, 0.725}},
 {{1, 0, 0.225}, {1, 1, 0.375}, {1, 2, 0.525}, {1, 3, 0.675}, {1, 4, 0.825}},
 {{2, 0, 0.325}, {2, 1, 0.475}, {2, 2, 0.625}, {2, 3, 0.775}, {2, 4, 0.925}},
 {{3, 0, 0.425}, {3, 1, 0.575}, {3, 2, 0.725}, {3, 3, 0.875}, {3, 4, 1.025}},
 {{4, 0, 0.525}, {4, 1, 0.675}, {4, 2, 0.825}, {4, 3, 0.975}, {4, 4, 1.125}}}
```

```
In[28]:= Sort[ArrayReshape[data[[;;,;;,3]],5*5]][[;;10]] // TableForm
Out[28]//TableForm=
0.125
0.225
0.275
0.325
0.375
0.425
0.425
0.475
0.525
0.525
```

```
In[29]:= Clear[m, ωx, ωy, V, data]
```

Harmonic oscillator 3D - symmetric

Harmonic oscillator with potential in form:

$$V(x, y, z) = \frac{1}{2} m \omega_x^2 x^2 + \frac{1}{2} m \omega_y^2 y^2 + \frac{1}{2} m \omega_z^2 z^2$$

Exact energies correspond to the following formula:

$$E_{n_x, n_y, n_z} = \hbar \omega \left(n_x + n_y + n_z + \frac{3}{2} \right), \quad n_x, n_y, n_z = 0, 1, 2, \dots$$

```
In[30]:= ωx = 0.1; ωy = 0.1; ωz = 0.1;
m = 1;
V = 1/2 * m * ωx^2 * x^2 + 1/2 * m * ωy^2 * y^2 + 1/2 * m * ωz^2 * z^2
Out[32]=
0.005 x^2 + 0.005 y^2 + 0.005 z^2
```

Energies for benchmark

```
In[33]:= data = Table[{nx, ny, nz, ħ * (ωx * (1/2 + nx) + ωy * (1/2 + ny) + ωz * (1/2 + nz))},
{nx, 0, 4}, {ny, 0, 4}, {nz, 0, 4}];

In[34]:= Sort[ArrayReshape[data[[;;,;;,4]],5*5*5]][[;;10]] // TableForm
Out[34]//TableForm=
0.15
0.25
0.25
0.25
0.35
0.35
0.35
0.35
0.35
0.35
```

```
In[35]:= Clear[m, ωx, ωy, ωz, V, data]
```

Harmonic oscillator 3D - asymmetric

Harmonic oscillator with potential in form:

$$V(x, y, z) = \frac{1}{2} m \omega_x^2 x^2 + \frac{1}{2} m \omega_y^2 y^2 + \frac{1}{2} m \omega_z^2 z^2$$

Exact energies correspond to the following formula:

$$E_{n_x, n_y, n_z} = \hbar \left[\omega_x \left(n_x + \frac{1}{2} \right) + \omega_y \left(n_y + \frac{1}{2} \right) + \omega_z \left(n_z + \frac{1}{2} \right) \right], \quad n_x, n_y, n_z = 0, 1, 2, \dots$$

```
In[36]:= ωx = 0.1; ωy = 0.07; ωz = 0.13;
```

```
m = 1;
```

```
V = 1 / 2 * m * ωx^2 * x^2 + 1 / 2 * m * ωy^2 * y^2 + 1 / 2 * m * ωz^2 * z^2
```

```
Out[38]=
```

```
0.005 x^2 + 0.00245 y^2 + 0.00845 z^2
```

Energies for benchmark

```
In[39]:= data = Table[{nx, ny, nz, ħ * (ωx * (1 / 2 + nx) + ωy * (1 / 2 + ny) + ωz * (1 / 2 + nz))},
  {nx, 0, 4}, {ny, 0, 4}, {nz, 0, 4}];
```

```
In[40]:= Sort[ArrayReshape[data[[;;, ;;, ;;, 4]], 5 * 5 * 5]][[;;, 10]] // TableForm
```

```
Out[40]//TableForm=
```

```
0.15
```

```
0.22
```

```
0.25
```

```
0.28
```

```
0.29
```

```
0.32
```

```
0.35
```

```
0.35
```

```
0.36
```

```
0.38
```

```
In[41]:= Clear[m, ωx, ωy, ωz, V, data]
```