

Abstract

These notes, which are unfinished, are based on Scott Weinstein’s “Topics in Logic: Set Theory” lectures at UPenn along with Thomas Jech’s *Set Theory - The Third Millennium Edition, revised and expanded*. Any mistake in what follows is my own.

Contents

1 Point-set topology of \mathbb{R}	3
1.1 Lecture 1	3
1.2 Lecture 2	4
1.3 Lecture 3	5
1.4 Lecture 4	7
1.5 Lecture 5	9
2 Zermelo-Fraenkel set theory	10
2.1 Lecture 6	10
2.2 Lecture 7	13
3 Ordinal numbers	14
3.1 Lecture 8	16
3.2 Lecture 9	18
3.3 Lecture 10	20
4 The axiom of choice	21
4.1 Lecture 11	23
4.2 Lecture 12	25
5 Relativization	27
5.1 Lecture 13	27
5.2 Lecture 14	30
5.3 Lecture 15	31
5.4 Lecture 16	33
5.5 Lecture 17	34
6 Gödel’s hierarchy of constructible sets	36
6.1 Lecture 18	36
6.2 Lecture 19	38
6.3 Lecture 20	38
6.4 Lecture 21	38
6.5 Lecture 22	38

6.6	Lecture 23	38
6.7	Lecture 24	38
6.8	Lecture 25	38
6.9	Lecture 26	38
6.10	Lecture 27	38
6.11	Lecture 28	38
6.12	Lecture 29	38
6.13	Lecture 30	38
6.14	Lecture 31	38
6.15	Lecture 32	38
7	Large cardinals	38
7.1	Lecture 33	38
7.2	Lecture 34	38
7.3	Lecture 35	38
7.4	Lecture 36	38
7.5	Lecture 37	38
7.6	Lecture 38	38
7.7	Lecture 39	38
7.8	Lecture 40	38
7.9	Lecture 41	39
7.10	Lecture 42	39
7.11	Lecture 43	39

1 Point-set topology of \mathbb{R}

1.1 Lecture 1

Throughout this section, we shall work tacitly in ZFC.

Notation. $\mathbb{N} := \{0, 1, 2, \dots\}$.

Definition 1.1.1. We say that two sets X and Y are *equipollent* if there exists a bijection from X onto Y .

Notation. If two sets X and Y are equipollent, then we shall write either $X \sim Y$ or $|X| = |Y|$.

Note 1.1.2. Recall that the *linear continuum* $(\mathbb{R}, <)$ is the unique ordered field in which every nonempty bounded (above) set has a supremum.

Theorem 1.1.3 (Cantor). *The set \mathbb{R} is uncountable, i.e., \mathbb{R} is not equipollent to \mathbb{N} or to any finite set.*

Proof. It's obvious that \mathbb{R} is not finite. Suppose, towards a contradiction, that \mathbb{R} is countable. Enumerate \mathbb{R} as

$$\mathbb{R} = \{x_0, x_1, \dots, x_n, \dots\}.$$

Define the sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ recursively as follows. Let $a_0 = x_0$ and $b_0 = x_{k_0}$ where k_0 denotes the least k such that $a_0 < x_k$. For each $n \in \mathbb{N}$, let $a_{n+1} = x_{i_n}$ where i_n denotes the least i such that $a_n < x_i < b_n$ and let $b_{n+1} = x_{k_n}$ where k_n denotes the least k such that $a_{n+1} < x_k < b_n$. In terms of our enumeration of \mathbb{R} , we have that

$$\begin{aligned} \mathbb{R} &= \{a_0, \dots, b_0, \dots, a_1, \dots, b_1, \dots, a_2, \dots, b_2, \dots, a_3, \dots, b_4, \dots\} \\ a_0 &< a_1 < a_2 < \dots < a_k < \dots < b_k < \dots < b_2 < b_1 < b_0. \end{aligned}$$

Note that $A := \{a_n \mid n \in \mathbb{N}\}$ is nonempty and bounded above by b_0 . Hence $\sup(A)$ exists in \mathbb{R} , and it belongs to $\bigcap_{n \in \mathbb{N}} (a_n, b_n)$. This implies that for each $n \in \mathbb{N}$, $\sup(A)$ does not equal any x_k that precedes a_n . But every x_k other than x_0 precedes some a_{n_k} . It follows that $\sup(A) \neq x_n$ for any n . Thus, $\mathbb{R} \neq \{x_0, x_1, \dots, x_n, \dots\}$, a contradiction. \square

Definition 1.1.4.

1. Let $a, b \in \mathbb{R}$ with $a < b$. We call the set

$$\begin{aligned} (a, b) &:= \{x \in \mathbb{R} \mid a < x < b\} \\ (\text{resp. } [a, b] &:= \{x \in \mathbb{R} \mid a \leq x \leq b\}) \end{aligned}$$

an *open* (resp. *closed*) *interval* (with endpoints a and b).

2. A set $X \subset \mathbb{R}$ is *open in \mathbb{R}* if it equals a union of open intervals.
3. A set $X \subset \mathbb{R}$ is *closed in \mathbb{R}* if there is some open set Y in \mathbb{R} such that $X = \mathbb{R} \setminus Y$.
4. The topology on \mathbb{R} generated by the set of open intervals is called the *order topology on \mathbb{R}* .

Note 1.1.5. A set X is open in \mathbb{R} if and only if for any $x \in X$, there is some open interval I such that $x \in I$ and $I \subset X$.

Proposition 1.1.6. *The set $\{(a, b) \mid a, b \in \mathbb{Q}\}$ of rational intervals forms a countable basis of \mathbb{R} .*

Definition 1.1.7. Let $X \subset \mathbb{R}$ and $c \in \mathbb{R}$.

1. We say that c is a *limit point* of X if for any open interval I with $I \ni c$, we have $(I \setminus \{c\}) \cap X \neq \emptyset$.
2. We say that c is an *isolated point* of X if $c \in X$ and c is not a limit point of X .

Example 1.1.8.

1. Let $X = \{0\} \cup [1, \frac{3}{2}) \cup (\frac{3}{2}, 2] \cup \{3\}$. Then the isolated points of X are precisely 0 and 3.
2. Let $X = \{0\} \cup \{\frac{1}{n} \mid n \in \mathbb{Z}_{>0}\}$. Then the isolated points of X are precisely $\frac{1}{n}$ for each $n \in \mathbb{Z}_{>0}$.

1.2 Lecture 2

Definition 1.2.1. Let $X \subset \mathbb{R}$.

1. The (*Cantor-Bendixson*) *derivative* X' of X is the set of limit points of X .
2. We say that X is *perfect* if $X \neq \emptyset$ and $X = X'$.

Lemma 1.2.2.

1. X is closed in \mathbb{R} if and only if $X' \subset X$.
2. If \mathcal{C} is any set of closed sets in \mathbb{R} , then $\bigcap \mathcal{C}$ is also closed in \mathbb{R} .

Proof.

1. Suppose that X is closed, so that $X = \mathbb{R} \setminus Y$ for some open Y . Let x be a limit point of X . If $x \in Y$, then there is some open interval I around x such that $I \subset Y$. In this case, $I \cap X = \emptyset$, so that x is not a limit point of X , a contradiction. Hence $x \notin Y$, i.e., $x \in X$. This proves that $X' \subset X$.

Conversely, suppose that $X' \subset X$. Let $x \notin X$. Then there is some open interval I around x such that $I \cap X = \emptyset$, i.e., $I \subset \mathbb{R} \setminus X$. This proves that $\mathbb{R} \setminus X$ is open, so that X is closed.

2. For each $C \in \mathcal{C}$, there is some open set Y_C such that $C = \mathbb{R} \setminus Y_C$. Then

$$\bigcap \mathcal{C} = \bigcap_{C \in \mathcal{C}} \mathbb{R} \setminus Y_C = \mathbb{R} \setminus \bigcup_{C \in \mathcal{C}} Y_C.$$

Since $\bigcup_{C \in \mathcal{C}} Y_C$ is open, it follows that $\bigcap \mathcal{C}$ is closed, as desired.

□

Corollary 1.2.3. *X is perfect if and only if it is nonempty, is closed, and has no isolated points.*

Definition 1.2.4. Let $X \subset \mathbb{R}$. Suppose that for any open interval $I \subset \mathbb{R}$, we have $I \cap X \neq \emptyset$. In this case, we say that X is *dense* in \mathbb{R} .

Example 1.2.5. \mathbb{Q} is dense in \mathbb{R} .

Note 1.2.6. Any set of pairwise disjoint open intervals must be countable because \mathbb{Q} is both countable and dense in \mathbb{R} .

Definition 1.2.7. Let $(P, <)$ be a (strict) linear ordering (also known as a (strict) total order).

1. We say that $(P, <)$ is *dense (in itself)* if for any $a, b \in P$ with $a < b$, there exists $c \in P$ such that $a < c < b$.
2. We say that $(P, <)$ is *complete* if every nonempty bounded subset of P has a supremum.
3. We say that $(P, <)$ is *unbounded* if it contains neither an upper bound nor a lower bound.

Definition 1.2.8. Let $(P, <)$ and $(Q, <')$ be any two linear orderings.

1. We say that $f : P \rightarrow Q$ is *order-preserving* if $x < y \implies f(x) <' f(y)$.
2. We say that P and Q are (*order*) *isomorphic*, written as $P \cong Q$, if there is some order-preserving bijection from P onto Q .

Proposition 1.2.9. *The following properties are invariant under isomorphism.*

- *Being dense in a set.*
- *Being complete.*
- *Being unbounded.*

Theorem 1.2.10. *Let $(C, <)$ be a complete linear ordering containing some countable dense subset $(P, <)$ such that $P \cong \mathbb{Q}$. Then $C \cong \mathbb{R}$.*

Proof. By hypothesis, there is some isomorphism $f : P \rightarrow \mathbb{Q}$. Extend f to $\tilde{f} : C \rightarrow \mathbb{R}$ as follows. For any $c \in C \setminus P$, let

$$\tilde{f}(c) = \sup \{f(p) \mid p \in P \wedge p < c\}.$$

It is easy to check that \tilde{f} is order-preserving. It remains to show that it is surjective. If $r \in \mathbb{R} \setminus \mathbb{Q}$, then there is some strictly increasing sequence of rationals $(q_n)_{n \in \mathbb{N}}$ that converges to r . In particular, $\sup\{q_n \mid n \in \mathbb{N}\} = r$. Then \tilde{f} sends $\sup\{f^{-1}(q_n) \mid n \in \mathbb{N}\}$ to r . \square

1.3 Lecture 3

Definition 1.3.1. Let $(P, <)$ be any linear ordering. A *Dedekind cut in P* is a pair (A, B) of disjoint nonempty subsets of P such that

- (i) $A \cup B = P$,
- (ii) $a < b$ for any $a \in A$ and $b \in B$, and
- (iii) A does not contain an upper bound.

Theorem 1.3.2. *Let $(P, <)$ be a dense unbounded linearly ordered set. Then there exist a linear ordering $(P', <')$ isomorphic to $(P, <)$ and a complete unbounded linearly ordered set (C, \prec) such that*

- $P' \subset C$,
- $<' equals \prec restricted to P' , and$

- P' is dense in C .

Proof. Set C equal to the set of all Dedekind cuts in P . Let $(A_1, B_1) \preceq (A_2, B_2)$ if $A_1 \subset A_2$. It's easy to check that (C, \prec) is a linear ordering. Also, since P is unbounded, so is C . If $T := \{(A_s, B_s) \mid s \in S\}$ is any nonempty bounded subset of C , then

$$\left(\bigcup_{s \in S} A_s, \bigcap_{s \in S} B_s \right) = \sup(T).$$

Thus, C is complete.

Now, for each $p \in P$, let

$$A_p = \{x \in P \mid x < p\}$$

$$B_p = \{x \in P \mid x \geq p\}.$$

Note that each (A_p, B_p) belongs to C and that

$$P' := \{(A_p, B_p) \mid p \in P\} \cong P.$$

Suppose that $(A_1, B_1) \prec (A_2, B_2)$, so that $A_1 \subsetneq A_2$. Then there is some $x \in A_2 \setminus A_1$. Pick any $x' \in A_2$ such that $x < x'$. Note that $x \in A_{x'} \setminus A_1$ and $x' \in A_2 \setminus A_{x'}$. Therefore, $A_1 \subsetneq A_{x'} \subset A_2$. This implies $(A_1, B_1) \prec (A_{x'}, B_{x'}) \prec (A_2, B_2)$. It follows that P' is dense in C . \square

Example 1.3.3. Given the ordered field $(\mathbb{Q}, <)$, we can construct \mathbb{R} as the set of all Dedekind cuts in \mathbb{Q} .

Lemma 1.3.4. *Every nonempty closed interval I is perfect.*

Proof. It suffices to show that I has no isolated points. But this follows from the fact that \mathbb{R} is dense in itself. \square

Notation. If I is an interval, then $|I|$ will denote the length of I .

Theorem 1.3.5 (Cantor's intersection theorem). *Let*

$$I_0 \supset I_1 \supset I_2 \supset \cdots$$

be a descending chain of nonempty closed intervals.

$$1. \bigcap_{n \in \mathbb{N}} I_n \neq \emptyset.$$

$$2. \text{ If } \lim_{n \rightarrow \infty} |I_n| = 0, \text{ then } \bigcap_{n \in \mathbb{N}} I_n \text{ consists of exactly one element.}$$

Example 1.3.6. The *Cantor set* \mathcal{C} is the set of all real numbers of the form

$$\sum_{n=1}^{\infty} \frac{a_n}{3^n}, \quad a_n \in \{0, 2\}.$$

That is, \mathcal{C} consists of those real numbers whose ternary expansions exclude the digit 1. Alternatively, we can obtain \mathcal{C} by the following procedure.

1. Take the interval

$$C_0 := [0, 1].$$

2. Remove $(\frac{1}{3}, \frac{2}{3})$ from I_0 to get

$$C_1 := \left[0, \frac{1}{3}\right] \cup \left[\frac{2}{3}, 1\right].$$

3. Remove $(\frac{1}{9}, \frac{2}{9})$ and $(\frac{7}{9}, \frac{8}{9})$ from I_1 to get

$$C_2 := \left[0, \frac{1}{9}\right] \cup \left[\frac{2}{9}, \frac{1}{3}\right] \cup \left[\frac{2}{3}, \frac{7}{9}\right] \cup \left[\frac{8}{9}, 1\right].$$

4. Continue in this fashion to get a sequence of sets $(C_n)_{n \in \mathbb{N}}$. Then

$$\mathcal{C} = \bigcap_{n \in \mathbb{N}} C_n.$$

Note that \mathcal{C} is closed as the intersection of closed sets. Moreover, since we have a nested sequence

$$C_0 \supset C_1 \supset C_2 \supset C_3 \supset \cdots,$$

Cantor's intersection theorem implies that \mathcal{C} is nonempty.

Claim. \mathcal{C} has no isolated points.

Proof. Let $\epsilon > 0$. Choose $n \in \mathbb{N}$ so large that $(\frac{1}{3})^n < \epsilon$. Let $x \in \mathcal{C}$. Then $x \in C_n$, so that x belongs to some closed interval of length $\frac{1}{3}^n$. Pick $y \neq x$ such that y is an endpoint of this interval. Then $y \in \mathcal{C}$, and $|y - x| < \epsilon$. This shows that any open interval around x contains a point in \mathcal{C} other than x . Thus, x is not an isolated point. \square

We now see that \mathcal{C} is, in fact, perfect. Moreover, since there is a bijection from $\{0, 2\}^{\mathbb{N}}$ onto \mathcal{C} , we see that $\mathcal{C} \sim 2^{\aleph_0}$.

1.4 Lecture 4

Note 1.4.1. Let \mathcal{O} denote the set of all open sets. Recall that \mathbb{R} has the set of rational intervals as a countable basis. Since $\underbrace{\mathbb{P}(\mathbb{Q} \times \mathbb{Q})}_{\text{power set}} \sim \mathbb{R}$, it follows by the Cantor-Schröder-Bernstein theorem that $\mathcal{O} \sim \mathbb{R}$.

Lemma 1.4.2. *If C is a closed interval, P is perfect, and $|C \cap P| > 2$, then for every $n \in \mathbb{N}$, there are closed intervals $C_1, C_2 \subset C$ each of length $\leq n^{-1}$ such that $|C_1 \cap P| > 2$, $|C_2 \cap P| > 2$, and $C_1 \cap C_2 = \emptyset$.*

Proof. By hypothesis, we can find a set $\{x, y, z\} \subset C \cap P$ of size three. There are two cases to consider.

First, suppose that at least two of x, y , and z belong to the interior $\text{Int}(C)$ of C , say, x and y . Choose $C_1 \subset C$ and $C_2 \subset C$ such that $\text{Int}(C_1) \ni x$ and $\text{Int}(C_2) \ni y$. We can also make them so small that $C_1 \cap C_2 = \emptyset$ and both are of length at most n^{-1} . Note that $\text{Int}(C_1)$ and $\text{Int}(C_2)$ contain at least two points other than x and y , respectively, because P has no isolated points.

Second, suppose that exactly one of x , y , and z belongs to $\text{Int}(C)$, say, z . Then there is some $w \neq z$ such that $w \in \text{Int}(C)$ because P has no isolated points. Assume, wlog, that $x < y$ and $w < z$. Choose $C_1 \subset C$ and $C_2 \subset C$ of length at most n^{-1} such that $C_1 \cap C_2 = \emptyset$, $x \in C_1$, $w \in \text{Int}(C_1)$, $y \in C_2$, and $z \in \text{Int}(C_2)$. Then, as before, $\text{Int}(C_1)$ and $\text{Int}(C_2)$ contain other points than w and z , respectively. \square

Theorem 1.4.3. *If P is a perfect set, then $P \sim \mathbb{R}$.*

Proof. Due to the Cantor-Schröder-Bernstein theorem, it suffices to exhibit an injection $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow P$. We can view $\{0, 1\}^{\mathbb{N}}$ as the set of all infinite binary strings. Let $\{0, 1\}^*$ denote the set of all (finite) binary strings including the empty string ϵ .

By induction on the length of string, we shall define a set of closed intervals $(I_w)_{w \in \{0, 1\}^*}$ such that

- $|I_w \cap P| > 2$,
- $I_{w\sigma} \subset I_w$,
- $I_{w0} \cap I_{w1} = \emptyset$, and
- $|I_s| < \frac{1}{|w|+1}$.

where $\sigma \in \{0, 1\}$. For the base case, there exists a closed interval I_ϵ such that $|I_\epsilon| < 1$ and $|I_\epsilon \cap P| > 2$ since P has no isolated points. In addition, by Lemma 1.4.2, we can find disjoint subintervals I_1 and I_0 each on length $< \frac{1}{2}$ such that $|I_1 \cap P| > 2$ and $|I_0 \cap P| > 2$. Moreover, the same lemma automatically completes our induction step.

Now, for each $f \in \{0, 1\}^{\mathbb{N}}$ and $n \in \mathbb{N}$, let $\bar{f}(n) = f(0)f(1)\cdots f(n)$. Note that

$$I_{\bar{f}(0)} \supset I_{\bar{f}(1)} \supset I_{\bar{f}(2)} \supset \cdots$$

is a descending chain of nonempty closed intervals with $\lim_{n \rightarrow \infty} |I_{\bar{f}(n)}| = 0$. It follows by Cantor's intersection theorem that

$$\bigcap_{n \in \mathbb{N}} I_{\bar{f}(n)} = \{c_f\}$$

for some $c_f \in \mathbb{R}$.

Claim. $c_f \in P$.

Proof. For each $I_{\bar{f}(k)}$, there is some $x_k \in P \cap I_{\bar{f}(k)}$. Then $\lim_{n \rightarrow \infty} |x_n - c_f| = 0$, so that $x_n \rightarrow c_f$. Since P is closed and c_f is a limit point of P , it follows that $c_f \in P$. \square

Therefore, we can let $\varphi(f) = c_f$. It remains to show that φ is injective. Suppose that $f, g \in \{0, 1\}^{\mathbb{N}}$ with $f \neq g$. Let m be the least $k \in \mathbb{N}$ such that $f(k) \neq g(k)$. Then $I_{\bar{f}(k)} \cap I_{\bar{g}(k)} = \emptyset$ by construction. Hence

$$\bigcap_{n \in \mathbb{N}} I_{\bar{f}(n)} \cap \bigcap_{n \in \mathbb{N}} I_{\bar{g}(n)} = \emptyset.$$

This means that $\varphi(f) \neq \varphi(g)$. \square

Corollary 1.4.4. $\mathcal{C} \sim \mathbb{R}$.

1.5 Lecture 5

Definition 1.5.1. Let $X \subset \mathbb{R}$. To define the *(transfinite) iteration of the derivative of X* by recursion on the ordinals, let

$$\begin{aligned} X^0 &= X \\ X^{\alpha+1} &= (X^\alpha)' \\ X^\lambda &= \bigcap_{\alpha < \lambda} X^\alpha \quad \text{when } \lambda > 0 \text{ is a limit ordinal.} \end{aligned}$$

If X is closed, then the *Cantor-Bendixson rank* $\text{cb}(X)$ of X is the least ordinal α such that $X^\alpha = X^{\alpha+1}$.

Lemma 1.5.2. *If $X \subset \mathbb{R}$, then X' is closed.*

Proof. We must show that $X'' \subset X'$. Let $x \in X''$. Then for any open interval I around x , there is some $y \in (I \setminus \{x\}) \cap X'$. Since y is a limit point of X , there exist an open subinterval $J \subset I$ around y and a $z \in J \cap X$ such that $z \notin \{x, y\}$. Hence x is a limit point of X , as required. \square

Corollary 1.5.3. *X^β is closed for any ordinal β .*

Theorem 1.5.4 (Cantor-Bendixson). *If C is an uncountable closed set, then there exists a perfect set $P \subset C$ such that $C \setminus P$ is at most countable.*

Proof. By Lemma 1.5.2, we get a descending chain

$$C = C^0 \supset C^1 \supset C^2 \supset \dots \supset C^\alpha \supset \dots$$

There must be some ordinal τ such that $C^\tau = C^{\tau+1}$. Otherwise, $\alpha \neq \beta \implies C^\beta \neq C^\alpha$. In this case, there is a definable injection F from the proper class of ordinals OR into the set $\mathbb{P}(C)$, which implies that $\text{im } F$ is a set and thus that the image of the definable inverse $F^{-1} : \text{im } F \rightarrow \text{OR}$ is a set. But this is impossible since $\text{im } F^{-1} = \text{OR}$. Now, let $P = C^\tau$. Note that P is perfect as long as it's nonempty. Since C is uncountable, this holds as long as $C \setminus P$ is at most countable.

Enumerate the rational intervals $(J_n)_{n \in \mathbb{N}}$. We have that

$$C \setminus P = \bigcup_{\alpha < \tau} C^\alpha \setminus (C^\alpha)'.$$

Since the sets $C^\alpha \setminus (C^\alpha)'$ are pairwise disjoint, for any $x \in C \setminus P$, there is some unique α_x such that x is an isolated point of C^{α_x} . Hence there is a least $k_x \in \mathbb{N}$ such that $J_{k_x} \cap C^{\alpha_x} = \{x\}$. If $y \in C \setminus P$ with $x \neq y$, then it's easy to check that $k_x \neq k_y$. Thus, the function $C \setminus P \rightarrow \mathbb{N}$ given by $x \mapsto k_x$ is an injection, which proves that $C \setminus P$ is at most countable. \square

Corollary 1.5.5. *Any uncountable closed set is equipollent to \mathbb{R} .*

Remark 1.5.6 (Continuum Hypothesis). Observe that any real number r equals $\sup\{q \in \mathbb{Q} \mid q < r\}$. Thus, there exists a surjection $\mathbb{R} \rightarrow \mathbb{P}(\mathbb{Q})$. Further, we have the inclusion $\mathcal{C} \subset \mathbb{R}$. By the Cantor-Schröder-Bernstein, it follows that

$$|\mathbb{R}| = 2^{\aleph_0}.$$

The *continuum hypothesis* (CH) asserts that every uncountable subset of \mathbb{R} is equipollent to \mathbb{R} (i.e., that $\aleph_1 = 2^{\aleph_0}$).

Hilbert placed the resolution of CH first on his famous 1900 research agenda for mathematics in the twentieth century. W. Hugh Woodin, a prominent set theorist, has recently argued that CH is true. It's clear that every nonempty open set is equipollent to \mathbb{R} . Moreover, Corollary 1.5.5 shows that any closed set is either finite, equipollent to \mathbb{N} , or equipollent to \mathbb{R} . Therefore, no closed or open subset of \mathbb{R} is a counterexample to CH. Our next result, however, shows that we *cannot* confirm CH by showing that every uncountable subset of \mathbb{R} contains a perfect set.

Theorem 1.5.7. *There exists a set $X \subset \mathbb{R}$ such that $X \sim \mathbb{R}$ and for any perfect set P , $P \not\subset X$.*

Proof. This is Exercise 3 of Homework 2. □

2 Zermelo-Fraenkel set theory

2.1 Lecture 6

Definition 2.1.1. Define the *rank hierarchy (of sets)* by ordinal recursion as the following sequence of objects (viewed, for now, as formal or primitive objects).¹

$$\begin{aligned} V_0 &= \emptyset \\ V_{\alpha+1} &= \mathbb{P}(V_\alpha) \\ V_\lambda &= \bigcup_{\alpha < \lambda} V_\alpha \text{ for limit } \lambda. \end{aligned}$$

Remark 2.1.2. According to Platonism, the rank hierarchy is not an arbitrary invention of our minds but rather an object independent of our conceptual scheme. Our project will be to discover properties of the rank hierarchy. As such, our project will be one of *metamathematics*.

Definition 2.1.3. The *language of set theory* is the language of first-order logic with equality that has a single binary predicate symbol \in (intended to mean set membership). Note that *set* is a primitive notion and that every object in the universe is considered to be a set.

Remark 2.1.4. Historically, there were three major paradoxes of naive set theory that pushed certain mathematicians of the twentieth century to axiomatize it in first-order logic.

1. *Burali-Forti:* The binary relation \in is meant to be a well-ordering of the family of all ordinals OR (see Lemma 3.1.2 below). Further, we have that $\alpha \in \text{OR} \implies \alpha \subset \text{OR}$. It follows that if OR is a set, then $\text{OR} \in \text{OR}$, which is a contradiction since \in is a strict ordering.²
2. *Cantor:* In 1891, Cantor showed that if X is a set, then $|X| < |\mathbb{P}(X)|$. Thus, if the universe \mathcal{U} of all sets is a set, then $|U| < |\mathbb{P}(U)| \leq |U|$, a contradiction.

¹This is also known as the *cumulative hierarchy*, but we shall use this term for a more general concept later.

²This version of the paradox is anachronistic as it uses the definition of ordinals formulated by von Neumann in 1923. The original, equivalent version was published in 1897.

3. *Russell*: Originally, Frege proposed the *axiom schema of unrestricted comprehension*: If P is a first-order property, then there exists a set of the form $\{x \mid P(x)\}$. In 1902, Russell found that unrestricted comprehension is false. Indeed, it implies that

$$S := \{x \mid x \notin x\}$$

is a set. But, in this case, $S \in S \iff S \notin S$, which is false.

Those mathematicians hoped that such an axiomatization would act as a consistent yet powerful enough foundation for the great advances being made in analysis, algebra, and geometry.

Definition 2.1.5. The following eight axioms constitute *Zermelo-Fraenkel set theory* (ZF).

1. *Extensionality* (Ext):

$$\forall u(u \in X \leftrightarrow u \in Y) \rightarrow X = Y.$$

2. *Pairing* (Pair):

$$\forall a \forall b \exists c \forall x (x \in c \leftrightarrow x = a \vee x = b).$$

This asserts that for any two sets a and b , the *unordered pair* $\{a, b\}$ exists.

3. *Union* (Union):

$$\forall X \exists Y \forall u (u \in Y \leftrightarrow \exists z (z \in X \wedge u \in z)).$$

Let $X \cup Y := \bigcup \{X, Y\}$ and $\{a, b, c\} := \{a, b\} \cup \{c\}$.

4. *Power set* (PowerSet):

$$\forall X \exists Y \forall u (u \in Y \leftrightarrow u \subset X)$$

where the expression $u \subset X$ stands for $\forall z (z \in u \rightarrow z \in X)$.

5. *Schema of separation* (Sep):

$$\forall X \forall p \exists Y \forall u (u \in Y \leftrightarrow u \in X \wedge \varphi(u, p))$$

for each formula $\varphi(u, p)$ (of the language of set theory with free variables among u and p). Given Pair, this is equivalent to saying that for each formula $\psi(u, p_1, \dots, p_n)$,

$$\forall X \forall p_1 \dots \forall p_n \exists Y \forall u (u \in Y \leftrightarrow u \in X \wedge \psi(u, p_1, \dots, p_n)).$$

6. *Infinity* (Inf):

$$\exists S (\exists x (x \in S \wedge \forall y (y \notin x) \wedge \forall z (z \in S \rightarrow \exists u (u \in S \wedge \forall w (w \in u \leftrightarrow u \in z \vee u = z)))).$$

This asserts the existence of an *inductive set*.

7. *Schema of replacement* (Rep):

$$\begin{aligned} & \forall p (\forall x \forall y \forall z (\varphi(x, y, p) \wedge \varphi(x, z, p) \rightarrow y = z) \\ & \rightarrow \forall X \exists Y \forall y (y \in Y \leftrightarrow (\exists x \in X) \varphi(x, y, p))) \end{aligned}$$

for each formula $\varphi(x, y, p)$. Equivalently,

$$\begin{aligned} \forall p_1 \cdots \forall p_n (\forall x \forall y \forall z (\varphi(x, y, p_1, \dots, p_n) \wedge \varphi(x, z, p_1, \dots, p_n) \rightarrow y = z) \\ \rightarrow \forall X \exists Y \forall y (y \in Y \leftrightarrow (\exists x \in X) \varphi(x, y, p_1, \dots, p_n))) \end{aligned}$$

for each formula $\varphi(x, y, p_1, \dots, p_n)$. This asserts that if the class F is a function and $\text{dom } F$ is a set, then $\text{im } F$ is also a set.

8. *Regularity* (Reg):³

$$\forall u ((\exists x (x \in u)) \rightarrow (\exists m (m \in u \wedge \forall y (y \in m \rightarrow \neg(y \in u))))).$$

Definition 2.1.6. A *class* is a collection (used informally) of sets that is definable (with parameters in the language of set theory). Any set S can be written as the class $S = \{x \mid x \in S\}$. Any class that is not a set is called a *proper class*.

Note 2.1.7.

1. The converse of **Ext** is an axiom of predicate logic, so that $X = Y$ if and only if X and Y consist of the same elements.
2. By **Pair** + **Union**, any class consisting of finitely many sets is a set.
3. The class of all sets is not a set. Otherwise, $\{x \mid x \notin x\}$ is a set by **Sep**, in which case we get Russell's paradox.
4. **Sep** asserts that any subclass of a set is a set.
5. **Rep** asserts that every nonempty set has an \in -minimal element.
6. There is no infinite descending chain $x_0 \ni x_1 \ni x_2 \ni x_3 \ni \cdots$, for otherwise $\{x_n \mid n \in \mathbb{N}\}$ (which is a set by **Rep**) violates **Reg**. In particular, if A is a set, then $A \notin A$.

Example 2.1.8 (The empty set and omega).

1. The sentence $(\exists x)x = x$ is an axiom of predicate logic. Hence there is at least one set S . By **Ext** + **Sep**, the empty set exists because it equals $\{x \mid x \neq x\}$, which is a subclass of S .
2. By **Inf**, there exists a set containing the class

$$\omega := \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \dots\}.$$

For any set x , let $\varphi(x)$ denote the (first-order) formula expressing that x is inductive. But note that

$$\omega = \{n \in S \mid \forall y (\varphi(y) \rightarrow n \in y)\},$$

which is a set by **Sep**, specifically, the smallest inductive set.

Notation. For each set x , let $V(x)$ denote the formula $\exists \alpha (x \in V_\alpha)$.

³The axiom of regularity is also called the *axiom of foundation*.

Definition 2.1.9. The *von Neumann universe*

$$V := \bigcup_{\alpha \in \text{OR}} V_\alpha$$

is precisely the proper class $\{x \mid V(x)\}$ of all sets belonging to some stage of the rank hierarchy.

Note 2.1.10.

1. The first five axioms of ZF are satisfied by the set $\text{HF} := V_\omega$, whose elements are called *hereditarily finite sets*. The axiom of infinity, however, requires $V_{\omega+1}$. Thus, we need to have $V_{\omega+\omega}$ since $\omega + \omega$ is the first limit ordinal after $\omega + 1$.
2. The set $V_{\omega+\omega}$ satisfies each of the first six axioms.
3. Consider the functional relation F on $V_{\omega+\omega}$ defined by $n \mapsto \omega + n$ where $n \in \omega$. Then $\text{dom } F = \omega$, which is a set in $V_{\omega+\omega}$. But $\text{im } F = \omega + \omega \notin V_{\omega+\omega}$. This shows that $V_{\omega+\omega}$ fails to satisfy Rep. For any $n \geq 2$, a similar argument shows that $\underbrace{V_\omega + \dots + \omega}_{n \text{ copies}}$ fails to satisfy Rep.
4. HF, however, satisfies Rep. This is because the image of any function with finite domain consists of finitely many sets and thus is itself a set.

Remark 2.1.11. In 1908, Zermelo introduced the first six axioms of ZF (along with the axiom of choice). For this reason, we shall denote the theory consisting of them by Z. In 1922, Fraenkel and Skolem together introduced the axiom schema of replacement to ensure that objects like $V_{\omega+\omega}$ counted as a set. Finally, in 1925, von Neumann introduced the axiom of regularity to enable proofs by induction on so-called well-founded proper classes, such as OR.

In ZF, we avoid each of the three paradoxes from Remark 2.1.4 by replacing unrestricted comprehension with Sep. In fact, they serve as proofs that the following classes are proper.

1. The class of all ordinals (see Corollary 3.3.4 below).
2. The class of all sets.
3. The class $\{x \mid x \notin x\}$.

2.2 Lecture 7

Definition 2.2.1. For any sets x and y , the *ordered pair* (x, y) is the set $\{\{x\}, \{x, y\}\}$, which exists by Pair.

Lemma 2.2.2. For any sets X and Y , the *Cartesian product* $X \times Y := \{(x, y) \mid x \in X \wedge y \in Y\}$ exists.

Proof. If $x \in X$ and $y \in Y$, then (x, y) is in $\mathbb{P}(\mathbb{P}(X \cup Y))$, which exists by Union + PowerSet. In particular,

$$\begin{aligned} X \times Y = \{z \in \mathbb{P}(\mathbb{P}(X \cup Y)) \mid & \exists x_1 \exists x_2 (\exists y_1 \in z)(\exists y_2 \in z)(x_1 \neq x_2 \wedge \forall c(c \in y_1 \leftrightarrow c = x_1) \\ & \wedge \forall d(d \in y_2 \leftrightarrow (d = x_1 \vee d = x_2)) \\ & \wedge (\forall k \in z)(k = y_1 \vee k = y_2))\}, \end{aligned}$$

which exists by Sep. □

Remark 2.2.3.

1. Let T denote the theory $(\mathbf{ZF} \setminus \{\text{Inf}\}) \cup \{\neg \text{Inf}\}$. Note that $V_\omega \models T$. It is known that T is bi-interpretable with PA (Peano arithmetic). This means that there is some *translation* τ from the language of HF to the language of PA such that $\mathbf{HF} \vdash \varphi \implies \mathbf{PA} \vdash \varphi^\tau$.
2. We can find some theory T in the language of set theory such that $T \vdash [\exists \lambda (V_\lambda \models \mathbf{ZF})]$. But $T \neq \mathbf{ZF}$, since Gödel's second incompleteness theorem implies that \mathbf{ZF} cannot prove its own consistency.

3 Ordinal numbers

Definition 3.0.1. Let R be a binary relation on a class X

1. . For each $x \in X$, the *extension of x* is

$$\text{ext}_R(x) = \{z \in X \mid zRx\}.$$

2. We say that R is *extensional* if $\text{ext}_R(x) \neq \text{ext}_R(y)$ for any distinct $x, y \in X$.
3. We say that R is *well-founded* if

- every nonempty subset of X has an R -minimal element and
- $\text{ext}_R(x)$ is a set for each $x \in X$ (i.e., R is *set-like*).

In this case, we also say that X is *well-founded (with respect to R)*.

Proposition 3.0.2. Let X be a well-founded class with respect to R . Then every non-empty subclass of X has an R -minimal element.

Definition 3.0.3. Let X be a class. A linear ordering $(X, <)$ is a *well-ordering* if it is well-founded. If X is a set, then we call it a *well-ordered set* or *woset*.

Example 3.0.4.

1. \mathbb{N} with its usual order is a woset.
2. \mathbb{N} ordered as $\{0, 2, 4, 6, \dots, 1, 3, 5, 7, \dots\}$ is a woset.

Non-example 3.0.5. $\mathbb{Q}_{>0}$ with its usual order is not a woset. For example, the subset $\{\frac{1}{n} \mid n \in \mathbb{Z}_{>0}\}$ has no minimal element.

Lemma 3.0.6. Let $(W, <)$ be a woset. If $f : W \rightarrow W$ is order-preserving, then $f(w) \geq w$ for each $w \in W$.

Proof. Suppose, towards a contradiction, that the set $M := \{w \in W \mid f(w) < w\}$ is nonempty. Let $m = \min(M)$. Then $f(f(m)) < f(m) < m$, a contradiction. \square

Corollary 3.0.7. Let $(W_1, <_1)$ and $(W_2, <_2)$ be two well-orderings.

- (1) The structure $(W_1, <_1)$ is rigid in the sense that id_{W_1} is the only automorphism of it.
- (2) Any isomorphism $\delta : W_1 \rightarrow W_2$ is unique.

Proof.

- (1) Let $f : (W, <) \rightarrow (W, <)$ be an automorphism. Let $x \in W$. By Lemma 3.0.6, we see that $f(x) \geq x$ and $x = f^{-1}(f(x)) \geq f(x)$. Hence $f(x) = x$.
- (2) Suppose, towards a contradiction, that δ and ψ are distinct isomorphisms $W_1 \rightarrow W_2$. Then $\psi^{-1} \circ \delta$ is an automorphism of W_1 . But $\delta^{-1} \neq \psi^{-1}$, and thus $\psi^{-1} \circ \delta \neq \text{id}_{W_1}$. This contradicts part (1).

□

Definition 3.0.8. Let $(W, <)$ be a woset and $x \in W$. The *initial segment of W determined by x* is the set

$$W_x := \{w \in W \mid w < x\}.$$

Note 3.0.9. Let $(W, <)$ be a woset. If $x < y$, then W_x is an initial segment of W_y .

Lemma 3.0.10. Let $(W, <)$ be a woset and $x \in W$. Then $W \not\cong W_x$.

Proof. Let $f : W \xrightarrow{\cong} W_x$. By Lemma 3.0.6, $f(x) > x$ since $x \notin W_x$. But since $f(x) \in W_x$, it follows that both $f(x) < x$ and $x > f(x)$, which is impossible. □

Theorem 3.0.11. Let $(W, <)$ and $(Y, <')$ be any two well-orderings. Then exactly one of the following scenarios occurs.

- (a) $W \cong Y$.
- (b) $W \cong Y_y$ for some $y \in Y$.
- (c) $W_w \cong Y$ for some $w \in W$.

Proof. First, note that Lemma 3.0.10 implies that no two of (a), (b), and (c) can occur simultaneously. We must show that at least one occurs. Consider the set

$$f := \{(w, y) \in W \times Y \mid W_w \cong Y_y\}.$$

If $Y_y \cong Y_{y'}$, then $y = y'$ by Lemma 3.0.10. Hence f is a partial function. Moreover, if $w, w' \in \text{dom } f$ and $w < w'$ with $h : W_{w'} \xrightarrow{\cong} Y_{f(w')}$, then $W_w \cong Y_{h(w)}$, so that $f(w) = h(w) <' f(w')$. It follows that f is order-preserving. There are three cases to consider.

- (i) Suppose that $\text{dom } f \subsetneq W$. Let $m = \min(W \setminus \text{dom } f)$. If $Y \setminus \text{im } f \neq \emptyset$, then $(m, n) \in f$ where $n = \min(Y \setminus \text{im } f)$, a contradiction. Hence $\text{im } f = Y$. Then $f \upharpoonright_{W_m} : W_m \rightarrow Y$ is an isomorphism. Thus, scenario (c) occurs.
- (ii) Suppose that $\text{dom } f = W$ but $\text{im } f \subsetneq Y$. Then scenario (b) occurs.
- (iii) Suppose that $\text{dom } f = W$ and $\text{im } f = Y$. Then f is an isomorphism, i.e., scenario (a) occurs.

□

Definition 3.0.12. A class x is *transitive* if for any $y \in x$, we have $y \subset x$.

Note 3.0.13. A set x is transitive if and only if $z \in y \in x \implies z \in x$. By Note 5(4), it follows that \in is a partial ordering of OR.

Definition 3.0.14. A set x is an *ordinal (number)*, written as $\text{ord}(x)$, if x is transitive and (x, \in) is a well ordering.

Notation. We shall use the symbols \in and $<$ interchangeably for the ordering of ordinals.

Lemma 3.0.15. Let α be an ordinal. Let $\alpha + 1 = \alpha \cup \{\alpha\}$. Then $\alpha + 1$ is the minimal ordinal greater than α .

Proof. It is easy to see that $\alpha + 1$ is transitive. Note that $\alpha = \max\{\alpha + 1\}$. Thus, it is well-ordered since for any nonempty $X \subset \alpha + 1$, we have $\min(X) = \min(X \setminus \{\alpha\})$. Moreover, if $\alpha \in \beta$ where β is an ordinal and $\beta \leq \alpha + 1$, then $\alpha + 1 \subset \beta \subset \alpha + 1$, in which case $\beta = \alpha + 1$. Therefore, $\alpha + 1$ is minimal, as desired. \square


Definition 3.0.16. We say that an ordinal α is a *successor (ordinal)* if $\alpha = \beta + 1$ for some ordinal β . We say that α is a *limit (ordinal)* if it is not a successor.

Note 3.0.17. Define $\sup(\emptyset) = 0$. Then for any limit ordinal α , we have $\alpha = \sup\{x \mid x \in \alpha\} = \bigcup \alpha$.

Example 3.0.18. The set ω is the smallest nonzero limit ordinal. Its elements are precisely the *natural numbers*.

Definition 3.0.19.

1. A set x is *finite* if there is an $n \in \omega$ such that $x \sim n$.
2. A set x is *Dedekind finite* if for any proper subset y of x , $y \not\sim x$.



Conversely, let $x \in \gamma$. Since $\gamma \subset \beta$, we get $x \in \beta$. By our choice of γ , it follows that $x \in \alpha$. Hence $\gamma \subset \alpha$. \square

Lemma 3.1.2. (OR, \in) is a well-ordered class.

Proof. It is clear that \in is set-like. Let $S \subset \text{OR}$ be a nonempty subset. We must show that S has a minimal element. Note that $\bigcap S \subset \alpha$ for every $\alpha \in S$. Therefore, $(\bigcap S, \in)$ is a well-ordering as the restriction of a well-ordering. Now, suppose that $\beta \in \bigcap S$. Then $\beta \in \alpha$ for every $\alpha \in S$. By transitivity, for any $x \in \beta$, we have $x \in \alpha$ for every $\alpha \in S$, so that $\beta \subset \bigcap S$. This implies that $\bigcap S$ is transitive. Hence $\bigcap S$ is an ordinal. By Lemma 3.1.1, we see that $\bigcap S$ is minimal.

It remains to show that OR is linearly ordered. By Lemma 3.1.1, it suffices to show that if α and β are ordinals, then either $\alpha \subset \beta$ or $\beta \subset \alpha$. Suppose, towards a contradiction, that both $\alpha \not\subset \beta$ and $\beta \not\subset \alpha$. Our last paragraph shows that $\alpha \cap \beta$ is an ordinal, which must be a proper subset both of α and of β . Lemma 3.1.1 implies that $\alpha \cap \beta \in \alpha$ and $\alpha \cap \beta \in \beta$. Hence $\alpha \cap \beta \in \alpha \cap \beta$, a contradiction. \square

Corollary 3.1.3. If S is a nonempty set of ordinals, then $\bigcap S = \min(S)$.

Lemma 3.1.4. Let $(W, <)$ be a woset. Suppose that for every $x \in W$, there is some ordinal $Z(x)$ such that $Z(x) \cong W_x$. Then W is isomorphic to some ordinal.

Proof. Lemma 3.0.10 together with Lemma 3.1.2 proves that such a $Z(x)$ must be unique. Therefore,

$$A := \{Z(x) \mid x \in W\}$$

is a set by Rep. Note that (A, \in) is a well-ordering. To see that A is transitive, let $v \in A$. Then $v = Z(y)$ for some $y \in W$. Let $x \in Z(y)$. Then

$$x \cong X_{g_y^{-1}(x)} \cong Z(g_y^{-1}(x))$$

where $g_y : W_y \xrightarrow{\cong} Z(y)$. This implies that $x = Z(g_y^{-1}(x))$ since both x and $Z(g_y^{-1}(x))$ are ordinals. Thus, $x \in A$, so that $v \subset A$.

Define $f : W \rightarrow A$ by $f(x) = Z(x)$.

Claim. Let $x, y \in W$. Then $x < y \implies Z(x) \subsetneq Z(y)$.

Proof. Let $x < y$. Consider the isomorphism $g_y : W_y \rightarrow Z(y)$. We have that

$$W_x = \{z \in W_y \mid z < x\} = (W_y)_x.$$

Hence $g_y \upharpoonright_{W_x}$ is an isomorphism from W_x onto $Z(y) \upharpoonright_{g_y(x)}$. It follows that $Z(y) \upharpoonright_{g_y(x)} \cong Z(x)$, so that

$$Z(x) = Z(y) \upharpoonright_{g_y(x)} \subsetneq Z(y),$$

as desired. \square

By Lemma 3.1.1, it follows that f is order-preserving. Since f is clearly surjective, it is an isomorphism. \square

Theorem 3.1.5. *For any woset $(W, <)$, there is a unique ordinal α such that $W \cong \alpha$.*

Proof. Lemma 3.0.10 together with Lemma 3.1.2 proves that such an α must be unique. We must prove that such an α exists.

Claim. *For any $x \in W$, there is some ordinal $Z(x)$ such that $W_x \cong Z(x)$.*

Proof. Now, suppose, towards a contradiction, that the set $D := \{x \in W \mid W_x \text{ is not isomorphic to an ordinal}\}$ is nonempty. Let $m = \min(D)$. Then for each $y < m$, there is some ordinal $Z(y)$ such that $(W_m)_y = W_y \cong Z(y)$. By Lemma 3.1.4, this implies that W_m is isomorphic to some ordinal, contrary to our choice of m . \square

It follows from Lemma 3.1.4 that $W \cong \alpha$ for some ordinal α . \square

3.2 Lecture 9

Lemma 3.2.1 (Transfinite induction). *Suppose that C is a class of ordinals. Suppose that*

- (a) $0 \in C$,
- (b) $\alpha + 1 \in C$ whenever $\alpha \in C$, and
- (c) for any nonzero limit λ , $\lambda \in C$ whenever $\{\beta \mid \beta < \lambda\} \subset C$.

Then $C = \text{OR}$.

Proof. Suppose, towards a contradiction, that $\text{OR} \setminus C \neq \emptyset$. Let $m = \min(\text{OR} \setminus C)$. It's easy to see that m is neither zero nor a successor. Hence m is a nonzero limit. By condition (c), there must be some $\alpha < m$ such that $\alpha \notin C$. Then $\alpha \in \text{OR} \setminus C$, so that m is not minimal, a contradiction. \square

Note 3.2.2. We can replace the condition “ $\alpha + 1 \in C$ whenever $\alpha \in C$ ” with the condition “ $\alpha + 1 \in C$ whenever $\{\beta \mid \beta \leq \alpha\} \subset C$.”

Theorem 3.2.3 (Transfinite recursion). *Let Seq denote the class $\{f : \alpha \rightarrow x \mid \alpha \in \text{OR} \wedge x \in V\}$ of all transfinite sequences. For any class functional relation $G : \text{Seq} \rightarrow V$, there exists a unique class function $F : \text{OR} \rightarrow V$ such that*

$$F(\alpha) = G(F \upharpoonright_\alpha)$$

for every $\alpha \in \text{OR}$.

Proof. We use transfinite induction to prove that there is a proper class $\langle f_\alpha \mid \alpha \in \text{OR} \rangle$ (viewed as a sequence) such that for each $\alpha \in \text{OR}$, f_α is the unique function such that

- (i) $\text{dom } f_\alpha = \alpha$,
- (ii) for any $\beta \in \alpha$, $f_\alpha(\beta) = G(f_\alpha \upharpoonright_\beta)$, and
- (iii) for any $\beta \in \alpha$, $f_\alpha \upharpoonright_\beta = f_\beta$.

There are three cases to consider for our induction.

- Suppose that $\alpha = 0$. Then $f_\alpha := \emptyset$ is the unique function satisfying all three conditions.

- Suppose that $\alpha = \beta + 1$ and that there exists a unique f_β satisfying all three conditions. Let

$$f_{\beta+1}(\gamma) = \begin{cases} G(f_\beta) & \gamma = \beta \\ f_\beta(\gamma) & \gamma < \beta \end{cases}.$$

Let $\delta \in \beta + 1$. If $\delta = \beta$, then $f_{\beta+1}(\delta) = G(f_\beta) = G(f_{\beta+1} \upharpoonright_\beta) = G(f_{\beta+1} \upharpoonright_\delta)$. If $\delta \in \beta$, then $f_{\beta+1}(\delta) = \underbrace{f_\beta(\delta)}_{\text{by induction}} = G(f_\beta \upharpoonright_\delta) = G(f_{\beta+1} \upharpoonright_\delta)$. Moreover, $f_{\beta+1} \upharpoonright_\beta = f_\beta$, and $f_{\beta+1} \upharpoonright_\gamma = f_\beta \upharpoonright_\gamma = f_\gamma$ for any $\gamma \in \beta$ by induction. We have shown that $f_{\beta+1}$ satisfies all three properties.

It remains to show that $f_{\beta+1}$ is unique. Suppose that there is another function g satisfying all three properties for $\beta + 1$. Suppose, towards a contradiction, that the set $E := \{\alpha \in \beta + 1 \mid g \neq f_{\beta+1}\}$ is nonempty. Let $m = \min(E)$. Then $g \upharpoonright_m = f_{\beta+1} \upharpoonright_m$, so that

$$g(m) = G(g \upharpoonright_m) = G(f_{\beta+1} \upharpoonright_m) = f_{\beta+1}(m),$$

a contradiction. It follows that $g = f_{\beta+1}$.

- Suppose that α is a nonzero limit and that, for each $\beta \in \alpha$, there exists a unique f_β satisfying all three conditions. Let

$$f_\alpha = \bigcup_{\beta \in \alpha} f_\beta.$$

By applying **Rep** followed by **Union**, we see that f_α is a set. By condition (iii) from our IH, we also see that f_α is a partial function on α . In fact, since α is a limit, f_α has domain equal to α . Hence f_α satisfies condition (i). Further, it is clear that f_α satisfies condition (iii).

Let $\beta \in \alpha$. Note that $\beta + 1 \in \alpha$. By applying our IH, we get

$$\begin{aligned} f_\alpha(\beta) &= f_{\beta+1}(\beta) \\ &= G(f_{\beta+1} \upharpoonright_\beta) \\ &= G(f_\beta) \\ &= G(f_\alpha \upharpoonright_\beta). \end{aligned}$$

This shows that f_α satisfies condition (ii). The fact that f_α is unique follows exactly as in our last case.

This completes our induction. Now, define $F : \mathbf{OR} \rightarrow V$ by $\alpha \mapsto G(f_\alpha)$. This is clearly a class function. Also, if $\alpha \in \mathbf{OR}$, then $F \upharpoonright_\alpha = \{(\beta, F(\beta)) \mid \beta \in \alpha\}$ is a set by **Rep** because each $F(\beta)$ is a set. Note that $\text{dom } f_\alpha = \alpha = \text{dom } F \upharpoonright_\alpha$ and that for any $\beta \in \alpha$,

$$f_\alpha(\beta) = G(f_\alpha \upharpoonright_\beta) = G(f_\beta) = F_\alpha(\beta).$$

Therefore, $f_\alpha = F \upharpoonright_\alpha$, so that $F(\alpha) = G(F \upharpoonright_\alpha)$, as required.

Let $F' : \mathbf{OR} \rightarrow V$ also have $F'(\alpha) = G(F' \upharpoonright_\alpha)$. Using transfinite induction, it is easy to check that $F'(\alpha) = F(\alpha)$ for any ordinal α . Thus, F is unique. \square

3.3 Lecture 10

Exercise 3.3.1. Show that a set x is an ordinal if and only if x is transitive and every $y \in x$ is transitive.

Lemma 3.3.2. If α is an ordinal and $x \in \alpha$, then x is an ordinal.

Proof. This follows from the (\implies) direction of Exercise 3.3.1. \square

Corollary 3.3.3. If X is a nonempty set of ordinals, then $\bigcup X$ is an ordinal and $\sup(X) = \bigcup X$.

Corollary 3.3.4 (Burali-Forti). OR is a proper class.

Proof. Otherwise, $\sup \text{OR}$ and $\sup \text{OR} + 1$ are ordinals by Corollary 3.3.3, in which case

$$\sup \text{OR} < \sup \text{OR} + 1 \leq \sup \text{OR},$$

a contradiction. \square

Definition 3.3.5. An ordinal κ is a *cardinal (number)* (written as $\text{card}(\kappa)$) if for every ordinal $\lambda < \kappa$, $\kappa \not\sim \lambda$.

Example 3.3.6. The finite cardinals are precisely the finite ordinals. In addition, ω is an infinite cardinal.

Note 3.3.7. Every infinite cardinal is a nonzero limit ordinal.

Definition 3.3.8. Let X and Y be sets.

- Let $|X| \leq |Y|$ if there is an injection from X into Y .
- Let $|X| < |Y|$ if $|X| \leq |Y|$ but $|Y| \not\leq |X|$.

Theorem 3.3.9 (Cantor-Schröder-Bernstein). If $|X| \leq |Y|$ and $|Y| \leq |X|$, then $|X| = |Y|$.

Theorem 3.3.10 (Cantor). $|X| < |\mathbb{P}(X)|$.

Proof. It's clear that $|X| \leq |\mathbb{P}(X)|$. Let $f : X \rightarrow \mathbb{P}(X)$. Consider the set

$$Z := \{x \in X \mid x \notin f(x)\}.$$

If $Z = f(y)$ for some $y \in X$, then $y \in Z \iff y \notin Z$, which is false. Thus, $Z \notin \text{im } f$. This shows that there is no surjection (and hence no bijection) from X onto $\mathbb{P}(X)$. By Theorem 3.3.9, it follows that $|\mathbb{P}(X)| \not\leq |X|$, so that $|X| < |\mathbb{P}(X)|$. \square

Remark 3.3.11. Recall that $\mathbb{R} \sim \mathbb{P}(\omega)$. The continuum hypothesis asserts that there is no set X such that $|\omega| < |X| < |\mathbb{P}(\omega)|$.

Definition 3.3.12. For any set X , let $H(X)$ denote the least ordinal α such that $\alpha \not\sim X$. We call $H(X)$ the *Hartogs number* of X .

Theorem 3.3.13. For every cardinal κ , there is some cardinal $\lambda > \kappa$.

Proof. Suppose that $H(\kappa)$ exists. Then $\kappa \sim \beta$ for some $\beta \in H(\kappa)$, so that $|\kappa| \leq |H(\kappa)|$. But clearly $|\kappa| \neq |H(\kappa)|$. Therefore, $|\kappa| < |H(\kappa)|$, which means that it suffices to show that $H(\kappa)$, in fact, exists.

Note that the class D of all well-orderings of κ is a subclass of the set $\mathbb{P}(\kappa \times \kappa)$. Hence D is a set by Sep. If $H(\kappa)$ does not exist, then we get an injective class function $F : \text{OR} \rightarrow D$. In this case, $\text{im } F^{-1} = \text{OR}$ is a set by Rep, a contradiction. Thus, $H(\kappa)$ exists. \square

Note 3.3.14. Our proof shows that $H(X)$ exists for any X in ZF.

Notation. If κ is a cardinal, then let κ^+ denote the least cardinal greater than κ .

Corollary 3.3.15. *The class C of all cardinals is a proper class.*

Proof. For any ordinal α , consider the cardinal $\kappa = \min\{\beta \in \text{OR} \mid \beta \sim \alpha\}$. By Cantor-Schröder-Bernstein, it's easy to check that $\alpha < \kappa^+$. Therefore, if C is a set, then $\bigcup C$ is a set and equals OR , a contradiction. Hence C is a proper class. \square

Definition 3.3.16. By transfinite recursion, define the sequence

$$\begin{aligned}\aleph_0 &= \omega \\ \aleph_{\alpha+1} &= \aleph_\alpha^+ \\ \aleph_\lambda &= \bigcup_{\beta \in \lambda} \aleph_\beta \text{ when } \lambda > 0 \text{ is a limit.}\end{aligned}$$

The symbol ω_α means the same thing as \aleph_α .

Lemma 3.3.17. *If $\lambda > 0$ is a limit ordinal, then $\text{card}(\aleph_\lambda)$.*

Proof. Suppose, towards a contradiction, that there exist an $x \in \aleph_\lambda$ and a bijection $f : x \rightarrow \aleph_\lambda$. Then $x \in \aleph_\beta$ for some successor $\beta \in \lambda$. It follows that

$$|\aleph_\lambda| = |x| \leq |\aleph_\beta| \leq |\aleph_\lambda|.$$

Hence $\aleph_\beta \sim x$. But we know that \aleph_β is a cardinal, a contradiction. \square

4 The axiom of choice

Definition 4.0.1 (The axiom of choice (AC)). Let I be a set and $C = \{A_i\}_{i \in I}$ where each A_i is nonempty. Then there is a function $f : I \rightarrow \bigcup C$ such that $f(i) \in A_i$ for each $i \in I$. (Such an f is called a *choice function*.)

Notation. Let ZFC denote the theory ZF + AC.

Definition 4.0.2 (Zermelo's well-ordering principle (WOP)). For any set x , there is some cardinal λ such that $x \sim \lambda$.

Theorem 4.0.3. $\text{AC} \iff \text{WOP}$.

Definition 4.0.4. (ZFC) Let X be a set, The *cardinality of X* is the cardinal number

$$|X| := \min\{\alpha \in \text{OR} \mid \alpha \sim X\}.$$

Remark 4.0.5.

1. One can show that $\text{ZF} \not\models \underbrace{\forall X \forall Y (|X| \leq |Y| \vee |Y| \leq |X|)}_\psi$. In fact, AC is equivalent to the sentence ψ . In particular, AC is independent of ZF.

2. Let φ denote the sentence “Every surjection has a right inverse.” Then $\mathbf{ZF} \not\models \varphi$, whereas $\mathbf{ZFC} \models \varphi$. Since every right inverse is injective, the following assertion is also provable in \mathbf{ZFC} : If there exists a surjection of X onto Y , then $|Y| \leq |X|$.
3. We can, however, prove in \mathbf{ZF} that every injection has a left inverse and that every left inverse is surjective. Therefore, in \mathbf{ZF} , if $|Y| \leq |X|$, then there is a surjection of X onto Y . It follows that, in \mathbf{ZFC} , $|Y| \leq |X|$ if and only if there is a surjection of X onto Y .

Note 4.0.6 (Linear ordering of cardinals). 1. Consider the binary relation $<$ on the class of all cardinal numbers where $|X| < |Y|$ if there is an injection from X into Y but $|X| \neq |Y|$. In light of the Cantor-Schröder-Bernstein, we see that $<$ is a partial ordering. By Remark 4.0.5, we see that $<$ is actually a linear ordering in \mathbf{ZFC} .

2. In \mathbf{ZFC} , if λ is an infinite cardinal, then $\lambda = \aleph_\alpha$ for some ordinal α . In this case, we get a transfinite enumeration of class of all cardinals

$$0 < 1 < 2 < 3 < \cdots < \aleph_0 < \aleph_1 < \aleph_2 < \aleph_3 < \cdots < \aleph_\omega < \cdots.$$

Definition 4.0.7 (Cardinal arithmetic (in \mathbf{ZFC})). Let κ and λ be cardinals. Define cardinal addition, multiplication, and exponentiation, respectively, as follows.

1. $\kappa + \lambda = |\kappa \amalg \lambda|$.
2. $\kappa \cdot \lambda = |\kappa \times \lambda|$.
3. $\kappa^\lambda = |\{f \mid f : \lambda \rightarrow \kappa\}|$.

Note 4.0.8. For any ordinal α , the lexicographic ordering of $\alpha \times \alpha$ is a well-ordering. Hence we do not need the axiom of choice for our definition of cardinal multiplication.

Lemma 4.0.9. (\mathbf{ZF}) $\aleph_\alpha \cdot \aleph_\alpha = \aleph_\alpha$ for any ordinal α .

Proof. Consider the binary class relation \triangleleft on $\mathbf{OR} \times \mathbf{OR}$ where $(\alpha, \beta) \triangleleft (\gamma, \delta)$ if at least one of the following conditions holds.

- $\max\{\alpha, \beta\} < \max\{\gamma, \delta\}$.
- $\max\{\alpha, \beta\} = \max\{\gamma, \delta\}$ and $\alpha < \delta$.
- $\max\{\alpha, \beta\} = \max\{\gamma, \delta\}$ and $\alpha = \delta$ and $\beta < \delta$.

It is straightforward to check that \triangleleft is a well-ordering. Now, define the class function $\Gamma : \mathbf{OR} \times \mathbf{OR} \rightarrow \mathbf{OR}$ by $\gamma(\alpha, \beta) = \delta$ where δ is the unique ordinal isomorphic to the initial segment $(\mathbf{OR} \times \mathbf{OR})_{(\alpha, \beta)}$. It is easy to see that Γ is injective. Thus, it suffices to show that

$$\Gamma(\omega_\alpha \times \omega_\alpha) = \omega_\alpha$$

for each ordinal α .

Note that for any $(n, m) \in \omega \times \omega$,

$$\Gamma(n, m) = m(n + 1).$$

Hence $\omega = \Gamma(\omega \times \omega)$, as desired.

Suppose, towards a contradiction, that the class $E := \{x > 0 \mid \Gamma(\omega_x \times \omega_x) \neq \omega_x\}$ of ordinals is nonempty. Let $\beta = \min(E)$. Then $\Gamma(\gamma, \delta) = \omega_\beta$ for some $\gamma, \delta \in \omega_\beta$. Since ω_β is a limit, we can find some $\eta \in \omega_\beta$ such that $\gamma, \delta \in \eta$. Note that $(\gamma, \delta) \in \eta \times \eta$. Since $\eta \times \eta = (\text{OR} \times \text{OR})_{(0, \eta)}$, it follows that

$$\Gamma(\eta \times \eta) \supset \underbrace{\Gamma(\{(x, y) \mid (x, y) \triangleleft (\gamma, \delta)\})}_{\text{transitive}} \supset \omega_\beta.$$

This implies that $|\aleph_\beta| \leq |\eta \times \eta|$. But

$$|\eta \times \eta| = ||\eta| \times |\eta|| = |\eta| < \aleph_\beta,$$

a contradiction. □

Corollary 4.0.10. (ZFC) *Let κ and λ be infinite cardinals. Then*

$$\kappa \cdot \lambda = \kappa + \lambda = \max\{\kappa, \lambda\},$$

4.1 Lecture 11

Remark 4.1.1. If \mathcal{U} denotes the universe of all sets, then $\text{ZF} \setminus \{\text{Reg}\} \vdash \text{Reg} \leftrightarrow \mathcal{U} = V$.

We now prove a theorem stated in Lecture 10.

Proof. (AC \iff WOP)

(\Leftarrow) Let I be any set and let $C := \{A_i\}_{i \in I}$ be a collection of nonempty sets. By Union, we see that $\bigcup C$ is a set. By WOP, there exists a well-ordering $<$ on $\bigcup C$. Note that each A_i is a nonempty subset of $\bigcup C$. Thus, A_i contains a least element a_i . Now, define $f : I \rightarrow \bigcup C$ by $f(i) = a_i$. As long as f is well-defined, it is a choice function. To show that it is well-defined, it suffices to show that the a_i are unique. Suppose that both a_i and b_i are minimal elements of A_i . Note that $<$ is a linear ordering of C . Thus, either $a_i \leq b_i$ or $b_i \leq a_i$. But both are minimal, so that, in either case, $a_i = b_i$. This shows that f is well-defined.

(\Rightarrow) Let X be a set. Let $E = \mathbb{P}(X) \setminus \{\emptyset\}$, which is a collection of nonempty sets. By AC, there is some choice function $f : E \rightarrow \bigcup E$. By transfinite recursion, we can define a sequence $F : \text{OR} \rightarrow V$ by

$$F(\alpha) = f(X \setminus \{F(\beta) \mid \beta \in \alpha\}).$$

Since f is a choice function, we have that $F(\alpha) \notin \{F(\beta) \mid \beta \in \alpha\}$ whenever $\{F(\beta) \mid \beta \in \alpha\} \neq X$. Hence if there is no $\delta \in \text{OR}$ such that $\{F(\gamma) \mid \gamma \in \delta\} = X$, then F is injective. But, in this case, $\text{im } F^{-1} = \text{OR}$ is a set by Rep, a contradiction. Therefore, there exists a least such δ . Then $F \upharpoonright_\delta$ is a bijection from δ onto X . □

Definition 4.1.2. Let $(P, <)$ be a partially ordered set. A *chain in P* is a nonempty subset $X \subset P$ such that $(X, < \upharpoonright_X)$ is a linear ordering.

Definition 4.1.3 (Zorn's lemma (Zorn)). If $(P, <)$ is a nonempty poset and every chain in P has an upper bound in P , then P has a maximal element.

Lemma 4.1.4. WOP \implies Zorn.

Proof. Let $(P, <_P)$ be a nonempty poset such that every chain in P has an upper bound in P . By assumption, we have a bijection $f : \lambda \rightarrow P$ for some cardinal λ . We see that $\lambda > 0$ for otherwise $P = \emptyset$. Note that $P = \{f(\delta) \mid \delta \in \lambda\}$. By transfinite recursion, define $F : \text{OR} \rightarrow \lambda + 1$ by

$$F(\alpha) = \begin{cases} \min\{\gamma \in \lambda \mid \beta \in \alpha \rightarrow f(F(\beta)) <_P f(\gamma)\} & \{\gamma \in \lambda \mid \beta \in \alpha \rightarrow f(F(\beta)) <_P f(\gamma)\} \neq \emptyset \\ \lambda & \text{otherwise} \end{cases}.$$

Note that $\alpha_1 < \alpha_2 \implies F(\alpha_1) <_P F(\alpha_2)$ provided that $F(\alpha_1), F(\alpha_2) \in \lambda$, in which case $\text{im } F^{-1} = \text{OR}$ is a set by **Rep**, a contradiction.

Therefore, there is some smallest nonzero $\delta \in \text{OR}$ such that $F(\delta) = \lambda$. This implies that $\{f(F(\beta)) \mid \beta \in \delta\}$ is a chain in P . If δ is a limit, then P has no upper bound, a contradiction. Thus, $\delta = \eta + 1$ for some ordinal η . Suppose that $f(F(\eta))$ is not maximal in P . Then there is some least $\tau \in \lambda$ such that $f(F(\eta)) <_P f(\tau)$. But this means

$$\lambda = F(\delta) = \tau \in \lambda,$$

a contradiction. It follows that $f(F(\eta))$ is a maximal element. \square

Proposition 4.1.5. $\text{Zorn} \implies \text{WOP}$.

Corollary 4.1.6. $\text{AC} \iff \text{WOP} \iff \text{Zorn}$.

Definition 4.1.7. Let S be a set. Let F be a subset of $\mathbb{P}(S)$. We say that F is a *filter on S* if

- (i) $F \neq \emptyset$,
- (ii) $\emptyset \notin F$,
- (iii) if $X \in F$ and $X \subset Y$, then $Y \in F$, and
- (iv) $X, Y \in F \implies X \cap Y \in F$.

Example 4.1.8. The *Fréchet filter* on S is the set of all cofinite sets in S .

Definition 4.1.9. Let F be a filter on S .

1. We say that F is an *ultrafilter* if for any $X \in \mathbb{P}(S)$, either $X \in F$ or $(S \setminus X) \in F$.
2. If F is an ultrafilter, then we say that F is *nonprincipal* if it contains the Fréchet filter on S .

Remark 4.1.10. Let F be an ultrafilter on S . Define $\mu : \mathbb{P}(S) \rightarrow \{0, 1\}$ by

$$\mu(A) = \begin{cases} 1 & A \in F \\ 0 & A \notin F \end{cases}.$$

Then μ is a measure on S so long as we substitute the condition of finite additivity for that of countable additivity.

Definition 4.1.11. A subset $G \subset \mathbb{P}(S)$ has the *finite intersection property* (FIP) if

$$Z_1, \dots, Z_k \in G \implies Z_1 \cap \dots \cap Z_k \neq \emptyset.$$

Lemma 4.1.12. *If $G \subset \mathbb{P}(S)$ has FIP, then there is some filter H on S such that $H \supset G$.*

Proof. Take $H = \{W \subset S \mid \text{there exist a } k \in \mathbb{N} \text{ and } Z_1, \dots, Z_k \in G \text{ such that } Z_1 \cap \dots \cap Z_k \subset W\}$. \square

Theorem 4.1.13. *Every filter E on S can be extended to an ultrafilter.*

Proof. Let $\mathcal{F} = \{F \subset \mathbb{P}(S) \mid F \text{ is a filter on } S, F \supset E\}$. Then (\mathcal{F}, \subset) is a poset with $E \in \mathcal{F}$.

Claim. *A filter $F \subset \mathbb{P}(S)$ is an ultrafilter if and only if it is maximal in \mathcal{F} .*

Proof. It is clear that if F is an ultrafilter, then it is maximal. For the converse, suppose that F is not an ultrafilter. Then there is some $X \in \mathbb{P}(S)$ such that both $X \notin F$ and $(S \setminus X) \notin F$. We must show that F is not maximal in \mathcal{F} . Note that for any $Y \in F$, the intersection $X \cap Y$ is nonempty. Therefore, $F \cup \{X\}$ has FIP. By our least lemma, $F \cup \{X\}$ can be extended to a filter H on S . But $H \supsetneq F \supset E$, so that F is not maximal in \mathcal{F} . \square

Thus, it suffices to show that \mathcal{F} has some maximal element. Let C be a chain in \mathcal{F} . It is easy to check that $\bigcup C$ is an upper bound of C in \mathcal{F} . It follows from Zorn that \mathcal{F} has some maximal element. \square

Corollary 4.1.14. *There exists a nonprincipal ultrafilter.*

Definition 4.1.15. Let κ be an infinite cardinal.

1. A filter $F \subset \mathbb{P}(S)$ is κ -complete if every subset $X \subset F$ with $|X| < \kappa$ satisfies $\bigcap X \in F$.
2. We say that κ is *measurable* if $\kappa > \omega$ and there exists a nonprincipal κ -complete ultrafilter on κ .

Definition 4.1.16. A filter $F \subset \mathbb{P}(S)$ is σ -complete if for any countable set C of elements of F , we have $\bigcap C \in F$.

Proposition 4.1.17. *The least cardinal κ with a σ -complete nonprincipal ultrafilter is measurable.*

4.2 Lecture 12

Definition 4.2.1. Let α and β be nonzero limit ordinals.

1. Let $f : \alpha \rightarrow \beta$. We say that f is *cofinal in β* , written as $f : \alpha \xrightarrow{\text{cofinally}} \beta$, if $\sup(\text{im } f) = \beta$.
2. The *cofinality* $\text{cf}(\alpha)$ of α is the least ordinal γ for which there exists a cofinal map $f : \gamma \rightarrow \alpha$.

Example 4.2.2.

1. $\text{cf}(\aleph_\alpha) = \text{cf}(\alpha)$ for any limit ordinal $\alpha > 0$.
2. $\text{cf}(\omega) = \omega$.

Definition 4.2.3. Let κ be a cardinal.

- κ is *regular* if $\text{cf}(\kappa) = \kappa$.
- κ is *singular* if it is not regular.
- κ is a *limit (cardinal)* if $\lambda < \kappa \implies \lambda^+ < \kappa$.

- κ is a *strong limit (cardinal)* if $\lambda < \kappa \implies 2^\lambda < \kappa$.
- κ is *weakly inaccessible* if it is an uncountable regular limit.
- κ is (*strongly*) *inaccessible* if it is an uncountable regular strong limit.

Proposition 4.2.4. *If κ is measurable, then κ is strongly inaccessible and $|\{\lambda < \kappa \mid \lambda \text{ is inaccessible}\}| = \kappa$.*

Remark 4.2.5.

1. $\text{ZF} \not\models \aleph_1$ is regular. In fact, in ZF , we can prove that there is some $f : \omega \rightarrow \aleph_1$ such that $\sup(\text{im } f) = \aleph_1$, so that $\aleph_1 = \bigcup_{n \in \omega} f(n)$ even though each $f(n)$ is countable.
2. If κ is a strongly inaccessible cardinal, then $V_\kappa \models \text{ZFC}$.
3. Let $\text{WI}(\kappa)$ denote the statement that κ is a weakly inaccessible cardinal. It is known that

$$\text{ZFC} \not\models \exists \kappa (\text{WI}(\kappa)).$$

Moreover, if $\text{Con}(\text{ZFC} + \exists \kappa (\text{WI}(\kappa)))$ where $\text{Con}(T)$ means that the theory T is consistent, then $\text{Con}(\text{ZFC} + \text{WI}(2^{\aleph_0}))$.

Lemma 4.2.6. *For any X , we have $|\bigcup X| \leq \lambda \cdot \kappa$ where λ denotes $|X|$ and κ denotes $\sup\{|y| : y \in x\}$.*

Proof. We can write $X = \{X_\alpha \mid \alpha < \lambda\}$. For each $\alpha < \lambda$, we have that $\kappa_\alpha := |X_\alpha| \leq \kappa$. Therefore, we can also write $X_\alpha = \{x_{\alpha\beta} \mid \beta < \kappa_\alpha\}$. It follows that

$$\bigcup X = \{x_{\alpha\beta} \mid \alpha < \lambda, \beta < \kappa_\alpha\},$$

which has cardinality at most $|\lambda \times \kappa|$. □

Corollary 4.2.7. *A countable union of countable sets is countable.*

Proof. Apply Lemma 4.2.6 with $\lambda = \omega$ and $\kappa = \omega$. Since $\omega \times \omega$ is countable, we're done. □

Lemma 4.2.8. *$\text{cf}(\alpha)$ is a cardinal for any limit ordinal $\alpha > 0$.*

Proof. We can find a cofinal map $g : \text{cf}(\alpha) \rightarrow \alpha$. Suppose, towards a contradiction, that $\text{cf}(\alpha)$ is not a cardinal. Then there exist a $\delta \in \text{cf}(\alpha)$ and a bijection $f : \delta \rightarrow \text{cf}(\alpha)$. Then $\sup(\text{im } g \circ f) = \sup(\text{im } g) = \alpha$, so that $g \circ f$ is a cofinal map. This contradicts the minimality of $\text{cf}(\alpha)$. □

Theorem 4.2.9. *For any infinite cardinal κ , κ^+ is regular.*

Proof. Note that $\text{cf}(\alpha) \leq \alpha$ for any α since id_α is a cofinal map. Thus, it suffices to show $\text{cf}(\kappa^+) \geq \kappa^+$.

Claim. *If $\lambda \leq \kappa$, then $\text{cf}(\kappa^+) > \lambda$.*

Proof. Suppose, towards a contradiction, that $\text{cf}(\kappa^+) \leq \lambda$ for some $\lambda \leq \kappa$. Then there is some $\eta \leq \lambda$ such that

$$\kappa^+ = \bigcup \{\alpha_i \mid i < \eta\}$$

for some $\alpha_i \in \kappa^+$. Note that $|\alpha_i| \leq \kappa$ for each i and that $|\eta| \leq \kappa$. By Lemma 4.2.6, it follows $\kappa^+ \leq \kappa \cdot \kappa = \kappa$, a contradiction. □

But $\text{cf}(\kappa^+)$ is a cardinal by Lemma 4.2.8. Hence $\text{cf}(\kappa^+) \geq \kappa^+$. \square

Definition 4.2.10. Let I be a set. Consider any collection of sets $\{X_i\}_{i \in I}$ indexed by I . The *infinite sum* of the X_i is $\sum_{i \in I} X_i := \prod_{i \in I} X_i$. Let $\iota_i : X_i \rightarrow \sum_{i \in I} X_i$ denote the canonical inclusion.

Theorem 4.2.11 (König). If $|X_i| < |Y_i|$ for each $i \in I$, then $|\sum_{i \in I} X_i| < |\prod_{i \in I} Y_i|$.

*Proof.*⁴ We must show that $|\sum_{i \in I} X_i| \not\geq |\prod_{i \in I} Y_i|$. Suppose, to the contrary, that there is some surjection $f : \sum_{i \in I} X_i \rightarrow \prod_{i \in I} Y_i$. Since $|X_i| < |Y_i|$ for each $i \in I$ by hypothesis, the function

$$f_i := \pi_i \circ f \circ \iota_i : X_i \rightarrow Y_i$$

cannot be surjective where π_i denotes the i -th projection map. For each i , we can thus choose some y_i such that $y_i \notin \text{im } f_i$. Since f is surjective, there is some $x \in \sum_{i \in I} X_i$ such that $f(x) = (y_i)_{i \in I}$. But $x \in X_j$ for some $j \in I$, so that $y_j = f_j(x)$. This contradicts the fact that $y_j \notin \text{im } f_j$. \square

Corollary 4.2.12. For any infinite cardinal κ , $\kappa < \text{cf}(2^\kappa)$.

Proof. Suppose, towards a contradiction, that there exist an $\eta \leq \kappa$ and a sequence $\langle \alpha_i \mid i < \eta \rangle$ in 2^κ such that

$$\sup\{\alpha_i \mid i < \eta\} = 2^\kappa.$$

Then we get

$$\begin{aligned} 2^\kappa &= (2^\kappa)^\kappa \geq \left| \prod_{i < \eta} 2^\kappa \right| > \left| \sum_{i < \eta} \alpha_i \right| \\ &\geq |\sup\{\alpha_i \mid i < \eta\}| = 2^\kappa, \end{aligned}$$

a contradiction. \square

5 Relativization

5.1 Lecture 13

Lemma 5.1.1. For any ordinal α , V_α is transitive and $V_\beta \subset V_\alpha$ for every $\beta \leq \alpha$.

Proof. Note that if a set x is transitive, then $\mathbb{P}(x)$ is transitive and $x \subset \mathbb{P}(x)$. In light of this, it's easy to use transfinite induction to complete the proof. \square

Corollary 5.1.2. Let x be a set. If every $y \in x$ belongs to V , then x belongs to V .

Proof. Using Lemma 5.1.1, we have that $\bigcup x \in V$. Thus, $\bigcup x \subset V_\beta$ for some $\beta \in \text{OR}$. This implies that $y \in V_\beta$ for each $y \in x$. Hence $x \subset \mathbb{P}(V_\beta) = V_{\beta+1}$, so that $x \in \mathbb{P}(V_{\beta+1}) = V_{\beta+2}$. \square

Definition 5.1.3. Let x be a set. The *transitive closure* $\text{TC}(x)$ of x is defined recursively by

$$\begin{aligned} x_0 &= x \\ x_{n+1} &= \bigcup x_n \\ \text{TC}(x) &= \bigcup_{n \in \omega} x_n. \end{aligned}$$

⁴The format of this proof is taken from *nLab*'s article "König's theorem" (as of 05/19).

Note 5.1.4. For any x , $\text{TC}(x)$ is the smallest transitive set containing x , i.e.,

$$\text{TC}(x) = \bigcap \{y \supset x \mid y \text{ is a transitive set}\}.$$

Lemma 5.1.5 (Regularity for classes). *Let C be a nonempty class. Then there is some $x \in C$ such that if $y \in C$, then $y \notin x$. In other words, C has some \in -minimal element.*

Proof. By assumption, there is some $u \in C$. Let $e = \{x \in C \mid x \in \text{TC}(u)\}$, which is a set by **Sep**. If $e = \emptyset$, then there is no element of $C \cap u$, in which case u is minimal in C . If $e \neq \emptyset$, then there is some \in -minimal element m of e . In this case, no element n of m can belong to C . For otherwise $n \in m \subset \text{TC}(u)$ and $n \in C$, so that $n \in e$, contrary to our choice of m . It follows that m is minimal in C . \square

Definition 5.1.6. The *rank* $\text{rank}(x)$ of a set x is the least ordinal α such that $x \in V_{\alpha+1}$.

Lemma 5.1.7. *In ZF, every set is in V . In other words, $\text{ZF} \vdash \forall x(V(x))$.*

Proof. Suppose, towards a contradiction, that there is some z such that $\neg V(z)$. By Lemma 5.1.5, there is some \in -minimal such set m . Then we have a functional relation $\text{rank} : m \rightarrow \text{OR}$. Hence im rank is a set, and we can form the set $\beta := \sup(\text{im rank})$, which is an ordinal. If $u \in m$, then $\text{rank}(u) \leq \beta$, so that $u \in V_{\text{rank}(u)+1} \subset V_{\beta+1}$ by Lemma 5.1.1. This shows that $m \subset V_{\beta+1}$, and thus $m \in \mathbb{P}(V_{\beta+1}) = V_{\beta+2}$. But this implies that $V(m)$, a contradiction. \square

Corollary 5.1.8. *A class c is a set if and only if it is bounded in rank, i.e., there is some $\beta \in \text{OR}$ such that for any $y \in c$, $\text{rank}(y) \leq \beta$.*

Notation.

- For any language \mathcal{L} , we shall write \bar{x} for a finite sequence of symbols (x_1, \dots, x_k) in \mathcal{L} .
- For any set A , a symbol of the form \bar{a} will refer to a finite sequence (a_1, \dots, a_k) in A . Abusing notation, we shall write $\bar{a} \in A$ to express that the components of \bar{a} belong to A . Also, we shall write $f(\bar{a})$ for the sequence $(f(a_1), \dots, f(a_k))$.

Definition 5.1.9 (Relativization). Let (M, \in_M) consist of a class M and the binary relation \in on M (i.e., $x \in_M y \iff x \in y$). In this case, we call M a *standard model (for the language of set theory)*. Let $\varphi(\bar{x})$ denote a formula of the language of set theory. The *relativization* $\varphi^M(\bar{x})$ of φ to M is the formula obtained inductively by

- fixing each atomic formula appearing in φ ,
- preserving each logical connective appearing in φ ,
- translating each formula of the form $\exists x \psi$ appearing in φ where ψ is quantifier-free to $(\exists x \in M) \psi^M$, and
- similarly translating $\forall x \psi$ to $(\forall x \in M) \psi^M$.

Part (b) says that the operator $(-)^M$ commutes with all logical connectives.

Remark 5.1.10. For any particular $\varphi(\bar{x})$, the expression

$$M \models \varphi(\bar{x})$$

(pronounced “ M satisfies φ ”) will mean the same thing as $\varphi^M(\bar{x})$. Note, however, that unless M is a set, we cannot formally define such a satisfaction relation \models in ZF. Indeed, due to Tarski’s undefinability theorem, if the existence of a satisfaction relation for (V, \in) were provable in ZF, then ZF could prove its own consistency, which is impossible.

Definition 5.1.11. A formula $\theta(\bar{x})$ of the language of set theory is a Δ_0 -formula (written as $\theta(\bar{x}) \in \Delta_0$) if

- (a) it is quantifier-free, or
- (b) it is of the form $\varphi \square \psi$ or $\neg \varphi$ where $\varphi, \psi \in \Delta_0$ and \square denotes a logical connective, or
- (c) it is of the form $(\exists x \in y)\varphi$ or $(\forall x \in y)\varphi$ where $\varphi \in \Delta_0$.

Lemma 5.1.12. *Let \mathfrak{U} be a standard model. TFAE.*

- (i) *For any $\varphi(\bar{x}) \in \Delta_0$ and any $\bar{v} \in \mathfrak{U}$, $\mathfrak{U} \models \varphi[\bar{v}]$ iff $\varphi[\bar{v}]$.*
- (ii) *\mathfrak{U} is transitive.*

Proof.

(ii) \implies (i):

We proceed by induction on the complexity of φ . Our equivalence holds immediately when φ is an atomic formula. For our induction step, it suffices to consider just the logical symbols \neg , \wedge , and \exists . In either of the first two cases, our equivalence holds immediately by induction. Finally, let

$$\varphi(\bar{x}) = (\exists u \in y)\psi(u, y, \bar{z})$$

and suppose that our equivalence is true of $\psi(u, y, \bar{z})$. Suppose that $\mathfrak{U} \models \varphi(\bar{x})[\bar{v}]$. Then we have

$$(\exists u \in \mathfrak{U})(u \in y \wedge \psi^{\mathfrak{U}}(u, y, \bar{z}))[\bar{v}].$$

By our IH, it follows that $(\exists u \in \mathfrak{U})(u \in y \wedge \psi(u, y, \bar{z}))[\bar{v}]$, which implies that $(\exists u \in y)\psi(u, y, \bar{z})[\bar{v}]$, as desired. Conversely, suppose $\varphi(\bar{x})[\bar{v}]$ with witness u . Then $u \in y[\bar{v}]$. Note that y is free in φ , and thus $y[\bar{v}]$ belongs to \mathfrak{U} by hypothesis. Since \mathfrak{U} is transitive, it follows that $u \in \mathfrak{U}$. Hence u is a witness for $\varphi^{\mathfrak{U}}$ as well.

\neg (ii) \implies \neg (i):

Since \mathfrak{U} is not transitive, we can find some $y \in \mathfrak{U}$ and some $u \in y$ such that $u \notin \mathfrak{U}$. Consider the formula

$$\psi(t) := (\exists b \in t)(\forall c \in t)c = b,$$

which expresses that t is a singleton. We see that $\psi \in \Delta_0$. But if $\psi(y)$, then $\mathfrak{U} \models y = \emptyset$, so that $\mathfrak{U} \not\models \psi(y)$. This shows that $\mathfrak{U} \models \psi[\bar{v}]$ is *not* equivalent to $\psi[\bar{v}]$. \square

Example 5.1.13. Let κ be a regular cardinal. Any element of the set

$$\text{HC}(\kappa) := \{x : |\text{TC}(x)| < \kappa\}$$

is said to have *hereditary cardinality* $< \kappa$. Then $x \in \text{HC}(\kappa)$ if and only if x has cardinality $< \kappa$, every element of x has cardinality $< \kappa$, every element of every element of x has cardinality $< \kappa$, and so on. Note that $(\text{HC}(\kappa), \in)$ is a transitive model for the language of set theory. One can show that $\text{HC}(\omega) = V_\omega$ and that

$$\text{HC}(\kappa) = V_\kappa \quad (\star)$$

when κ is inaccessible.

Theorem 5.1.14. $\text{Con}(\text{ZFC} - \text{Reg}) \implies \text{Con}(\text{ZFC})$.

Proof sketch. It suffices to verify that $\text{ZFC} - \text{Reg} \vdash (\text{ZFC})^V \vdash (\emptyset \neq \emptyset)^V \leftrightarrow (\emptyset \neq \emptyset)$. □

5.2 Lecture 14

Notation.

- Let Sl denote the first-order assertion that there exists a strongly inaccessible cardinal.
- Let $\text{Sl}(x)$ denote the first-order assertion that x is a strongly inaccessible cardinal.

Lemma 5.2.1. *Suppose that $\text{Sl}(\kappa)$. Then $|V_\alpha| < \kappa$ for every $\alpha < \kappa$.*

Proof. Let us induct on α . The base case is obvious. Suppose that $|V_\alpha| < \kappa$. Then

$$|V_{\alpha+1}| = |\mathbb{P}(V_\alpha)| = 2^{|V_\alpha|} < \kappa$$

since κ is a strong limit. Next, suppose that $\lambda < \kappa$ is a limit ordinal such that $|V_\beta| < \kappa$ for each $\beta < \lambda$. As the supremum of a set of cardinals is again a cardinal, we have that

$$|V_\lambda| = \left| \bigcup_{\beta < \lambda} V_\beta \right| = \sup_{\beta < \lambda} |V_\beta|.$$

If this equals κ , then we have a cofinal map $\lambda \rightarrow \kappa$, contrary to the fact that κ is regular. Hence $|V_\lambda| < \kappa$, as required. □

Proposition 5.2.2. $\text{Sl}(x) \iff V_\kappa \models \text{Sl}(x)$.

Theorem 5.2.3. *Suppose that κ is the least λ such that $\text{Sl}(\lambda)$. Then*

- (a) $V_\kappa \models \text{ZFC}$, and
- (b) $V_\kappa \models \neg \text{Sl}$.

Proof.

- (a) It is easy to see that $V_\kappa \models \text{Z} + \text{Reg}$. It remains to check that V_κ satisfies both Rep and AC .

- **Rep** : Suppose that $X \in V_\kappa$ and F is a function $X \rightarrow V_\kappa$. We must show that $F(X) \in V_\kappa$. As κ is a limit, we have that $X \in V_\alpha$ for some $\alpha < \kappa$. By Lemma 5.2.1, it follows that

$$|X| \leq |V_\alpha| < \kappa.$$

Let $\beta = \sup\{\text{rank}(F(y)) \mid y \in X\}$, so that $F(X) \in V_{\beta+1}$. If $\beta = \kappa$, then the map $X \rightarrow \kappa$ defined by $y \mapsto \text{rank}(F(y))$ is cofinal. But this is impossible, because κ is regular. Hence $\beta < \kappa$, and thus $F(X) \in V_\kappa$.

- **AC** : Suppose that $X \in V_\kappa$ with each element of X nonempty. There exists a choice function $F : X \rightarrow V$. Note that F consists of ordered pairs (x, y) where $x \in X$ and $y \in x$. Both of these belong to V_α where $\alpha := \text{rank}(X) + 1 < \kappa$, and thus (x, y) belongs to $V_{\alpha+1}$. This implies that $\text{rank}(F) \leq \text{rank}(X) + 2 < \kappa$ because κ is a limit. This proves that $F \in V_\kappa$.

- (b) Let α be any ordinal belonging to V_κ . By Proposition 5.2.2, we see that $V_\kappa \models \text{SI}(\alpha)$ if and only if $\text{SI}(\alpha)$. But $\alpha < \kappa$, and thus $\neg \text{SI}(\alpha)$ holds by our choice of κ . This means that $V_\kappa \not\models \text{SI}(\alpha)$. As α was arbitrary, it follows that $V_\kappa \models \neg \text{SI}$.

□

Corollary 5.2.4. $\text{HC}(\kappa) \models \text{ZFC}$ in light of (\star) .

Corollary 5.2.5. $\text{ZFC} \not\models \text{SI}$.

Proof. If there exists an inaccessible cardinal, then we can find a model of $\text{ZFC} + \neg \text{SI}$ via Theorem 5.2.3. Otherwise, we can take V to be a model of $\text{ZFC} + \neg \text{SI}$. □

Remark 5.2.6. In light of part (a) of Theorem 5.2.3, we have another way of proving Corollary 5.2.5. Indeed, if $\text{ZFC} \models \text{SI}$, then $\text{ZFC} \models (V_\kappa \models \text{ZFC})$ where κ denotes the minimal strongly inaccessible cardinal according to ZFC. Such a situation, however, contradicts Gödel's second incompleteness theorem (provided that ZFC is consistent).

Corollary 5.2.7. $\text{Con}(\text{ZFC}) \implies \text{Con}(\text{ZFC} + \neg \text{SI})$.

Proof. Suppose that $\text{Con}(\text{ZFC})$, so that there is some set model \mathfrak{A} of ZFC. If there is no inaccessible cardinal in \mathfrak{A} , then just take \mathfrak{A} to be a model of $\text{ZFC} + \neg \text{SI}$. Otherwise, let κ be the least such cardinal. Then it is provable that $V_\kappa \models \text{ZFC} + \neg \text{SI}$. In either case, $\text{ZFC} + \neg \text{SI}$ is consistent because it has a set model. □

5.3 Lecture 15

Definition 5.3.1. Let \mathfrak{A} and \mathfrak{B} be two models for a given language \mathcal{L} .

1. We say that \mathfrak{B} is an *elementary submodel* of \mathfrak{A} (written as $\mathfrak{B} \preceq \mathfrak{A}$) if \mathfrak{B} is a submodel of \mathfrak{A} such that for any first-order formula $\varphi(\bar{x})$ of \mathcal{L} and any $\bar{b} \in \mathfrak{B}$,

$$\mathfrak{A} \models \varphi[\bar{b}] \iff \mathfrak{B} \models \varphi[\bar{b}].$$

2. An embedding $f : \text{dom}(A) \rightarrow \text{dom}(B)$ (i.e., an isomorphism onto its image) is an *elementary embedding* if $f(\mathfrak{A}) \preceq \mathfrak{B}$. Equivalently, f is an elementary embedding if

$$\mathfrak{A} \models \varphi[\bar{a}] \iff \mathfrak{B} \models \varphi[f(\bar{a})]$$

for any first-order formula $\varphi(\bar{x})$ of \mathcal{L} and any $\bar{a} \in \mathfrak{A}$.

Lemma 5.3.2 (Tarski-Vaught elementary submodel criterion). *Let \mathfrak{A} be a model for \mathcal{L} such that its domain A is a set. (Such an \mathfrak{A} is sometimes called an \mathcal{L} -structure.) Let $B \subset A$. Suppose that for every formula of the form*

$$\exists y \theta(x_1, \dots, x_n, y)$$

and every $b_1, \dots, b_n \in B$, if $\mathfrak{A} \models \exists y \theta[b_1, \dots, b_n]$, then there is some $b \in B$ such that $\mathfrak{A} \models \theta[b_1, \dots, b_n, b]$. Then B forms an elementary submodel of \mathfrak{A} .

Theorem 5.3.3 ((Downward) Löwenheim-Skolem (L-S)). *Let \mathcal{L} be a countable language. For any infinite set model \mathfrak{A} for \mathcal{L} and any set $X \subset \text{dom}(\mathfrak{A})$, we can use AC to find a set model \mathfrak{B} for \mathcal{L} such that*

1. $X \subset \text{dom}(\mathfrak{B})$,
2. $\mathfrak{B} \preceq \mathfrak{A}$, and
3. $\text{dom}(\mathfrak{B})$ has cardinality at most $|X| \cdot |\mathcal{L}| \cdot \aleph_0$.

In particular, there is a countable \mathcal{L} -structure \mathfrak{B} such that $\mathfrak{B} \preceq \mathfrak{A}$.

Lemma 5.3.4 (Mostowski collapse). *Consider a class model (A, E^A) of ZF, where E^A is a (strict) binary relation interpreting \in . If E^A is both well-founded and extensional, then there is some transitive class model (B, \in_B) such that $A \cong B$.*

Proof. Suppose that E^A is both well-founded and extensional. We can extend transfinite induction (and thus recursion) on OR to any well-founded class. Thus, we can define $f : A \rightarrow V$ by

$$f(a) = \{f(b) \mid b \in A \wedge bE^A a\}.$$

Let $x \in f(A)$, so that $x = \{f(y) \mid y \in A \wedge yE^A z\}$ for some $z \in A$. Hence any element of x belongs to the image of f , which shows that $f(A)$ is a transitive class.

Now, we want to show that f is an embedding. Let $a, b \in A$ with $a \neq b$. Suppose, towards a contradiction, that the class

$$D := \{z \in f(A) \mid (\exists a, b \in A)(a \neq b \wedge z = f(a) = f(b))\}$$

is nonempty. Choose $m \in D$ of least rank and let a and b witness that $m \in D$. Since $a \neq b$, we have $\text{ext}_{E^A}(a) \neq \text{ext}_{E^A}(b)$. Assume, wlog, that there is some $u \in \text{ext}_{E^A}(a) \setminus \text{ext}_{E^A}(b)$. Then

$$f(u) \in f(a) = m = f(b),$$

and thus there is some $v \in \text{ext}_{E^A}(b)$ such that $s := f(u) = f(v)$. Since $u \neq v$, we have $s \in D$. But $s \in m$, so that $\text{rank}(s) < \text{rank}(m)$, a contradiction. It follows that f is injective.

It remains to show that $(x, y) \in E^A \iff f(x) \in f(y)$. The forward direction is obvious. For the reverse direction, suppose that $f(x) \in f(y)$. Then $f(x) = f(c)$ for some $cE^A y$. Since f is injective, we have $x = c$. Thus, $(x, y) \in E^A$. \square

Terminology. We call (B, E^B) from Lemma 5.3.4 the *transitive collapse* of A .

Proposition 5.3.5. *Provided that $\text{Con}(\text{ZF})$, there exists a set model of ZF that is not well-founded.*

Proof. In general, as a result of the compactness theorem, if an \mathcal{L} -structure (A, E^A) has a chain

$$x_1 E^A x_2 E^A \dots E^A x_n$$

for each $n \in \mathbb{N}$, then there is some \mathcal{L} -structure (B, E^B) such that $\underbrace{A \equiv B}_{\text{elem. equiv.}}$ but E^B is not well-founded. \square

Theorem 5.3.6 (Skolem's paradox). *Let $\varphi(x)$ denote the formula*

$$\exists f(f : \omega \rightarrow x \wedge f \text{ is a bijection}),$$

where ω denotes the smallest infinite ordinal. Provided that ZF has at least one well-founded set model \mathfrak{A} ,⁵ there is a transitive set model \mathfrak{B} of ZF such that $\mathfrak{B} \not\models \varphi$ but φ holds in the universe of all sets.

Proof. By the L-S theorem, there is some countable elementary submodel \mathfrak{A}' of \mathfrak{A} . Note that \mathfrak{A}' is both extensional and well-founded as a submodel of \mathfrak{A} . By Mostowski, we have that $\mathfrak{A}' \cong \mathfrak{B}$ for some transitive model \mathfrak{B} . Then \mathfrak{B} is a countable transitive model of ZF (written as $\text{ctm}(\mathfrak{B})$). Since $\mathfrak{B} \models \text{ZF}$, we can take the Hartogs number $H(\omega)$ to get a cardinal $\kappa^{\mathfrak{B}} \in \text{dom}(\mathfrak{B})$ with $\kappa^{\mathfrak{B}} > \omega$. Therefore,

$$\mathfrak{B} \models \neg\varphi(\kappa^{\mathfrak{B}}).$$

That is, $\kappa^{\mathfrak{B}}$ is not countable inside \mathfrak{B} . But since \mathfrak{B} is transitive, we get $\kappa^{\mathfrak{B}} \subset \text{dom}(\mathfrak{B})$, which implies that $\kappa^{\mathfrak{B}}$ is countable outside \mathfrak{B} . That is, $\varphi(\kappa^{\mathfrak{B}})$ holds in the universe of all sets. \square

5.4 Lecture 16

Definition 5.4.1 (Absoluteness). Let $\varphi(\bar{x})$ be a formula of set theory. Suppose that B is a class and that $A \subset B$. We say that φ is *absolute for A and B* , written as $\text{Abs}(A, B, \varphi)$, if for every $\bar{a} \in A$, we have

$$\varphi^A[\bar{a}] \leftrightarrow \varphi^B[\bar{a}].$$

When B denotes the universe of all sets, we say that φ is *absolute for A* .

Example 5.4.2. Any quantifier-free formula is absolute for any class A .

Note 5.4.3. Let M be a transitive class. Lemma 5.1.12 states that any Δ_0 -formula is absolute for M .

Example 5.4.4. The following expressions can be written as Δ_0 -formulas and thus are absolute for all transitive models.

1. Being an ordinal.
2. Being a limit/successor ordinal.
3. “ $x = \omega$ ”.
4. Being a function.
5. Being an injection/surjection.

Example 5.4.5. Proposition 5.2.2 states that $\text{SI}(x)$ is absolute for V_κ when κ is strongly inaccessible.

Non-example 5.4.6.

1. Skolem's paradox states that if there exists a well-founded set model of ZF, then there exists a transitive model M such that being countable is not absolute for M .

⁵This assumption is strictly stronger than the assumption that $\text{Con}(\text{ZF})$ but is weaker than the assumption that $\exists \kappa(\text{SI}(\kappa))$.

2. Being a cardinal.⁶

The following is, in essence, a re-wording of Lemma 5.3.2.

Lemma 5.4.7 (Tarski-Vaught (alternative formulation)). *Suppose that $A \subset B$ and that $\varphi(\bar{x}, y)$ is a formula such that $\text{Abs}(A, B, \varphi(\bar{x}, y))$. Further, suppose that for any $\bar{a} \in A$, if there exists $b \in B$ such that $\varphi^B[\bar{a}, b]$, then there exists $b' \in A$ such that $\varphi^B[\bar{a}, b']$. Then $\text{Abs}(A, B, \exists y \varphi(\bar{x}, y))$.*

Proof. By induction on the complexity φ , let us show that

$$A \models \exists y \varphi(\bar{a}, y) \iff B \models \exists y \varphi(\bar{a}, y)$$

for every $\bar{a} \in A$. The case where φ is atomic is obvious, as is the case where $\varphi = \varphi_1 \square \varphi_2$ where $\square \in \{\neg, \wedge\}$. It remains to consider the case where $\varphi = \exists z \psi$.

In this case, suppose that $A \models \psi[\bar{a}, c, c']$ for some $c, c' \in A$. By our induction hypothesis, B also must satisfy $\psi[\bar{a}, c, c']$. This means that $B \models \exists y \varphi(\bar{a}, y)$, as required. Conversely, suppose that $B \models \psi[\bar{a}, d, d']$ for some $d, d' \in B$. Then $B \models \varphi[\bar{a}, d']$. By hypothesis, there is some $e \in A$ such that $B \models \varphi[\bar{a}, e]$. By our induction hypothesis, we have that $A \models \varphi[\bar{a}, \tilde{e}]$ for some $\tilde{e} \in A$. Thus, $A \models \exists y \varphi(\bar{a}, y)$, as required. \square

Theorem 5.4.8 (Reflection principle). *For any set M and any formulas $\varphi_1, \dots, \varphi_k$, there exists a set $M' \supset M$ such that $\text{Abs}(M', V, \varphi_i)$ for each $1 \leq i \leq k$.*

5.5 Lecture 17

Our next result is a strong version of Theorem 5.4.8 and is provable in ZF. For simplicity, we shall state it under the assumption that $k = 1$.

Theorem 5.5.1. *For every formula $\varphi(\bar{x})$ and every ordinal α , there is some limit ordinal $\beta > \alpha$ such that for any $\bar{b} \in V_\beta$, we have $\varphi^{V_\beta}[\bar{b}] \iff \varphi[\bar{b}]$.*

Proof. Let us induct on the complexity of φ . The case where φ is atomic is obvious, as is the case where $\varphi = \varphi_1 \square \varphi_2$ where $\square \in \{\neg, \wedge\}$. It remains to consider the case where $\varphi = \exists z \psi(z, x_1, \dots, x_n)$. For each n -tuple \bar{w} , let

$$F(\bar{w}) = \{s \mid \psi(s, \bar{w}) \wedge \forall t (\psi(t, \bar{w}) \rightarrow \text{rank}(s) \leq \text{rank}(t))\}.$$

Now, let us construct a sequence $\{G_i\}_{i \in \omega}$ of sets inductively as follows.

$$\begin{aligned} G(0) &:= V_\alpha \\ G(i+1) &:= G(i) \cup \bigcup_{\bar{w} \in G(i)} F(\bar{w}) \end{aligned} \tag{*}$$

Let α_i denote the least limit ordinal $\delta > \alpha$ such that $G(i) \subset V_\delta$. We have that

$$\bigcup_{i \in \omega} V_{\alpha_i} = V_{\sup_{i \in \omega} \{\alpha_i\}},$$

⁶According to http://cantorsattic.info/Hereditary_Cardinality, for any large regular λ , the property of being a cardinal is absolute for $\text{HC}(\lambda)$.

where $\sup_{i \in \omega} \{\alpha_i\}$ must be a limit ordinal greater than α , which we take to be β . Note that for any $\bar{b} \in V_\beta$, there is some $m \in \mathbb{N}$ such that $b_1, \dots, b_n \in V_{\alpha_m}$. If there exists a witness of $\varphi[\bar{b}]$ in V , then there exists one in $V_{\alpha_{m+1}} \subset V_\beta$ in light of (*). It follows that

$$\begin{aligned} V_\beta \models \varphi[\bar{b}] &\leftrightarrow (\exists z \in V_\beta) \psi^{V_\beta}[\bar{b}] \\ &\leftrightarrow (\exists z \in V_\beta) \psi[\bar{b}] \\ &\leftrightarrow \exists z \psi[\bar{b}], \end{aligned} \tag{IH}$$

as desired. □

Corollary 5.5.2. *ZF is not finitely axiomatizable.*

Proof. Suppose that there is some formula φ such that $\text{Con}(\mathbf{Z} + \varphi)$. Suppose, towards a contradiction, that $\mathbf{Z} + \varphi \models \text{ZF}$. In light of Theorem 5.5.1, take γ as the least limit ordinal $\beta > \omega$ such that φ is absolute for V_β . Note that Theorem 5.5.1 is provable in $\mathbf{Z} + \varphi$ and that $V_\gamma \vdash \mathbf{Z}$ since γ is an uncountable limit ordinal. Therefore, we have that

$$V_\gamma \models \exists \alpha (\alpha \text{ is an uncountable limit ordinal} \wedge (\varphi \leftrightarrow \varphi^{V_\alpha})).$$

Claim. *Being uncountable is absolute for V_γ .*

Proof. Let $x \in V_\gamma$. We must show that x is uncountable iff $V_\gamma \models “x \text{ is uncountable}”$. Suppose that x is uncountable, i.e., there is no injection $x \rightarrow \omega$. Due to Example 5.4.4, this means that there is no such injection in V_γ .

Conversely, suppose that there is no injection $x \rightarrow \omega$ in V_γ , so that $\gamma > \omega$. Suppose, however, that there is such an injection f in V . Note that $f \subset x \times \omega \subset \mathbb{P}(\mathbb{P}(x \cup \omega))$, and thus $\text{rank}(f) \leq \max\{\text{rank}(x), \omega\} + 2$. Hence $\text{rank}(f) \leq \alpha + 2$ for some $\alpha < \gamma$ as γ is a limit ordinal. This implies that $\text{rank}(f) < \gamma$, i.e., $f \in V_\gamma$, a contradiction. □

Moreover, “ $x = V_\alpha$ ” is absolute for V_γ . It follows that there is some uncountable limit ordinal $\alpha < \gamma$ such that φ is absolute for V_α , contrary to our choice of γ . □

The following notion generalizes the rank hierarchy V .

Definition 5.5.3. A *cumulative hierarchy* is a sequence of sets $\{W_\alpha\}_{\alpha \in \text{OR}}$ such that

- $W_0 = \emptyset$,
- $W_\alpha \subset W_{\alpha+1} \subset \mathbb{P}(W_\alpha)$, and
- $W_\lambda = \bigcup_{\beta < \lambda} W_\beta$ for limit λ .

Note that each stage W_α is transitive and satisfies $W_\alpha \subset V_\alpha$.

Remark 5.5.4. Our proof of Theorem 5.5.1 is easily adapted to *any* cumulative hierarchy.

6 Gödel's hierarchy of constructible sets

6.1 Lecture 18

Let (A, E^A) be a set model. Recall that a set x is *definable with parameters* over an \mathcal{L} -structure (A, E^A) if there exist a formula $\varphi(y, \bar{z})$ of \mathcal{L} and a finite sequence $\bar{b} \in A$ such that

$$x = \{a \in A \mid A \models \varphi[a, \bar{b}]\}.$$

Let $\text{Def}(A)$ denote the set consisting of all sets definable with parameters over (A, E^A) .

Note 6.1.1. Let $\kappa = |A|$. We have that

$$|\text{Def}(A)| \leq \left| \bigcup_{n \in \omega} A^n \right| = \kappa < 2^\kappa.$$

Therefore, $\text{Def}(A) \subsetneq \mathbb{P}(A)$.

The *generalized continuum hypothesis* (GCH) asserts that $2^{\aleph_\alpha} = \aleph_{\alpha+1}$ for every ordinal α . In order to prove that $\text{ZF} + \text{GCH}$ is consistent, Gödel defined a certain cumulative hierarchy L , called the *constructible universe*, as follows.

$$\begin{aligned} L_0 &= \emptyset \\ L_{\alpha+1} &= \text{Def}(L_\alpha) \\ L_\lambda &= \bigcup_{\beta < \lambda} L_\beta, \quad \lambda \text{ limit.} \end{aligned}$$

Note 6.1.2.

1. Any ordinal α is an element of $L_{\alpha+1}$.
2. $|L_{\aleph_1} \cap \mathbb{P}(\omega)| = \aleph_1$.

Notation. For any $x \in L$, let $\text{order}(x)$ denote the least ordinal α such that $x \in L_{\alpha+1}$.

In 1939, Gödel proved that $\text{ZF} \vdash (\text{ZF})^L + (V = L)^L$, which means that if ZF is consistent, then so is $\text{ZF} + V = L$. The assertion “ $V = L$ ” is known as the *axiom of constructibility*.

In order to prove this relative consistency result, we should introduce a classification of first-order definable properties that expands Definition 5.1.11. Define the *Lévy hierarchy* by mutual recursion as follows.

- A formula is Σ_0 if it is Δ_0 .
- A formula is Π_0 if it is Δ_0 .
- A formula is Σ_{n+1} if it has the form $\exists x \varphi$ where φ is Π_n .
- A formula is Π_{n+1} if it has the form $\forall x \varphi$ where φ is Σ_n .
- A formula is Δ_n if it both Σ_n and Π_n .

Let T be a theory in the language of set theory. We say that a first-order property P of \bar{x} is Σ_n^T (resp. Π_n^T) if the formula $P(\bar{x}) \leftrightarrow \varphi(\bar{x})$ is provable in T for some Σ_n (resp. Π_n) formula φ .

Note 6.1.3. If $\varphi(\bar{x})$ is Δ_1 , then $\varphi(\bar{x})$ is absolute for all transitive models. This follows easily from the fact that any Δ_0 formula is absolute for all transitive models.

Example 6.1.4. “ R is a well-founded relation on A ” is a Δ_1^{ZF} property.

Proof. Consider the formulas

$$\varphi := \forall X \left(\overbrace{X \subset A \wedge X \neq \emptyset \rightarrow (\exists y \in X) (\forall z \in A) (\neg Rzy)}^{\tau(R, A, X)} \right)$$

$$\psi := \exists f (f \text{ is a function} \wedge \text{dom}(f) = A \wedge \text{range}(f) \in \text{OR} \wedge (\forall x, y \in A) (Rxy \rightarrow f(x) < f(y))) .$$

Note that τ is Δ_0 and that “ R is a relation on A ” is a Δ_0^{ZF} property $\eta(R, A)$, and thus $\eta \wedge \varphi$ is Π_1 . Moreover, $\eta \wedge \psi$ is Σ_1 . But both of these formulas express that R is a well-founded relation on A , which must be Δ_1^{ZF} as a result. \square

6.2 Lecture 19

6.3 Lecture 20

6.4 Lecture 21

6.5 Lecture 22

6.6 Lecture 23

6.7 Lecture 24

6.8 Lecture 25

6.9 Lecture 26

6.10 Lecture 27

6.11 Lecture 28

6.12 Lecture 29

6.13 Lecture 30

6.14 Lecture 31

6.15 Lecture 32

7 Large cardinals

7.1 Lecture 33

7.2 Lecture 34

7.3 Lecture 35

7.4 Lecture 36

7.5 Lecture 37

7.6 Lecture 38

7.7 Lecture 39

7.8 Lecture 40

Remark 7.8.1. Cohen proved that

$$\text{Con}(\text{ZF}) \implies \text{Con}(\text{ZFC} + \neg\text{CH}),$$

thereby proving that CH (as well as GCH) is independent of ZFC. In fact, assuming that ZF is consistent, we have not only that $\text{ZFC} + (2^{\aleph_0} = \aleph_{\alpha+1})$ is consistent for any α but also that $\text{ZFC} + (2^{\aleph_0} = \aleph_\lambda)$ is consistent for any λ with $\text{cf}(\lambda) > \omega$.

7.9 Lecture 41

Discussion of Homework 3 solutions.

7.10 Lecture 42

Discussion of Homework 3 solutions.

7.11 Lecture 43

Discussion of Homework 3 solutions.