

Endotyping asthma: new insights into key pathogenic mechanisms in a complex, heterogeneous disease

Gary P Anderson

Clinical asthma is very widely assumed to be the net result of excessive inflammation driven by aberrant T-helper-2 (Th2) immunity that leads to inflamed, remodelled airways and then functional derangement that, in turn, causes symptoms. This notion of disease is actually poorly supported by data, and there are substantial discrepancies and very poor correlation between inflammation, damage, functional impairment, and degree of symptoms. Furthermore, this problem is compounded by the poor understanding of the heterogeneity of clinical disease. Failure to recognise and discover the underlying mechanisms of these major variants or endotypes of asthma is, arguably, the major intellectual limitation to progress at present. Fortunately, both clinical research and animal models are very well suited to dissecting the cellular and molecular basis of disease endotypes. This approach is already suggesting entirely novel pathways to disease—eg, alternative macrophage specification, steroid refractory innate immunity, the interleukin-17–regulatory T-cell axis, epidermal growth factor receptor co-amplification, and Th2-mimicking but non-T-cell, interleukins 18 and 33 dependent processes that can offer unexpected therapeutic opportunities for specific patient endotypes.

Introduction

In recent years, the morbidity and mortality of asthma have decreased, probably as a result of improved management. Some evidence suggests that the relentless rise in disease incidence and prevalence is now also reaching a plateau.¹ However, although contemporary treatment approaches are indisputably effective, many patients have substantial residual disease and some, with very severe asthma, respond suboptimally even to high-dose oral steroids (figure 1).^{2–6} Furthermore, asthma—the most common serious chronic lung disease afflicting around 150 million people worldwide—remains both unpreventable and incurable.⁷ Despite decades of intensive research, little progress in identification of new treatments has been made since the introduction of inhaled β_2 adrenoceptor selective agonists (1969) and inhaled glucocorticosteroids (1974).




Figure 1: Residual disease burden in optimally treated asthma

The diagram shows that as severity increases (purple), the number of afflicted individuals decreases (represented as an inverse triangle). Exacerbations are represented as a pie chart. Severe asthma (red triangle), which is defined as failure of optimised treatment, is rare and estimated at about 5% of all disease. Notional best possible treatment (green) shows (1) residual disease in patients with moderate disease, (2) refractory severe asthma, and (3) breakthrough acute exacerbations. The purple and red regions represent the targets for future therapies. Th2-directed therapies will most probably compete with present therapies for mild asthma.

This Review aims to advance the argument that the way in which we think about the pathogenesis of asthma is flawed (or incomplete), which in turn is preventing the discovery of better treatments, preventions, and cures. Clear evidence now suggests that asthma is a heterogeneous and genetically complex disease (>100 genes have already been implicated) that cannot be explained by one mechanism alone. To order this heterogeneity and the volume and complexity of clinical and basic research data, the new notion of disease endotypes (panel 1), identifying definable subpopulations of asthma with discrete pathogenic pathways, is introduced and a conceptual framework to model endotypes is presented.

This Review is structured into four sections: weaknesses of the current T-helper-2 (Th2)-inflammation

Search strategy and selection criteria

National Center for Biotechnology Information (NCBI) public domain databases, Pubmed, and OMIM (On-line mendelian inheritance in man) were searched by combining the key term "asthma" sequentially with each of the following descriptors: "phenotype", "subtype", "factor analysis", "principal component analysis", "cluster", "clade", "classification", "pattern recognition", "severe", "severity", "refractory", "complex disease", "heterogeneity", "variant", "inflammation", "sputum", "biopsy", "eosinophil", "neutrophil", "lymphocyte", "Th2", "IL4", "IL13", "GATA3", "TBET", "macrophage", "immunity", "immunity-adaptive", "immunity-innate", "Treg", "IL17", "IL23", "biomarker", "exhaled", "condensate", "exacerbation", "virus", "COPD", "atopy", "epidemiology", "genome-wide scan", "candidate gene", "ontology", "childhood", "remission", "lung function", "FEV1 decline", "airway reactivity", "bronchial hyperreactivity", "airway smooth muscle", "inhomogeneity", "resolution", "repair", "resolving", "lipoxin", "obesity", "comorbidity", "perception", and "dyspnea". Very recent papers within the past 3 years have been preferentially selected. Human asthma candidate genes, or mouse genes with human homologues, identified in the primary search from gene profiling studies, linkage studies, or from basic and clinical biology were assigned to endotype component categories on the basis of gene ontology (GO) categories, OMIM, and published work proximity to relevant processes with the iHoP (Information hyperlinked over Protein), search tools, and biological plausibility.

Lancet 2008; 372: 1107–19

See [Editorial](#) page 1009

Lung Disease Research Group, Departments of Medicine and Pharmacology, University of Melbourne, Parkville, VIC, Australia

(Prof G P Anderson PhD)

Correspondence to:

Prof Gary P Anderson, Lung Disease Research Group, Departments of Medicine and Pharmacology, University of Melbourne, Parkville, 3010 VIC, Australia

gpa@unimelb.edu.au

Panel 1: What is an endotype?

Endotype—a contraction of endophenotype—is a subtype of disease defined functionally and pathologically by a molecular mechanism or by treatment response. Asthma, like many chronic disorders, is a heterogeneous and genetically complex disease, meaning that many genes (>100 have been identified) are likely to contribute, variably, to its different manifestations. Asthma is likely to have several specific endotypes associated with distinct clinical features, divergent underlying molecular causes, and distinct treatment responses.




Figure 2: Linear representation of the Th2-inflammation hypothesis

The diagram shows the most common view of asthma, which is that allergen exposure in sensitive individuals induces Th2 immune deviation. Th2 immunity drives eosinophilic inflammation and tissue damage, resulting in airway hyper-responsiveness and mediator release. These, in turn, cause symptoms. The hypothesis is simple but does not accord well with current evidence and cannot explain the heterogeneity of asthma.

hypothesis; asthma heterogeneity in terms of ontogeny, clinical phenotypes, and molecular patterns; asthma endotypes; and novel mechanisms, with particular emphasis on alternative macrophage specification programmes and the role of innate immunity as determinants of more severe and steroid-refractory asthma endotypes. By understanding asthma endotypes, and their molecular determinants, effective therapies, and possibly cures, can be developed that are highly effective in targeted patient subgroups.

Limitations of the Th2-inflammation hypothesis

Since the mid-1990s, asthma research has been propelled forward by innovations stemming from the Th2-inflammation hypothesis, providing a molecular framework for understanding the well known associations of atopy or IgE and eosinophilic lung inflammation with asthma. A helper T-cell population induced by interleukin 4 is able to produce a panel of cytokines—such as interleukin 4 or 13 (causing B-cell IgE production, mucus secretion, and fibrosis), interleukin 5 (causing eosinophilic inflammation and

damage), and interleukin 9 (promoting mast cell growth)—which induce traits associated with classic asthma (figure 2). The Th2-inflammation hypothesis^{8,9} coincided with the rise of genetically modified mouse technology and molecular profiling methods. Thus, lung Th2 immunity is now understood in fine molecular detail: from the nature of antigen, through the co-stimulation topology of antigen-presenting dendritic cells, to the language of transcription factors and chromatin reshaping that controls gene programmes governing the emergence and persistence of Th2-biased lymphocytes and their trafficking patterns *in vivo*.^{10–12}

Th2 immunity is undoubtedly important in some asthma endotypes. But even from its inception, concerns have arisen about whether the Th2-inflammation hypothesis would lead to improved treatments.^{8,9} These concerns have now been heightened. The main reasons for questioning the Th2-inflammation hypothesis is that it cannot explain why airway hyper-responsiveness and tissue remodelling are not clearly linked to inflammation; why existing T-cell immunosuppressives and new Th2-targeted treatments, which often worked well in Th2 disease models, have no or marginal effectiveness in the clinic; why many patients have recurrent exacerbations; why substantial residual disease remains when anti-inflammatory therapy is optimised; why asthma shares some genetic risk factors with chronic obstructive pulmonary disease (COPD); and why some patients have severe asthma. Moreover, the Th2-inflammation model cannot account for the substantial clinical and molecular heterogeneity that has now been unequivocally documented in human asthma. Therefore, this intellectual framework needs to be revised. Indeed, that the entry criteria for patients into almost all clinical trials for asthma does not reflect the pattern of actual asthma in the community is remarkable.¹³

Several problems exist. If T cells were fundamentally important, T-cell inhibitors should be very effective treatments; however, T-cell-directed therapies have uniformly failed in clinical trials.¹⁴ Th2 immunity is fundamental to atopy, but although atopy is a risk factor for asthma in populations, it has poor sensitivity and specificity as a predictor of disease. Eosinophilic inflammation is the Th2 driven trait that most consistently tracks with disease activity, exacerbation susceptibility, treatment responses, and as a useful biomarker to guide treatment.^{15–17} However, airway inflammation is much the same between non-asthmatic atopics, allergic rhinitics, and atopic asthmatics.¹⁸ An almost identical pattern of response occurs after allergen challenge in people allergic to house-dust mites with and without asthma.^{19–21} The Th2-inflammation model predicts that eosinophilic inflammation should drive airway hyper-responsiveness, but no clear relation exists;^{22,23} furthermore, population studies show atopy and airway hyper-responsiveness are not concordant.²⁴ Neutrophils, mast cell infiltration of

airway smooth muscle, intensity of inflammation, and inflammation of airway smooth muscle, might discriminate between inflammation in atopy versus that in asthma, but the sensitivity and specificity of these putative co-determinants has not been formally proven.^{25,26} Lung Th2 cytokines are found as often in atopy as in asthma, and interferon γ (a Th1 cytokine) is actually upregulated in human asthma together with interleukins 4 and 5 in sputum but not in blood.²⁷

T-cell immunosuppressive drugs (eg, ciclosporine and methotrexate) have measurable but very weak effects in asthma, and immunosuppressive therapy after allograft transplantation does not prevent asthma or allergy in children and adolescents.²⁸ Furthermore, glucocorticosteroids (and β_2 agonists), which are highly effective in atopic asthma with eosinophilic inflammation, paradoxically consolidate and intensify Th2 immunity and increase IgE.²⁹ At the molecular level, steroids preferentially suppress interleukin 12 and T-bet (negative regulators of Th2 immunity) and spare STAT6 (which induces Th2 genes).^{30,31} Steroids also dampen expression of T-bet, the transcription factor controlling Th1 (but not Th2) immunity.³² Steroid sensitivity wanes in severe asthma; however, genetic manipulation to enhance steroid effectiveness does not suppress experimental asthma and instead favours Th2 immunity.³³ In children, steroids worsen Th2 immunity and increase IgE, which has been related to the genetic association of the low affinity IgE receptor (FCER2) with risk of severe exacerbations.³⁴ Considered together with the weak effects of therapies of anti-interleukins 4 and 13 (eg, pitakinra) and anti-interleukin 5 (eg, mepolizumab) that are in trials,^{35,36} it is reasonable to suggest that steroids exert their beneficial effects in asthma at loci other than Th2 immunity. Steroids suppress the end effects of Th2 immunity but consolidate the underlying aberration.

Despite these caveats, Th2 immunity is clinically important, specifically for childhood asthma with atopy and mild allergic adult asthma, and Th2 directed therapy will be effective in some asthma endotypes.³⁶ However, present treatments are effective in mild atopic asthma (figure 1), and Th2-directed therapies are unlikely to ameliorate the residual disease burden of more severe disease.

Heterogeneity of clinical asthma and treatment responses

Asthma continues to elude specific definition and can therefore currently only be characterised in functional terms (panel 2). Although evidence has suggested the pronounced heterogeneity of asthma for decades, little interest has focused on understanding its basis. Most asthma trials and research protocols have used inclusion criteria—typically predicted forced expiratory volume in 1 second (FEV₁), degree of reversibility, inflammation, eosinophilia, and often IgE, because they can be measured objectively and accord well with disease

Panel 2: Definitions of asthma and indices of severity

The first systematic attempts to define asthma were made in the 1970s but, despite decades of effort, there is still no specific definition of, or validated diagnostic algorithm for, the disease. Instead, asthma is defined functionally as an inflammatory disorder linked to hyper-responsiveness that causes symptoms:

“Asthma is a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role. The chronic inflammation is associated with airway hyper-responsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness, and coughing, particularly at night or in the early morning. These episodes are usually associated with widespread, but variable, airflow obstruction within the lung that is often reversible either spontaneously or with treatment.”³⁷

Similarly, no current methods exist to objectively define disease severity other than by functional criteria:

- day-time symptoms
- limitations of activity
- nocturnal symptoms and waking
- need for rescue/reliever drugs
- lung function as percentage predicted
- exacerbations
- treatment responses (success or failure to improve or control these factors)³⁷

These functional or operational criteria are, however, practical, pragmatic, and very well thought out because they are simple, work in the context of global asthma guidelines, and allow the disease to be managed even in health-care systems with constrained resources.

notions. However, because asthma is heterogeneous, these criteria have resulted in patients being selected to trial new asthma drugs who are not representative of asthma in general practice.¹³ Advances in asthma epidemiology including longitudinal outcomes studies, population genetics, and molecular profiling methods; application of statistical methods, such as clustering and principal component techniques, to show distinct asthma patterns and subsets; and compelling new clinical research, have all contributed to a reawakening of interest in understanding disease heterogeneity.

Paediatricians have defined three major patterns of wheezing in infants that have also been assessed for their effect on subsequent persistent asthma in adulthood: transient infant wheeze, non-atopic wheezing in toddlers, and IgE-mediated wheeze or asthma. A fourth category has also been introduced: late-onset childhood asthma.³⁸ Evidence now suggests that transdermal sensitisation associated with polymorphism in filaggrin, a molecule important in maintaining cutaneous integrity, rather than aeroallergen links asthma risk with atopy and atopic dermatitis.^{39,40}




Figure 3: Open framework asthma endotype model

The diagram represents processes that are known to be involved in the diverse presentations of asthma as colour-coded elements. Distinct endotypes arise through the variable interplay of these components. In the model, asthma symptoms almost always arise through the interaction of altered lung function (forced expiratory volume in 1 second [FEV₁], airway hyper-responsiveness [AHR], and airway smooth muscle [ASM]) and immunopathologies (remodelling/inflammation) but these symptoms arise and interact independently. Modifying process such as body-mass index (BMI), diet, and in utero exposures; CNS sensitivity to dyspnoea (perception); acute asthma exacerbations (AEA); cigarette smoke (smoke) and environmental irritants are shown impinging on lung function on remodelling and inflammation. Both divisions of the immune system, (adaptive and innate), are shown closely interconnected (arrowed dotted line) and regulated by immune tolerance (Treg) in equilibrium with Th17 immune activation. Defective healing is represented by resolution/repair. Immunity, together with smoke and environmental contaminants, and also exacerbations, are shown interacting with inflammation and remodelling, but no assumption is made that remodelling is dependent on inflammation. Because asthma is defined by symptoms, defects in lung function are essential, and their necessary interaction with other components is indicated by +. Candidate genes linked to these processes are listed in the table.

Wheeze in early life in the absence of atopic sensitisation almost invariably resolves to normal lung function by 13 years of age, whereas atopic sensitisation, particularly before 3 years of age, is associated with a much higher chance of reduced lung function and worsened airway hyper-responsiveness,⁴¹ which occur despite the airway biopsy pathology of non-atopic asthma being almost indistinguishable from atopic asthma in children.⁴² This finding suggests that very early intervention with steroids would be beneficial, but this is not the case,⁴³ perhaps because steroids actually worsen Th2 immune deviation.²⁹ Long-term follow-up studies also suggest that up to half of asthma in adolescence or early adult life represents a relapse of previously quiescent disease.⁴⁴ Asthma affects more boys than girls in childhood, but more women in later life.⁴⁵ At the cellular and molecular level, children who are predisposed to life-long asthma might have substantial changes early in disease, including oxidative stress and acetylation,^{46,47} IRAK-M (a negative regulator of innate immunity),^{48,49} epidermal growth factor (EGF), interleukin 6, and prostaglandin E2.⁵⁰

The heterogeneity of asthmatic inflammation is well documented; Golash first suggested sputum eosinophilia as a hallmark of disease in the 1890s. Woolcock and Peat²⁴ noted that airway hyper-responsiveness was

unimodally distributed in the general population and only partially overlapped with atopy, precluding atopy as a cause of all airway hyper-responsiveness. Wardlaw and colleagues²⁵ have reported the absence of a clear relation between intensity of inflammation (eosinophils) and severity of asthma. They also noted that only airway smooth muscle infiltration by mast cells differentiated the airway pathology of eosinophilic bronchitis from asthma^{25,26} and that this trait, not airway remodelling, is associated with airway hyper-responsiveness.⁵¹

More recently, Simpson and colleagues⁵² have identified distinct inflammatory endotypes in sputum samples from patients with clinical asthma, forming the basis of a simple classification schema: (1) eosinophilic; (2) neutrophilic; (3) mixed (ie, both neutrophils and eosinophils found); and (4) paucigranulocytic (few or no granulocytes in the sputum).

Since these investigators noted evidence of upregulated toll-like receptors, they have specifically linked neutrophilic asthma to innate immunity.⁵³ Halder and Pavord⁵⁴ have independently replicated the identification of neutrophil variant asthma (which has been suggested for many years). These heterogeneous inflammatory patterns are entirely consistent with the identification of a distinct cytokine profile in patients with asthma.⁵⁵ Similarly, application of factor analysis to molecular genetics studies tends to segregate, rather than cluster, atopic and asthmatic disease traits.⁵⁶ Segregation of inflammatory subtypes by non-invasive exhaled breath methods is not yet possible.⁵⁷

The biology of very severe asthma is almost certainly distinct from milder forms of disease, and severe asthma is heterogeneous.¹⁴ Severity should be viewed as a separately regulated biology rather than as one end of the spectrum of disease or the result of an inexorably progressive process. Results from long-term epidemiological studies in asthma have shown that the severity grade of asthma tends to be established early in life, and disease seldom progresses to a more severe grade.⁵⁸ Two major subdivisions of severe asthma have been proposed on the basis of discordant inflammatory patterns.^{59,60} Furthermore, Pavord and colleagues⁶¹ have reported severe asthma in the absence of eosinophilic inflammation and, by inference, Th2 immunity. Brasier and co-workers⁵⁵ applied mathematical pattern-recognition methods to compare cytokine patterns in bronchoalveolar lavage from 43 patients with mild or moderate asthma with 43 patients with severe disease, and noted four distinct profiles to predict methacholine responsiveness. However, few molecular mechanisms exist that unequivocally distinguish severe asthma. Possibilities include CREB (cyclic AMP response element binding protein 1), which regulates gene expression in responses to increases in cyclic AMP;⁶² RIP-2 (receptor-interacting serine-threonine kinase 2), an intermediate in toll-like receptor signalling;⁶³ reduced generation of pro-healing lipoxin A4; or anti-inflammatory interleukin 10.^{64,65}

Combination therapy (bronchodilator plus inhaled glucocorticosteroid) achieves good outcomes for many, but not all, patients with mild to moderately severe asthma and can reduce the rate of severe exacerbations by up to a third (figure 1). Other therapies have also proven to be of measurable, but lesser and variable benefit—eg, cysteinyl-leukotriene receptor antagonists and anti-IgE monoclonal antibodies.^{2–6} Other therapeutic options, such as selective phosphodiesterase isozyme inhibitors, selective chemokine and cytokine antagonists, vaccines, and T-cell-directed or general immunosuppressives, are ineffective or of marginal benefit.⁶⁶ Primary prevention has not proven possible so far, since a meta-analysis shows that avoidance of house-dust mites is not effective.⁷ The β_2 adrenoceptor is polymorphic and its Arg16 variant might adversely affect response to regular short-acting drugs but not long-acting β agonists.⁶⁷ About 15% of patients with asthma respond well to leukotriene antagonists and, although urinary concentrations of leukotriene do not predict responses, polymorphism in 5 lipoxygenase (ALOX) and the receptor (CYSLTR2) can establish drug sensitivity.^{68,69} Steroid responses are very heterogeneous in childhood and adult asthma.^{31,70}

An asthma endotype model

In view of the manifest problems with the Th2-inflammation model and the pronounced heterogeneity of asthma, how is the disease to be modelled and how will new treatments be found? Gibson's four inflammatory patterns provide one simple and useful model.⁵² A more extensive rational approach is to assess

components of disease that could be considered in the definition of endotypes. The asthma endotype model shown in figure 3 maps inter-relations between clinical determinants that are known to be important in the manifestation and expression of asthma across its diverse patterns and severities. The table lists candidate genes linked to endotype determinants. The model is non-linear: it does not make a-priori assumption about the weight of a specific factor—ie, inflammation—to expression of disease. Some of the evidence base for the model is summarised below.

Airway smooth muscle

Airway narrowing, which is largely caused by contraction of airway smooth muscle, is the mechanism that can be most clearly linked to symptoms, as can be inferred from the additional benefit of combining long-acting bronchodilators with steroids and the remarkably effective benefits of killing muscle by bronchial thermoplasty in severe disease.^{71,72} More smooth muscle exists in airways of asthmatic patients, but alone it might not be a cause;⁵¹ modelling studies suggest that increased bulk might even protect against excessive closure. Airway smooth muscle in asthmatic patients also secretes inflammatory cytokines including stem-cell factor (or kit-ligand), a mast cell growth and activation protein that is now also implicated in dendritic cell activation.⁷³ Increased mast cell number and degranulation have been suggested as pathological indicators of changed function.^{25,51}

Specific molecular changes affecting contractility—such as changes in myosin light chain kinase isoforms or cross talk between signalling G proteins—remain controversial,

	Genetic	Biology
Lung function: basal FEV ₁ , airway hyper-responsiveness, airway smooth muscle	EDN1, ADAM33, B2ADR, CREB, CCR5, COL29A1, CSTA, CYSLTR1, CYSLTR2, EP2, FCER2, GSTM, HNMT, KCNS1, LELP1, MMP, MUC7, MLCK, NK2R, PDGFRA, PLA2, PLAU, PTGDR, PTGER, PTGIR, TBX21, VDR	CREB, GSNOR, NOS, NR3C1
Immunity	FLG, IL17F, TGF β , IL6, ROR α , ROR γ , BDNF, chemokines, CD14, CD40, CD86, DPP10, FCER2, FLG, HLA-G, ICOS, IGHG, IL12B, IL2, IL4, IL6, IL9, IL10, IL13, IL16, IL17, IL18, IL27, IL33, IRAKM, ITK, MICB, MMP, MRP1, MUC1, NOD, PHF11, PLA2, PPARG, PTGDR, PTGER, RIP2, RUNX1, SFTPC, SOCS, SPP1, STAT6, TBX21, TIM1, VDR, VEGFR	CREB, HCK, IL23, IL33, LYN, NFATc, NOS, NR3C1, PTEN, RIP, ROR, SHIP, SHP, TSLP
Inflammation and remodelling	EDN1, ADAM33, IL17A, IL17F, NRF2, SOD, CREB, VDR, CAT, chemokines, COL29A1, CSTA, DPP10, ECP, EP2, FYN, GSTM, IGHG, IL2, IL5, IL9, IL13, IL17, IL18, IL33, PLA2, PLAU, SOCS, STAT6, TNFA, UTG, VEGFR	AMCase, ARG, C3AR1, c-kit, C3 β , EGFR, CSF2(=GM-CSF), HCK, HMGB1, LYN, NOS, NR3C1, NRF2, PTEN, RAGE, RIP, SCF, SHIP, SHP, SOD, TIMP, TSLP
Resolution and repair	VDR, LEP	IL-10, FAS, NR3C1, RAGE, TIMP, Lipoxin A4 (15LOX, 5LOX), presqualene phosphates
Exacerbations, smoke, and environmental irritants	VDR, AOA, CAT, CYP24A1, GSNOR, HLA-G, IL12B, IL2, IL6, IL12, IL17, IL23, IL33, IRAKM, MMP, MRP1, NOD, SFTPC, UTG, NRF2	CD200, EGFR, IFN, NR3C1, SOD, TIMP
BMI and nutrition	ADRB2, FABP, NR3C1, VDR, FABP4, NR3C1	
Perception	KCNS1, GAD65	
Genes of unknown function from linkage	DCNP1, GCLM, ORMDL3, SCGB3A2	

Candidate genes selected from linkage and microarray studies (genetic) or from the known biology of established disease processes (biology) are shown as putative determinants of asthma endotype trait elements (see also figure 3). Definitions of these genes can be found via the NCBI website OMIM database (<http://www.ncbi.nlm.nih.gov/sites/entrez>).

Table: Putative contribution of asthma candidate genes to endotype elements

and microarray profiling and genetic linkage studies have not suggested strong candidates for changed function.⁷⁴ However, increased rate of shortening has been shown, and although the specific mechanism of this effect is disputed, faster or excessive shortening will probably be the major functional defect causing airway narrowing and symptoms.⁷⁵ Clinical asthma is unlikely to occur in almost all cases without the contribution of airway smooth muscle. All known bronchodilators are functional antagonists, or direct pharmacological antagonists of contractile mediators, and they are not fundamentally able to prevent contraction to intense stimuli.

Airway hyper-responsiveness and dynamic inhomogeneity

Alexander and Paddock first described airway hyper-responsiveness to systemic pilocarpine in 1921.⁷⁶ People with asthma often show pronounced airway hyper-responsiveness with loss of plateau that is orders of magnitude greater than in healthy controls. Airway smooth muscle responds differently to inflammation in patients with asthma, with substantial variation in degree and site of bronchoconstriction during the late-phase rise in inflammation that follows antigen challenge.^{77,78} This heterogeneity of constriction has been linked directly to the loss of bronchoprotective response to deep inspiration,⁷⁹ which might indicate a defect in responses of airway smooth muscles to cyclic stretch, leading to risk of catastrophic bronchoconstriction.⁸⁰ Dynamic hyperinflation caused by patchy constriction of large airways is also thought to contribute greatly to airway hyper-responsiveness.⁸¹ This dynamic inhomogeneity has been directly recorded by tomography (with hyperpolarised 3He-MRI or CT).

Steroids are able to reverse this defect only in the mildest disease.⁸² Airway hyper-responsiveness can be worsened by inflammation, but the main component is inherited separately.⁸³ Palmer and colleagues⁸⁴ noted that serum IgE concentrations, blood eosinophil counts, and airways responsiveness to inhaled agonist were inherited separately in human beings. Inflammation actually correlates poorly with airway hyper-responsiveness. Perhaps the greatest misunderstanding in basic (mouse) and applied asthma research is that intrinsic (inherited) and antigen/inflammation-induced airway hyper-responsiveness arise from different mechanisms.⁸⁵ This notion is particularly ironic in asthma models in mice because they clearly show that basal airway reactivity is a heritable trait, whereas the small and transient labile component indicates changes in access of agonists to airways smooth muscle that is secondary to inflammation.^{84,85}

FEV₁ decline and fixed obstruction risk

Long-term outcome studies show that basal FEV₁ is set early in life and few patients with asthma have excessive rates of decrease.^{58,86,87} However, some patients—particularly those with adult onset asthma, smokers, and those with persistent uncontrolled eosinophilic

inflammation, airway hyper-responsiveness, or with inherited polymorphisms in *ADAM33*—can have rapid decline and progress to fixed obstruction.⁸⁸ Some evidence suggests that inhaled steroid use can reduce an excessive decline, but this notion is controversial.⁸⁷ Linkage studies have shown that inheritance of FEV₁ has no known genetic determinants in common with asthma severity or symptom score.⁸⁹ Bisgaard and colleagues⁹⁰ have noted that the rate of lung function decline and airway hyper-responsiveness in children is associated with lower lung function, delayed use of steroids for symptomatic disease, smoking, and positive allergic skin-prick test,⁹⁰ which accords with Grol and co-workers' earlier work.⁸⁷ In childhood, lower lung function and lower increase in FEV₁ predicts worse airway hyper-responsiveness in adulthood,⁹¹ but the molecular basis for this finding remains unknown.

Immunity, inflammation and remodelling, resolution, and repair

The interplay of immunity, inflammation, and remodelling has been a central theme in asthma research for decades. Eosinophilic inflammation is the trait that is best linked to symptoms and treatment responses, but alone it is not enough to cause asthma, which is absent in atopy, eosinophilic bronchitis, and Crohn's disease (for which airway inflammation also occurs).⁹² Similarly, atopic rhinitis with allergen exposure produces pathological changes similar to asthma in airways and cytokine release without asthma symptoms,²¹ and anti-interleukin-5 antibodies seem to have some benefit but only in a patient subset with a high eosinophil load. Lung eosinophilia is a useful biomarker to titrate steroid responsiveness, and decreasing eosinophils reduce the risk of exacerbations. Inflammation is heterogeneous in asthma. Steroids have little effect in neutrophilic asthma, and basal eosinophilia predicts steroid effect.⁹³ Airway pathological changes are established early in life as basement membrane pseudo-thickening and angiogenesis are evident in children with asthma (and in atopic children without asthma).⁹⁴ Similar early changes probably occur in lung nerves that undergo remarkable plastic changes in asthma, which could partly relate to increased cough. Good evidence also suggests that resolution processes are defective in asthma. Resolvin E1 (18R-trihydroeicosapentaenoic acid) suppresses experimental asthma *in vivo*,⁹⁵ and pro-resolving lipoxins are diminished as asthma severity worsens.⁹⁶ Production of interleukin 10, which exerts an inflammation-suppression effect under some conditions, is also dampened.⁶⁵

Exacerbations

Acute asthma exacerbations—usually but not uniformly caused by rhinovirus infection—are a major cause of morbidity in asthma. In severe asthma, five risk factors have been identified for recurrent exacerbations: severe nasal sinus disease, gastro-oesophageal reflux, recurrent

respiratory infections, psychological affective disorders, and obstructive sleep apnoea.⁹⁷ Rhinovirus infection affects lower airways even in healthy people, causing narrowing and inflammation.⁹⁸ Eosinophilia is a known risk factor for asthma exacerbations.¹⁶ Patients with asthma do not have more frequent infections but rather more intense reactions. This finding has been linked to interplay of the EGF receptor (EGFR) with matrix metalloproteases and external regulated kinase (ERK) signalling,⁹⁹ and primary or acquired interferon α and λ deficiency.^{100,101} Conventional plasmacytoid-dendritic cells govern lung viral immunity. The dendritic cell growth factor FLT3 induces conventional plasmacytoid and plasmacytoid-dendritic cells, and suppresses respiratory syncytial virus infection;¹¹ however, little is known about variations across endotypes or the contribution of inhibitory pathways such as CD200.^{97,102}

Body-mass index, nutrition, and obesity

Thomas Platts-Mills is credited with proposing that watching television was the cause of the asthma epidemic in the 1980s.¹⁰³ Analysis of 20016 children (aged 6–7 years) showed that high bodyweight, salty diet, and time spent watching television were independent risk factors for asthma.¹⁰⁴ This finding is supported by a prospective multiple logistic regression analysis of 932 children in Boston.¹⁰⁵ Body-mass index affects treatment outcome.¹⁰⁶ Vitamin D deficiency, due to less sun exposure and indoor inactivity, has been proposed as an asthma cause.¹⁰⁷ Vitamin D receptor expression in the lung is needed for inflammation and expression of experimental asthma, suggesting that vitamin D might have a stronger effect on host defense than inflammation does.^{108,109} Importantly, both the β_2 adrenoceptor and glucocorticosteroid receptor, which are closely associated with asthma, are also obesity candidate genes. Furthermore, asthma risk almost certainly begins in utero. Prenatal stress, smoke, and exposure to air pollutants all increase asthma risk.^{110–112} Asthma is defined by symptoms, and patients who have poor perception of the severity of their disease (under-perceivers) and patients who suffer overt symptoms but show only small changes in lung function (over-perceivers) have been described. Because dyspnoea is perceived in the anterior insula and amygdala,¹¹³ perception modifying neuronal mechanisms will probably emerge in coming years.

Asthma-COPD overlap

Many asthmatics smoke, which worsens their disease and impairs steroids responses. Lapperre and colleagues¹¹⁴ used factor analysis statistics in 117 patients with COPD by measuring lung function, DLCO (the single-breath diffusing lung capacity for carbon monoxide), PC₂₀ MeCh (the concentration of inhaled methacholine causing a 20% fall in FEV₁), total IgE, exhaled nitric oxide, and differential cell counts in induced sputum. They noted




Figure 4: The unexpected importance of innate immunity in asthma severity and reduced steroid sensitivity

Innate immunity is associated with more severe disease and steroid insensitivity. The diagram shows that as disease severity worsens and becomes less steroid sensitive (red), the contribution of innate immunity increases (black). Classic Th2 immunity is shown in blue. Two novel processes are represented: (1) Auto-innate immunity is shown activating macrophages (tissue is also activated), particularly by factors released from damaged tissue (DAMPs) that trigger innate immunity locally; and (2) T-cell independent processes mediated by innate immunity that resemble Th2 immunity. M ϕ =macrophage.

that airflow inflammation and inflammation of the airways, and systemic features commonly associated with asthma (eg, IgE and eosinophils) were separate and predominantly independent contributors to COPD.¹¹⁴ These data are entirely consistent with genome-wide scans of COPD susceptibility and severity, and underscore the certainty that shared co-determinates of asthma and COPD exist in some patients.^{115–118} Interleukin-13 promoter polymorphism is associated with adverse effects of smoke on lung function,¹¹⁹ and smoking, airway hyper-responsiveness, and eosinophils interact positively for respiratory symptoms.¹²⁰ As in asthma, lung eosinophilia predicts smokers who will benefit from steroids; smoking greatly impedes the activity of steroids.^{121,122} The newly discovered intermediate penderin, like EGF, induces mucin in both asthma and COPD.¹²³ Furthermore, the interplay of interleukin 1 β and tumour growth factor β —recently identified as a cause of small airways disease in COPD—could be a cause of fixed airflow restriction in asthma.¹²⁴ Additionally, vitamin D biology, immune defense mediated by serpins and collectins, dysregulation of oxidative stress and apoptosis with decreased clearance of senescent cells, and secondary necrosis and impaired repair capacity are probably co-determinants of asthma and COPD.

Novel disease mechanisms

Asthma has a heritable component that is estimated to be between 36% and 94%. More than 100 plausible candidate genes have been suggested, but each has a

very low attributable risk (<5%), with often poor replication¹²⁵ or lack of plausible biology or functional single nucleotide polymorphisms. Of this very large information set, two broad new notions are emerging: auto-innate immunity driven by tissue damage and Th2-like responses that occur without T cells (figure 4).

Until recently, innate immunity was thought of only as a rapid front-line defender against infections. It works by triggering host defence after recognition of pathogen associated molecular patterns (PAMPs) on invading pathogens. However, endogenous ligands released from damage associated molecular patterns (DAMPs—ie, ligands such as heat shock proteins, RAGE ligands, and HMGB1) are very able activators of innate immunity. Like PAMPs, DAMPS often signal via the toll-like receptor system and its characteristic MyD88 transduction pathway to directly promote inflammation. Innate immunity is intrinsically resistant to steroids,⁴⁰ which could help to explain why as asthma severity worsens, steroid sensitivity decreases (the molecular basis for this notion is an amalgam of processes including HDAC nitrosylation, MAP kinase phosphatase induction, and inhibitory signalling from altered matrix). This notion has a profound implication: wherever DAMPs and innate immunity contribute to disease, the process will be intrinsically steroid insensitive, which might be why more severe asthma endotypes with established tissue damage are steroid resistant. Furthermore, the main source of ligands to drive this auto-innate immunity is damaged tissue, which is normally removed by macrophages. However, as oxidative stress worsens (with severity or smoking) the ability of macrophages to recognise and remove effete cells decreases, leading to much greater so-called spill of DAMPs.^{126,127}

A second major emerging idea is that there are mechanisms to induce Th2-like effects in the absence of Th2 cells, which is probably the reason why interleukin 13 is so widely associated with highly divergent asthma endotypes. In both cases, the existence of these pathways most probably is an indicator of phylogenetic evolution in that ancient innate immunity, and many of its effector cytokines, predate the evolution of lymphocytes and glucocorticosteroids by million of years. Molecular dissection of these processes is suggesting new intervention points for asthma endotypes.

Interleukin 17 came to attention as an indirect mediator of sustained neutrophilic inflammation.^{128,129} Evidence suggests that interleukin 17 is a mediator of neutrophil variant and severe neutrophilic asthma endotypes.¹³⁰ A defined T-cell subset—Th17—has been identified, but interleukin-17 family cytokine production is not constrained to T cells. Th17 differentiation is dependent on the retinoid receptors ROR α and ROR γ (implicating diet and lipid metabolism),¹³¹ and signalling intermediates STAT3 and STAT4.¹³² Interleukin 17 is induced in human beings by interleukin 23, which is

related to interleukin 12 (a negative regulator of Th2 immunity), especially in the context of interleukin 1 β , interleukin 6, and tumour growth factor β . This mixture is important because tumour growth factor β and interleukin 6 exert counter-balancing control of the regulatory T cells (Tregs) that suppress inflammation, indicating a fine balance between suppression and disease.^{130,133,134}

Dendritic cells are poised at the interface between innate and adaptive immunity, and are essential for both the induction and maintenance of allergic inflammation. Coactivation of dendritic cells by antigen in the presence of stem cell factor—a potent mast cell growth factor—induces sustained interleukin-6 production, triggering concurrent Th2 and Th17 induction via the c-kit receptor.¹³⁵ Because stem cell factor is also implicated in mast cell infiltration of airways smooth muscle, this axis predicts the usefulness of imatinib mesylate (Gleevec, or related compounds), which has potent c-kit receptor kinase blocking activity in some asthma endotypes, especially when mast cells are concurrently implicated.

Viral bronchiolitis in infancy, especially with respiratory syncytial virus, is a risk factor for persistent asthma. In rodents, similar exposure leads to life-long inflammation and phenotypic alteration of the lung. Kim and colleagues¹³⁶ have discovered that activation of macrophages by CD1d expressing natural killer T (NKT) cells induces production of interleukin 13 and its pathological sequelae entirely independently of adaptive immunity. This mechanism operates in both human asthma and COPD,¹³⁶ and defines an adaptive immunity (ie, T cell) independent endotype. Smoke directly and substantially worsens viral inflammation and remodelling.^{137,138}

Interleukin-18 polymorphisms have been replicated in several genetic linkage studies. This cytokine is known to induce IgE by causing NKT cells to upregulate the co-stimulation molecule CD40 and interleukin 4. Interleukin 18, which potently induces interferon γ , has been implicated in processes relevant to neutrophilic and mixed inflammatory patterns because exogenous interleukin 18 triggers bystander memory cells—the types of cells that would reflect past viral infections—to release not only interferon γ , causing neutrophilia, but also interleukin 13, inducing airway remodelling.¹³⁹ This mechanism might relate to asthma endotypes in which recurrent infection drives accelerated lung function decline or which show extensive neutrophilia. However, the role of NKT cells in asthma, where they have been linked to allergic (via interleukin 4) and neutrophilic (via interleukin 17) endotypes, is controversial.

Interleukin 33 is an interleukin-1-like cytokine that was identified initially as the ligand to an orphan receptor called T1/ST2, which is preferentially expressed on Th2 cells. However, interleukin 33 induces airway hyper-responsiveness and goblet hyperplasia and

eosinophilic inflammation dependent on ST2 binding and transduction via the innate immunity transducer, MyD88, concurrently with induction of interleukins 4, 5, and 13. These effects occur in Rag^{-/-} mice that do not have a functional adaptive immune system. These data suggest that Th2 mimicking pathological changes can be induced entirely in the absence of an adaptive immune system defining a second adaptive immunity independent endotype.¹⁴⁰

Rhinovirus infection is the main cause of asthma exacerbations. Epidemiological studies have shown that patients with asthma do not have more frequent infections but rather more severe inflammation. Such patients also overexpress the EGFR. Liu and colleagues⁹⁹ have identified a new mechanism governing the intensity of inflammatory responses to rhinovirus. Rhinovirus binds to ICAM-1, inducing upregulation of EGF and triggering an inflammatory response in the infected cell. In the presence of raised matrix metalloproteinase, excessive EGF is cleaved from the cell surface and binds to unregulated EGFR, sending a signal via ERKs that synergises with the direct response to rhinovirus, greatly enhancing inflammation.⁹⁹ This mechanism is an example of how an asthma candidate gene (matrix metalloproteinase) with weak attributable risk can exert a stronger effect in an altered disease context. Because EGFR also mediates mucus induction, these results suggest the use of EGFR-tyrosine kinase inhibitors (eg, gefitinib or erlotinib) or antihuman epidermal growth factor receptor 2 family antibodies or blockers in some asthma endotypes. Whether EGFR affects the macrophage CD200 pathway, which limits viral inflammation, is not known.¹⁰²

The role of macrophages in asthma has probably been greatly underestimated, and this cell lineage is increasingly researched as the role of innate immunity emerges. Macrophages do not always undergo classic activation, but might adopt alternative phenotypes^{141,142} associated with hallmark features such as induction of AMCases (chitinases), which are linked to asthma severity.¹⁴³ Alternatively, specified macrophages are the main candidates for asthma endotypes that do not need T cells.

Much of the evidence that alternatively specified macrophages can achieve these effects comes from genetically manipulated mice. SHIP-1 is a negative regulator (ie, turns off) inflammatory cytokine and surface receptor signalling. SHIP-1 deficiency causes spontaneous asthma in mice with a Th2-like pattern,¹⁴⁴⁻¹⁴⁶ including AMCases induction.¹⁴³ Similarly, mice with activated *Hck*—a *Src* family kinase—develop an aggressive T-cell-independent eosinophilic lung inflammation associated with progressive airway fibrosis.¹⁴⁵⁻¹⁴⁷ *Lyn* is related to *Hck*. A profound and multi-trait severe asthma syndrome develops in *Lyn*-deficient mice who display hyper-IgE, enhanced bronchoconstriction, mast cell, and eosinophil degranulation; very persistent inflammation associated

with deficient apoptosis; and Th2-like cytokines together with enhanced interferon γ .¹⁴⁸

Biochemically, alternative macrophage activation has been linked to tumour growth factor β and interleukin 13.¹⁴¹ Because these responses can occur in the complete absence of T cells, interleukin 13 probably exerts its effect via the type II interleukin-4 receptor.¹⁴⁹ These findings might explain why interleukin 13 is a candidate gene in both asthma and COPD.¹⁵⁰

Summary and implications

This Review has developed the argument that the Th2-inflammation hypothesis, although useful, is not adequate to understand the substantial heterogeneity of asthma. The first iteration of an open-frame asthma endotype model has been presented and discussed in the context of entirely novel disease pathways, many of which are independent of adaptive immunity. The role of innate immunity, which is intrinsically insensitive to steroids and can be driven by tissue damage, has been emphasised. There are important implications. Definition of asthma endotypes opens the possibility of much more precise disease classification and definition of biomarkers that meet formal diagnostic or prognostic criteria. From analysis of several asthma candidates across many endotype components, effective future therapies will be used as an adjunct to existing medicines or will be combinations of activities, since few known candidates affect enough crucial endotype components to be effective in their own right. Some asthma endotypes will be reclassified as orphan diseases. Because the inclusion criteria and endpoints for clinical trials directed at some novel endotypes are likely to be unvalidated, we will probably increasingly use adaptive clinical trial methods to identify responding patient endotypes. In adaptive clinical trials, responsive patients (and non-responsive patients) are identified during the course of the trial itself (allowing the trial design to be adapted while running) rather than during retrospective statistical analysis after the trial has closed. Specific definition of asthma endotypes should also spur and redirect basic research to discover the mechanisms and highly innovative ideas that are required to improve asthma therapy and ultimately prevent or cure the disorder or disorders.

Conflict of interest statement

I declare that I have no specific conflict of interest with the material presented in this Review. Currently, or within the past 3 years, I have received consultancy, travel, and speaker fees from AstraZeneca Pharmaceuticals in relation to β -agonist steroid combination products. I have received consultancy fees from Roche Pharmaceuticals in relation to review of preclinical investigational concepts and compounds. My laboratory has several full-time staff who undertake fee-for-service testing of compounds in preclinical animal models of asthma and COPD, for which the surplus arising is used to support basic research. The University of Melbourne has licensed patented intellectual property arising partly from research in my laboratory on GM-CSF to MorphoSys AG, Germany, for the treatment of chronic inflammatory disorders including lung disease.

Acknowledgments

The author's research is funded by the National Health and Medical Research Council of Australia (NHMRC) via project and programme grants.

References

- Anderson HR, Gupta R, Strachan DP, Limb ES. 50 years of asthma: UK trends from 1955 to 2004. *Thorax* 2007; **62**: 85–90.
- Bateman ED, Clark TJ, Frith L, Bousquet J, Busse WW, Pedersen SE. Rate of response of individual asthma control measures varies and may overestimate asthma control: an analysis of the goal study. *J Asthma* 2007; **44**: 667–73.
- Bateman ED, Jacques L, Goldfrad C, Atienza T, Mihaescu T, Duggan M. Asthma control can be maintained when fluticasone propionate/salmeterol in a single inhaler is stepped down. *J Allergy Clin Immunol* 2006; **117**: 563–70.
- Tee AK, Koh MS, Gibson PG, Lasserson TJ, Wilson AJ, Irving LB. Long-acting beta2-agonists versus theophylline for maintenance treatment of asthma. *Cochrane Database Syst Rev* 2007; **3**: CD001281.
- Ducharme FM, Lasserson TJ, Cates CJ. Long-acting beta2-agonists versus anti-leukotrienes as add-on therapy to inhaled corticosteroids for chronic asthma. *Cochrane Database Syst Rev* 2006; **4**: CD003137.
- Walker S, Monteil M, Phelan K, Lasserson TJ, Walters EH. Anti-IgE for chronic asthma in adults and children. *Cochrane Database Syst Rev* 2006; **2**: CD003559.
- MacDonald C, Sternberg A, Hunter PR. A systematic review and meta-analysis of interventions used to reduce exposure to house dust and their effect on the development and severity of asthma. *Environ Health Perspect* 2007; **115**: 1691–95.
- Anderson GP, Coyle AJ. TH2 and 'TH2-like' cells in allergy and asthma: pharmacological perspectives. *Trends Pharmacol Sci* 1994; **15**: 324–32.
- Coyle AJ, Le Gros G, Bertrand C, et al. Interleukin-4 is required for the induction of lung Th2 mucosal immunity. *Am J Respir Cell Mol Biol* 1995; **13**: 54–59.
- Holloway AF, Rao S, Chen X, Shannon MF. Changes in chromatin accessibility across the GM-CSF promoter upon T cell activation are dependent on nuclear factor kappaB proteins. *J Exp Med* 2003; **197**: 413–23.
- Smit JJ, Lindell DM, Boon L, Kool M, Lambrecht BN, Lukacs NW. The balance between plasmacytoid DC versus conventional DC determines pulmonary immunity to virus infections. *PLoS ONE* 2008; **3**: e1720.
- van Rijt LS, Jung S, Kleinjan A, et al. In vivo depletion of lung CD11c+ dendritic cells during allergen challenge abrogates the characteristic features of asthma. *J Exp Med* 2005; **201**: 981–91.
- Travers J, Marsh S, Williams M, et al. External validity of randomised controlled trials in asthma: to whom do the results of the trials apply? *Thorax* 2007; **62**: 219–23.
- Chanez P, Wenzel SE, Anderson GP, et al. Severe asthma in adults: what are the important questions? *J Allergy Clin Immunol* 2007; **119**: 1337–48.
- Brown HM, Storey G, George WH. Beclomethasone dipropionate: a new steroid aerosol for the treatment of allergic asthma. *BMJ* 1972; **1**: 585–90.
- Green RH, Brightling CE, McKenna S, et al. Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002; **360**: 1715–21.
- Jackola DR, Miller MB, Liebeler CL, Blumenthal MN. Search for quantitative trait loci of atopy-associated immune responses using allergen-specific IgG1 as an "endophenotype". *Hum Immunol* 2007; **68**: 839–43.
- Brown JL, Behndig AF, Sekerel BE, et al. Lower airways inflammation in allergic rhinitis: a comparison with asthmatics and normal controls. *Clin Exp Allergy* 2007; **37**: 688–95.
- Braunstaal GJ, Overbeek SE, Fokkens WJ, et al. Segmental bronchoprovocation in allergic rhinitis patients affects mast cell and basophil numbers in nasal and bronchial mucosa. *Am J Respir Crit Care Med* 2001; **164**: 858–65.
- Braunstaal GJ, Fokkens WJ, Overbeek SE, Kleinjan A, Hoogsteden HC, Prins JB. Mucosal and systemic inflammatory changes in allergic rhinitis and asthma: a comparison between upper and lower airways. *Clin Exp Allergy* 2003; **33**: 579–87.
- Lopuhaa CE, Out TA, Jansen HM, Aalberse RC, van der Zee JS. Allergen-induced bronchial inflammation in house dust mite-allergic patients with or without asthma. *Clin Exp Allergy* 2002; **320**: 1720–27.
- Alvarez MJ, Olaguibel JM, Garcia BE, Tabar AI, Urbiola E. Comparison of allergen-induced changes in bronchial hyperresponsiveness and airway inflammation between mildly allergic asthma patients and allergic rhinitis patients. *Allergy* 2000; **55**: 531–39.
- Alvarez MJ, Olaguibel JM, Garcia BE, Rodriguez A, Tabar AI, Urbiola E. Airway inflammation in asthma and perennial allergic rhinitis. Relationship with nonspecific bronchial responsiveness and maximal airway narrowing. *Allergy* 2000; **55**: 355–62.
- Woolcock AJ, Peat J. What is the relationship between airway hyperresponsiveness and atopy? *Am J Respir Crit Care Med* 2000; **161**: 2155–217.
- Wardlaw AJ, Brightling CE, Green R, Woltmann G, Bradding P, Pavord ID. New insights into the relationship between airway inflammation and asthma. *Clin Sci (Lond)* 2002; **103**: 201–11.
- Brightling CE, Bradding P, Symon FA, Holgate ST, Wardlaw AJ, Pavord ID. Mast-cell infiltration of airway smooth muscle in asthma. *N Engl J Med* 2002; **346**: 1699–705.
- Cho SH, Stanciu LA, Holgate ST, Johnston SL. Increased interleukin-4, interleukin-5, and interferon-gamma in airway CD4+ and CD8+ T cells in atopic asthma. *Am J Respir Crit Care Med* 2005; **171**: 224–30.
- Dehlink E, Gruber S, Eiwegger T, et al. Immunosuppressive therapy does not prevent the occurrence of Immunoglobulin E-mediated allergies in children and adolescents with organ transplants. *Pediatrics* 2006; **118**: e764–70.
- Anderson GP. Interactions between corticosteroids and beta-adrenergic agonists in asthma disease induction, progression, and exacerbation. *Am J Respir Crit Care Med* 2000; **161**: S188–96.
- Miyaura H, Iwata M. Direct and indirect inhibition of Th1 development by progesterone and glucocorticoids. *J Immunol* 2002; **168**: 1087–94.
- Tantisira KG, Hwang ES, Raby BA, et al. TBX21: a functional variant predicts improvement in asthma with the use of inhaled corticosteroids. *Proc Natl Acad Sci USA* 2004; **101**: 18099–104.
- Raby BA, Hwang ES, Van Steen K, et al. T-bet polymorphisms are associated with asthma and airway hyperresponsiveness. *Am J Respir Crit Care Med* 2006; **173**: 64–70.
- Brandt Jvd, Luhder F, McPherson KG, et al. Enhanced glucocorticoid receptor signaling in T cells impacts thymocyte apoptosis and adaptive immune responses. *Am J Pathol* 2007; **170**: 1041–53.
- Tantisira KG, Silverman ES, Mariani TJ, et al. FCER2: a pharmacogenetic basis for severe exacerbations in children with asthma. *J Allergy Clin Immunol* 2007; **120**: 1285–91.
- Leckie MJ, ten Brinke A, Khan J, et al. Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000; **356**: 2144–48.
- Wenzel S, Wilbraham D, Fuller R, Getz EB, Longphre M. Effect of an interleukin-4 variant on late phase asthmatic response to allergen challenge in asthmatic patients: results of two phase 2a studies. *Lancet* 2007; **370**: 1422–31.
- Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J* 2008; **31**: 143–78.
- Bel EH. Clinical phenotypes of asthma. *Curr Opin Pulm Med* 2004; **10**: 44–50.
- McLean WH, Palmer CN, Henderson J, Kabesch M, Weidinger S, Irvine AD. Filaggrin variants confer susceptibility to asthma. *J Allergy Clin Immunol* 2008; **121**: 1294–95.
- Zhang N, Truong-Tran QA, Tancowny B, Harris KE, Schleimer RP. Glucocorticoids enhance or spare innate immunity: effects in airway epithelium are mediated by CCAAT/enhancer binding proteins. *J Immunol* 2007; **179**: 578–89.
- Illi S, von Mutius E, Lau S, Niggemann B, Gruber C, Wahn U. Perennial allergen sensitisation early in life and chronic asthma in children: a birth cohort study. *Lancet* 2006; **368**: 763–70.
- Turato G, Barbato A, Baraldo S, et al. Non-atopic Children with multi-trigger wheezing have airway pathology comparable to atopic asthma. *Am J Respir Crit Care Med* 2008.

- 43 den Otter JJ, van Schayck CP, Folgering HT, van den Boom G, Akkermans RP, van Weel C. Early intervention with inhaled corticosteroids in subjects with rapid decline in lung function and signs of bronchial hyperresponsiveness: results from the DIMCA programme. *Eur J Gen Pract* 2007; **13**: 89–91.
- 44 Piippo-Savolainen E, Korppi M. Wheezy babies—wheezy adults? Review on long-term outcome until adulthood after early childhood wheezing. *Acta Paediatr* 2008; **97**: 5–11.
- 45 Postma DS. Gender differences in asthma development and progression. *Gen Med* 2007; **4** (suppl B): S133–46.
- 46 Carraro S, Rezzi S, Reniero F, et al. Metabolomics applied to exhaled breath condensate in childhood asthma. *Am J Respir Crit Care Med* 2007; **175**: 986–90.
- 47 Zanconato S, Carraro S, Corradi M, et al. Leukotrienes and 8-isoprostane in exhaled breath condensate of children with stable and unstable asthma. *J Allergy Clin Immunol* 2004; **113**: 257–63.
- 48 Balaci L, Spada MC, Olla N, et al. IRAK-M is involved in the pathogenesis of early-onset persistent asthma. *Am J Hum Genet* 2007; **80**: 1103–14.
- 49 Nakashima K, Hirota T, Obara K, et al. An association study of asthma and related phenotypes with polymorphisms in negative regulator molecules of the TLR signaling pathway. *J Hum Genet* 2006; **51**: 284–91.
- 50 Kicic A, Sutanto EN, Stevens PT, Knight DA, Stick SM. Intrinsic biochemical and functional differences in bronchial epithelial cells of children with asthma. *Am J Respir Crit Care Med* 2006; **174**: 1110–18.
- 51 Siddiqui S, Mistry V, Doe C, et al. Airway hyperresponsiveness is dissociated from airway wall structural remodeling. *J Allergy Clin Immunol* 2008; **122**: 335–41.
- 52 Simpson JL, Scott R, Boyle MJ, Gibson PG. Inflammatory subtypes in asthma: assessment and identification using induced sputum. *Respirology* 2006; **11**: 54–61.
- 53 Simpson JL, Grissell TV, Douwes J, Scott RJ, Boyle MJ, Gibson PG. Innate immune activation in neutrophilic asthma and bronchiectasis. *Thorax* 2007; **62**: 211–18.
- 54 Haldar P, Pavord ID. Noneosinophilic asthma: a distinct clinical and pathologic phenotype. *J Allergy Clin Immunol* 2007; **119**: 1043–52.
- 55 Brasier AR, Victor S, Boetticher G, et al. Molecular phenotyping of severe asthma using pattern recognition of bronchoalveolar lavage-derived cytokines. *J Allergy Clin Immunol* 2008; **121**: 30–37.
- 56 Pillai SG, Tang Y, van den Oord E, et al. Factor analysis in the Genetics of Asthma International Network family study identifies five major quantitative asthma phenotypes. *Clin Exp Allergy* 2008; **38**: 421–29.
- 57 Dragonieri S, Schot R, Mertens BJ, et al. An electronic nose in the discrimination of patients with asthma and controls. *J Allergy Clin Immunol* 2007; **120**: 856–62.
- 58 Phelan PD, Robertson CF, Olinsky A. The Melbourne Asthma Study: 1964–1999. *J Allergy Clin Immunol* 2002; **109**: 189–94.
- 59 Wenzel SE, Schwartz LB, Langmack EL, et al. Evidence that severe asthma can be divided pathologically into two inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J Respir Crit Care Med* 1999; **160**: 1001–08.
- 60 Lemiere C, Ernst P, Olivenstein R, et al. Airway inflammation assessed by invasive and noninvasive means in severe asthma: eosinophilic and noneosinophilic phenotypes. *J Allergy Clin Immunol* 2006; **118**: 1033–39.
- 61 Pavord ID, Brightling CE, Woltmann G, Wardlaw AJ. Non-eosinophilic corticosteroid unresponsive asthma. *Lancet* 1999; **353**: 2213–14.
- 62 Chiappara G, Chanez P, Bruno A, et al. Variable p-CREB expression depicts different asthma phenotypes. *Allergy* 2007; **62**: 787–94.
- 63 Nakashima K, Hirota T, Suzuki Y, et al. Association of the RIP2 gene with childhood atopic asthma. *Allergol Int* 2006; **55**: 77–83.
- 64 Levy BD, Bonnans C, Silverman ES, Palmer LJ, Marigowda G, Israel E. Diminished lipoxin biosynthesis in severe asthma. *Am J Respir Crit Care Med* 2005; **172**: 824–30.
- 65 Xystrakis E, Kusumakar S, Boswell S, et al. Reversing the defective induction of IL-10-secreting regulatory T cells in glucocorticoid-resistant asthma patients. *J Clin Invest* 2006; **116**: 146–55.
- 66 Barnes PJ. New therapies for asthma. *Trends Mol Med* 2006; **11**: 515–20.
- 67 Hawkins GA, Weiss ST, Bleecker ER. Clinical consequences of ADRbeta2 polymorphisms. *Pharmacogenomics* 2008; **9**: 349–58.
- 68 Klotzman M, York TP, Pillai SG, et al. Pharmacogenetics of the 5-lipoxygenase biosynthetic pathway and variable clinical response to montelukast. *Pharmacogenet Genomics* 2007; **17**: 189–96.
- 69 Pillai SG, Cousens DJ, Barnes AA, et al. A coding polymorphism in the CYSLT2 receptor with reduced affinity to LTD4 is associated with asthma. *Pharmacogenetics* 2004; **14**: 627–33.
- 70 Szeffler SJ, Martin RJ, King TS, et al. Significant variability in response to inhaled corticosteroids for persistent asthma. *J Allergy Clin Immunol* 2002; **109**: 410–18.
- 71 Pauwels RA, Lofdahl CG, Postma DS, et al. Effect of inhaled formoterol and budesonide on exacerbations of asthma. Formoterol and Corticosteroids Establishing Therapy (FACET) International Study Group. *N Engl J Med* 1997; **337**: 1405–11.
- 72 Pavord ID, Cox G, Thomson NC, et al. Safety and efficacy of bronchial thermoplasty in symptomatic, severe asthma. *Am J Respir Crit Care Med* 2007; **176**: 1185–91.
- 73 Kassel O, Schmidlin F, Duvernelle C, Gasser B, Massard G, Frossard N. Human bronchial smooth muscle cells in culture produce stem cell factor. *Eur Respir J* 1999; **13**: 951–54.
- 74 Woodruff PG. Gene expression in asthmatic airway smooth muscle. *Proc Am Thorac Soc* 2008; **5**: 113–18.
- 75 An SS, Bai TR, Bates JHT, et al. Airway smooth muscle dynamics: a common pathway of airway obstruction in asthma. *Eur Respir J* 2007; **29**: 834–60.
- 76 Alexander HL, Paddock R. Bronchial asthma: response to pilocarpine and epinephrine. *Arch Intern Med* 1921; **27**: 184–91.
- 77 Henderson AC, Ingenito EP, Atileh H, Israel E, Suki B, Lutchen KR. How does airway inflammation modulate asthmatic airway constriction? An antigen challenge study. *J Appl Physiol* 2003; **95**: 873–82.
- 78 Winkler T, Venegas JG. Complex airway behavior and paradoxical responses to bronchoprovocation. *J Appl Physiol* 2007; **103**: 655–63.
- 79 Lutchen KR, Jensen A, Atileh H, et al. Airway constriction pattern is a central component of asthma severity: the role of deep inspirations. *Am J Respir Crit Care Med* 2001; **164**: 207–15.
- 80 Venegas JG, Winkler T, Musch G, et al. Self-organized patchiness in asthma as a prelude to catastrophic shifts. *Nature* 2005; **434**: 777–82.
- 81 Brown RH, Pearse DB, Pyrgos G, Liu MC, Togias A, Permutt S. The structural basis of airways hyperresponsiveness in asthma. *J Appl Physiol* 2006; **101**: 30–39.
- 82 Scichilone N, Permutt S, Bellia V, Togias A. Inhaled corticosteroids and the beneficial effect of deep inspiration in asthma. *Am J Respir Crit Care Med* 2005; **172**: 693–99.
- 83 Denham S, Koppelman GH, Blakey J, et al. Meta-analysis of genome-wide linkage studies of asthma and related traits. *Respir Res* 2008; **9**: 38.
- 84 Palmer LJ, Burton PR, James AL, Musk AW, Cookson WO. Familial aggregation and heritability of asthma-associated quantitative traits in a population-based sample of nuclear families. *Eur J Hum Genet* 2000; **8**: 853–60.
- 85 Wagers SS, Haverkamp HC, Bates JHT, et al. Intrinsic and antigen-induced airway hyperresponsiveness are the result of diverse physiological mechanisms. *J Appl Physiol* 2007; **102**: 221–30.
- 86 Sears MR, Greene JM, Willan AR, et al. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med* 2003; **349**: 1414–22.
- 87 Grol MH, Gerritsen J, Vonk JM, et al. Risk factors for growth and decline of lung function in asthmatic individuals up to age 42 years. A 30-year follow-up study. *Am J Respir Crit Care Med* 1999; **160**: 1830–37.
- 88 ten Brinke A. Risk factors associated with irreversible airflow limitation in asthma. *Curr Opin Allergy Clin Immunol* 2008; **8**: 63–69.
- 89 Bouzigon E, Siroux V, Dizier MH, et al. Scores of asthma and asthma severity reveal new regions of linkage in EGEA study families. *Eur Respir J* 2007; **30**: 253–59.
- 90 Bisgaard H, Pedersen S, Anhoj J, et al. Determinants of lung function and airway hyperresponsiveness in asthmatic children. *Respir Med* 2007; **101**: 1477–82.

- 91 Grol MH, Postma DS, Vonk JM, et al. Risk factors from childhood to adulthood for bronchial responsiveness at age 32–42 yr. *Am J Respir Crit Care Med* 1999; **160**: 150–56.
- 92 Louis E, Louis R, Shute J, et al. Bronchial eosinophilic infiltration in Crohn's disease in the absence of pulmonary disease. *Clin Exp Allergy* 1999; **29**: 660–66.
- 93 Gauvreau GM, Inman MD, Kelly M, Watson RM, Dorman SC, O'Byrne PM. Increased levels of airway neutrophils reduce the inhibitory effects of inhaled glucocorticosteroids on allergen-induced airway eosinophils. *Can Respir J* 2002; **9**: 26–32.
- 94 Barbato A, Turato G, Baraldo S, et al. Epithelial damage and angiogenesis in the airways of children with asthma. *Am J Respir Crit Care Med* 2006; **174**: 975–81.
- 95 Aoki H, Hisada T, Ishizuka T, et al. Resolvin E1 dampens airway inflammation and hyperresponsiveness in a murine model of asthma. *Biochem Biophys Res Commun* 2008; **367**: 509–15.
- 96 Bonnans C, Levy BD. Lipid mediators as agonists for the resolution of acute lung inflammation and injury. *Am J Respir Cell Mol Biol* 2007; **36**: 201–05.
- 97 ten Brinke A, Sterk PJ, Masclée AA, et al. Risk factors of frequent exacerbations in difficult-to-treat asthma. *Eur Respir J* 2005; **26**: 812–18.
- 98 de Kluijver J, Grunberg K, Sont JK, et al. Rhinovirus infection in nonasthmatic subjects: effects on intrapulmonary airways. *Eur Respir J* 2002; **20**: 274–79.
- 99 Liu K, Gualano RC, Hibbs ML, Anderson GP, Bozinovski S. Epidermal growth factor receptor signaling to Erk1/2 and STATs control the intensity of the epithelial inflammatory responses to rhinovirus infection. *J Biol Chem* 2008; **283**: 9977–85.
- 100 Contoli M, Message SD, Laza-Stanca V, et al. Role of deficient type III interferon-lambda production in asthma exacerbations. *Nat Med* 2006; **12**: 1023–26.
- 101 Wark PA, Johnston SL, Bucchieri F, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med* 2005; **201**: 937–47.
- 102 Snelgrove RJ, Goulding J, Didierlaurent AM, et al. A critical function for CD200 in lung immune homeostasis and the severity of influenza infection. *Nat Immunol* 2008; **9**: 1074–83.
- 103 Platts-Mills TAE, Perzanowski M, Carter MC, Woodfolk JA. The rising prevalence and severity of asthma in western society: are the causes of asthma the cause of the increase? In: Platts-Mills TAE, ed. *Asthma: causes and mechanisms of an epidemic inflammatory disease*. Boca Raton: CRC Press, 1999: 1–22.
- 104 Corbo GM, Forastiere F, Sario MD, et al. Wheeze and asthma in children: associations with body mass index, sports, television viewing, and diet. *Epidemiology* 2008; **19**: 747–55.
- 105 Taveras EM, Rifas-Shiman SL, Camargo CA Jr, et al. Higher adiposity in infancy associated with recurrent wheeze in a prospective cohort of children. *J Allergy Clin Immunol* 2008; **121**: 1161–66 e3.
- 106 Peters-Golden M, Swern A, Bird SS, Hustad CM, Grant E, Edelman JM. Influence of body mass index on the response to asthma controller agents. *Eur Respir J* 2006; **27**: 495–503.
- 107 Devereux G, Litonjua AA, Turner SW, et al. Maternal vitamin D intake during pregnancy and early childhood wheezing. *Am J Clin Nutr* 2007; **85**: 853–59.
- 108 Wittke A, Weaver V, Mahon BD, August A, Cantorna MT. Vitamin D receptor-deficient mice fail to develop experimental allergic asthma. *J Immunol* 2004; **173**: 3432–36.
- 109 Wittke A, Chang A, Froicu M, et al. Vitamin D receptor expression by the lung micro-environment is required for maximal induction of lung inflammation. *Arch Biochem Biophys* 2007; **460**: 306–13.
- 110 Pincus-Knackstedt MK, Joachim RA, Blois SM, et al. Prenatal stress enhances susceptibility of murine adult offspring toward airway inflammation. *J Immunol* 2006; **177**: 8484–92.
- 111 Hamada K, Suzuki Y, Leme A, et al. Exposure of pregnant mice to an air pollutant aerosol increases asthma susceptibility in offspring. *J Toxicol Environ Health A* 2007; **70**: 688–95.
- 112 Wang L, Joad JP, Zhong C, Pinkerton KE. Effects of environmental tobacco smoke exposure on pulmonary immune response in infant monkeys. *J Allergy Clin Immunol* 2008; **122**: 400–06.
- 113 von Leupoldt A, Sommer T, Kegat S, et al. The unpleasantness of perceived dyspnea is processed in the anterior insula and amygdala. *Am J Respir Crit Care Med* 2008; **177**: 1026–32.
- 114 Lapperre TS, Snoeck-Stroband JB, Gosman MME, et al. Dissociation of lung function and airway inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004; **170**: 499–504.
- 115 Silverman EK, Speizer FE, Weiss ST, et al. Familial aggregation of severe, early-onset COPD: candidate gene approaches. *Chest* 2000; **117** (suppl 1): 273S–4S.
- 116 Silverman EK, Palmer LJ, Mosley JD, et al. Genomewide linkage analysis of quantitative spirometric phenotypes in severe early-onset chronic obstructive pulmonary disease. *Am J Hum Genet* 2002; **70**: 1229–39.
- 117 Silverman EK, Mosley JD, Palmer LJ, et al. Genome-wide linkage analysis of severe, early-onset chronic obstructive pulmonary disease: airflow obstruction and chronic bronchitis phenotypes. *Hum Mol Genet* 2002; **11**: 623–32.
- 118 Patel BD, Loo WJ, Tasker AD, et al. Smoking related COPD and facial wrinkling: is there a common susceptibility? *Thorax* 2006; **61**: 568–671.
- 119 Sadeghnejad A, Meyers DA, Bottai M, Sterling DA, Bleecker ER, Ohar JA. IL13 promoter polymorphism 1112C/T modulates the adverse effect of tobacco smoking on lung function. *Am J Respir Crit Care Med* 2007; **176**: 748–52.
- 120 Jansen DF, Schouten JP, Vonk JM, et al. Smoking and airway hyperresponsiveness especially in the presence of blood eosinophilia increase the risk to develop respiratory symptoms: a 25-year follow-up study in the general adult population. *Am J Respir Crit Care Med* 1999; **160**: 259–64.
- 121 Pedersen SE, Bateman ED, Bousquet J, Busse WW, Yoxall S, Clark TJ. Determinants of response to fluticasone propionate and salmeterol/fluticasone propionate combination in the Gaining Optimal Asthma control study. *J Allergy Clin Immunol* 2007; **120**: 1036–42.
- 122 Bel EH. Smoking: a neglected cause of glucocorticoid resistance in asthma. *Am J Respir Crit Care Med* 2003; **168**: 1265–66.
- 123 Nakao I, Kanaji S, Ohta S, et al. Identification of pendrin as a common mediator for mucus production in bronchial asthma and chronic obstructive pulmonary disease. *J Immunol* 2008; **180**: 6262–69.
- 124 Araya J, Cambier S, Markovics JA, et al. Squamous metaplasia amplifies pathologic epithelial-mesenchymal interactions in COPD patients. *J Clin Invest* 2007; **117**: 3551–62.
- 125 Hersh CP, Raby BA, Soto-Quiros ME, et al. Comprehensive testing of positionally cloned asthma genes in two populations. *Am J Respir Crit Care Med* 2007; **176**: 849–57.
- 126 Bhavsar P, Hew M, Khorasani N, et al. Relative corticosteroid insensitivity of alveolar macrophages in severe asthma compared to non-severe asthma. *Thorax* 2008; **63**: 784–90.
- 127 Hodge S, Hodge G, Ahern J, Jersmann H, Holmes M, Reynolds PN. Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2007; **37**: 748–55.
- 128 Ivanov S, Bozinovski S, Bossios A, et al. Functional relevance of the IL-23-IL-17 axis in lungs in vivo. *Am J Respir Cell Mol Biol* 2007; **36**: 442–51.
- 129 Linden A, Laan M, Anderson GP. Neutrophils, interleukin-17A and lung disease. *Eur Respir J* 2005; **25**: 159–72.
- 130 Pene J, Chevalier S, Preisser L, et al. Chronically inflamed human tissues are infiltrated by highly differentiated Th17 lymphocytes. *J Immunol* 2008; **180**: 7423–30.
- 131 Yang XO, Pappu BP, Nurieva R, et al. T helper 17 lineage differentiation is programmed by orphan nuclear receptors ROR alpha and ROR gamma. *Immunity* 2008; **28**: 29–39.
- 132 Mathur AN, Chang H-C, Zisoulis DG, et al. Stat3 and Stat4 direct development of IL-17-secreting Th cells. *J Immunol* 2007; **178**: 4901–07.
- 133 Lewkowich IP, Herman NS, Schleifer KW, et al. CD4+CD25+ T cells protect against experimentally induced asthma and alter pulmonary dendritic cell phenotype and function. *J Exp Med* 2005; **202**: 1549–61.
- 134 Yang XO, Chang SH, Park H, et al. Regulation of inflammatory responses by IL-17F. *J Exp Med* 2008; **205**: 1063–75.
- 135 Krishnamoorthy N, Oriss TB, Paglia M, et al. Activation of c-Kit in dendritic cells regulates T helper cell differentiation and allergic asthma. *Nat Med* 2008; **14**: 565–73.

- 136 Kim EY, Battaile JT, Patel AC, et al. Persistent activation of an innate immune response translates respiratory viral infection into chronic lung disease. *Nat Med* 2008; **14**: 633–40.
- 137 Gualano RC, Hansen MJ, Vlahos R, et al. Cigarette smoke worsens lung inflammation and impairs resolution of influenza infection in mice. *Respir Res* 2008; **9**: 53.
- 138 Kang MJ, Lee CG, Lee JY, et al. Cigarette smoke selectively enhances viral PAMP- and virus-induced pulmonary innate immune and remodeling responses in mice. *J Clin Invest* 2008; **118**: 2771–84.
- 139 Hayashi N, Yoshimoto T, Izuhara K, Matsui K, Tanaka T, Nakanishi K. T helper 1 cells stimulated with ovalbumin and IL-18 induce airway hyperresponsiveness and lung fibrosis by IFN- γ and IL-13 production. *Proc Natl Acad Sci USA* 2007; **104**: 14765–70.
- 140 Kondo Y, Yoshimoto T, Yasuda K, et al. Administration of IL-33 induces airway hyperresponsiveness and goblet cell hyperplasia in the lungs in the absence of adaptive immune system. *Int Immunol* 2008; **20**: 791–800.
- 141 Muller U, Stenzel W, Kohler G, et al. IL-13 induces disease-promoting type 2 cytokines, alternatively activated macrophages and allergic inflammation during pulmonary infection of mice with *Cryptococcus neoformans*. *J Immunol* 2007; **179**: 5367–77.
- 142 Tiemessen MM, Jagger AL, Evans HG, van Herwijnen MJC, John S, Taams LS. CD4⁺CD25⁺Foxp3⁺ regulatory T cells induce alternative activation of human monocytes/macrophages. *Proc Natl Acad Sci USA* 2007; **104**: 19446–51.
- 143 Chupp GL, Lee CG, Jarjour N, et al. A chitinase-like protein in the lung and circulation of patients with severe asthma. *N Engl J Med* 2007; **357**: 2016–27.
- 144 Oh SY, Zheng T, Bailey ML, et al. Src homology 2 domain-containing inositol 5-phosphatase 1 deficiency leads to a spontaneous allergic inflammation in the murine lung. *J Allergy Clin Immunol* 2007; **119**: 123–31.
- 145 Sly LM, Ho V, Antignano F, et al. The role of SHIP in macrophages. *Front Biosci* 2007; **12**: 2836–48.
- 146 Rauh MJ, Ho V, Pereira C, et al. SHIP represses the generation of alternatively activated macrophages. *Immunity* 2005; **23**: 361–74.
- 147 Ernst M, Inglese M, Scholz GM, et al. Constitutive activation of the SRC family kinase Hck results in spontaneous pulmonary inflammation and an enhanced innate immune response. *J Exp Med* 2002; **196**: 589–604.
- 148 Beavitt SJ, Harder KW, Kemp JM, et al. Lyn-deficient mice develop severe, persistent asthma: Lyn is a critical negative regulator of Th2 immunity. *J Immunol* 2005; **175**: 1867–75.
- 149 Munitz A, Brandt EB, Mingler M, Finkelman FD, Rothenberg ME. Distinct roles for IL-13 and IL-4 via IL-13 receptor $\alpha 1$ and the type II IL-4 receptor in asthma pathogenesis. *Proc Natl Acad Sci USA* 2008; **105**: 7240–45.
- 150 Meyers DA, Larj MJ, Lange L. Genetics of asthma and COPD. Similar results for different phenotypes. *Chest* 2004; **26** (suppl 2): 105S–10S.