

- 6 Illi S, von Mutius E, Lau S, Niggemann B, Gruber C, Wahn U, on behalf of the Multicentre Allergy Study (MAS) group. Perennial allergen sensitisation early in life and chronic asthma in children: a birth cohort study. *Lancet* 2006; **368**: 763–70.
- 7 Kusel MM, de Klerk NH, Keadze T, et al. Early-life respiratory viral infections, atopic sensitization, and risk of subsequent development of persistent asthma. *J Allergy Clin Immunol* 2007; **119**: 1105–10.
- 8 Bonness S, Bieber T. Molecular basis of atopic dermatitis. *Curr Opin Allergy Clin Immunol* 2007; **7**: 382–86.
- 9 Weidinger S, O'Sullivan M, Illig T, et al. Filaggrin mutations, atopic eczema, hay fever, and asthma in children. *J Allergy Clin Immunol* 2008; **121**: 1203–09.
- 10 Novembre E, Galli E, Landi F, et al. Coseasonal sublingual immunotherapy reduces the development of asthma in children with allergic rhinoconjunctivitis. *J Allergy Clin Immunol* 2004; **114**: 851–57.
- 11 Moller C, Dreborg S, Ferdousi HA, et al. Pollen immunotherapy reduces the development of asthma in children with seasonal rhinoconjunctivitis (the PAT study). *J Allergy Clin Immunol* 2002; **109**: 251–56.
- 12 Niggemann B, Jacobsen L, Dreborg S, et al, the PAT Investigator Group. Five-year follow-up on the PAT study: specific immunotherapy and long-term prevention of asthma in children. *Allergy* 2006; **61**: 855–59.
- 13 Jacobsen L, Niggemann B, Dreborg S, et al, the PAT Investigator Group. Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy* 2007; **62**: 943–48.
- 14 Grembale RD, Camporota L, Naty S, Tranfa CM, Djukanovic R, Marsico SA. Effects of specific immunotherapy in allergic rhinitic individuals with bronchial hyperresponsiveness. *Am J Respir Crit Care Med* 2000; **162**: 2048–52.

## Transition from childhood to adult asthma

See [Articles](#) page 1058


If a child has more than one episode of bronchitis in infancy, parents and doctors are concerned about whether the child will outgrow the disease or develop chronic obstructive airway disease—ie, asthma. In today's *Lancet*, Debra Stern and colleagues<sup>1</sup> report data on the prediction of adult asthma at 22 years of age by evaluation of early-life factors at 6 years of age in the Tucson Children's Respiratory Study, one of the oldest birth-cohort studies. Longitudinal studies analyse time courses in disease development and progression and can find relations between exposure factors and outcome (figure).<sup>2</sup>

Although no real cure for asthma exists, early identification of patients at risk of disease progression could lead to better treatment opportunities and, hopefully, improved outcomes in adulthood. Nevertheless, early intervention has not yet been shown to be beneficial in terms of disease modification, which might be because of the failure to identify appropriate subgroups of responders. Studies during early childhood

have not shown a clear benefit of early anti-inflammatory treatment in terms of symptom-free days after the treatment period in preschool children.<sup>3</sup> However, inhaled steroids seem to reduce the accelerated decline in forced expiratory volume in 1 s and airway remodelling in adulthood.<sup>4</sup> A kind of tracking (those who start low, remain low) in terms of lung function seems to exist,<sup>5,6</sup> and children with wheezing disorder before 2 years of age compared with those who started wheezing at school age and girls compared with boys probably have poorer lung function in early adulthood.<sup>7</sup>

Cohort studies have tried to identify different wheezing phenotypes, which were mainly defined by onset and prolongation of symptoms (early onset, late onset, persistent) and presence and absence of atopy.<sup>8,9</sup> One study used latent class analysis, a statistical method developed for the social sciences to identify distinct subsets or classes underlying the observed heterogeneity in a population, to distinguish phenotypes of childhood wheeze and cough.<sup>10</sup> The investigators described five phenotypes: persistent and transient cough, atopic and non-atopic persistent wheeze, and transient viral wheeze. The most common phenotype was atopic wheeze at 8 years of age; however, no prediction for adulthood has been made so far.

In an unselected birth cohort study from New Zealand, 25% of the children had wheezing that persisted from childhood to adulthood or that relapsed after remission.<sup>11</sup> Factors predicting persistence or relapse were sensitisation to house-dust mites, airway hyper-responsiveness, female sex, smoking, and early age at onset. Nevertheless, prediction of outcome seems to be difficult for childhood asthma.



**Figure: Incidence of childhood asthma**  
Data are from German MAS study.<sup>2</sup>

In their new paper, Stern and colleagues try to define phenotypes of asthma with special features early in life, favouring ongoing airway disease later in life on the one hand and characterisation of patients with asthma who developed the disease after puberty on the other. Most cases with current asthma in early adult life already had episodes of wheezing during the first 3 years of life. In 27% of the individuals with asthma at 22 years, asthma was newly diagnosed and most of them were female (71%). Predictive factors for chronic asthma in adulthood were late onset of wheezing (6 years and older), persistent wheeze in early life (before 3 years, and at 6 years and later), sensitisation to the fungus *Alternaria alternata*, and low airway-function and bronchial hyper-responsiveness to cold dry air at 6 years. These findings identify a population at risk of chronic obstructive airway disease in early adulthood, and they already showed a predisposition during preschool years. Whether therapeutic approaches at early preschool age can affect progression of the disease is yet to be established.

Susanne Lau

Department of Paediatric Pneumology and Immunology,  
Charité University Medicine, 13353 Berlin, Germany  
Susanne.lau@charite.de

I declare that I have no conflict of interest.

- 1 Stern DA, Morgan WJ, Halonen M, Wright AL, Martinez FD. Wheezing and bronchial hyper-responsiveness in early childhood as predictors of newly diagnosed asthma in early adult life: a longitudinal birth-cohort study. *Lancet* 2008; **372**: 1058–64.
- 2 Matricardi PM, Illi S, Grüber C, et al. Wheezing in childhood: incidence, longitudinal patterns and factors predicting persistence. *Eur Respir J* 2008; **32**: 585–92.
- 3 Guilbert TW, Morgan WJ, Zeiger RS, et al. Long-term inhaled corticosteroids in preschool children at high risk for asthma. *N Engl J Med* 2006; **117**: 1985–97.
- 4 Dijkstra A, Vonk JM, Jongepier H, et al. Lung function decline in asthma: association with inhaled corticosteroids, smoking and sex. *Thorax* 2006; **61**: 105–10.
- 5 Morgan WJ, Stern DA, Sherill DL, et al. Outcome of asthma and wheezing in the first 6 years of life: follow-up through adolescence. *Am J Respir Crit Care Med* 2005; **172**: 1253–58.
- 6 Håland G, Carlsen KC, Sandvik L, et al. Reduced lung function at birth and the risk of asthma at 10 years of age. *N Engl J Med* 2006; **355**: 1682–89.
- 7 Goksör E, Gustafsson PM, Alm B, Amak M, Wennergren G. Reduced airway function in early adulthood among subjects with wheezing disorders before two years of age. *Pediatr Pulmonol* 2008; **43**: 396–403.
- 8 Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995; **332**: 133–38.
- 9 Stein RT, Holberg CJ, Morgan WJ, et al. Peak flow variability, methacholine responsiveness and atopy as markers for detecting different wheezing phenotypes in childhood. *Thorax* 1997; **52**: 946–52.
- 10 Spycher B, Silverman M, Brooke AM, Minder CE, Kuehni CE. Distinguishing phenotypes of childhood wheeze and cough using latent class analysis. *Eur Respir J* 2008; **31**: 974–81.
- 11 Sears MR, Greene JM, Willan AR, et al. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med* 2003; **349**: 1414–22.

## Exhaled nitric oxide in guideline-based asthma management

The importance of good asthma control has been amply documented and new guidelines focus on achievement and maintenance of good asthma control.<sup>1</sup> However, although validated scoring systems have been developed to facilitate asthma control, several studies have found that assessment of asthma control varies markedly between health-care professionals as well as patients.<sup>2</sup> The result is high risk of too little or too much treatment, or both. Therefore the medical community has been hoping for a single, easy-to-measure, and reliable biomarker (such as haemoglobin A<sub>1c</sub> in diabetes) that could facilitate the assessment of asthma control and help physicians to appropriately increase or decrease treatment. In this respect, exhaled nitric oxide (FE<sub>NO</sub>)—an indirect marker of airway inflammation—has generated much enthusiasm. FE<sub>NO</sub> is easy to measure, correlates with eosinophilic airway inflammation, and is increased during periods of uncontrolled asthma and reduced during treatment

with anti-inflammatory agents.<sup>3–5</sup> However, before a new biomarker is implemented in daily practice, it should be assessed in light of the words of Albert Einstein: “What counts cannot always be measured and what can be measured does not always count.”

The measurement of FE<sub>NO</sub> is relatively easy and has shown promising results, but is not cheap. Therefore the important question is: does addition of FE<sub>NO</sub> to guideline-recommended clinical assessments and treatment algorithms reduce over-treatment or under-treatment (or both), and does it improve daily asthma control? In today's *Lancet*, Stanley Szefer and colleagues report the findings of a large well-conducted study designed to address these questions.<sup>6</sup> The investigators conclude that adding measurements of FE<sub>NO</sub> to guideline-recommended asthma management did not provide any important clinical benefit. No differences were found in clinical scores for asthma control or other asthma

See [Comment](#) page 1017

See [Articles](#) page 1065