

In their new paper, Stern and colleagues try to define phenotypes of asthma with special features early in life, favouring ongoing airway disease later in life on the one hand and characterisation of patients with asthma who developed the disease after puberty on the other. Most cases with current asthma in early adult life already had episodes of wheezing during the first 3 years of life. In 27% of the individuals with asthma at 22 years, asthma was newly diagnosed and most of them were female (71%). Predictive factors for chronic asthma in adulthood were late onset of wheezing (6 years and older), persistent wheeze in early life (before 3 years, and at 6 years and later), sensitisation to the fungus *Alternaria alternata*, and low airway-function and bronchial hyper-responsiveness to cold dry air at 6 years. These findings identify a population at risk of chronic obstructive airway disease in early adulthood, and they already showed a predisposition during preschool years. Whether therapeutic approaches at early preschool age can affect progression of the disease is yet to be established.

Susanne Lau

Department of Paediatric Pneumology and Immunology,  
Charité University Medicine, 13353 Berlin, Germany  
Susanne.lau@charite.de

I declare that I have no conflict of interest.

- 1 Stern DA, Morgan WJ, Halonen M, Wright AL, Martinez FD. Wheezing and bronchial hyper-responsiveness in early childhood as predictors of newly diagnosed asthma in early adult life: a longitudinal birth-cohort study. *Lancet* 2008; **372**: 1058–64.
- 2 Matricardi PM, Illi S, Grüber C, et al. Wheezing in childhood: incidence, longitudinal patterns and factors predicting persistence. *Eur Respir J* 2008; **32**: 585–92.
- 3 Guilbert TW, Morgan WJ, Zeiger RS, et al. Long-term inhaled corticosteroids in preschool children at high risk for asthma. *N Engl J Med* 2006; **117**: 1985–97.
- 4 Dijkstra A, Vonk JM, Jongepier H, et al. Lung function decline in asthma: association with inhaled corticosteroids, smoking and sex. *Thorax* 2006; **61**: 105–10.
- 5 Morgan WJ, Stern DA, Sherill DL, et al. Outcome of asthma and wheezing in the first 6 years of life: follow-up through adolescence. *Am J Respir Crit Care Med* 2005; **172**: 1253–58.
- 6 Håland G, Carlsen KC, Sandvik L, et al. Reduced lung function at birth and the risk of asthma at 10 years of age. *N Engl J Med* 2006; **355**: 1682–89.
- 7 Goksör E, Gustafsson PM, Alm B, Amak M, Wennergren G. Reduced airway function in early adulthood among subjects with wheezing disorders before two years of age. *Pediatr Pulmonol* 2008; **43**: 396–403.
- 8 Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995; **332**: 133–38.
- 9 Stein RT, Holberg CJ, Morgan WJ, et al. Peak flow variability, methacholine responsiveness and atopy as markers for detecting different wheezing phenotypes in childhood. *Thorax* 1997; **52**: 946–52.
- 10 Spycher B, Silverman M, Brooke AM, Minder CE, Kuehni CE. Distinguishing phenotypes of childhood wheeze and cough using latent class analysis. *Eur Respir J* 2008; **31**: 974–81.
- 11 Sears MR, Greene JM, Willan AR, et al. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med* 2003; **349**: 1414–22.

## Exhaled nitric oxide in guideline-based asthma management

The importance of good asthma control has been amply documented and new guidelines focus on achievement and maintenance of good asthma control.<sup>1</sup> However, although validated scoring systems have been developed to facilitate asthma control, several studies have found that assessment of asthma control varies markedly between health-care professionals as well as patients.<sup>2</sup> The result is high risk of too little or too much treatment, or both. Therefore the medical community has been hoping for a single, easy-to-measure, and reliable biomarker (such as haemoglobin A<sub>1c</sub> in diabetes) that could facilitate the assessment of asthma control and help physicians to appropriately increase or decrease treatment. In this respect, exhaled nitric oxide (FE<sub>NO</sub>)—an indirect marker of airway inflammation—has generated much enthusiasm. FE<sub>NO</sub> is easy to measure, correlates with eosinophilic airway inflammation, and is increased during periods of uncontrolled asthma and reduced during treatment

with anti-inflammatory agents.<sup>3–5</sup> However, before a new biomarker is implemented in daily practice, it should be assessed in light of the words of Albert Einstein: “What counts cannot always be measured and what can be measured does not always count.”

The measurement of FE<sub>NO</sub> is relatively easy and has shown promising results, but is not cheap. Therefore the important question is: does addition of FE<sub>NO</sub> to guideline-recommended clinical assessments and treatment algorithms reduce over-treatment or under-treatment (or both), and does it improve daily asthma control? In today's *Lancet*, Stanley Szefer and colleagues report the findings of a large well-conducted study designed to address these questions.<sup>6</sup> The investigators conclude that adding measurements of FE<sub>NO</sub> to guideline-recommended asthma management did not provide any important clinical benefit. No differences were found in clinical scores for asthma control or other asthma

See [Comment](#) page 1017

See [Articles](#) page 1065



Science Photo Library

outcomes, and the occurrence of under-treatment was not improved. These findings agree with the results from smaller trials.<sup>7-10</sup> So, it seems unlikely that adding FE<sub>NO</sub> measurement to guideline-recommended asthma management is going to improve asthma control or reduce under-treatment. This finding is highly relevant to daily practice.

Even if daily asthma control was not improved by the use of FE<sub>NO</sub>, the measurement might still have improved individualised dosing, such that the same level of asthma control could be achieved with less treatment. However, Szeffler and colleagues did not see that result. By contrast, the use of FE<sub>NO</sub> resulted in higher doses of inhaled corticosteroids and more frequent use of long-acting  $\beta_2$  agonists. At first glance, this finding could lead to the conclusion that, in the population studied, use of FE<sub>NO</sub> to help manage asthma could lead to over-treatment because of the use of a "measure that did not count". This finding could also be due to the design of the study. Although the study followed the recommendations of guidelines to regularly assess asthma-control status and tailor and adjust treatment in regular cycles, the approach did not use FE<sub>NO</sub> concentrations to reduce the amount of treatment. FE<sub>NO</sub> measurement was mainly used to increase the amount of treatment in patients with increased concentrations, even if clinical variables were stable. The conclusions in earlier trials about the potentially treatment-saving effects of adding FE<sub>NO</sub> to clinical assessment have varied from no change in the daily dose of inhaled corticosteroid,<sup>9,10</sup> to an increase,<sup>8</sup> and to a reduction in daily dose of inhaled corticosteroid.<sup>7</sup>

Although Szeffler and colleagues' study was not designed to assess the value of the guideline recommendations, the substantial clinical benefits achieved during the run-in period, when guideline-recommended treatment was introduced, agree with findings from other studies.<sup>11,12</sup> This finding emphasises, once again, the importance of even low doses of effective anti-inflammatory drugs in asthma management.

It takes more than one good study to completely rule out potential benefits of a new biomarker. Although Szeffler and colleagues' findings convincingly show that addition of FE<sub>NO</sub> measurement to guideline-recommended asthma management will not improve daily asthma control or under-treatment in the population studied, more studies are needed to explore the usefulness of FE<sub>NO</sub> in subgroups of patients with asthma and its treatment-saving potential. Until such data are available, a recommendation to use FE<sub>NO</sub> measurements routinely in patients treated according to guidelines is not ready to be made yet.

\*Søren Pedersen, Paul M O'Byrne

Paediatric Research Unit, Kolding Hospital, DK 6000 Kolding, Denmark (SP); and Department of Medicine, Michael G DeGroot School of Medicine, McMaster University, Hamilton, ON, Canada (PMO'B)

spconsult@post1.tele.dk

SP has received research funding, consultancies, and lecture fees from Nycomed, AstraZeneca, and GlaxoSmithKline. PMO'B has been a member of advisory boards for AstraZeneca, Biopix, GlaxoSmithKline, Merck, Nycomed, Topigen, Resistentia, and Wyeth, has received lecture fees from AstraZeneca, Chiesi, GlaxoSmithKline, and Nycomed, and grants for research from AstraZeneca, Boehringer Genentech, GlaxoSmithKline, Medimmune, Merck, Nycomed, Pfizer, and Wyeth.

- 1 Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J* 2008; **31**: 143-78.
- 2 Rabe KF, Vermeire PA, Soriano JB, Maier WC. Clinical management of asthma in 1999: the Asthma Insights and Reality in Europe (AIRE) study. *Eur Respir J* 2000; **16**: 802-07.
- 3 Silkoff PE, Carlson M, Bourke T, Katial R, Ogren E, Szeffler SJ. The Aerocrine exhaled nitric oxide monitoring system NIOX is cleared by the US Food and Drug Administration for monitoring therapy in asthma. *J Allergy Clin Immunol* 2004; **114**: 1241-56.
- 4 Silkoff PE, McClean P, Spino M, Erlich L, Slutsky AS, Zamel N. Dose-response relationship and reproducibility of the fall in exhaled nitric oxide after inhaled beclomethasone dipropionate therapy in asthma patients. *Chest* 2001; **119**: 1322-28.
- 5 Stirling RG, Kharitonov SA, Campbell D, et al, for the Asthma and Allergy Group. Increase in exhaled nitric oxide levels in patients with difficult asthma and correlation with symptoms and disease severity despite treatment with oral and inhaled corticosteroids. *Thorax* 1998; **53**: 1030-34.
- 6 Szeffler SJ, Mitchell H, Sorkness CA, et al. Management of asthma based on exhaled nitric oxide in addition to guideline-based treatment for inner-city adolescents and young adults: a randomised controlled trial. *Lancet* 2008; **372**: 1065-72.
- 7 Smith AD, Cowan JO, Brassett KP, Herbison GP, Taylor DR. Use of exhaled nitric oxide measurements to guide treatment in chronic asthma. *N Engl J Med* 2005; **352**: 2163-73.

- 8 Fritsch M, Uxa S, Horak F Jr, et al. Exhaled nitric oxide in the management of childhood asthma: a prospective 6-months study. *Pediatr Pulmonol* 2006; **41**: 855–62.
- 9 Pijnenburg MW, Bakker EM, Hop WC, de Jongste JC. Titrating steroids on exhaled nitric oxide in children with asthma: a randomized controlled trial. *Am J Respir Crit Care Med* 2005; **172**: 831–36.
- 10 Shaw DE, Berry MA, Thomas M, et al. The use of exhaled nitric oxide to guide asthma management: a randomized controlled trial. *Am J Respir Crit Care Med* 2007; **176**: 231–37.
- 11 Bateman ED, Boushey HA, Bousquet J, et al, for the GOAL Investigators Group. Can guideline-defined asthma control be achieved? The Gaining Optimal Asthma Control study. *Am J Respir Crit Care Med* 2004; **170**: 836–44.
- 12 Pauwels RA, Pedersen S, Busse WW, et al, on behalf of the START Investigators Group. Early intervention with budesonide in mild persistent asthma: a randomised, double-blind trial. *Lancet* 2003; **361**: 1071–76.

## Inflammometry to assess airway diseases

The diagnostic labels used to characterise common airway diseases have always been a problem. The term asthma implies the presence of variable airflow obstruction; however, objective demonstration can be difficult. Commonly used tests, such as spirometry or serial peak-flow measurements, are neither sensitive nor specific,<sup>1</sup> especially in patients with mild disease and normal or near-normal lung function, or in those with fixed airflow obstruction. Conversely, the term chronic obstructive pulmonary disease (COPD) implies largely irreversible airflow obstruction, yet clinicians may attempt to confirm the presence of reversibility, and having done so, label the disease as having an “asthmatic” component. Furthermore, the use of specific diagnostic labels implies a probable natural history, and influences expectations about treatment outcomes.

Although causative factors in the various airway diseases are different, the natural history and treatment responsiveness are less distinct. For example, accelerated decline in lung function and fixed airflow obstruction are features of COPD, but not exclusively so; they also occur in some patients with asthma, and the mechanisms might be similar.<sup>2</sup> In view of this picture, diagnostic labelling on the basis of physiological data can be confusing and misleading.

Treatment with corticosteroids, usually by inhalation, is arguably the most important therapeutic intervention in patients with airway disease. Judicious use of these agents improves symptoms and reduces exacerbations. However, the response to inhaled corticosteroids is heterogeneous. The identification of clinical or physiological features that predict corticosteroid responsiveness in patients with symptoms that suggest airway disease is difficult, irrespective of the final diagnosis. A common approach is to base long-term use of corticosteroid on the response to a short-term trial of treatment. This approach is potentially flawed for several reasons.

First, symptoms that suggest asthma are non-specific, and are mimicked by acute conditions such as postviral bronchial hyper-responsiveness, anxiety hyperventilation syndrome, vocal cord dysfunction, and gastro-oesophageal reflux, and by chronic conditions such as COPD or bronchiectasis. Most of these conditions do not respond to corticosteroids, but spontaneous improvement over time leads to the mistaken belief that such treatment has been beneficial. The correct diagnosis is thus delayed, or inappropriate treatment might be increased when symptoms worsen. Second, it is not valid to draw inferences about the longer-term benefits of treatment (ie, reduction in exacerbation frequency) from the outcome of a short-term trial. Third, expectation, observer or ascertainment biases, and incomplete adherence to the prescribed treatment can also influence results. Most of these problems, together with the natural tendency of clinicians to be cautious in borderline cases, increase the likelihood that patients may be started on inappropriate corticosteroid therapy, with associated cost and potential toxicity.

An alternative approach is to identify the need for corticosteroids in relation to the underlying inflammation. It is logical that both the indications and the outcomes for anti-inflammatory treatment should be related to the presence of airway inflammation. There is now consistent evidence that eosinophilic airway inflammation is the most reliable predictor of a response to corticosteroids in patients with airway disease,<sup>3–7</sup> irrespective of which diagnostic label applies. The long-term benefits of corticosteroids on exacerbation frequency also occur predominantly in patients with evidence of eosinophilic airway inflammation.<sup>8–10</sup> Management strategies that seek to minimise eosinophilic airway inflammation substantially reduce the frequency of severe exacerbations, both in asthma<sup>8,9</sup> and COPD,<sup>10</sup> and hence result in more efficient use of corticosteroids.

See [Comment](#) page 1015