

New targets for drug development in asthma

Ian M Adcock, Gaetano Caramori, K Fan Chung

Asthma is a chronic inflammatory disease that affects about 300 million people worldwide, a total that is expected to rise to about 400 million over the next 15–20 years. Most asthmatic individuals respond well to the currently available treatments of inhaled corticosteroids and β -adrenergic agonists; however, 5–10% have severe disease that responds poorly. Improved knowledge of asthma mechanisms has led to the recognition of different asthma phenotypes that might reflect distinct types of inflammation, explaining the effectiveness of anti-leucotrienes and the anti-IgE monoclonal antibody omalizumab in some patients. However, more knowledge of the inflammatory mechanisms within the airways is required. Improvements in available therapies—such as the development of fast-onset, once-a-day combination drugs with better safety profiles—will occur. Other drugs, such as inhaled p38 MAPK inhibitors and anti-oxidants, that target specific pathways or mediators could prove useful as monotherapies, but could also, in combination with corticosteroids, reduce the corticosteroid insensitivity often seen in severe asthma. Biological agents directed against the interleukin-13 pathway and new immunoregulatory agents that modulate functions of T-regulatory and T-helper-17 cells are likely to be successful. Patient-specific treatments will depend on the development of discriminatory handprints of distinct asthma subtypes and are probably over the horizon. Although a cure is unlikely to be developed in the near future, a greater understanding of disease mechanisms could bring such a situation nearer to reality.

Introduction

Asthma is one of the most common chronic inflammatory diseases, affecting about 300 million people worldwide, a total that is expected to rise by an additional 100 million—mainly in children—over the next 15–20 years.¹ Asthma accounts for about one out of every 250 deaths worldwide and has profound health-care costs in terms of emergency room visits and hospitalisations. Asthma also has enormous indirect costs and is one of the leading causes of work and school absenteeism.² Most patients with asthma respond well to current treatments; however, 5–10% of patients have severe disease that often fails to respond to conventional therapy; these patients account for more than 50% of the total health-care costs associated with asthma.³

By understanding the different types of airway inflammation in various subtypes of asthma, it should be possible to address some of the important questions in asthma research: which triggers or factors underlie airway smooth muscle hyper-responsiveness? What are the processes (genetic or environmental) that underlie different subtypes of asthma? Which aspects of airway remodelling are important in disease subtypes? What are the best biomarkers of disease progression or treatment response? Why are some patients less responsive to conventional therapies than are others? It is now recognised that there are distinct asthma phenotypes⁴ and that distinct therapeutic approaches may only impinge on some aspects of the disease process, or at least outcome measures, within each subgroup. Thus, treatments might affect exacerbation rates without altering day-to-day symptoms or lung function, reflecting the fact that distinct cells or mediators in the lung could drive airway hyper-responsiveness or specific inflammatory components, and that treatments directed at a single cell or mediator

might only affect a single aspect of disease.⁵ The current reductionist approach to understanding the disease has led to the development of drugs that target specific pathways or mediators. In the future, we may need to target and assess several outcome measures and biomarkers simultaneously, and undertake subgroup analyses of the responses obtained when assessing combinations of new drugs.

Current asthma treatments

Initial approaches to treat asthma emphasised the relief of bronchoconstriction with bronchodilators, particularly β_2 -adrenergic agonists, but the discovery of airway inflammation as an important pathophysiological component of asthma has led to the use of inhaled corticosteroids as the mainstay of asthma therapy⁶—these drugs are the most effective anti-inflammatory treatment available for asthma. Asthmatic inflammation is characterised by eosinophilia, mast cell infiltration,

Lancet 2008; 372: 1073–87

Airways Disease Section, National Heart and Lung Institute, Imperial College London, London, UK (Prof I M Adcock PhD, Prof K F Chung DSc); MRC-Asthma UK Centre for Allergic Mechanisms in Asthma, Kings and Imperial College, London, UK (Prof I M Adcock, Prof K F Chung); and Centre of Research on Asthma and COPD, University of Ferrara, Ferrara, Italy (G Caramori MD)

Correspondence to: Prof Ian M Adcock, Airways Disease Section, National Heart and Lung Institute, Imperial College London, Dovehouse Street, London SW3 6LY, UK Ian.Adcock@imperial.ac.uk

Search strategy and selection criteria

We undertook a detailed appraisal of peer-reviewed publications over the past 10 years with the NCBI PubMed website for English language publications with the keywords: “asthma”, in combination with “treatment”, “novel therapy”, “glucocorticoid insensitivity”, “new drugs”, “steroid-sparing”, “severe”, “immunomodulation”, and “unmet need”. We also searched the reference lists of articles identified by this search strategy and selected those we judged relevant. We also had source publications that we have accumulated because of our association with asthma therapy and research in the past 15 years. Review articles are cited to provide readers with more details and more references than this Review has room for. Our reference list was modified on the basis of comments from peer reviewers.




Figure 1: Potential targets for new asthma drugs

Even before asthma becomes symptomatic, exposure to an allergen produces structural changes to the airways, including subepithelial fibrosis and epithelial denudation. Inhaled allergens encounter antigen-presenting cells (dendritic cells, macrophages) that line the airways and mast cells which release mediators, such as histamine. The antigen-presenting cells migrate to the lymph nodes, where they induce the proliferation and activation of CD4 T lymphocytes (Th2 cells). These cells express a variety of cytokines and chemokines, under the control of intracellular kinases and downstream transcription factors, which lead to the differentiation, migration, and pathobiological effects of eosinophils. These inflammatory processes lead to pathological effects, such as airway hyper-responsiveness and airway remodelling including vasodilation and angiogenesis. Neutrophils are increasingly important in severe asthma. As such asthma is a highly complex disease that involves many inflammatory cells, mediators, inflammatory proteins, and intracellular pathways. ICAM=intercellular adhesion molecule. JNK=c-Jun N-terminal kinase. MAPK=mitogen activated protein kinase. NFκB=nuclear factor κB. NF-AT=nuclear factor activated T cells. PDE=phosphodiesterase. PI3K=phosphoinositide-3 kinase. PPAR=peroxisome proliferator-activated receptor. STAT=signal transducer and activator of transcription. TNF=tumour necrosis factor. Treg=T-regulatory cells. VLA=very late antigen.

and activation of T-helper (Th) 2 cells that express interleukins 4, 5, and 13. Structural changes such as goblet cell hyperplasia, airway smooth muscle hypertrophy and hyperplasia, along with subepithelial fibrosis are also present, even in mild disease, although these features may be more prominent in severe disease (figure 1). The functional effects of these structural changes are unclear, but they might contribute to chronic airflow obstruction and reduced airway responses to bronchodilators or inhaled corticosteroids.⁷

Long-term treatment with inhaled corticosteroids reverses airflow obstruction, reduces exacerbations and the need for hospitalisation, and improves quality of life. Inhaled corticosteroids may have also contributed to the reduction in asthma-related deaths over the past few years.⁸ Concerns about the long-term detrimental effects of high-dose inhaled corticosteroids therapy include cataracts, osteoporosis in elderly patients, and stunting of growth in children,⁹ but there is a high safety margin for most patients on low to moderate doses of inhaled corticosteroids. Compliance with inhaled corticosteroids

is poor, possibly due to a misunderstanding of the side-effects and an inability to maintain a daily regime, especially when symptoms are well controlled.¹⁰ However, inhaled corticosteroids may not modify the progression of asthma and are not curative, because asthma symptoms and inflammation recur on discontinuation of treatment.¹¹

The effectiveness of inhaled corticosteroids—particularly at low to moderate doses—in controlling asthma and reducing exacerbations is improved by combination with long-acting β₂ agonists and is better than that shown by higher doses of inhaled corticosteroids alone.¹² Thus, combinations of inhaled corticosteroids and long-acting β₂ agonists are used in the treatment of moderate-to-severe asthma¹² and such combinations have now become established as the most efficacious treatment.¹ Because of the rapid action of formoterol compared with salmeterol, it is possible to use the combination of formoterol and budesonide or formoterol and beclomethasone as both maintenance and reliever medication. An alternative approach to asthma control

has been taken with the combination of salmeterol and fluticasone, which has been shown to achieve comprehensive asthma control in most patients.¹³ It must be emphasised that long-acting β_2 agonists are not recommended as monotherapy maintenance treatment for asthma.

The need for new therapies

Combinations of inhaled corticosteroids and long-acting β_2 agonists are effective in most (about 90%), but not all, asthmatic individuals.¹⁴ Indeed, even patients whose asthma is apparently well controlled by existing therapies might benefit from more efficacious therapies that are easier to comply with.¹¹ Surprisingly, telephone surveys of asthmatic patients have reported a high degree of morbidity among patients who report that their asthma was controlled,¹¹ which could reflect either a lack of compliance or that existing medications are not as efficacious in day-to-day practice as they are in controlled trials. Improved compliance in the mild-to-moderate asthmatic patient could be achieved by the development of safe oral versions of conventional treatments or of new, more efficacious treatments, particularly if these agents altered the course of the disease or pointed towards a cure.

The drive to find new drug targets has led to the introduction of two new classes of anti-mediator agents that are now in clinical practice, namely the leucotriene inhibitors and anti-IgE monoclonal antibodies, which are used in mild-to-moderate and severe allergic asthma, respectively.¹ The development of these drugs reflect the emphasis on specific anti-mediator agents over the past two decades and the discovery that many classes of mediators are seemingly not important in the pathogenesis of asthma. However, there has been a widening of potential targets for asthma treatments, which now include inflammatory and immune cytokines and chemokines, transcription factors, enzymes, and immune cells (figure 1).⁴

Furthermore, different subtypes of asthma occur in children and in adults and even within children and adults,¹⁵ resulting in distinct patterns of treatment sensitivity. A complete analysis of each patient's genome, proteome, and kinome is beyond current technology but could eventually lead to patient-centred rational therapy and highlights why some treatments are only effective in small subsets of patients.

Patients with poorly controlled severe asthma who do not respond well to combination therapy are another major unmet need. One characteristic of this group is relative corticosteroid insensitivity.⁷ These patients may have corticosteroid-dependent, or more rarely, corticosteroid-resistant asthma.⁷ These corticosteroid-resistant patients are a subset of those patients with severe asthma, but one must note that these terms are not interchangeable because some corticosteroid-resistant patients do not have severe disease and some patients with severe asthma

are not treatment-insensitive.¹⁶ The US Severe Asthma Research Program (SARP)¹⁶ and the European Network For Understanding Mechanisms Of Severe Asthma (ENFUMOSA)¹⁷ have highlighted the need to understand the molecular and cellular mechanisms present in severe asthma to enable the identification of novel targets for asthma and to speed up the drug development process.¹⁸ Clarification of the mechanisms that relate to disease subtypes in corticosteroid-resistant patients might not only indicate selective responsiveness to novel therapies in severe asthma but also point to similar subtypes and treatment responsiveness in patients with mild or moderate asthma.

It has been observed recently that the thickness of the airway epithelium, of the sub-basement membrane, and of the airway smooth muscle in the airways of patients with severe asthma is greater than in those with non-severe asthma.^{19,20} These differences were associated with altered expression of markers of epithelial proliferation.¹⁹ Simultaneously, a number of unbiased techniques such as hierarchical clustering of bronchoalveolar lavage cytokine expression,²¹ cluster analysis of inflammation and responses to treatment,²² the analysis of volatile organic components of exhaled breath using an electronic nose,²³ and the analysis of bronchial biopsy samples²⁴ have supported the possibility of several distinct phenotypes of patients with severe asthma. Targeting nodal points in these clusters may highlight potential novel sites for drug intervention.

Improvement of current therapies

New longer-acting bronchodilators

The long-acting β_2 agonists salmeterol and formoterol have a 12-h bronchodilator effect and, in conjunction with inhaled corticosteroids, improve asthma control and reduce exacerbation rates.²⁵ Several ultra-long-acting β_2 agonists that are under development, including indacaterol, carmoterol (Chiesi Farmaceutici, Italy), and GSK159797, act for more than 24 h, are fast acting, and are suitable for once-daily dosing.²⁶ It is also possible to increase the duration of action of inhaled corticosteroids which, in combination with once-a-day fast-onset β_2 agonists, will result in a combination therapy with prolonged action that, although possibly no more efficacious than current drugs, will improve compliance.²⁶ Although other bronchodilators exist—eg, vasoactive intestinal peptide, neuropeptide analogues, and K^+ -channel openers—these drugs have proved difficult to develop because of their potent vasodilator effects.²⁷

New corticosteroids with reduced systemic side-effects

Lung absorption of inhaled corticosteroids contributes to the systemic side-effects that are seen with such drugs, particularly with higher doses. Various means to reduce these systemic side-effects have been developed, including systemic or local inactivation, administration of an inactive prodrug that is only converted to active




Figure 2: Approaches to reduce the side-effects of inhaled corticosteroids

Several approaches are being taken to reduce systemic absorption including the use of pro-drugs such as ciclesonide, which is activated to release the active form des-ciclesonide only in the lungs, and the use of corticosteroids that are able to repress inflammatory gene expression without affecting the induction of corticosteroid-inducible genes, many of which are responsible for the side-effects. Alternatively, it may be possible to deliver monodispersed corticosteroid particles to specific parts of the airway using new delivery devices, thereby reducing the dose of inhaled drug required to obtain the equivalent clinical benefit.

drug in the airways (eg, ciclesonide²⁸), or targeted airway deposition using monodispersed particles (figure 2).²⁹ Another approach has been to develop dissociated corticosteroids that do not activate the mechanisms that lead to side-effects while preserving therapeutic anti-inflammatory effects. Many of the detrimental side-effects of corticosteroids have been linked to trans-activation (ie, activation of gene transcription through DNA binding of the activated glucocorticoid receptor) whereas most beneficial effects are associated with transrepression (ie, activated glucocorticoid receptor binding to other transcription factors rather than to DNA sites), although this mechanism requires confirmation in human beings.^{9,30,31} Dissociated corticosteroids may be just as effective as conventional inhaled corticosteroids, have a better safety profile, and might even lead to safer oral corticosteroids.³⁰ Linking the development of dissociated corticosteroids to drugs with a non-steroidal backbone such as AL-438 (Abbott Ligand) and ZK 216348 (Bayer-Schering Pharma) may further improve the therapeutic index, because these drugs will also lack activity at other nuclear hormone receptors which also cause side-effects.³² Lastly, the addition of an NO-donating group to prednisolone (NCX1015, NiCox Research Institute, Italy) and budesonide (NCX1020, NiCox Research Institute) has resulted in improved corticosteroid efficacy against lipopolysaccharide-induced inflammatory responses and prevented lipopolysaccharide-induced airway hyper-responsiveness, eosinophilia, and neutrophilia in guinea-pigs. By contrast, the parent compounds had only

a small effect on eosinophilia.^{33,34} In a similar manner, NO groups have been tagged to β_2 agonists.³⁵ The NO moiety is probably donated to specific residues within the respective ligand binding domains, thereby affecting receptor function.

Lipid mediator blockade

Many inflammatory mediators have been implicated in asthma and specific inhibitors and antagonists against many of these have been developed (figure 1, table). However, the only mediator antagonists that are currently used in asthma therapy are antileucotrienes, which block cysteinyl-leucotriene receptors or the synthesis of leucotriene.³⁶ Despite initial promise, inhibitors of other pro-inflammatory mediators, including histamine, prostaglandins, platelet-activating factor, bradykinin, and tachykinin, have proved ineffective.²⁷

Cysteinyl-leucotriene receptor antagonists, such as montelukast, improve lung function and asthma symptoms in patients with mild to moderate asthma, but their efficacy is reduced compared with low-dose inhaled corticosteroids. Moreover, such agents are less effective in severe asthma,³⁷ although a recent study suggests that they may be beneficial in asthmatic patients who smoke.³⁸ Zileuton, a 5-lipoxygenase inhibitor that blocks the synthesis of cysteinyl-leucotrienes and leucotriene B₄, provides similar efficacy to montelukast in asthma. These compounds need to be tested in neutrophilic asthma, because they would also inhibit neutrophilic chemotaxis and activation. Levels of lipoxin A₄, an endogenous anti-inflammatory eicosanoid, are reduced in severe

	Function	Drug and stage of development
β_2 adrenergic receptor	Ultra-long bronchodilation	Indacaterol (phase II), carmoterol (phase II), GSK159797 (phase II)
Glucocorticoid receptor	Anti-inflammatory	GSK685689 (phase II), GSK870086 (phase II), AL-438 (phase I), ZK 216348 (phase I)
PGD2/CRT2 inhibitors	Th2 cell recruitment and activation	TM30089, ODC9101 (phase II), AZD1981 (phase II), ramatroban (phase II)
BLT1 antagonist	Mononuclear/granulocyte recruitment	CP-105696 (phase I), LY293111 (phase II, no effect against allergen challenge)
CCL11	Blocks eosinophil recruitment/activation	CAT-213 (preclinical)
CCR3	Blocks eosinophil recruitment/activation	Met-RANTES (phase II, moderate/severe asthma)
CXCR4	Blocks Th2 activation	AMD070, AMD3100, SP01A (all preclinical for asthma, all phase II HIV, AMD3100 phase III for multiple myeloma)
CXCR1/2	Blocks neutrophil recruitment/activation	Repertaxin (preclinical, phase II for graft vs host disease)
Interleukin 5	Blocks eosinophil recruitment/activation	MEDI-563 (phase I, severe asthma), mepolizumab (phase II)
Interleukin 12	..	Interleukin 12 (phase II, no effect on lung function, adverse side-effects, not developed further)
Interleukin 10	Endogenous anti-inflammatory agent	Interleukin 10 (preclinical for asthma, approved for psoriasis/Crohn's disease, recruited in 1999 for asthma)
Interferon γ	..	Interferon γ (phase II, no effect on lung function in severe asthma, not developed further)
Interleukin 13	Key driver of asthmatic inflammation	Pitrakinra (interleukin-4/13 mutein), CAT-354, IMA-638 (both in phase II)
VLA4 antagonist	Adhesion molecule blocker	GW-559090, IVL745, CDP323 (CDP323 phase II, not developed)
PDE4	Anti-inflammatory	GSK256066 (phase II)
p38 MAPK	Anti-inflammatory	GSK681323, GSK856553, VX-745, BIRB-796, Ro-320-1195, Scio-469 (all in phase II), SB2439063, RWJ-67657
JNK	Anti-inflammatory	SP600125, CC-401, CNI-1493 (dual JNK/p38 MAPK) (all in preclinical for asthma; CC-401 and CNI-1493 in phase II in rheumatoid arthritis and Crohn's disease)
SYK	Mast cell degranulation, T-cell and B-cell function	Antisense (preclinical), BAY61-3606 (preclinical), R343 (phase I)
IKK2	Anti-inflammatory	AS206868, SC-514, BMS345541, TPCA-1 (all preclinical, MLN0415 [phase I])
CD23	Reduces IgE	Lumiliximab (phase I)
Sphingosine-1 phosphate receptor	Prevents dendritic cell activity	FTY720 (preclinical for asthma, Phase II for multiple sclerosis and transplant rejection)
DP1	Prevents dendritic cell activity	BW245C (preclinical)
VDR	Increased interleukin-10 expression in Treg cells	Vitamin D3 (phase II, steroid sparing)

Table: Summary of some of the compounds in development for asthma

asthma³⁹ and treatment with this compound might be considered as an anti-inflammatory agent for these patients.³⁹

Recently, the gene that encodes 5-lipoxygenase-activating protein (FLAP) has been linked to risk for myocardial infarction, stroke, and restenosis of the coronary artery. FLAP inhibitors showed some efficacy in early clinical trials in asthma but were not developed commercially because of efficacy and safety concerns, but new safer FLAP inhibitors such as DG031 (DeCode Genetics, Iceland) could be useful in patients with distinct genetic or phenotypic backgrounds, especially as a steroid add-on therapy in the treatment of asthma exacerbations.⁴⁰

Prostaglandin PGD₂, produced by mast cells, acts mainly through the G-protein-coupled receptor DP2 (also known as CRT2, chemoattractant receptor-homologous molecule expressed on Th2 lymphocytes) to mediate Th2 recruitment and activation.⁴¹ Ramatroban, a thromboxane receptor antagonist and a partial DP2 antagonist, is used to treat perennial rhinitis in Japan⁴¹

and more selective and potent DP2 antagonists are being developed for asthma and allergic rhinitis. For example, TM30089 (7TM Pharma, Denmark) is effective in animal models of asthma and a once daily oral molecule, ODC9101 (Oxagen, UK), is now in phase IIa clinical trials for asthma.⁴¹

Leucotriene B₄, acting through its receptor BLT1, is chemoattractive for neutrophils, eosinophils, monocytes, and macrophages. The number of BLT1-expressing T cells is increased in asthma and BLT1 antagonists prevent airway inflammation and airway hyper-responsiveness in animal models.⁴² However, the BLT1 antagonist LY293111 (Eli Lilly, USA) had no effect on allergen challenge in a small group of asthmatic patients,⁴³ although this has not been tested in patients with neutrophilic asthma.

Targeting chemokines and their receptors

Recruitment of inflammatory cells into the airway by chemokines is a crucial process in the development of asthma. Chemokines are classified into four families




Figure 3: Chemokine receptor modulation in asthma

Prevention of inflammatory cell migration into the airways might be achieved by modulation of chemokine receptors. CCR3 antagonists would be predicted to prevent recruitment and activation of T cells, eosinophils, and mast cells in the airways, whereas CXCR2 antagonism would target monocytes/macrophage and neutrophil infiltration. Antagonism of the CCR2 receptor or biological agents against its ligands would prevent mast cell, monocytes/macrophage, and T-cell effects in asthma. Selective targeting of Th1/Tc1 or Th2/Tc2 cells could be achieved with CXCR3 and CCR4/8 or CRTh2 (chemoattractant receptor-homologous molecule expressed on Th2 lymphocytes) antagonism, respectively, or by activation of CCR5 in the case of Th1/Tc1 cells.

based on a conserved cysteine motif—C, CC, CXC, or CX3C—and act through specific receptors according to the ligand class.⁴² The major focus of interest in asthma has been CCR3 and its ligands (CCL11, CCL24, and CCL26), which are increased in asthma⁴⁴ and mediate eosinophil recruitment⁴² (figure 3). CCR3 is also expressed on mast cells and some Th2 cells. CCL11 is increased temporally before induction of CCL24 and CCL26 after allergen challenge.⁴⁵ Targeting of CCR3 would therefore be more reasonable than targeting its ligands, as is the case with most chemokine receptors. Antisense oligonucleotides that inhibit CCR3 and interleukin-5 receptor led to a reduction in sputum eosinophilia after allergen challenge associated with an inhibition of the early response, but only a trend for a reduced late response was noted.⁴⁶ Furthermore, inhibitors of CCR3 have been effective in inhibiting eosinophilic inflammation, airway hyper-responsiveness, and goblet cell hyperplasia in mouse models^{47,48} and are currently undergoing clinical trials for asthma.

Other chemokines may be important in asthma. The expression of CCL17 and CCL22⁴⁹ and the number of CCR4-expressing Th2 cells is increased in asthmatic airways after allergen challenge,⁵⁰ and antibodies directed against CCL17 and CCL22 are effective in animal models of asthma.⁴² Additionally, CXCL8, acting through CXCR1, is important for neutrophil chemotaxis and activation; the expression of CXCR1/2 and its ligands is increased in biopsy samples obtained from asthmatic patients during severe exacerbations that required mechanical ventilation.⁵¹ CXCR1/2 antagonists or biological agents could therefore be particularly effective in the treatment of severe exacerbations or in patients with severe asthma with evidence of neutrophilia.^{51,52} CCR2, which is expressed on monocytes and T cells, and CCR4 and

CCR8, which are expressed on Th2 cells, are also targets for asthma therapy⁴² (figure 3).

Lastly, the glycosaminoglycan heparin is co-released with histamine from mast cells and can, in addition to its well-described anticoagulant properties, combine with chemokines to prevent their action and prevent exercise-induced bronchoconstriction.⁵³ A phase II study in patients with mild asthma using IVX 0142 (IVAX Research, USA), a novel heparin-derived oligosaccharide, has just been completed (ClinicalTrials.gov, number NCT00232999).

Targeting Th2-derived and inflammatory cytokines

Cytokines are major targets for new asthma therapies because of their key role in chronic inflammation and in remodelling of the airway.²⁷ The concept of asthma as a Th2-driven disease and the demonstration that inhibition of Th2-derived cytokines in many animal models of asthma prevented all aspects of disease drove the development of antagonists and antibodies directed against interleukins 4 and 5 (figure 1). However, these findings have not been replicated in asthmatic patients, illustrating the limitations of the animal models used.

Interleukin 5 is critical for terminal differentiation of eosinophils and for eosinophilic inflammation; however, a blocking antibody to interleukin 5 had no effect on airway hyper-responsiveness, lung function, or exacerbation frequency in asthmatic patients, despite depletion of blood and sputum eosinophilia.^{54,55} In a subsequent study, an interleukin-5 antibody failed to prevent eosinophilia within the airway submucosa and resulted in some beneficial effects on aspects of airway remodelling.⁵⁶ It is possible that anti-interleukin-5 therapy may be more effective in patients with high levels of circulating and sputum eosinophils, as it has been shown to be effective in the treatment of hyper-eosinophilic syndrome.⁵⁷

Both interleukin 4 and interleukin 13 are important in B-cell IgE isotype switching, and interleukin 4 is also important in maintaining the Th2 phenotype. Interleukin 13 is also involved in the modulation of eosinophilic inflammation and airway smooth muscle hyperplasia, the induction of goblet-cell hyperplasia, the recruitment of monocytes and T cells, and the induction of a corticosteroid-insensitive airway inflammation.⁵⁸ Early studies with a soluble recombinant human interleukin-4 receptor (altrakinept, Immunex [Amgen], USA) in patients with mild-to-moderate asthma showed some efficacy in maintaining asthma control when inhaled corticosteroids were being withdrawn,⁵⁹ but this effect was not subsequently confirmed and development stopped. Humanised interleukin-4-specific antibodies, antibodies that block the alpha subunit of the interleukin-4 receptor, and peptide-based vaccines against interleukin 4 are now being tested.⁶⁰ A recent study showed that interleukin-4 muteins (eg, pitrakinra) can inhibit the binding of interleukins 4 and 13 to the alpha subunit of

the interleukin-4 receptor, reducing the allergen-induced late-phase response in asthmatic patients.⁶¹ Several monoclonal antibodies against interleukin 13, including CAT-354 and IMA-638, are undergoing clinical trials for asthma and these may have a better therapeutic index than small molecule inhibitors because the large size of the binding pocket could prevent selectivity.⁶⁰

Suplatast tosilate selectively inhibits interleukin 4 and 5 production from T cells in vitro, attenuates allergen-induced goblet-cell metaplasia, and reduces blood and sputum eosinophilia and airway hyper-responsiveness in mild-to-moderate asthmatic patients.⁶⁰ In a small placebo-controlled clinical trial, suplatast was as effective as inhaled beclomethasone in improving the forced expiratory volume in one second (FEV₁) and mean morning peak expiratory flow and was corticosteroid-sparing in patients with moderate asthma.⁶⁰

Thymic stromal lymphopoietin is highly expressed in airway epithelial cells of asthmatic patients and can either activate dendritic cells to orchestrate an allergic pattern of inflammation through the activation of Th2 cells or directly stimulate Th2 cytokine expression from T cells.^{62,63} Additionally, mice in which the gene for thymic stromal lymphopoietin has been knocked out fail to develop an antigen-specific Th2-mediated inflammatory response in the airways. Clinical studies are now needed in patients with all types of asthma.

Some cytokines are intrinsically anti-inflammatory and are therefore potential therapeutic agents. The expression of interleukins 10 and 12 is reduced in patients with severe asthma;^{64,65} restoration of levels of these cytokines has been proposed to restore asthma control. However, repeated injections of interleukin 12 have no effect on airway hyper-responsiveness despite marked effects on blood eosinophilia.⁶⁶ Administration of interleukin 10 has proved effective in animal models of asthma but no studies have been reported in asthmatic patients despite interleukin 10 being approved for psoriasis.⁶⁷ These cytokines have unacceptable side-effects and an alternative strategy to enhance endogenous expression of interleukin 10 through immunomodulatory pathways is preferable.

Tumour necrosis factor (TNF) α has been implicated as a pro-inflammatory cytokine in asthma and blood monocytes from patients with refractory asthma have increased expression of membrane-bound TNF α , TNF α receptor 1, and TNF α -converting enzyme.⁶⁸ In two small studies, patients with severe asthma on conventional therapy exhibited improved asthma control and bronchial hyper-responsiveness in response to TNF α blockade,^{68,69} although a slightly larger study with etanercept, a soluble TNF receptor, showed no clinical efficacy despite effects on sputum cell numbers.⁷⁰

Targeting adhesion molecules

Considerable effort has been made to exploit the importance of adhesion molecules in several facets of asthma—eg, leucocyte migration, exocytosis, cytokine

production, and respiratory burst—by developing drugs that target these molecules⁴² (figure 1). Much interest has been generated by the very late antigen-4 (VLA4, $\alpha 4 \beta 1$ integrin), which is involved in the recruitment of eosinophils and T cells.⁴² However, the clinical outcome of trials of these agents in patients with asthma have been disappointing^{71–73} and indeed the development of some classes of these drugs (eg, $\alpha 4 \beta 1$ integrin antagonists) was placed on hold by the US Food and Drug Administration because of reports of progressive multifocal leucoencephalopathy in patients taking natalizumab. Although now lifted, the ban delayed approval causing widespread implications for the development of adhesion molecule antagonists for the treatment of asthma.⁴²

Other therapeutic areas

Infections

Respiratory viral and bacterial infections are a major cause of asthma exacerbations.¹ Therefore, antibiotics and antivirals may be beneficial for asthma exacerbations. Telithromycin, a macrolide antibiotic, caused a small but significant reduction in asthma symptoms without changes in lung function compared with placebo when administered to patients with acute exacerbations of asthma.⁷⁴ Additionally, clarithromycin can reduce levels of interleukin 8 and sputum neutrophilia and can improve asthma-related quality of life in patients with refractory non-eosinophilic asthma.⁷⁵ Whether these antibiotics act by directly inhibiting infections or by inhibiting neutrophil-based inflammation is unclear. Therapeutic interventions aimed at treating viral or bacterial infections by inhibition of Toll-like receptors (TLRs) are also under investigation, although evidence in animal models is inconclusive.⁷⁶

Rhinovirus infection can reduce nuclear translocation of the glucocorticoid receptor and corticosteroid function;⁷⁷ conversely, inhibition of the glucocorticoid receptor-associated heat shock protein 90 (hsp90) by geldanamycin or its analogues can attenuate rhinovirus replication without the formation of drug-resistant strains,⁷⁸ implying mutual antagonism between viral infection and corticosteroids.

Intracellular signalling pathways

Phosphodiesterase 4 (PDE4) inhibitors such as roflumilast and cilomilast prevent eosinophilic inflammation in ovalbumin-sensitised Brown Norway rats with an efficacy similar to that seen with budesonide.⁷⁹ Moreover, PDE4 inhibitors are able to suppress neutrophilic inflammation and exert distinct anti-inflammatory effects compared with those seen with corticosteroids,⁸⁰ suggesting that PDE4 inhibitors could be useful in the treatment of severe asthma. The suppressive activity of PDE4 inhibitors on inflammatory cells in asthma involves the inhibition of the generation of cytokines, oxidants, pro-inflammatory mediators, the reduction or inhibition

of eosinophilic migration, and the mitigation of degranulation; the compounds can also modify airway hyper-responsiveness and remodelling by reducing smooth muscle cell proliferation and migration, the pro-inflammatory activity of epithelial cells, and plasma exudation that can lead to oedema.⁸⁰ Both cilomilast and roflumilast show dose-dependent inhibition of early and late phase responses to allergen and exercise challenges.⁸⁰ In a parallel-group trial, cilomilast caused small improvements in FEV₁ at 6 weeks, although this was lost by 12 months.⁸⁰ By contrast, roflumilast (500 µg daily) produced greater effects and caused improvements in FEV₁, morning and evening peak flow, and asthma symptoms over 12 weeks similar to those seen with beclomethasone 400 µg.⁸¹ The problem of nausea and vomiting that occurs at the top of the dose–effect curve has not been overcome despite improved isoform selectivity, and administration of these compounds by inhalation is being considered.

Kinase inhibitors

Kinases have a critical role in the expression and activation of inflammatory mediators in the airway, in both resident and infiltrating cell function and airway remodelling^{82,83} (figure 4). Although different kinase pathways can activate specific downstream transcription factors, there is considerable cross-talk between pathways.^{82,83} Changes in kinase activation status have

been reported in all asthmatic patients, but particularly in those with severe asthma where an association with reduced glucocorticoid responsiveness has been proposed.⁸⁴ Thus, enhanced activation of extracellular signal-related kinase (ERK), c-Jun N-terminal kinase (JNK), p38 mitogen-activated protein kinase (MAPK), and the janus kinase/signal transducers and activators of transcription (JAK/STAT) signalling pathway have all been proposed to have a role in steroid-insensitive asthma in a stimulus-dependent manner.^{7,85,86}

p38 MAPK is involved in many inflammatory processes in the lower airways and in tissue remodelling. Selective second generation p38 MAPK inhibitors such as SB2439063 (GlaxoSmithKline, UK) and the p38 MAPK antisense ISIS 101757 (ISIS Pharmaceuticals, USA) reduce the release of inflammatory mediators and some characteristics of allergic inflammation in animal models with no hepatic or neurological toxicity.⁸² Interestingly, the corticosteroid insensitivity seen in peripheral blood cells and macrophages obtained from bronchoalveolar lavage fluid in patients with severe asthma can be overcome by the combination of a p38 MAPK inhibitor and dexamethasone.^{7,87} Safety issues remain a concern for long-term use, although delivery to the airways by aerosol may reduce side-effects. An alternative approach is to target downstream substrates such as MAPKAPK2 since, by contrast with p38 MAPK knockout mice, MAPKAPK2 knockouts are viable and exhibit an anti-inflammatory phenotype.⁸⁸

JNK activity is increased in corticosteroid-resistant asthma⁸⁹ and SP600125 (Celgene, USA), a JNK inhibitor, reduces accumulation of eosinophils and lymphocytes in bronchoalveolar lavage fluid, cytokine release, serum IgE production, and smooth muscle proliferation after repeated allergen exposure in acute and chronic animal models of asthma.^{90,91}

Spleen tyrosine kinase (SYK, p72Syk) has a pivotal role in triggering mast cell degranulation through the high affinity IgE receptor (FcεRI).⁹² SYK is also involved in B and T lymphocyte antigen receptor signalling and in eosinophil survival, suggesting that this might be an important potential target for the development of new anti-asthma drugs.⁹³ BAY 61-3606 (Bayer, Japan), a potent and selective inhibitor of SYK inhibited lipid mediator release, cytokine synthesis, and mast cell degranulation, had inhibitory effects on human basophils, eosinophils, and monocytes,⁹⁴ and attenuated ovalbumin-induced airway inflammation in rats.⁹⁴ The SYK inhibitor R112 (Rigel Pharmaceuticals, USA), given topically, rapidly reduced symptoms of allergic rhinitis⁹⁵ and its follow-up R343 (Rigel-Pfizer, USA) began phase I studies in asthma in late 2007.

Phosphoinositide 3-kinases (PI3Ks) catalyse the phosphorylation of phosphoinositides and regulate a number of distinct cellular responses including cell growth and division, cell apoptosis and survival, and activation in response to cytokines, antigens, and co-stimulatory molecules.^{83,96} PI3Ks may also contribute




Figure 4: Major pro-inflammatory kinases in asthma

Many stimuli, including cytokines, allergens, chemokines, and infectious agents known to be important in asthma and asthma exacerbations, can activate key intracellular signalling pathways in a cell and stimulus dependent manner. Activation of these pathways results in enhanced recruitment and activation of infiltrating and resident cell types and increases the expression of inflammatory mediators, including other cytokines and chemokines that can act, in turn, in an either an autocrine or paracrine manner to further drive the inflammatory response in asthma. Selective inhibitors of these pathways are under development with the intention of dampening the inflammatory response. Some of these pathways (eg, IKK2 and MAPK) are also implicated in corticosteroid function and suppression of these may be corticosteroid-sparing. Additionally, inhibitors of SYK are able to prevent mast cell degranulation. AP-1=activator protein-1. ATF=activating transcription factor. cAMP=cyclic adenosine monophosphate. IKK=inhibitor of κB kinase. JNK=c-Jun N-terminal kinase. NFκB=nuclear factor κB; MAPK=mitogen-activated protein kinase. MAPKAPK=MAPK-activated protein kinase. PDE=phosphodiesterase. PI3K=phosphoinositide-3 kinase. SYK=spleen tyrosine kinase.




Figure 5: Stimulation of cell surface or intracellular receptors can activate a number of signalling cascades that alter the DNA binding or transcriptional activation status of key pro-inflammatory transcription factors

The activity of the transcription factors can be modulated by crosstalk between factors at the level of DNA binding or transactivation—eg, AP-1 and c-myc can associate with GATA-3 or NF-AT to enhance transcription. By contrast, activation of nuclear hormone (steroid) receptors in the cytoplasm leads to nuclear import and either gene induction after DNA binding or gene repression by crosstalk between other factors such as NFκB and AP-1. IKK=IκB kinase. GATA-3=GATA binding protein 3. IκB=inhibitor of κB. JAK=Janus kinase. NF-AT=nuclear factor of activated T cells. NFκB=nuclear factor κB. p38 MAPK=p38 mitogen-activated protein kinase. PPARγ=peroxisome proliferator-activated receptor γ. STAT=signal transducers and activators of transcription.

to the pathogenesis of asthma by affecting airway smooth muscle proliferation and eosinophil recruitment.^{83,96} The need for isoform specificity may limit the usefulness of current PI3K inhibitors in asthma.

Inhibition of transcription factors

Nuclear factor κB (NFκB) is induced by many factors involved in asthmatic inflammation including allergen challenge, cytokines, chemokines, and bacterial and viral infection, and induces the expression of many mediators, growth factors, receptors, and enzymes important in the inflammatory cascade, leading to a feed-forward enhancement of inflammation.⁸² Activation of NFκB is enhanced in mild asthma and further increased in severe asthma,^{97,98} genetic studies also implicate the NFκB pathway in corticosteroid-resistant asthma.⁹⁹ Small molecule inhibitors that target the stimulating kinase (IKK2) completely suppressed inflammatory responses in animal models of asthma and in macrophages from bronchoalveolar lavage of asthmatic patients^{100,101} (figure 5). Targeting IKK2 may also have additional benefits since it seems that current IKK2 inhibitors can also modulate the corticosteroid-insensitive release of CXCL10 that is induced by interferon γ.^{102,103} There are concerns that inhibition of NFκB may cause side-effects such as increased susceptibility

to infections.⁸² Delivery by inhalation could decrease the risk of serious side-effects and these drugs may be particularly effective for exacerbations.

NFκB is a major target for corticosteroids and downstream co-activators or co-repressors may also be potential therapeutic targets as steroid-sparing agents. Recruitment of histone deacetylase 2 (HDAC2) is involved in the glucocorticoid receptor-mediated suppression of NFκB and its expression and activity is reduced in some corticosteroid-insensitive diseases.^{104,105} Importantly, over-expression of HDAC2 in corticosteroid-insensitive cells can restore corticosteroid responsiveness.¹⁰⁶ The suppression of HDAC2 activity may be due to tyrosine nitration¹⁰⁶ further implicating a potential therapeutic role for anti-oxidants or inhibitors of nitric oxide synthase 2 in restoring corticosteroid responsiveness. Theophylline and curcumin both enhance HDAC2 activity under conditions of oxidative stress, leading to a restoration of corticosteroid responsiveness,^{107,108} which might explain why adding a low, sub-bronchodilator dose of theophylline is more effective than increasing the dose of inhaled corticosteroids in patients with poorly controlled disease and why theophylline withdrawal worsens control of patients with severe asthma.¹⁰⁹

Expression of the transcription factor GATA-3 is markedly increased in T cells and bronchial biopsies from asthmatic patients^{110,111} and in cells from bronchoalveolar lavage after allergen challenge.¹¹² GATA-3 is also involved in the regulation of early T-cell development in the thymus, the differentiation of invariant natural killer T cells,¹¹³ and the control of regulatory T (Treg) cells.¹¹⁴ GATA-3 has a critical role in Th2 differentiation from naive CD4 T cells¹¹³ and knockdown of GATA-3 expression using small interfering RNA in human T cells results in the loss of anti-CD3/CD28-mediated Th2 cytokine expression.¹¹⁵ These findings suggest that antisense oligonucleotides may be an excellent novel anti-asthma therapy.¹¹⁶

The nuclear factor of activated T cells (NF-AT) regulates the release of cytokines from Th2 cells by forming complexes with GATA-3 and AP-1, another transcription factor (figure 5). The immunosuppressive drugs ciclosporin A, tacrolimus, and pimecrolimus block calcineurin-dependent dephosphorylation of NF-AT, thereby preventing its activation.¹¹⁶ Systemic administration of ciclosporin A to patients with severe asthma has provided some improvement in lung function but has unacceptable side-effects.¹¹⁶ It is hoped that the development of the locally active T-cell modulator MLD987 (Novartis, Switzerland), an inhaled derivative, may reduce the side-effects.¹¹⁶ Alternative strategies for inhibiting NF-AT include the use of peptides known as inhibitors of NF-AT-calcineurin association (INCA) that block docking of calcineurin to NF-AT.¹¹⁶

STATs are the primary signal-specific mediators of cytokine-regulated gene expression activated by receptor-associated JAKs. For example, interleukins 4 and 12 drive the differentiation of Th2 and Th1 cells, respectively, through activation of STAT6 and STAT4.¹¹⁷ Thus, targeting JAK/STAT pathways could be an effective therapeutic strategy for asthma. Indeed, a STAT1 decoy oligonucleotide (AVT-01, Avontec, Germany) has proved successful in an animal model of asthma¹¹⁸ and is now in phase IIa clinical trials.⁶⁰ Of the many STAT proteins, only STAT6 is unique to the asthma-related cytokines interleukin 4 and interleukin 13.⁶⁰

The characteristics of Treg cells have made these cells attractive candidates for immunotherapy.¹¹⁹ The activity of Treg cells is under the control of the transcription factor forkhead box P3 (FOXP3). Stable over-expression or induction of FOXP3 by retinoic acid drives the conversion of naive T cells to Treg cells and this has been proposed to modulate asthmatic inflammation.¹¹⁹

Peroxisome proliferator-activated receptors (PPARs) are nuclear receptors activated by polyunsaturated fatty acid derivatives, oxidised fatty acids, and phospholipids¹²⁰ (figure 5). PPAR γ ligands decrease antigen-induced airway hyper-responsiveness, lung inflammation and eosinophilia, cytokine production, and GATA-3 expression, as well as serum levels of antigen-specific IgE in different animal models of asthma.¹²⁰ Activation of

PPAR γ also alters the maturation of dendritic cells and rosiglitazone reduces the proliferation of antigen-specific T cells while increasing the production of interleukin 10 by these cells.¹²⁰ Interestingly, the panel of inflammatory genes regulated by PPAR γ agonists is distinct from that regulated by corticosteroids and the combination of both a PPAR γ agonist and a corticosteroid might have a greater anti-inflammatory effect than either drug alone,¹²¹ particularly since corticosteroids can induce PPAR γ expression.¹²²

Oxidative stress

Oxidative stress has been implicated as a driving force behind the inflammatory response and lack of corticosteroid sensitivity in severe asthma.¹²³ Moreover, oxidative stress and its byproducts may drive a Th2-dependent immune response.¹²⁴ Anti-oxidants including N-acetylcysteine, n-acetylcysteinyl, and the superoxide dismutase mimetic AEOL10150 (Aeolus Pharmaceuticals, USA) are able to restore corticosteroid functions that were reduced in response to cigarette smoke or other oxidative stresses in both primary human cells and in animal models,¹²³ but current anti-oxidants do not seem to affect the redox balance in the airways of human beings. Therefore, smarter and more potent drugs are required to target the correct cellular compartment. Oxidative stress in combination with the high levels of nitric oxide seen in asthma¹²⁵ will result in the formation of peroxynitrite, tyrosine nitration, and lipid peroxidation products,¹²⁶ which have been linked to corticosteroid insensitivity. Inhibitors of nitric oxide synthase 2 have been shown to be safe, but ineffective, in mild asthma¹²⁷ but they may have an important role in treatment of patients with severe asthma and in asthmatic patients who smoke, who have increased levels of oxidative or nitrosative stress.¹²³

Resveratrol (3,5,4'-trihydroxystilbene), a component of red wine, has anti-inflammatory and antioxidant properties.¹²³ Although resveratrol inhibits cytokine release by alveolar macrophages from patients with chronic obstructive pulmonary disease, it is not clear if this is due to its anti-oxidant properties. Nevertheless, this compound may also be beneficial for patients with severe asthma.¹²⁸ Curcumin has many therapeutic properties as a result of its anti-oxidant, anti-inflammatory, and anti-cancer effects.^{108,129} Despite the low bioavailability of curcumin after oral application, it is now in phase I clinical studies for several respiratory diseases.¹³⁰

Immunomodulation and anti-allergy treatments

Since asthmatic patients are often atopic, much effort has been directed at modulating the allergic response (figure 1). IgE is the immunoglobulin that mediates the acute allergic response in mast cells and basophils through cross-linking of high-affinity IgE receptors, and may increase allergen uptake by dendritic cells. A

humanised monoclonal antibody that binds to IgE (omalizumab) has been introduced for the treatment of severe allergic asthma. Omalizumab is a useful add-on therapy in some patients who are affected by frequent exacerbations since it reduces the rate of exacerbations,¹³¹ but it is expensive and its cost-effectiveness is debated.¹³² Other more potent anti-IgE antibodies are in development and other strategies also being considered include the development of peptide-based vaccines or immunotherapy to induce IgE-specific antibodies. An antibody directed against the low-affinity IgE receptor (FcεRII or CD23), lumiliximab (Biogen Idec, UK), reduces IgE concentrations in atopic patients.¹³³ A similar effect has been reported in mild asthma from the same group but the results have not been published in full.¹³⁴ Development of therapies directed at T-cell co-stimulatory molecules such as CD23, ICOS, and OX40 is also progressing.⁶⁰

Dendritic cells mount and maintain immune responses to inhaled allergen and modulation of their function represents a new approach to asthma treatment. The sphingosine-1 phosphate receptor antagonist, FTY720 (fingolimod, Novartis, Switzerland), strongly attenuates established lung inflammation in mice through inhibition of activation of dendritic cells,¹³⁵ and is currently in clinical trials for the treatment of multiple sclerosis and transplant rejection.¹³⁵ Selective prostaglandin D receptor agonists such as BW245C (Wellcome Research Laboratories, UK) also suppresses dendritic cell function, leading to decreased airway inflammation and bronchial hyper-reactivity in a mouse model of asthma through the induction of FOXP3-expressing Treg cells.¹³⁶ A similar mechanism may account for the actions of the prostacyclin analogue, iloprost.⁶²

Immunomodulatory therapies offer the opportunity to reverse the abnormal immune function observed in asthma by enhancing Treg expression or function or altering T-cell class switching away from the Th2 response.¹³⁷ Sublingual immunotherapy has been used in asthma, although questions remain about effective doses, treatment schedules, and treatment duration.¹³⁸

Specific subcutaneous immunotherapy increases the production of interleukin 10 from Treg cells^{137,139} and allergen-specific T-cell peptides that enhance Treg function and increase the release of interleukin 10 are now under development.⁷ Treating Treg cells from corticosteroid-resistant patients with vitamin D3 in combination with dexamethasone restores the ability of these cells to release interleukin 10.¹⁴⁰ This, in turn, allowed interleukin 10 to upregulate the expression of the glucocorticoid receptor and reverse the dexamethasone-induced reduction in expression of this receptor. Impressively, oral administration of vitamin D3 (0.5 µg daily) for 7 days to corticosteroid-resistant asthmatic patients enhanced *ex vivo* Treg responses to dexamethasone.¹⁴⁰ This observation suggests that vitamin D3 could potentially increase the therapeutic response to glucocorticoids in corticosteroid-resistant patients.

The potential of non-selective skewing of the T-cell response in asthma using non-pathogenic bacterial products, such as immunostimulatory oligodeoxynucleotides, including CpG oligodeoxynucleotides, which target TLR9, is under active investigation.¹⁴¹ However, early indications in asthma are not encouraging despite evidence of a robust increase in interferon γ and genes that are induced by interferon γ.¹⁴² The long-term consequences of this approach need to be carefully assessed, especially since they would probably need to be used in young children before the onset of asthma.

Conclusions

Several new treatments are now under development for mild or moderate asthma but many of them are highly specific, targeting a single receptor, enzyme, or mediator, and are unlikely to have a major clinical impact. New treatments also have a high barrier to overcome in that combination therapy is efficacious and it is likely that once-a-day, fast-onset combinations of long-acting β agonists and steroids will be available soon. Because of the overexpression of interleukin 13 in asthma and its role in many immuno-inflammatory processes in asthma, it is possible that biological agents targeted against the interleukin-13 receptor may buck this trend. Compliance is an issue with current therapy and the development of an effective oral therapy for mild or moderate asthma may be an advantage, but it is likely that such treatments will have major side-effects. Although we have not discussed these issues here, drug delivery and comorbidities, particularly in patients with severe asthma, will affect the pharmacopoeia used to treat all asthmatic patients.

Prospects for prevention or cure are currently remote but might arise from the development of vaccines and immune therapies directed against Treg and Th17 cells, such as combinations of vitamin D3 and steroids. It is important to recognise that distinct subgroups of patients may respond better to particular therapies, such as those targeting the leucotriene pathway, although it is unlikely that these will be successful as monotherapies.

There are several approaches that can be taken to treat those patients with severe and treatment-refractory asthma. The type of inflammation in these patients may be distinct and targeting this inflammation with selective therapeutic agents such as inhaled p38 MAPK and IKK2 inhibitors may be beneficial. More selective targeting of drugs to patients with particular subtypes of asthma might be possible in the future with the development of discriminatory handprints of clinical phenotypes, biomarkers, and genetic profiles. Rapid tests that distinguish these subtypes may make the advent of selective therapy closer and more effective.

An alternative approach is to try to restore steroid sensitivity rather than prevent inflammation *per se*. Understanding the multiple mechanisms underlying

corticosteroid-resistant asthma might indicate patient-specific abnormalities in signalling pathways that could be targeted to restore asthma control—eg, inhaled p38 MAPK and JAK inhibitors—which could result in a reduced need for inhaled or systemic corticosteroids.

Finally, the fact that combination therapies are more effective than monotherapy, as has been previously observed in the treatment of rheumatoid arthritis, emphasises the need to examine multidrug approaches to asthma that are tailored to the genotype or phenotype of the particular asthma subtype.

Contributors

IMA prepared the initial draft; KFC and GC revised and corrected the subsequent versions and helped to prepare the figures. All authors saw and approved the final version.

Conflict of interest statement

IMA has received consultancy fees from Chiesi, GSK, and Novartis, lecture fees from Altana, GSK, Novartis, Pfizer, and Sanofi-Aventis and contributions towards attending ATS and ERS meetings from Boehringer-Ingelheim and GSK over the past 3 years. KFC has been remunerated for attending advisory board meetings and for delivering lectures at meetings sponsored by pharmaceutical companies marketing respiratory drugs. GC has received lecture fees from GSK (Italy) and MSD (Italy), and contributions towards attending ATS and ERS meetings from Chiesi and Novartis (Italy) over the past 3 years.

Acknowledgments

We thank Paul A Kirkham who wrote the section on oxidative stress. PAK is a Royal Society Industrial Fellow on secondment from Novartis and holds shares in Novartis. Novartis do not work in the area of oxidative stress. We also thank members of the Department of Airways Disease, NHLI, Imperial College London, for helpful discussions. Work in the authors' laboratories in this area is funded by Asthma UK, the UK Medical Research Council, the Royal Society, the University of Ferrara, and the Wellcome Trust.

References

- Masoli M, Fabian D, Holt S, Beasley R. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004; **59**: 469–78.
- Barnes PJ, Jonsson B, Klim JB. The costs of asthma. *Eur Respir J* 1996; **9**: 636–42.
- Chung KF, Godard P, Adelroth E, et al. Difficult/therapy-resistant asthma: the need for an integrated approach to define clinical phenotypes, evaluate risk factors, understand pathophysiology and find novel therapies. *Eur Respir J* 1999; **13**: 1198–208.
- Wenzel SE. Asthma: defining of the persistent adult phenotypes. *Lancet* 2006; **368**: 804–13.
- Holgate ST. Pathogenesis of asthma. *Clin Exp Allergy* 2008; **38**: 872–97.
- Barnes PJ. Inhaled glucocorticoids for asthma. *N Engl J Med* 1995; **332**: 868–75.
- Ito K, Chung KF, Adcock IM. Update on glucocorticoid action and resistance. *J Allergy Clin Immunol* 2006; **117**: 522–43.
- Suissa S, Ernst P. Inhaled corticosteroids: impact on asthma morbidity and mortality. *J Allergy Clin Immunol* 2001; **107**: 937–44.
- Schacke H, Docke WD, Asadullah K. Mechanisms involved in the side effects of glucocorticoids. *Pharmacol Ther* 2002; **96**: 23–43.
- Horne R. Compliance, adherence, and concordance: implications for asthma treatment. *Chest* 2006; **130**: 65–72S.
- Rabe KF, Adachi M, Lai CK, et al. Worldwide severity and control of asthma in children and adults: the global asthma insights and reality surveys. *J Allergy Clin Immunol* 2004; **114**: 40–47.
- Chung KF, Adcock IM. Combination therapy of long-acting beta₂-adrenoceptor agonists and corticosteroids for asthma. *Treat Respir Med* 2004; **3**: 279–89.
- Postma DS, Kerstjens HA, ten Hacken NH. Inhaled corticosteroids and long-acting beta-agonists in adult asthma: a winning combination in all? *Naunyn Schmiedeberg Arch Pharmacol* 2008; **378**: 203–15.
- Bateman ED, Boushey HA, Bousquet J, et al. Can guideline-defined asthma control be achieved? The Gaining Optimal Asthma Control study. *Am J Respir Crit Care Med* 2004; **170**: 836–44.
- Moffatt MF, Kabisch M, Liang L, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* 2007; **448**: 470–73.
- Moore WC, Bleecker ER, Curran-Everett D, et al. Characterization of the severe asthma phenotype by the National Heart, Lung, and Blood Institute's Severe Asthma Research Program. *J Allergy Clin Immunol* 2007; **119**: 405–13.
- Anon. The ENFUMOSA cross-sectional European multicentre study of the clinical phenotype of chronic severe asthma. European Network for Understanding Mechanisms of Severe Asthma. *Eur Respir J* 2003; **22**: 470–77.
- Kamel N, Compton C, Middelvelld R, Higenbottam T, Dahlen SE. The Innovative Medicines Initiative (IMI): a new opportunity for scientific collaboration between academia and industry at the European level. *Eur Respir J* 2008; **31**: 924–26.
- Cohen L, Xueping E, Tarsi J, et al. Epithelial cell proliferation contributes to airway remodeling in severe asthma. *Am J Respir Crit Care Med* 2007; **176**: 138–45.
- Bourdin A, Neveu D, Vachier I, Paganin F, Godard P, Chanez P. Specificity of basement membrane thickening in severe asthma. *J Allergy Clin Immunol* 2007; **119**: 1367–74.
- Brasier AR, Victor S, Boetticher G, et al. Molecular phenotyping of severe asthma using pattern recognition of bronchoalveolar lavage-derived cytokines. *J Allergy Clin Immunol* 2008; **121**: 30–37.
- Haldar P, Pavord ID, Shaw DE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med* 2008; **178**: 218–24.
- Dragonieri S, Schot R, Mertens BJ, et al. An electronic nose in the discrimination of patients with asthma and controls. *J Allergy Clin Immunol* 2007; **120**: 856–62.
- Wenzel SE, Schwartz LB, Langmack EL, et al. Evidence that severe asthma can be divided pathologically into two inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J Respir Crit Care Med* 1999; **160**: 1001–08.
- Johnson M. Molecular mechanisms of beta(2)-adrenergic receptor function, response, and regulation. *J Allergy Clin Immunol* 2006; **117**: 18–24.
- Matera MG, Cazzola M. Ultra-long-acting beta₂-adrenoceptor agonists: an emerging therapeutic option for asthma and COPD? *Drugs* 2007; **67**: 503–15.
- Barnes PJ. New therapies for asthma. *Trends Mol Med* 2006; **12**: 515–20.
- Derendorf H. Pharmacokinetic and pharmacodynamic properties of inhaled ciclesonide. *J Clin Pharmacol* 2007; **47**: 782–89.
- Usmani OS, Ito K, Maneechotesuwan K, et al. Glucocorticoid receptor nuclear translocation in airway cells after inhaled combination therapy. *Am J Respir Crit Care Med* 2005; **172**: 704–12.
- Schacke H, Berger M, Rehwinkel H, Asadullah K. Selective glucocorticoid receptor agonists (SEGRAs): novel ligands with an improved therapeutic index. *Mol Cell Endocrinol* 2007; **275**: 109–17.
- Reichardt HM, Tuckermann JP, Gottlicher M, et al. Repression of inflammatory responses in the absence of DNA binding by the glucocorticoid receptor. *EMBO J* 2001; **20**: 7168–73.
- Miner JN, Ardecky B, Benbatoul K, et al. Antiinflammatory glucocorticoid receptor ligand with reduced side effects exhibits an altered protein-protein interaction profile. *Proc Natl Acad Sci USA* 2007; **104**: 19244–49.
- Paul-Clark MJ, Roviezzo F, Flower RJ, et al. Glucocorticoid receptor nitration leads to enhanced anti-inflammatory effects of novel steroid ligands. *J Immunol* 2003; **171**: 3245–52.
- Nevin BJ, Broadley KJ. Comparative effects of inhaled budesonide and the NO-donating budesonide derivative, NCX 1020, against leukocyte influx and airway hyperreactivity following lipopolysaccharide challenge. *Pulm Pharmacol Ther* 2004; **17**: 219–32.
- Lagente V, Advenier C. New nitric oxide-donating drugs for the treatment of airway diseases. *Curr Opin Investig Drugs* 2004; **5**: 537–41.
- Dahlen SE. Treatment of asthma with antileukotrienes: first line or last resort therapy? *Eur J Pharmacol* 2006; **533**: 40–56.

- 37 Jayaram L, Duong M, Pizzichini MM, et al. Failure of montelukast to reduce sputum eosinophilia in high-dose corticosteroid-dependent asthma. *Eur Respir J* 2005; **25**: 41–46.
- 38 Lazarus SC, Chinchilli VM, Rollings NJ, et al. Smoking affects response to inhaled corticosteroids or leukotriene receptor antagonists in asthma. *Am J Respir Crit Care Med* 2007; **175**: 783–90.
- 39 Levy BD, Lukacs NW, Berlin AA, et al. Lipoxin A4 stable analogs reduce allergic airway responses via mechanisms distinct from CysLT1 receptor antagonism. *FASEB J* 2007; **21**: 3877–84.
- 40 Peters-Golden M, Henderson WR Jr. Leukotrienes. *N Engl J Med* 2007; **357**: 1841–54.
- 41 Pettipher R, Hansel TT, Armer R. Antagonism of the prostaglandin D2 receptors DP1 and CRTH2 as an approach to treat allergic diseases. *Nat Rev Drug Discov* 2007; **6**: 313–25.
- 42 Medina-Tato DA, Watson ML, Ward SG. Leukocyte navigation mechanisms as targets in airway diseases. *Drug Discov Today* 2006; **11**: 866–79.
- 43 Evans DJ, Barnes PJ, Spaethe SM, van Alstyne EL, Mitchell MI, O'Connor BJ. Effect of a leukotriene B4 receptor antagonist, LY293111, on allergen induced responses in asthma. *Thorax* 1996; **51**: 1178–84.
- 44 Ying S, Robinson DS, Meng Q, et al. Enhanced expression of eotaxin and CCR3 mRNA and protein in atopic asthma. Association with airway hyperresponsiveness and predominant co-localization of eotaxin mRNA to bronchial epithelial and endothelial cells. *Eur J Immunol* 1997; **27**: 3507–16.
- 45 Ravensberg AJ, Ricciardolo FL, van Schadewijk A, et al. Eotaxin-2 and eotaxin-3 expression is associated with persistent eosinophilic bronchial inflammation in patients with asthma after allergen challenge. *J Allergy Clin Immunol* 2005; **115**: 779–85.
- 46 Gauvreau GM, Boulet LP, Cockcroft DW, et al. Antisense therapy against CCR3 and the common beta chain attenuates allergen-induced eosinophilic responses. *Am J Respir Crit Care Med* 2008; **177**: 952–58.
- 47 Das AM, Vaddi KG, Solomon KA, et al. Selective inhibition of eosinophil influx into the lung by small molecule CC chemokine receptor 3 antagonists in mouse models of allergic inflammation. *J Pharmacol Exp Ther* 2006; **318**: 411–17.
- 48 Wegmann M, Goggel R, Sel S, et al. Effects of a low-molecular-weight CCR3 antagonist on chronic experimental asthma. *Am J Respir Cell Mol Biol* 2007; **36**: 61–67.
- 49 Panina-Bordignon P, Papi A, Mariani M, et al. The C-C chemokine receptors CCR4 and CCR8 identify airway T cells of allergen-challenged atopic asthmatics. *J Clin Invest* 2001; **107**: 1357–64.
- 50 Kallinich T, Schmidt S, Hamelmann E, et al. Chemokine-receptor expression on T cells in lung compartments of challenged asthmatic patients. *Clin Exp Allergy* 2005; **35**: 26–33.
- 51 Qiu Y, Zhu J, Bandi V, Guntupalli KK, Jeffery PK. Bronchial mucosal inflammation and upregulation of CXC chemoattractants and receptors in severe exacerbations of asthma. *Thorax* 2007; **62**: 475–82.
- 52 Barnes PJ. New molecular targets for the treatment of neutrophilic diseases. *J Allergy Clin Immunol* 2007; **119**: 1055–62.
- 53 Lever R, Page CP. Novel drug development opportunities for heparin. *Nat Rev Drug Discov* 2002; **1**: 140–48.
- 54 Leckie MJ, ten Brinke A, Khan J, et al. Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000; **356**: 2144–48.
- 55 Kips JC, O'Connor BJ, Langley SJ, et al. Effect of SCH55700, a humanized anti-human interleukin-5 antibody, in severe persistent asthma: a pilot study. *Am J Respir Crit Care Med* 2003; **167**: 1655–59.
- 56 Flood-Page P, Menzies-Gow A, Phipps S, et al. Anti-IL-5 treatment reduces deposition of ECM proteins in the bronchial subepithelial basement membrane of mild atopic asthmatics. *J Clin Invest* 2003; **112**: 1029–36.
- 57 Rothenberg ME, Klion AD, Roufosse FE, et al. Treatment of patients with the hyper-eosinophilic syndrome with mepolizumab. *N Engl J Med* 2008; **358**: 1215–28.
- 58 Therien AG, Bernier V, Weicker S, et al. Adenovirus IL-13-induced airway disease in mice: a corticosteroid-resistant model of severe asthma. *Am J Respir Cell Mol Biol* 2008; **39**: 26–35.
- 59 Borish LC, Nelson HS, Corren J, et al. Efficacy of soluble IL-4 receptor for the treatment of adults with asthma. *J Allergy Clin Immunol* 2001; **107**: 963–70.
- 60 Caramori G, Groneberg D, Ito K, Casolari P, Adcock IM, Papi A. New drugs targeting Th2 lymphocytes in asthma. *J Occup Med Toxicol* 2008; **3**: S6.
- 61 Wenzel SE, Balzar S, Ampleford E, et al. IL4R alpha mutations are associated with asthma exacerbations and mast cell/IgE expression. *Am J Respir Crit Care Med* 2007; **175**: 570–76.
- 62 Hammad H, Lambrecht BN. Dendritic cells and epithelial cells: linking innate and adaptive immunity in asthma. *Nat Rev Immunol* 2008; **8**: 193–204.
- 63 Liu YJ, Soumelis V, Watanabe N, et al. TSLP: an epithelial cell cytokine that regulates T cell differentiation by conditioning dendritic cell maturation. *Annu Rev Immunol* 2007; **25**: 193–219.
- 64 Tomita K, Lim S, Hanazawa T, et al. Attenuated production of intracellular IL-10 and IL-12 in monocytes from patients with severe asthma. *Clin Immunol* 2002; **102**: 258–66.
- 65 Lim S, Crawley E, Woo P, Barnes PJ. Haplotype associated with low interleukin-10 production in patients with severe asthma. *Lancet* 1998; **352**: 113.
- 66 Bryan SA, O'Connor BJ, Matti S, et al. Effects of recombinant human interleukin-12 on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000; **356**: 2149–53.
- 67 Numerof RP, Asadullah K. Cytokine and anti-cytokine therapies for psoriasis and atopic dermatitis. *BioDrugs* 2006; **20**: 93–103.
- 68 Berry MA, Hargadon B, Shelley M, et al. Evidence of a role of tumor necrosis factor alpha in refractory asthma. *N Engl J Med* 2006; **354**: 697–708.
- 69 Howarth PH, Babu KS, Arshad HS, et al. Tumour necrosis factor (TNFalpha) as a novel therapeutic target in symptomatic corticosteroid dependent asthma. *Thorax* 2005; **60**: 1012–18.
- 70 Morjaria JB, Chauhan AJ, Babu KS, Polosa R, Davies DE, Holgate ST. The role of a soluble TNFalpha receptor fusion protein (etanercept) in corticosteroid refractory asthma: a double blind, randomised, placebo controlled trial. *Thorax* 2008; **63**: 584–91.
- 71 Ravensberg AJ, Luijk B, Westers P, et al. The effect of a single inhaled dose of a VLA-4 antagonist on allergen-induced airway responses and airway inflammation in patients with asthma. *Allergy* 2006; **61**: 1097–103.
- 72 Norris V, Choong L, Tran D, et al. Effect of IVL745, a VLA-4 antagonist, on allergen-induced bronchoconstriction in patients with asthma. *J Allergy Clin Immunol* 2005; **116**: 761–67.
- 73 Diamant Z, Kuperus J, Baan R, et al. Effect of a very late antigen-4 receptor antagonist on allergen-induced airway responses and inflammation in asthma. *Clin Exp Allergy* 2005; **35**: 1080–87.
- 74 Johnston SL, Blasi F, Black PN, Martin RJ, Farrell DJ, Nieman RB. The effect of telithromycin in acute exacerbations of asthma. *N Engl J Med* 2006; **354**: 1589–600.
- 75 Simpson JL, Powell H, Boyle MJ, Scott RJ, Gibson PG. Clarithromycin targets neutrophilic airway inflammation in refractory asthma. *Am J Respir Crit Care Med* 2008; **177**: 148–55.
- 76 Feleszko W, Jaworska J, Hamelmann E. Toll-like receptors—novel targets in allergic airway disease (probiotics, friends and relatives). *Eur J Pharmacol* 2006; **533**: 308–18.
- 77 Bellettato C, Adcock IM, Ito K, et al. Rhinovirus infection reduces glucocorticoid receptor nuclear translocation in airway epithelial cells. *Eur Respir J* 2003; **22**: 565S.
- 78 Solit DB, Chiosis G. Development and application of Hsp90 inhibitors. *Drug Discov Today* 2008; **13**: 38–43.
- 79 Bundschuh DS, Eltze M, Barsig J, Wollin L, Hatzelmann A, Beume R. In vivo efficacy in airway disease models of roflumilast, a novel orally active PDE4 inhibitor. *J Pharmacol Exp Ther* 2001; **297**: 280–90.
- 80 Chung KF, Fan CK. Phosphodiesterase inhibitors in airways disease. *Eur J Pharmacol* 2006; **533**: 110–17.
- 81 Bousquet J, Aubier M, Sastre J, et al. Comparison of roflumilast, an oral anti-inflammatory, with beclomethasone dipropionate in the treatment of persistent asthma. *Allergy* 2006; **61**: 72–78.
- 82 Adcock IM, Chung KF, Caramori G, Ito K. Kinase inhibitors and airway inflammation. *Eur J Pharmacol* 2006; **533**: 118–32.

- 83 Ito K, Caramori G, Adcock IM. Therapeutic potential of phosphatidylinositol 3-kinase inhibitors in inflammatory respiratory disease. *J Pharmacol Exp Ther* 2007; **321**: 1–8.
- 84 Adcock IM, Lane SJ. Corticosteroid-insensitive asthma: molecular mechanisms. *J Endocrinol* 2003; **178**: 347–55.
- 85 Tsitoura DC, Rothman PB. Enhancement of MEK/ERK signaling promotes glucocorticoid resistance in CD4+ T cells. *J Clin Invest* 2004; **113**: 619–27.
- 86 Li LB, Goleva E, Hall CF, Ou LS, Leung DY. Superantigen-induced corticosteroid resistance of human T cells occurs through activation of the mitogen-activated protein kinase kinase/extracellular signal-regulated kinase (MEK-ERK) pathway. *J Allergy Clin Immunol* 2004; **114**: 1059–69.
- 87 Bhavsar P, Hew M, Khorasani N, et al. Relative corticosteroid insensitivity of alveolar macrophages in severe asthma compared to non-severe asthma. *Thorax* 2008; **39**: 312–23.
- 88 Kotlyarov A, Neiningner A, Schubert C, et al. MAPKAP kinase 2 is essential for LPS-induced TNF- α biosynthesis. *Nat Cell Biol* 1999; **1**: 94–97.
- 89 Sousa AR, Lane SJ, Soh C, Lee TH. In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation. *J Allergy Clin Immunol* 1999; **104**: 565–74.
- 90 Nath P, Eynott P, Leung SY, Adcock IM, Bennett BL, Chung KF. Potential role of c-Jun NH2-terminal kinase in allergic airway inflammation and remodelling: effects of SP600125. *Eur J Pharmacol* 2005; **506**: 273–83.
- 91 Eynott PR, Nath P, Leung SY, Adcock IM, Bennett BL, Chung KF. Allergen-induced inflammation and airway epithelial and smooth muscle cell proliferation: role of Jun N-terminal kinase. *Br J Pharmacol* 2003; **140**: 1373–80.
- 92 Costello PS, Turner M, Walters AE, et al. Critical role for the tyrosine kinase Syk in signalling through the high affinity IgE receptor of mast cells. *Oncogene* 1996; **13**: 2595–605.
- 93 Yousefi S, Hoessli DC, Blaser K, Mills GB, Simon HU. Requirement of Lyn and Syk tyrosine kinases for the prevention of apoptosis by cytokines in human eosinophils. *J Exp Med* 1996; **183**: 1407–14.
- 94 Yamamoto N, Takeshita K, Shichijo M, et al. The orally available spleen tyrosine kinase inhibitor 2-[7-(3,4-dimethoxyphenyl)-imidazo[1,2-c]pyrimidin-5-ylamino]nicotinamide dihydrochloride (BAY 61-3606) blocks antigen-induced airway inflammation in rodents. *J Pharmacol Exp Ther* 2003; **306**: 1174–81.
- 95 Meltzer EO, Berkowitz RB, Grossbard EB. An intranasal Syk-kinase inhibitor (R112) improves the symptoms of seasonal allergic rhinitis in a park environment. *J Allergy Clin Immunol* 2005; **115**: 791–96.
- 96 Finan PM, Thomas MJ. PI 3-kinase inhibition: a therapeutic target for respiratory disease. *Biochem Soc Trans* 2004; **32**: 378–82.
- 97 Hart LA, Krishnan VL, Adcock IM, Barnes PJ, Chung KF. Activation and localization of transcription factor, nuclear factor- κ B, in asthma. *Am J Respir Crit Care Med* 1998; **158**: 1585–92.
- 98 Gagliardo R, Chanez P, Mathieu M, et al. Persistent activation of nuclear factor- κ B signaling pathway in severe uncontrolled asthma. *Am J Respir Crit Care Med* 2003; **168**: 1190–98.
- 99 Hakonarson H, Bjornsdottir US, Halapi E, et al. Profiling of genes expressed in peripheral blood mononuclear cells predicts glucocorticoid sensitivity in asthma patients. *Proc Natl Acad Sci USA* 2005; **102**: 14789–94.
- 100 Birrell MA, Hardaker E, Wong S, et al. Ikappa-B kinase-2 inhibitor blocks inflammation in human airway smooth muscle and a rat model of asthma. *Am J Respir Crit Care Med* 2005; **172**: 962–71.
- 101 Cosio BG, Mann B, Ito K, et al. Histone acetylase and deacetylase activity in alveolar macrophages and blood mononocytes in asthma. *Am J Respir Crit Care Med* 2004; **170**: 141–47.
- 102 Tudhope SJ, Catley MC, Fenwick PS, et al. The role of IkappaB kinase 2, but not activation of NF- κ B, in the release of CXCR3 ligands from IFN- γ -stimulated human bronchial epithelial cells. *J Immunol* 2007; **179**: 6237–45.
- 103 Tliba O, Damera G, Banerjee A, et al. Cytokines induce an early steroid resistance in airway smooth muscle cells: novel role of IRF-1. *Am J Respir Cell Mol Biol* 2007; **38**: 463–72.
- 104 Ito K, Ito M, Elliott WM, et al. Decreased histone deacetylase activity in chronic obstructive pulmonary disease. *N Engl J Med* 2005; **352**: 1967–76.
- 105 Bilodeau S, Vallette-Kasic S, Gauthier Y, et al. Role of Brg1 and HDAC2 in GR trans-repression of the pituitary POMC gene and misexpression in Cushing disease. *Genes Dev* 2006; **20**: 2871–86.
- 106 Ito K, Yamamura S, Essilfie-Quaye S, et al. Histone deacetylase 2-mediated deacetylation of the glucocorticoid receptor enables NF- κ B suppression. *J Exp Med* 2006; **203**: 7–13.
- 107 Cosio BG, Tsaprouni L, Ito K, Jazrawi E, Adcock IM, Barnes PJ. Theophylline restores histone deacetylase activity and steroid responses in COPD macrophages. *J Exp Med* 2004; **200**: 689–95.
- 108 Meja KK, Rajendrasozhan S, Adenuga D, et al. Curcumin restores corticosteroid function in monocytes exposed to oxidants by maintaining HDAC2. *Am J Respir Cell Mol Biol* 2008; **63**: 784–90.
- 109 Barnes PJ. How corticosteroids control inflammation: Quintiles Prize Lecture 2005. *Br J Pharmacol* 2006; **148**: 245–54.
- 110 Nakamura Y, Ghaffar O, Olivenstein R, et al. Gene expression of the GATA-3 transcription factor is increased in atopic asthma. *J Allergy Clin Immunol* 1999; **103**: 215–22.
- 111 Caramori G, Lim S, Ito K, et al. Expression of GATA family of transcription factors in T-cells, monocytes and bronchial biopsies. *Eur Respir J* 2001; **18**: 466–73.
- 112 Erpenbeck VJ, Hagenberg A, Krentel H, et al. Regulation of GATA-3, c-maf and T-bet mRNA expression in bronchoalveolar lavage cells and bronchial biopsies after segmental allergen challenge. *Int Arch Allergy Immunol* 2006; **139**: 306–16.
- 113 Ho IC, Pai SY. GATA-3—not just for Th2 cells anymore. *Cell Mol Immunol* 2007; **4**: 15–29.
- 114 Mantel PY, Kuipers H, Boyman O, et al. GATA3-driven Th2 responses inhibit TGF- β 1-induced FOXP3 expression and the formation of regulatory T cells. *PLoS Biol* 2007; **5**: e329.
- 115 Manechotesuwan K, Xin Y, Ito K, et al. Regulation of Th2 cytokine genes by p38 MAPK-mediated phosphorylation of GATA-3. *J Immunol* 2007; **178**: 2491–98.
- 116 Cousins DJ, McDonald J, Lee TH. Therapeutic approaches for control of transcription factors in allergic disease. *J Allergy Clin Immunol* 2008; **121**: 803–09.
- 117 O'Shea JJ, Murray PJ. Cytokine signaling modules in inflammatory responses. *Immunity* 2008; **28**: 477–87.
- 118 Quarcoo D, Weixler S, Groneberg D, et al. Inhibition of signal transducer and activator of transcription 1 attenuates allergen-induced airway inflammation and hyperactivity. *J Allergy Clin Immunol* 2004; **114**: 288–95.
- 119 Chatila TA, Li N, Garcia-Lloret M, Kim HJ, Nel AE. T-cell effector pathways in allergic diseases: transcriptional mechanisms and therapeutic targets. *J Allergy Clin Immunol* 2008; **121**: 812–23.
- 120 Belvisi MG, Hele DJ, Birrell MA. Peroxisome proliferator-activated receptor gamma agonists as therapy for chronic airway inflammation. *Eur J Pharmacol* 2006; **533**: 101–09.
- 121 Ogawa S, Lozach J, Benner C, et al. Molecular determinants of crosstalk between nuclear receptors and toll-like receptors. *Cell* 2005; **122**: 707–21.
- 122 Usami A, Ueki S, Ito W, et al. Theophylline and dexamethasone induce peroxisome proliferator-activated receptor- γ expression in human eosinophils. *Pharmacology* 2006; **77**: 33–37.
- 123 Kirkham P, Rahman I. Oxidative stress in asthma and COPD: antioxidants as a therapeutic strategy. *Pharmacol Ther* 2006; **111**: 476–94.
- 124 King MR, Ismail AS, Davis LS, Karp DR. Oxidative stress promotes polarization of human T cell differentiation toward a T helper 2 phenotype. *J Immunol* 2006; **176**: 2765–72.
- 125 Kharitonov SA, Barnes PJ. Effects of corticosteroids on noninvasive biomarkers of inflammation in asthma and chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 2004; **1**: 191–99.
- 126 Sugiura H, Komaki Y, Koarai A, Ichinose M. Nitrate stress in refractory asthma. *J Allergy Clin Immunol* 2008; **121**: 355–60.
- 127 Singh D, Richards D, Knowles RG, et al. Selective inducible nitric oxide synthase inhibition has no effect on allergen challenge in asthma. *Am J Respir Crit Care Med* 2007; **176**: 988–93.
- 128 Culpitt SV, Rogers DF, Fenwick PS, et al. Inhibition by red wine extract, resveratrol, of cytokine release by alveolar macrophages in COPD. *Thorax* 2003; **58**: 942–46.

- 129 Rahman I, Biswas SK, Kirkham PA. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem Pharmacol* 2006; **72**: 1439–52.
- 130 Hsu CH, Cheng AL. Clinical studies with curcumin. *Adv Exp Med Biol* 2007; **595**: 471–80.
- 131 Sthoeger ZM, Eliraz A, Asher I, Berkman N, Elbirt D. The beneficial effects of Xolair (omalizumab) as add-on therapy in patients with severe persistent asthma who are inadequately controlled despite best available treatment (GINA 2002 step IV)—the Israeli arm of the INNOVATE study. *Isr Med Assoc J* 2007; **9**: 472–75.
- 132 Wu AC, Paltiel AD, Kuntz KM, Weiss ST, Fuhlbrigge AL. Cost-effectiveness of omalizumab in adults with severe asthma: results from the Asthma Policy Model. *J Allergy Clin Immunol* 2007; **120**: 1146–52.
- 133 Poole JA, Meng J, Reff M, Spellman MC, Rosenwasser LJ. Anti-CD23 monoclonal antibody, lumiliximab, inhibited allergen-induced responses in antigen-presenting cells and T cells from atopic subjects. *J Allergy Clin Immunol* 2005; **116**: 780–88.
- 134 Rosenwasser LJ, Meng J. Anti-CD23. *Clin Rev Allergy Immunol* 2005; **29**: 61–72.
- 135 Idzko M, Hammad H, van Nimwegen M, et al. Local application of FTY720 to the lung abrogates experimental asthma by altering dendritic cell function. *J Clin Invest* 2006; **116**: 2935–44.
- 136 Hammad H, Kool M, Soullie T, et al. Activation of the D prostanoid 1 receptor suppresses asthma by modulation of lung dendritic cell function and induction of regulatory T cells. *J Exp Med* 2007; **204**: 357–67.
- 137 Akdis CA, Blaser K, Akdis M. Mechanisms of allergen-specific immunotherapy. *Chem Immunol Allergy* 2006; **91**: 195–203.
- 138 Frew AJ. Sublingual immunotherapy. *N Engl J Med* 2008; **358**: 2259–64.
- 139 Radulovic S, Jacobson MR, Durham SR, Nouri-Aria KT. Grass pollen immunotherapy induces Foxp3-expressing CD4(+)CD25(+) cells in the nasal mucosa. *J Allergy Clin Immunol* 2008; **121**: 1467–72.
- 140 Xystrakis E, Kusumakar S, Boswell S, et al. Reversing the defective induction of IL-10-secreting regulatory T cells in glucocorticoid-resistant asthma patients. *J Clin Invest* 2006; **116**: 146–55.
- 141 Krieg AM. Therapeutic potential of Toll-like receptor 9 activation. *Nat Rev Drug Discov* 2006; **5**: 471–84.
- 142 Gauvreau GM, Hessel EM, Boulet LP, Coffman RL, O'Byrne PM. Immunostimulatory sequences regulate interferon-inducible genes but not allergic airway responses. *Am J Respir Crit Care Med* 2006; **174**: 15–20.