

The report from ISAAC Phase Three is the largest and most important contribution to date on the growing literature, summarised well by Beasley and colleagues, on paracetamol use and childhood asthma. It ends, appropriately, with a question rather than a conclusion and that question is about causality. The authors, and I, can offer informed opinion—or educated guesses—on the causal effects of paracetamol on incident childhood asthma. The studies to date are suggestive but not definitive enough to recommend a wholesale change in antipyretic use in children. Paracetamol has known benefits for paediatric febrile illness as well as known toxicities. The drug might contribute to asthma incidence and it might be prudent to minimise casual use of this—and all—drugs in otherwise healthy children. However, we need to take the guess-work out of recommending and prescribing antipyretic drugs for children. I agree with Beasley that a population-based randomised trial of adequate power and duration to examine childhood asthma incidence, with paracetamol compared with an active control such as ibuprofen and placebo, is warranted. In view of the heterogeneous nature of asthma, the pharmacogenetics of such a study is likely to be fascinating.

R Graham Barr

Columbia University Medical Center, New York, NY 10032, USA  
rgb9@columbia.edu

I have been a Generalist Physician Faculty Scholar, funded by the Robert Wood Johnson Foundation, on a project about acetaminophen, inflammation, and asthma.

- 1 Morse NM. Ueber eine neue Darstellungsmethode der Acetylamidophenole. *Berichte deutsch chemischen Gesellschaft* 1878; **11**: 232–33.
- 2 Hawton K, Simkin S, Deeks J, et al. UK legislation on analgesic packs: before and after study of long term effect on poisonings. *BMJ* 2004; **329**: 1076.
- 3 Beasley R, Clayton T, Crane J, et al, for the ISAAC Phase Three Study Group. Association between paracetamol use in infancy and childhood, and risk of asthma, rhinoconjunctivitis, and eczema in children aged 6–7 years: analysis from Phase Three of the ISAAC programme. *Lancet* 2008; **372**: 1039–49.
- 4 Jenkins C, Costello J, Hodge L. Systematic review of prevalence of aspirin induced asthma and its implications for clinical practice. *BMJ* 2004; **328**: 434.
- 5 Park JS, Chang HS, Park CS, et al. Association analysis of cysteinyl-leukotriene receptor 2 (CYS2LTR2) polymorphisms with aspirin intolerance in asthmatics. *Pharmacogenet Genomics* 2005; **15**: 483–92.
- 6 Kim SH, Choi JH, Park HS, et al. Association of thromboxane A2 receptor gene polymorphism with the phenotype of acetyl salicylic acid-intolerant asthma. *Clin Exp Allergy* 2005; **35**: 585–90.
- 7 Jinnai N, Sakagami T, Sekigawa T, et al. Polymorphisms in the prostaglandin E2 receptor subtype 2 gene confer susceptibility to aspirin-intolerant asthma: a candidate gene approach. *Hum Mol Genet* 2004; **13**: 3203–17.
- 8 Kordansky D, Adkinson NF Jr, Norman PS, Rosenthal RR. Asthma improved by nonsteroidal anti-inflammatory drugs. *Ann Intern Med* 1978; **88**: 508–11.
- 9 Barr RG, Wentowski CC, Curhan GC, et al. Prospective study of acetaminophen use and risk of newly diagnosed asthma among women. *Am J Respir Crit Care Med* 2004; **169**: 836–41.
- 10 Varner AE, Busse WW, Lemanske RFJ. Hypothesis: decreasing use of pediatric aspirin has contributed to the increasing prevalence of childhood asthma. *Ann Allergy Asthma Immunol* 1998; **81**: 347–51.
- 11 Barr RG, Kurth T, Stampfer MJ, Buring JE, Hennekens CH, Gaziano JM. Low dose aspirin and decreased adult-onset asthma: randomized comparisons from the Physicians' Health Study. *Am J Respir Crit Care Med* 2007; **175**: 120–25.
- 12 Kurth T, Barr RG, Gaziano JM, Buring JE. Randomized aspirin assignment and risk of adult-onset asthma in the Women's Health Study. *Thorax* 2008; **63**: 514–18.
- 13 Lesko SM, Mitchell AA. The safety of acetaminophen and ibuprofen among children younger than two years old. *Pediatrics* 1999; **104**: e39.
- 14 Lesko SM, Louik C, Vezina RM, Mitchell AA. Asthma morbidity after the short-term use of ibuprofen in children. *Pediatrics* 2002; **109**: e20.

## Rhinitis as predictor of adult-onset asthma

See [Perspectives](#) page 1029  
See [Articles](#) page 1049

Sneezing and wheezing are common bedfellows.<sup>1</sup> Because a large proportion of people with asthma have symptoms of rhinitis either attributable to allergic sensitisation or to viral infections, it is hard to know whether sneezing leads to wheezing or whether wheezing and sneezing are part of the same process. However, several studies indicate that rhinitis can precede new onset of asthma.<sup>2–4</sup> Rafea Shaaban and colleagues,<sup>5</sup> in today's *Lancet*, reinforce this idea by showing that rhinitis, even in the absence of atopy, was a strong predictor of adult-onset asthma in the European Community Respiratory Health Survey, which included more than 6000 patients in 14 countries. The

strongest risk occurred in patients with allergic rhinitis who were sensitised to several inhalant allergens, particularly house-dust mites.

These data have interesting parallels to the findings of the Multicentre Allergy Study (MAS) birth cohort,<sup>6</sup> in which the development of sensitisation to perennial but not seasonal allergens in the first years of life determined the development of persistent asthma, which was also mediated by increases in airway responsiveness. The role of rhinitis was not assessed in the MAS cohort, but there are data from the longitudinal Tucson Respiratory Study.<sup>3</sup> In that study, children who developed rhinitis in the first year of life

were more likely to have physician-diagnosed asthma by age 6 years, although they were no more likely to have a positive skin-prick test.

These findings could merely indicate the close association between viral infections and nasal symptoms. Alternatively, rhinitis apart from colds and atopy could be a clinical manifestation of impaired mucosal function. Such loss of function might entail a weakened mucosal barrier leading to increased allergen uptake and mounting of IgE responses, eventually resulting in allergic airway inflammation, airway hyper-responsiveness, and asthma. Conversely, an impaired mucosa of the upper and lower airways might also be more susceptible to viral infections which could interact with the process of allergic sensitisation and thereby increase the risk of asthma.<sup>7</sup> In this context it is noteworthy that mutations in the gene that codes for filaggrin, a protein known to have an important role in the integrity of the epidermal barrier,<sup>8</sup> were also associated with the prevalence of allergic rhinitis in independent populations.<sup>9</sup>

The idea that allergic rhinitis could cause asthma raises the possibility of preventing asthma by preventing atopic sensitisation, which could in turn prevent allergic rhinitis. The potential of subcutaneous and sublingual immunotherapy for asthma prevention in children with allergic rhinitis has been investigated in two open studies.<sup>10–13</sup> The uncontrolled study design of both trials urges caution in interpretation. Of greater concern is that, in the PAT study,<sup>11</sup> the therapy had no effect in two out of six participating centres, and airway hyper-responsiveness was not affected during 5-year and 10-year follow-up.<sup>10–12</sup> Moreover, other factors that potentially influence asthma risk, such as a family history of asthma and exposure to environmental tobacco smoke, were not taken into account in the randomisation. In the SLIT study, objective markers of airway hyper-responsiveness were missing and the findings were only based on subjective reports.<sup>10</sup>

Even if the long-term preventive effect of immunotherapy is still unknown, immunotherapy in patients with allergic rhinitis can improve nasal symptom scores and reduce airway responsiveness and therefore help reduce asthma burden.<sup>14</sup> By contrast, treatment of nasal symptoms with topical steroids or leukotriene antagonists,<sup>1</sup> because they are symptomatic rather than immunomodulatory, will probably have no role



House-dust mite

in the secondary prevention of asthma. Lastly, even if immunotreatments work, the fairly low population-attributable risk might diminish the overall effect of this therapeutic approach. This risk is the incidence (new cases) of a disease that would be avoided if the exposure, in this case rhinitis, would be eliminated. The population-attributable risk in Shaaban and colleagues' study ranged from 27% to 32%, and was only higher in men with allergic rhinitis (48%). Therefore a cure of rhinitis would prevent only some cases of new-onset asthma. In the end, potential risks and management options should be discussed cautiously with patients with allergic rhinitis.

*Erika von Mutius*

University Children's Hospital, D-80337 Munich, Germany  
Erika.von.Mutius@med.uni-muenchen.de

I am consultant for GlaxoSmithKline, UCB, and ProtectImmune.

- 1 Bousquet J, Khaltaev N, Cruz AA, et al. Allergic rhinitis and its impact on asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy* 2008; **63** (suppl): 8–160.
- 2 Guerra S, Sherrill DL, Martinez FD, Barbee RA. Rhinitis as an independent risk factor for adult-onset asthma. *J Allergy Clin Immunol* 2002; **109**: 419–25.
- 3 Wright AL, Holberg CJ, Martinez FD, Halonen M, Morgan W, Taussig LM. Epidemiology of physician-diagnosed allergic rhinitis in childhood. *Pediatrics* 1994; **94**: 895–901.
- 4 Leynaert B, Bousquet J, Neukirch C, Liard R, Neukirch F. Perennial rhinitis: an independent risk factor for asthma in nonatopic subjects: results from the European Community Respiratory Health Survey. *J Allergy Clin Immunol* 1999; **104**: 301–04.
- 5 Shaaban R, Zureik M, Soussan D, et al. Rhinitis and onset of asthma: a longitudinal population-based study. *Lancet* 2008; **372**: 1049–57.

- 6 Illi S, von Mutius E, Lau S, Niggemann B, Gruber C, Wahn U, on behalf of the Multicentre Allergy Study (MAS) group. Perennial allergen sensitisation early in life and chronic asthma in children: a birth cohort study. *Lancet* 2006; **368**: 763–70.
- 7 Kusel MM, de Klerk NH, Keadze T, et al. Early-life respiratory viral infections, atopic sensitization, and risk of subsequent development of persistent asthma. *J Allergy Clin Immunol* 2007; **119**: 1105–10.
- 8 Bonness S, Bieber T. Molecular basis of atopic dermatitis. *Curr Opin Allergy Clin Immunol* 2007; **7**: 382–86.
- 9 Weidinger S, O'Sullivan M, Illig T, et al. Filaggrin mutations, atopic eczema, hay fever, and asthma in children. *J Allergy Clin Immunol* 2008; **121**: 1203–09.
- 10 Novembre E, Galli E, Landi F, et al. Coseasonal sublingual immunotherapy reduces the development of asthma in children with allergic rhinoconjunctivitis. *J Allergy Clin Immunol* 2004; **114**: 851–57.
- 11 Moller C, Dreborg S, Ferdousi HA, et al. Pollen immunotherapy reduces the development of asthma in children with seasonal rhinoconjunctivitis (the PAT study). *J Allergy Clin Immunol* 2002; **109**: 251–56.
- 12 Niggemann B, Jacobsen L, Dreborg S, et al, the PAT Investigator Group. Five-year follow-up on the PAT study: specific immunotherapy and long-term prevention of asthma in children. *Allergy* 2006; **61**: 855–59.
- 13 Jacobsen L, Niggemann B, Dreborg S, et al, the PAT Investigator Group. Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy* 2007; **62**: 943–48.
- 14 Grembale RD, Camporota L, Naty S, Tranfa CM, Djukanovic R, Marsico SA. Effects of specific immunotherapy in allergic rhinitic individuals with bronchial hyperresponsiveness. *Am J Respir Crit Care Med* 2000; **162**: 2048–52.

## Transition from childhood to adult asthma

See [Articles](#) page 1058


If a child has more than one episode of bronchitis in infancy, parents and doctors are concerned about whether the child will outgrow the disease or develop chronic obstructive airway disease—ie, asthma. In today's *Lancet*, Debra Stern and colleagues<sup>1</sup> report data on the prediction of adult asthma at 22 years of age by evaluation of early-life factors at 6 years of age in the Tucson Children's Respiratory Study, one of the oldest birth-cohort studies. Longitudinal studies analyse time courses in disease development and progression and can find relations between exposure factors and outcome (figure).<sup>2</sup>

Although no real cure for asthma exists, early identification of patients at risk of disease progression could lead to better treatment opportunities and, hopefully, improved outcomes in adulthood. Nevertheless, early intervention has not yet been shown to be beneficial in terms of disease modification, which might be because of the failure to identify appropriate subgroups of responders. Studies during early childhood

have not shown a clear benefit of early anti-inflammatory treatment in terms of symptom-free days after the treatment period in preschool children.<sup>3</sup> However, inhaled steroids seem to reduce the accelerated decline in forced expiratory volume in 1 s and airway remodelling in adulthood.<sup>4</sup> A kind of tracking (those who start low, remain low) in terms of lung function seems to exist,<sup>5,6</sup> and children with wheezing disorder before 2 years of age compared with those who started wheezing at school age and girls compared with boys probably have poorer lung function in early adulthood.<sup>7</sup>

Cohort studies have tried to identify different wheezing phenotypes, which were mainly defined by onset and prolongation of symptoms (early onset, late onset, persistent) and presence and absence of atopy.<sup>8,9</sup> One study used latent class analysis, a statistical method developed for the social sciences to identify distinct subsets or classes underlying the observed heterogeneity in a population, to distinguish phenotypes of childhood wheeze and cough.<sup>10</sup> The investigators described five phenotypes: persistent and transient cough, atopic and non-atopic persistent wheeze, and transient viral wheeze. The most common phenotype was atopic wheeze at 8 years of age; however, no prediction for adulthood has been made so far.

In an unselected birth cohort study from New Zealand, 25% of the children had wheezing that persisted from childhood to adulthood or that relapsed after remission.<sup>11</sup> Factors predicting persistence or relapse were sensitisation to house-dust mites, airway hyper-responsiveness, female sex, smoking, and early age at onset. Nevertheless, prediction of outcome seems to be difficult for childhood asthma.



**Figure: Incidence of childhood asthma**  
Data are from German MAS study.<sup>2</sup>