

Groupwork 8 Problems

1. Commit Tea Party [15 points]

Two committees are having a meeting. If there are 12 people in each committee, how many different ways can they sit around a table given the following restrictions? Note that two orderings are considered equal if each person has the same two neighbors (without distinguishing their left and right neighbors).

- (a) There are no restrictions on seating.
- (b) Two people in the same committee cannot be neighbors.
- (c) Everybody must have exactly two neighbors from their committee.
- (d) Everybody must have exactly one neighbor from their committee.

Solution:

There are 24 people in total and 2 committees.

a) No restrictions on seating.

Therefore, it is a combination problem.

$$C(24, 12) = \frac{24!}{12!12!} = 2704156 \text{ ways.}$$

b) Two people in the same committee cannot be neighbors.

Therefore, it is a permutation problem.

$$P(24, 2) = 24 \cdot 23 = 552 \text{ ways.}$$

c) Everybody must have exactly two neighbors from their committee.

Therefore, it is a permutation problem.

$$P(24, 3) = 24 \cdot 23 \cdot 22 = 12144 \text{ ways.}$$

d) Everybody must have exactly one neighbor from their committee.

Therefore, it is a permutation problem.

$$P(24, 2) = 24 \cdot 23 = 552 \text{ ways.}$$

2. Hiking Extravaganza [15 points]

Prove that every complete n -node weighted graph (with all possible edges) with $n \geq 1$ and all distinct edge weights has a (possibly non-simple) path of $n - 1$ edges along which the edge weights are strictly increasing.

Hint: Start by placing a hiker on each node. Try to show that the hikers can walk paths of *total* length $n(n - 1)$, each along increasing-weight paths.

Solution:

Let G be a complete n -node weighted graph with all distinct edge weights.

Let v_1, v_2, \dots, v_n be the nodes of G .

Let w_{ij} be the weight of the edge between v_i and v_j .

Let H be a complete n -node weighted graph with all distinct edge weights.

Let u_1, u_2, \dots, u_n be the nodes of H .

Let x_{ij} be the weight of the edge between u_i and u_j .

Let h_{ij} be the weight of the edge between v_i and u_j .

Let $h_{ij} = w_{ij} + \sum_{k=1}^{j-1} x_{k(k+1)}$ for $1 \leq i \leq n$ and $1 \leq j \leq n$.