

Making Do with Less: An Introduction to Compressed Sensing 3

Kurt Bryan

July 7, 2023

Imperfect Solutions

Consider a linear system $\mathbf{Ax} = \mathbf{b}$.

Imperfect Solutions

Consider a linear system $\mathbf{Ax} = \mathbf{b}$.

For a vector \mathbf{w} the quantity $\|\mathbf{Aw} - \mathbf{b}\|$ is called the *residual error* and provides one way to measure the extent to which \mathbf{w} is a solution.

Imperfect Solutions

Consider a linear system $\mathbf{Ax} = \mathbf{b}$.

For a vector \mathbf{w} the quantity $\|\mathbf{Aw} - \mathbf{b}\|$ is called the *residual error* and provides one way to measure the extent to which \mathbf{w} is a solution.

If $\|\mathbf{Aw} - \mathbf{b}\| = 0$ then $\mathbf{x} = \mathbf{w}$ is a solution to $\mathbf{Ax} = \mathbf{b}$.

Imperfect Solutions

Consider a linear system $\mathbf{Ax} = \mathbf{b}$.

For a vector \mathbf{w} the quantity $\|\mathbf{Aw} - \mathbf{b}\|$ is called the *residual error* and provides one way to measure the extent to which \mathbf{w} is a solution.

If $\|\mathbf{Aw} - \mathbf{b}\| = 0$ then $\mathbf{x} = \mathbf{w}$ is a solution to $\mathbf{Ax} = \mathbf{b}$.

If we have two approximate solutions to $\mathbf{Ax} = \mathbf{b}$, the solution with smaller residual is generally preferred.

Example: Finding a 1-Sparse Solution

Consider the linear system $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

Example: Finding a 1-Sparse Solution

Consider the linear system $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

Conditions for uniqueness of 1-sparse solutions are met ($\mu(\mathbf{A}) \approx 0.95 < 1$, so no two columns are parallel).

Example: Finding a 1-Sparse Solution

Consider the linear system $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

Conditions for uniqueness of 1-sparse solutions are met ($\mu(\mathbf{A}) \approx 0.95 < 1$, so no two columns are parallel).

We're looking for a single nonzero x_k so that

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

Example: Finding a 1-Sparse Solution

Consider the linear system $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

Conditions for uniqueness of 1-sparse solutions are met ($\mu(\mathbf{A}) \approx 0.95 < 1$, so no two columns are parallel).

We're looking for a single nonzero x_k so that

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}.$$

The true solution is $\mathbf{x}^* = \langle 0, 0, 3, 0 \rangle$.

Example: Finding a 1-Sparse Solution

If a 1-sparse solution exists then \mathbf{b} is parallel to some column \mathbf{a}_k of \mathbf{A} . Then $\mu(\mathbf{a}_k, \mathbf{b}) = 1$.

Example: Finding a 1-Sparse Solution

If a 1-sparse solution exists then \mathbf{b} is parallel to some column \mathbf{a}_k of \mathbf{A} . Then $\mu(\mathbf{a}_k, \mathbf{b}) = 1$.

Also since no two columns of \mathbf{A} are parallel, $\mu(\mathbf{a}_j, \mathbf{b}) < 1$ for $j \neq k$.

Example: Finding a 1-Sparse Solution

If a 1-sparse solution exists then \mathbf{b} is parallel to some column \mathbf{a}_k of \mathbf{A} . Then $\mu(\mathbf{a}_k, \mathbf{b}) = 1$.

Also since no two columns of \mathbf{A} are parallel, $\mu(\mathbf{a}_j, \mathbf{b}) < 1$ for $j \neq k$.

Solution Strategy: Compute $\mu(\mathbf{a}_j, \mathbf{b})$ for each $j = 1, \dots, n$, choose the $j = k$ so that $\mu(\mathbf{a}_k, \mathbf{b}) = 1$, then find x_k so that $x_k \mathbf{a}_k = \mathbf{b}$.

Example: Finding a 1-Sparse Solution

With

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

and $\mathbf{b} = \langle 3, 3 \rangle$ we find coherences $0.707, 0.707, 1.00, 0.949$.

Example: Finding a 1-Sparse Solution

With

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

and $\mathbf{b} = \langle 3, 3 \rangle$ we find coherences $0.707, 0.707, 1.00, 0.949$.

Then \mathbf{b} is a multiple of \mathbf{a}_3 , so solve $x_3 \mathbf{a}_3 = \mathbf{b}$ to find $x_3 = 3$.

Example: Finding a 1-Sparse Solution

With

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

and $\mathbf{b} = \langle 3, 3 \rangle$ we find coherences $0.707, 0.707, 1.00, 0.949$.

Then \mathbf{b} is a multiple of \mathbf{a}_3 , so solve $x_3 \mathbf{a}_3 = \mathbf{b}$ to find $x_3 = 3$.

The recovered 1-sparse solution is then

$$\mathbf{x} = 3\mathbf{e}_3 = \langle 0, 0, 3, 0 \rangle$$

where \mathbf{e}_k means the k th standard basis vector.

Example: Finding a 1-Sparse Solution

But suppose the data is a bit off, say $\mathbf{b} = \langle 3.1, 3 \rangle$, and we want the best possible 1-sparse solution.

Example: Finding a 1-Sparse Solution

But suppose the data is a bit off, say $\mathbf{b} = \langle 3.1, 3 \rangle$, and we want the best possible 1-sparse solution.

Which column of

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

should we use?

Example: Finding a 1-Sparse Solution

But suppose the data is a bit off, say $\mathbf{b} = \langle 3.1, 3 \rangle$, and we want the best possible 1-sparse solution.

Which column of

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

should we use?

It seems natural to use the column \mathbf{a}_3 (again) since it is “most parallel” to \mathbf{b} .

Example: Finding a 1-Sparse Solution

But suppose the data is a bit off, say $\mathbf{b} = \langle 3.1, 3 \rangle$, and we want the best possible 1-sparse solution.

Which column of

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

should we use?

It seems natural to use the column \mathbf{a}_3 (again) since it is “most parallel” to \mathbf{b} .

We can put this reasoning on a firmer footing.

Example: Finding a 1-Sparse Solution with Imperfect Data

Suppose no 1-sparse solution to $\mathbf{Ax} = \mathbf{b}$ exists. That is, in

$$x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$$

no \mathbf{a}_j is parallel to \mathbf{b} .

Example: Finding a 1-Sparse Solution with Imperfect Data

Suppose no 1-sparse solution to $\mathbf{Ax} = \mathbf{b}$ exists. That is, in

$$x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$$

no \mathbf{a}_j is parallel to \mathbf{b} .

We can still construct a “best” 1-sparse solution.

Example: Finding a 1-Sparse Solution with Imperfect Data

Suppose no 1-sparse solution to $\mathbf{Ax} = \mathbf{b}$ exists. That is, in

$$x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$$

no \mathbf{a}_j is parallel to \mathbf{b} .

We can still construct a “best” 1-sparse solution.

Suppose we decide to use a solution with support $S = \{k\}$. That is, $\mathbf{x} = \langle 0, 0, \dots, 0, x_k, 0, \dots, 0 \rangle$. This means we need

$$x_k \mathbf{a}_k = \mathbf{b}.$$

Example: Finding a 1-Sparse Solution with Imperfect Data

Suppose no 1-sparse solution to $\mathbf{Ax} = \mathbf{b}$ exists. That is, in

$$x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$$

no \mathbf{a}_j is parallel to \mathbf{b} .

We can still construct a “best” 1-sparse solution.

Suppose we decide to use a solution with support $S = \{k\}$. That is, $\mathbf{x} = \langle 0, 0, \dots, 0, x_k, 0, \dots, 0 \rangle$. This means we need

$$x_k \mathbf{a}_k = \mathbf{b}.$$

We will choose x_k to minimize the squared residual $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$.

Example: Finding a 1-Sparse Solution with Imperfect Data

To minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ with respect to x_k write

Example: Finding a 1-Sparse Solution with Imperfect Data

To minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ with respect to x_k write

$$\begin{aligned}\|x_k \mathbf{a}_k - \mathbf{b}\|^2 &= (x_k \mathbf{a}_k - \mathbf{b}) \cdot (x_k \mathbf{a}_k - \mathbf{b}) \\ &= \|\mathbf{a}_k\|^2 x_k^2 - 2(\mathbf{a}_k \cdot \mathbf{b})x_k + \|\mathbf{b}\|^2.\end{aligned}$$

Example: Finding a 1-Sparse Solution with Imperfect Data

To minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ with respect to x_k write

$$\begin{aligned}\|x_k \mathbf{a}_k - \mathbf{b}\|^2 &= (x_k \mathbf{a}_k - \mathbf{b}) \cdot (x_k \mathbf{a}_k - \mathbf{b}) \\ &= \|\mathbf{a}_k\|^2 x_k^2 - 2(\mathbf{a}_k \cdot \mathbf{b})x_k + \|\mathbf{b}\|^2.\end{aligned}$$

Minimizing with respect to x_k is pure Calc 1: Differentiate with respect to x_k , set the derivative to 0, then solve to find minimizer

$$x_k = \frac{\mathbf{a}_k \cdot \mathbf{b}}{\|\mathbf{a}_k\|^2}.$$

Example: Finding a 1-Sparse Solution with Imperfect Data

To minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ with respect to x_k write

$$\begin{aligned}\|x_k \mathbf{a}_k - \mathbf{b}\|^2 &= (x_k \mathbf{a}_k - \mathbf{b}) \cdot (x_k \mathbf{a}_k - \mathbf{b}) \\ &= \|\mathbf{a}_k\|^2 x_k^2 - 2(\mathbf{a}_k \cdot \mathbf{b})x_k + \|\mathbf{b}\|^2.\end{aligned}$$

Minimizing with respect to x_k is pure Calc 1: Differentiate with respect to x_k , set the derivative to 0, then solve to find minimizer

$$x_k = \frac{\mathbf{a}_k \cdot \mathbf{b}}{\|\mathbf{a}_k\|^2}.$$

If you substitute this back into $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ and simplify you find the minimized squared residual can be written as

$$\text{residual} = \|\mathbf{b}\|^2 \left(1 - \left(\frac{\mathbf{a}_k \cdot \mathbf{b}}{\|\mathbf{a}_k\| \|\mathbf{b}\|} \right)^2 \right)$$

Example: Finding a 1-Sparse Solution with Imperfect Data

To minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ with respect to x_k write

$$\begin{aligned}\|x_k \mathbf{a}_k - \mathbf{b}\|^2 &= (x_k \mathbf{a}_k - \mathbf{b}) \cdot (x_k \mathbf{a}_k - \mathbf{b}) \\ &= \|\mathbf{a}_k\|^2 x_k^2 - 2(\mathbf{a}_k \cdot \mathbf{b})x_k + \|\mathbf{b}\|^2.\end{aligned}$$

Minimizing with respect to x_k is pure Calc 1: Differentiate with respect to x_k , set the derivative to 0, then solve to find minimizer

$$x_k = \frac{\mathbf{a}_k \cdot \mathbf{b}}{\|\mathbf{a}_k\|^2}.$$

If you substitute this back into $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$ and simplify you find the minimized squared residual can be written as

$$\text{residual} = \|\mathbf{b}\|^2 \left(1 - \left(\frac{\mathbf{a}_k \cdot \mathbf{b}}{\|\mathbf{a}_k\| \|\mathbf{b}\|} \right)^2 \right) = \|\mathbf{b}\|^2 (1 - \mu^2(\mathbf{a}_k, \mathbf{b})).$$

Example: Finding a 1-Sparse Solution with Imperfect Data

In summary, if we construct a 1-sparse solution \mathbf{x} to $x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$ with support index k , the best choice for x_k is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$.

Example: Finding a 1-Sparse Solution with Imperfect Data

In summary, if we construct a 1-sparse solution \mathbf{x} to $x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$ with support index k , the best choice for x_k is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$. The resulting squared residual is

$$\|\mathbf{b}\|^2(1 - \mu^2(\mathbf{a}_k, \mathbf{b})).$$

Example: Finding a 1-Sparse Solution with Imperfect Data

In summary, if we construct a 1-sparse solution \mathbf{x} to $x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$ with support index k , the best choice for x_k is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$. The resulting squared residual is

$$\|\mathbf{b}\|^2(1 - \mu^2(\mathbf{a}_k, \mathbf{b})).$$

Which column $1 \leq k \leq n$ gives the best 1-sparse solution (smallest residual)? The column that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.

Example: Finding a 1-Sparse Solution with Imperfect Data

In summary, if we construct a 1-sparse solution \mathbf{x} to $x_1 \mathbf{a}_1 + \cdots + x_n \mathbf{a}_n = \mathbf{b}$ with support index k , the best choice for x_k is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$. The resulting squared residual is

$$\|\mathbf{b}\|^2(1 - \mu^2(\mathbf{a}_k, \mathbf{b})).$$

Which column $1 \leq k \leq n$ gives the best 1-sparse solution (smallest residual)? The column that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.

Note that if $\mu(\mathbf{a}_k, \mathbf{b}) = 1$ then the residual is 0 and we obtain a perfect solution, since \mathbf{b} is parallel to \mathbf{a}_k .

Example: Finding a 1-Sparse Solution with Imperfect Data

Consider the system

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3.10 \\ 3.00 \end{bmatrix}.$$

Example: Finding a 1-Sparse Solution with Imperfect Data

Consider the system

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3.10 \\ 3.00 \end{bmatrix}.$$

First, compute each coherence $\mu(\mathbf{a}_j, \mathbf{b})$ to find the column “most parallel” to \mathbf{b} . We find coherences 0.719, 0.695, 0.999, 0.943 for $1 \leq j \leq 4$.

Example: Finding a 1-Sparse Solution with Imperfect Data

Consider the system

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3.10 \\ 3.00 \end{bmatrix}.$$

First, compute each coherence $\mu(\mathbf{a}_j, \mathbf{b})$ to find the column “most parallel” to \mathbf{b} . We find coherences 0.719, 0.695, 0.999, 0.943 for $1 \leq j \leq 4$.

The best 1-sparse solution will come from support index 3. Also, from $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$ we find optimal choice $x_3 = 3.05$.

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

- 1 Find the column index k that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

- 1 Find the column index k that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.
- 2 The solution will have support index k , and so $\mathbf{x}^1 = x_k \mathbf{e}_k$ where \mathbf{e}_k is the k th standard basis vector.

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

- 1 Find the column index k that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.
- 2 The solution will have support index k , and so $\mathbf{x}^1 = x_k \mathbf{e}_k$ where \mathbf{e}_k is the k th standard basis vector.
- 3 Choose x_k to minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$; this value is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$.

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

- 1 Find the column index k that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.
- 2 The solution will have support index k , and so $\mathbf{x}^1 = x_k \mathbf{e}_k$ where \mathbf{e}_k is the k th standard basis vector.
- 3 Choose x_k to minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$; this value is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$.

This is a “greedy” algorithm: We choose the index support k to make $\|x_k \mathbf{a}_k - \mathbf{b}\|$ as small as possible.

Summary: Computing the Best 1-sparse solution

To find the best 1-sparse solution $\mathbf{x} = \mathbf{x}^1$ to $\mathbf{Ax} = \mathbf{b}$ we

- 1 Find the column index k that maximizes $\mu(\mathbf{a}_k, \mathbf{b})$.
- 2 The solution will have support index k , and so $\mathbf{x}^1 = x_k \mathbf{e}_k$ where \mathbf{e}_k is the k th standard basis vector.
- 3 Choose x_k to minimize $\|x_k \mathbf{a}_k - \mathbf{b}\|^2$; this value is $x_k = (\mathbf{a}_k \cdot \mathbf{b}) / \|\mathbf{a}_k\|^2$.

This is a “greedy” algorithm: We choose the index support k to make $\|x_k \mathbf{a}_k - \mathbf{b}\|$ as small as possible.

We can extend this greedy approach to iteratively construct 2-, 3-, 4-, or higher-sparsity solutions.

Example: Finding a 2-Sparse Solution

Consider finding a 2-sparse solution to $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 2 & -1 & 3 & 0 \\ 3 & 1 & -1 & 1 & 1 & 3 \\ 0 & 1 & -1 & 2 & 4 & -2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} -3 \\ 5 \\ 2 \end{bmatrix}.$$

Example: Finding a 2-Sparse Solution

Consider finding a 2-sparse solution to $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 2 & -1 & 3 & 0 \\ 3 & 1 & -1 & 1 & 1 & 3 \\ 0 & 1 & -1 & 2 & 4 & -2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} -3 \\ 5 \\ 2 \end{bmatrix}.$$

The sparsest solution is $\mathbf{x} = \langle 1, 0, -2, 0, 0, 0 \rangle$, support set $S = \{1, 3\}$ (but we don't know this).

Example: Finding a 2-Sparse Solution

Consider finding a 2-sparse solution to $\mathbf{Ax} = \mathbf{b}$ with

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 2 & -1 & 3 & 0 \\ 3 & 1 & -1 & 1 & 1 & 3 \\ 0 & 1 & -1 & 2 & 4 & -2 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} -3 \\ 5 \\ 2 \end{bmatrix}.$$

The sparsest solution is $\mathbf{x} = \langle 1, 0, -2, 0, 0, 0 \rangle$, support set $S = \{1, 3\}$ (but we don't know this).

If we knew even the support of the 2-sparse solution things would be easy—but we don't.

Example: Finding a 2-Sparse Solution

Start by finding the best 1-sparse solution. The coherence of each column of \mathbf{A} with \mathbf{b} is

0.6156, 0.3746, 0.8609, 0.7947, 0.1273, 0.4949.

Example: Finding a 2-Sparse Solution

Start by finding the best 1-sparse solution. The coherence of each column of \mathbf{A} with \mathbf{b} is

0.6156, 0.3746, 0.8609, 0.7947, 0.1273, 0.4949.

Since column 3 is most highly correlated with \mathbf{b} , we will take a 1-sparse solution of the form $\mathbf{x}^1 = x_3 \mathbf{e}_3$.

Example: Finding a 2-Sparse Solution

Start by finding the best 1-sparse solution. The coherence of each column of \mathbf{A} with \mathbf{b} is

$$0.6156, 0.3746, 0.8609, 0.7947, 0.1273, 0.4949.$$

Since column 3 is most highly correlated with \mathbf{b} , we will take a 1-sparse solution of the form $\mathbf{x}^1 = x_3 \mathbf{e}_3$.

Choosing x_3 to minimize $\|x_3 \mathbf{a}_3 - \mathbf{b}\|^2$ yields $x_3 = -13/6$. The best 1-sparse solution is $\mathbf{x}^1 = \langle 0, 0, -13/6, 0, 0, 0 \rangle$ with support set $S = \{3\}$.

Example: Finding a 2-Sparse Solution

Start by finding the best 1-sparse solution. The coherence of each column of \mathbf{A} with \mathbf{b} is

$$0.6156, 0.3746, 0.8609, 0.7947, 0.1273, 0.4949.$$

Since column 3 is most highly correlated with \mathbf{b} , we will take a 1-sparse solution of the form $\mathbf{x}^1 = x_3 \mathbf{e}_3$.

Choosing x_3 to minimize $\|x_3 \mathbf{a}_3 - \mathbf{b}\|^2$ yields $x_3 = -13/6$. The best 1-sparse solution is $\mathbf{x}^1 = \langle 0, 0, -13/6, 0, 0, 0 \rangle$ with support set $S = \{3\}$.

We'll now boost \mathbf{x}^1 to a 2-sparse solution \mathbf{x}^2 .

Example: Finding a 2-Sparse Solution

Try $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$ (j and x_j to be determined).

Example: Finding a 2-Sparse Solution

Try $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$ (j and x_j to be determined). Note that

$$\mathbf{A}\mathbf{x}^2 = \mathbf{A}\mathbf{x}^1 + x_j \mathbf{A}\mathbf{e}_j = \mathbf{A}\mathbf{x}^1 + x_j \mathbf{a}_j.$$

Example: Finding a 2-Sparse Solution

Try $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$ (j and x_j to be determined). Note that

$$\mathbf{A}\mathbf{x}^2 = \mathbf{A}\mathbf{x}^1 + x_j \mathbf{A}\mathbf{e}_j = \mathbf{A}\mathbf{x}^1 + x_j \mathbf{a}_j.$$

If we want $\mathbf{A}\mathbf{x}^2 = \mathbf{b}$ then we need $\mathbf{A}\mathbf{x}^1 + x_j \mathbf{a}_j = \mathbf{b}$ or

$$x_j \mathbf{a}_j = \mathbf{b} - \mathbf{A}\mathbf{x}^1.$$

Example: Finding a 2-Sparse Solution

Try $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$ (j and x_j to be determined). Note that

$$\mathbf{Ax}^2 = \mathbf{Ax}^1 + x_j \mathbf{Ae}_j = \mathbf{Ax}^1 + x_j \mathbf{a}_j.$$

If we want $\mathbf{Ax}^2 = \mathbf{b}$ then we need $\mathbf{Ax}^1 + x_j \mathbf{a}_j = \mathbf{b}$ or

$$x_j \mathbf{a}_j = \mathbf{b} - \mathbf{Ax}^1.$$

This is just the problem of finding the best 1-sparse solution to $\mathbf{Ax} = \mathbf{b} - \mathbf{Ax}^1$ (instead of $\mathbf{Ax} = \mathbf{b}$), which we know how to do!

Example: Finding a 2-Sparse Solution

So we seek a 1-sparse solution to $\mathbf{Ax} = \mathbf{b} - \mathbf{Ax}^1$. Note that

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 2 & -1 & 3 & 0 \\ 3 & 1 & -1 & 1 & 1 & 3 \\ 0 & 1 & -1 & 2 & 4 & -2 \end{bmatrix}, \quad \mathbf{b} - \mathbf{Ax}^1 = \begin{bmatrix} 4/3 \\ 17/6 \\ -1/6 \end{bmatrix}.$$

Example: Finding a 2-Sparse Solution

So we seek a 1-sparse solution to $\mathbf{Ax} = \mathbf{b} - \mathbf{Ax}^1$. Note that

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 2 & -1 & 3 & 0 \\ 3 & 1 & -1 & 1 & 1 & 3 \\ 0 & 1 & -1 & 2 & 4 & -2 \end{bmatrix}, \quad \mathbf{b} - \mathbf{Ax}^1 = \begin{bmatrix} 4/3 \\ 17/6 \\ -1/6 \end{bmatrix}.$$

These coherence of $\mathbf{b} - \mathbf{Ax}^1$ with each column of \mathbf{A} is

$$0.9916, 0.7365, 0.0000, 0.1519, 0.3857, 0.7813$$

so we'll take $k = 1$ and take $\mathbf{x}^2 = x_1\mathbf{e}_1 + x_3\mathbf{e}_3$ where we already have $x_3 = -13/6$.

Example: Finding a 2-Sparse Solution

Choose x_1 to minimize $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$ or equivalently,
 $\|x_1\mathbf{a}_1 - (\mathbf{b} - x_3\mathbf{a}_3)\|^2$, which leads to $x_1 = 59/60$.

Example: Finding a 2-Sparse Solution

Choose x_1 to minimize $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$ or equivalently, $\|x_1\mathbf{a}_1 - (\mathbf{b} - x_3\mathbf{a}_3)\|^2$, which leads to $x_1 = 59/60$.

The best 2-sparse solution (by this algorithm) is estimated to be

$$\mathbf{x}^2 = \langle 59/60, 0, -13/6, 0, 0, 0 \rangle.$$

Example: Finding a 2-Sparse Solution

Choose x_1 to minimize $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$ or equivalently, $\|x_1\mathbf{a}_1 - (\mathbf{b} - x_3\mathbf{a}_3)\|^2$, which leads to $x_1 = 59/60$.

The best 2-sparse solution (by this algorithm) is estimated to be

$$\mathbf{x}^2 = \langle 59/60, 0, -13/6, 0, 0, 0 \rangle.$$

Compare to the correct sparsest solution

$$\mathbf{x}^* = \langle 1, 0, -2, 0, 0, 0 \rangle.$$

Example: Finding a p -Sparse Solution

This process can be continued and yields an algorithm called *matching pursuit*.

Example: Finding a p -Sparse Solution

This process can be continued and yields an algorithm called *matching pursuit*.

- 1 Construct a 1-sparse solution \mathbf{x}^1 using the above procedure.

Example: Finding a p -Sparse Solution

This process can be continued and yields an algorithm called *matching pursuit*.

- 1 Construct a 1-sparse solution \mathbf{x}^1 using the above procedure.
- 2 Let $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$; we'd like $\mathbf{A}\mathbf{x}^2 = \mathbf{b}$, or

$$x_j \mathbf{a}_j = \mathbf{b} - \mathbf{A}\mathbf{x}^1.$$

This is just the problem of finding a 1-sparse solution to $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{A}\mathbf{x}^1$, so proceed as before.

Example: Finding a p -Sparse Solution

This process can be continued and yields an algorithm called *matching pursuit*.

- 1 Construct a 1-sparse solution \mathbf{x}^1 using the above procedure.
- 2 Let $\mathbf{x}^2 = \mathbf{x}^1 + x_j \mathbf{e}_j$; we'd like $\mathbf{A}\mathbf{x}^2 = \mathbf{b}$, or

$$x_j \mathbf{a}_j = \mathbf{b} - \mathbf{A}\mathbf{x}^1.$$

This is just the problem of finding a 1-sparse solution to $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{A}\mathbf{x}^1$, so proceed as before.

- 3 Let $\mathbf{x}^3 = \mathbf{x}^2 + x_j \mathbf{e}_j$; we'd like $\mathbf{A}\mathbf{x}^2 = \mathbf{b}$, or

$$x_j \mathbf{a}_j = \mathbf{b} - \mathbf{A}\mathbf{x}^2.$$

This is just the problem of finding a 1-sparse solution to $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{A}\mathbf{x}^2$, so proceed as before.

Observation

In the 2-sparse example the true solution was

$$\mathbf{x}^* = \langle 1, 0, -2, 0, 0, 0 \rangle$$

with support $S = \{1, 3\}$.

Observation

In the 2-sparse example the true solution was

$$\mathbf{x}^* = \langle 1, 0, -2, 0, 0, 0 \rangle$$

with support $S = \{1, 3\}$.

The recovered solution was

$$\mathbf{x}^2 = \langle 59/60, 0, -13/6, 0, 0, 0 \rangle,$$

also with support $S = \{1, 3\}$.

Observation

In the 2-sparse example the true solution was

$$\mathbf{x}^* = \langle 1, 0, -2, 0, 0, 0 \rangle$$

with support $S = \{1, 3\}$.

The recovered solution was

$$\mathbf{x}^2 = \langle 59/60, 0, -13/6, 0, 0, 0 \rangle,$$

also with support $S = \{1, 3\}$.

Note that \mathbf{x}^2 has the correct support, but the component entries are off.

Improving Matching Pursuit

An improvement is this: we've identified a 2-sparse solution with support $S = \{1, 3\}$ so we think the best 2-sparse solution is of the form

$$\mathbf{x}^2 = x_1 \mathbf{e}_1 + x_3 \mathbf{e}_3.$$

Improving Matching Pursuit

An improvement is this: we've identified a 2-sparse solution with support $S = \{1, 3\}$ so we think the best 2-sparse solution is of the form

$$\mathbf{x}^2 = x_1 \mathbf{e}_1 + x_3 \mathbf{e}_3.$$

We don't settle for the x_1 and x_3 values obtained from matching pursuit. Instead, we go back and "re-choose" x_1 and x_3 simultaneously to minimize the residual quantity $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$.

Improving Matching Pursuit

An improvement is this: we've identified a 2-sparse solution with support $S = \{1, 3\}$ so we think the best 2-sparse solution is of the form

$$\mathbf{x}^2 = x_1 \mathbf{e}_1 + x_3 \mathbf{e}_3.$$

We don't settle for the x_1 and x_3 values obtained from matching pursuit. Instead, we go back and "re-choose" x_1 and x_3 simultaneously to minimize the residual quantity $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$.

This leads to minimizing $\|x_1 \mathbf{a}_1 + x_3 \mathbf{a}_3 - \mathbf{b}\|^2$, an easy quadratic minimization in variables x_1 and x_3 (it also yields the correct 2-sparse solution here).

Improving Matching Pursuit

An improvement is this: we've identified a 2-sparse solution with support $S = \{1, 3\}$ so we think the best 2-sparse solution is of the form

$$\mathbf{x}^2 = x_1 \mathbf{e}_1 + x_3 \mathbf{e}_3.$$

We don't settle for the x_1 and x_3 values obtained from matching pursuit. Instead, we go back and "re-choose" x_1 and x_3 simultaneously to minimize the residual quantity $\|\mathbf{A}\mathbf{x}^2 - \mathbf{b}\|^2$.

This leads to minimizing $\|x_1 \mathbf{a}_1 + x_3 \mathbf{a}_3 - \mathbf{b}\|^2$, an easy quadratic minimization in variables x_1 and x_3 (it also yields the correct 2-sparse solution here).

This modification yields *Orthogonal Matching Pursuit* (OMP), a standard computational algorithm in CS.

Convergence of OMP

Theorem

Suppose $\mu(\mathbf{A}) < 1/(2k - 1)$ and let $\mathbf{x} = \mathbf{x}^$ be the (unique) k -sparse solution to $\mathbf{Ax} = \mathbf{b}$. Then OMP will recover \mathbf{x}^* exactly, and will recover the components of \mathbf{x}^* in descending order of magnitude.*

Basis Pursuit

In the first lecture we saw some motivation that made it plausible that finding a sparse solution to an underdetermined linear system $\mathbf{Ax} = \mathbf{b}$ (which probably has many solutions) might be accomplished by solving the minimization problem

$$\min_{\mathbf{x}} \|\mathbf{x}\|_1$$

where $\|\mathbf{x}\|_1 = \sum_{j=1}^n |x_j|$, subject to the constraint $\mathbf{Ax} = \mathbf{b}$.

Basis Pursuit

In the first lecture we saw some motivation that made it plausible that finding a sparse solution to an underdetermined linear system $\mathbf{Ax} = \mathbf{b}$ (which probably has many solutions) might be accomplished by solving the minimization problem

$$\min_{\mathbf{x}} \|\mathbf{x}\|_1$$

where $\|\mathbf{x}\|_1 = \sum_{j=1}^n |x_j|$, subject to the constraint $\mathbf{Ax} = \mathbf{b}$.

This would be a Calculus 3 problem if not for the fact that the objective function is not differentiable.

Linear Programming

One of the great accomplishments of 20th century mathematics was the development of techniques for solving optimization problems of the form

$$\min_{\mathbf{y}} \sum_{j=1}^n c_j y_j$$

(this function is linear) subject to equality constraints $\mathbf{A}\mathbf{y} = \mathbf{b}$ and inequality constraints $\mathbf{B}\mathbf{y} \leq \mathbf{d}$ (inequality taken component-by-component).

Linear Programming

One of the great accomplishments of 20th century mathematics was the development of techniques for solving optimization problems of the form

$$\min_{\mathbf{y}} \sum_{j=1}^n c_j y_j$$

(this function is linear) subject to equality constraints $\mathbf{A}\mathbf{y} = \mathbf{b}$ and inequality constraints $\mathbf{B}\mathbf{y} \leq \mathbf{d}$ (inequality taken component-by-component).

Two main classes of methods are the *simplex method* and *interior point methods*.

Conversion to Linear Programming

The problem of minimizing $\|\mathbf{x}\|_1$ subject to constraint $\mathbf{Ax} = \mathbf{b}$ can be cast as a linear programming problem as follows:

Conversion to Linear Programming

The problem of minimizing $\|\mathbf{x}\|_1$ subject to constraint $\mathbf{Ax} = \mathbf{b}$ can be cast as a linear programming problems as follows:

- 1 Introduce new variables x_{n+1}, \dots, x_{2n} (doubling the size of \mathbf{x}).

Conversion to Linear Programming

The problem of minimizing $\|\mathbf{x}\|_1$ subject to constraint $\mathbf{Ax} = \mathbf{b}$ can be cast as a linear programming problems as follows:

- 1 Introduce new variables x_{n+1}, \dots, x_{2n} (doubling the size of \mathbf{x}).
- 2 Add inequality constraints $x_j \leq x_{j+n}$ and $-x_j \leq x_{j+n}$ for $1 \leq j \leq n$, which are equivalent to $-x_{j+n} \leq x_j \leq x_{j+n}$ or $|x_j| \leq x_{j+n}$.

Conversion to Linear Programming

The problem of minimizing $\|\mathbf{x}\|_1$ subject to constraint $\mathbf{Ax} = \mathbf{b}$ can be cast as a linear programming problem as follows:

- 1 Introduce new variables x_{n+1}, \dots, x_{2n} (doubling the size of \mathbf{x}).
- 2 Add inequality constraints $x_j \leq x_{j+n}$ and $-x_j \leq x_{j+n}$ for $1 \leq j \leq n$, which are equivalent to $-x_{j+n} \leq x_j \leq x_{j+n}$ or $|x_j| \leq x_{j+n}$.
- 3 We replace minimizing $\|\mathbf{x}\|_1$ with the problem of minimizing $x_{n+1} + \dots + x_{2n}$ subject to $\mathbf{Ax} = \mathbf{b}$ and the new inequality constraints in step 2 above.

Conversion to Linear Programming

The problem of minimizing $\|\mathbf{x}\|_1$ subject to constraint $\mathbf{Ax} = \mathbf{b}$ can be cast as a linear programming problem as follows:

- 1 Introduce new variables x_{n+1}, \dots, x_{2n} (doubling the size of \mathbf{x}).
- 2 Add inequality constraints $x_j \leq x_{j+n}$ and $-x_j \leq x_{j+n}$ for $1 \leq j \leq n$, which are equivalent to $-x_{j+n} \leq x_j \leq x_{j+n}$ or $|x_j| \leq x_{j+n}$.
- 3 We replace minimizing $\|\mathbf{x}\|_1$ with the problem of minimizing $x_{n+1} + \dots + x_{2n}$ subject to $\mathbf{Ax} = \mathbf{b}$ and the new inequality constraints in step 2 above.
- 4 Then x_1, \dots, x_n provide the solution to the original problem.

Small Example

Consider minimizing $|x_1| + |x_2|$ subject to $x_1 + 2x_2 = 6$. We

Small Example

Consider minimizing $|x_1| + |x_2|$ subject to $x_1 + 2x_2 = 6$. We

- 1 Introduce new variables x_3, x_4 .

Small Example

Consider minimizing $|x_1| + |x_2|$ subject to $x_1 + 2x_2 = 6$. We

- 1 Introduce new variables x_3, x_4 .
- 2 Add inequality constraints $x_1 \leq x_3$ and $-x_1 \leq x_3$ (equivalent to $|x_1| \leq x_3$), and $x_2 \leq x_4$ and $-x_2 \leq x_4$ (equivalent to $|x_2| \leq x_4$).

Small Example

Consider minimizing $|x_1| + |x_2|$ subject to $x_1 + 2x_2 = 6$. We

- 1 Introduce new variables x_3, x_4 .
- 2 Add inequality constraints $x_1 \leq x_3$ and $-x_1 \leq x_3$ (equivalent to $|x_1| \leq x_3$), and $x_2 \leq x_4$ and $-x_2 \leq x_4$ (equivalent to $|x_2| \leq x_4$).
- 3 The new problem is to minimize $x_3 + x_4$ subject to the inequalities $x_1 \leq x_3$, $-x_1 \leq x_3$, $x_2 \leq x_4$, and $-x_2 \leq x_4$, and the equality constraint $x_1 + 2x_2 = 6$.

Convergence of BP

Seeking a sparse solution to $\mathbf{Ax} = \mathbf{b}$ by minimizing $\|\mathbf{x}\|_1$ is called *basis pursuit* (BP).

Theorem

Suppose $\mu(\mathbf{A}) < 1/(2k - 1)$ and let $\mathbf{x}^ = \mathbf{x}^*$ be the (unique) k -sparse solution to $\mathbf{Ax} = \mathbf{b}$. Then BP will recover \mathbf{x}^* exactly.*

Other Algorithms and Issues

Another class of algorithms for finding sparse solutions has also been developed, *iterative hard thresholding*.

Other Algorithms and Issues

Another class of algorithms for finding sparse solutions has also been developed, *iterative hard thresholding*.

If the vector \mathbf{b} is noisy then enforcing $\mathbf{Ax} = \mathbf{b}$ may be too restrictive (you're forcing a fit to noise). In this case we might minimize $\|\mathbf{x}\|_1$ subject to the constraint $\|\mathbf{Ax} - \mathbf{b}\|_2 \leq \epsilon$ for some ϵ . This is called *basis pursuit denoising*.

Other Algorithms and Issues

Another class of algorithms for finding sparse solutions has also been developed, *iterative hard thresholding*.

If the vector \mathbf{b} is noisy then enforcing $\mathbf{Ax} = \mathbf{b}$ may be too restrictive (you're forcing a fit to noise). In this case we might minimize $\|\mathbf{x}\|_1$ subject to the constraint $\|\mathbf{Ax} - \mathbf{b}\|_2 \leq \epsilon$ for some ϵ . This is called *basis pursuit denoising*.

In OMP we would iterate only until the residual $\|\mathbf{Ax}^p - \mathbf{b}\|$ is comparable to the noise level of \mathbf{b} .