# **Shanghai Pinghe School Academic Year 2022-2023**

### 1st Semester Final Exam for Grade 10

Subject: Chemistry		
Time: 90 min		
Total Page: 12		
NAME: CLASS:		
Instructions:		
<ol> <li>Do not open this examination paper until instructed to do so.</li> <li>Do not use correction pen or liquid.</li> <li>Answer all the questions in the spaces provided.</li> </ol>		
Part I – Multiple choices:	Score:	
Part II – Short answer questions:	/ 62	

## Part I. Multiple choices (1\*20)

[Total: 20]

Directions: Read the following questions carefully and choose the letter of the correct answer (one letter for each question) and fill them into the table below.

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20

18	2 <b>He</b>	00	0	Ne	18	8	5	92	9	۲.	90	4	Xe	.29	9	_	(2)	8	윽	(294)
7	" I	.4	1	z	20.	1	_	39.95	3	<u>×</u>	83.90	2	×	131	8	~	(222)	11	5	(28
17			6	ш	19.00	17	ច	35.45	32	Ā	79.90	23	-	126.90	82	¥	(210)	117	Nns	(294)
16			8	0	16.00	16	ဟ	32.07	34	Se	78.96	52	<u>e</u>	127.60	8	S.	(508)	116	Uuh	(293)
15			7	z	14.01	15	۵	30.97	33	As	74.92	19	Sp	121.76	83	ö	208.98	115	Uup	(288)
4			9	ပ	12.01	14	S	28.09	32	ලී	72.63	20	Sn	118.71	82	Pb	207.20	114	Oug	(289)
13			2	ω	10.81	13	₹	26.98	31	Ğa	69.72	49	딤	114.82	81	F	204.38	113	Out	(286)
12		,							30	Zu	65.38	48	ၓ	112.41	80	Нg	200.59	112	ت د	(285)
7									59	D C	63.55	47	Ag	107.87	79	Ρ'n	196.97	111	Rg	(281)
10									28	Ż	58.69	46	Pd	106.42	78	ĭ	195.08	110	Ds	(281)
6									27	ပိ	58.93	45	R	102.91	77	<u>1</u>	192.22	109	Ĕ	(278)
æ	number	ent		atomic	SS				56	Рe	55.85	4	Ru	101.07	9/	SO	190.23	108	Hs	(569)
7	Atomic number	Element		Relative atomic	mass				25	Ā	54.94	43	ပ	(86)	75	Re	186.21	107	В	(270)
9									24	ပ်	52.00	42	ŝ	92.96	74	>	183.84	106	Sg	(269)
r2									23	>	50.94	41	g	92.91	73	Тa	180.95	105	В	(268)
4									22	F	47.87	40	Ż	91.22	72	¥	178.49	104	꿆	(267)
ო									21	သွ	44.96	33	>	88.91	57 🕇	Ľa	138.91	<b>\$</b> 68	¥	(227)
8			4	Be	9.01	12	Mg	24.31	20	Ca	40.08	38	Š	87.62	26	Ва	137.33	88	Ra	(226)
-	<b>- エ</b>	1.01	3	=	6.94	11	Na	22.99	19	×	39.10	37	Rb	85.47	55	S	132.91	87	ŗ	(223)
'	-			7			က			4			2			9			7	

28	26	9	61	62	63	49	65	99	29	89	69	2	71
ဝီ	<u>-</u>	Š	P	Sm	П	ဗွ	<b>P</b>	ò	운	ш	٣	χ	Ľ
140.12	140.12 140.91	144.24	(145)	150.36	151.96	157.25	158.93	162.50	164.93	167.26	168.93	173.05	174.97
06	91	92	93	98	92	96	26	86	66	100	101	102	103
ᆮ	Ра	<b>-</b>	Ν	Pu	Am	Ę,	æ	ర	Es	F	βd	Ŷ	בֿ
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)



**1.** Which contains the most atoms of oxygen?

A. 
$$64 g of O_2$$

B.  $1.2 \times 10^{24}$  molecules of  $O_2$ 

C. 64 g of C<sub>3</sub>H<sub>5</sub>O<sub>3</sub>

D.  $1.2 \times 10^{24}$  molecules of C<sub>3</sub>H<sub>5</sub>O<sub>3</sub>



**2.** Burette readings for a titration are shown.

Burette readings / cm $^3\pm 0.05\mathrm{cm}^3$	Trial 1	Trial 2	Trial 3
Final	11.35	24.60	11.70
Initial	0.20	13.50	0.50

What is the mean titre?

A. 
$$11.1 \text{ cm}^3 \pm 0.1 \text{ cm}^3$$

B. 
$$11.15 \text{ cm}^3 \pm 0.05 \text{ cm}^3$$

C. 
$$11.2 \text{ cm}^3 \pm 0.05 \text{ cm}^3$$

D. 
$$11.2 \text{ cm}^3 \pm 0.1 \text{ cm}^3$$

**3.** What volume of oxygen, in dm<sup>3</sup> at STP, is needed when 5.8 g of butane undergoes complete combustion?

$$2C_4H_{10}(g) + 13O_2(g) \to 8\mathrm{CO}_2(g) + 10H_2\mathrm{O}(g)$$


A. 
$$2 \times \frac{5.8}{12.01 \times 4 + 1.01 \times 10} \times 13 \times 22.7$$

B. 
$$\frac{5.8}{12.01\times4+1.01\times10} \times \frac{13}{2} \times 22.7$$

C. 
$$\frac{5.8}{12.01\times4+1.01\times10} \times \frac{2}{13} \times 22.7$$

D. 
$$\frac{5.8}{12.01\times4+1.01\times10} \times \frac{13}{2} \times \frac{22.7}{1000}$$

4. A liquid was added to a graduated cylinder. What can be deduced from the graph?



	Gradient	y-intercept
A.	density of liquid	amount of liquid
B.	density of liquid	mass of empty cylinder
C.	rate of adding liquid	amount of liquid
D.	rate of adding liquid	mass of empty cylinder

- **5.** What is represented by "2–" in  ${}^{A}Z^{2-}$ ?
- A. loss of electron
- B. gain of electron
- C. loss of proton
- D. gain of proton
- **6.** A sample of a compound contains approximately  $24.0\,\mathrm{g}$  C,  $3.0\,\mathrm{g}$  H, and  $1.6\,\mathrm{g}$  O. What is the empirical formula of the compound?
- A. C<sub>20</sub>H<sub>30</sub>O
- B. C<sub>84</sub>H<sub>10</sub>O<sub>6</sub>
- C. C<sub>2</sub>H<sub>3</sub>O
- D. C<sub>24</sub>H<sub>30</sub>O<sub>2</sub>
- $\textbf{7.} \ Which series shows the correct order of metallic bond strength from strongest to weakest?$


A. 
$$Na > K > Rb > Mg$$

$$\mathsf{B.}\ \mathrm{Mg}>\mathrm{Rb}>\mathrm{K}>\mathrm{Na}$$

C. 
$$Rb > K > Na > Mg$$

D. 
$$Mg > Na > K > Rb$$

**8.** What is the relative atomic mass,  $A_r$ , of an element with this mass spectrum?



- A. 24.0
- B. 24.3
- C. 24.9
- D. 25.0

#### **9.** Which is correct?

- A. Mixtures are either homogeneous or heterogeneous and their chemical properties are an average of the individual component properties.
- B. Mixtures are never heterogeneous and their chemical properties are an average of the individual component properties.
- C. Mixtures are either homogeneous or heterogeneous and the components retain their individual chemical properties.
- D. Mixtures are never homogeneous and the components retain their individual chemical properties.



- **10.** Which is correct for all solid ionic compounds?
- A. High volatility
- B. Poor electrical conductivity
- C. Low melting point
- $D. \ Good \ solubility \ in \ water$

### **11.** The rate of a reaction is studied at different temperatures.

Which is the best way to plot the data?

(îndequadent, deq	render is 12	场
-------------------	--------------	---

Ī	x-axis	Type of variable on x-axis
A.	rate	dependent
B.	rate	independent
C.	temperature	independent
D.	temperature	dependent

#### 12. In which set do all the species contain more electrons than neutrons?

- A. <sup>14</sup>N, <sup>16</sup>O, <sup>11</sup>C
- B. <sup>14</sup>N, <sup>16</sup>O, <sup>11</sup>C<sup>4</sup>-
- C. <sup>14</sup>N<sup>3</sup>-, <sup>16</sup>O<sup>2</sup>-, <sup>11</sup>C
- D. <sup>14</sup>N<sup>3</sup>-, <sup>16</sup>O<sup>2</sup>-, <sup>11</sup>C<sup>4</sup>+

### 13. What is the order of decreasing ionic radius?

- A.  $S^{2-} > Cl^- > Al^{3+} > Mg^{2+}$
- B.  $Cl^- > S^{2-} > Al^{3+} > Mg^{2+}$
- C.  $S^{2-} > Cl^- > Mg^{2+} > Al^{3+}$
- D.  $Mg^{2+} > Al^{3+} > Cl^{-} > S^{2-}$

### $\textbf{14.}\ \textbf{X}, \textbf{Y}\ and\ \textbf{Z}\ represent\ the\ successive\ elements,\ \textbf{Ne},\ \textbf{Na}\ and\ \textbf{Mg},\ \textbf{but}\ not\ necessarily\ in\ that\ order.$

	First ionization energy / kJ mol <sup>-1</sup>
X	2081
Y	496
Z	738


### What is the order of increasing atomic number?


- A. X < Y < Z
- B. X < Z < Y
- C. Y < Z < X
- D. Y < X < Z


<b>15.</b> Which compound contains both ionic and covalent bonds?
A. MgO
B. $CH_2Cl_2$




- D. NaOH
- **16.** Which of the following is a mixture?
- A. CaCO3
- B. Ozone
- C. CuSO4·5H2O
- D. Brass
- 17. Which diagram represents a heterogeneous mixture?











- **18.** Which statements about the periodic table are correct?
- I. The elements Mg, Ca and Sr have similar chemical properties.
- II. Elements in the same period have the same number of main energy levels.

decrease

- III. The atomic radius of Na, Mg and P increases as atomic number increases.
  - A. I and II only B. I and III only
  - C. II and III only D. I, II and III

<b>19.</b> A measuring cylinder was used to obtain a known volume of a liquid. The volume was read from the
top of the meniscus and the liquid completely emptied into a flask. The exact same process was then
repeated. Which statement is correct about the overall described procedure and the volumes measured?

- A. There is a systematic error and the volumes measured are accurate.
- B. There is a random error and the volumes measured are accurate.
- C. There is a random error and the volumes measured are inaccurate.
- D. There is a systematic error and the volumes measured are inaccurate.
- 20. How many bonding electrons are there in the urea molecule?

- A. 8
- B. 16
- C. 20
- D. 24

#### Part II. Short-answer and data based questions

[Total: 42 marks]

**Directions:** Read the following questions carefully and answer the questions in the spaces provided.

1. Complete the following tables.

[3]

Formula	IUPAC Name
Mn <sub>3</sub> (PO <sub>3</sub> ) <sub>2</sub>	Manganese (1) phosphite. (phosphote 等, 場)
P <sub>4</sub> O <sub>10</sub>	
HBrO	

<b>2.</b> a) <b>State</b> the electron arrangement for the P <sup>3</sup> ion.	[1]	
	• • • • • • • • • • • • • • • • • • • •	

b) <b>Explain</b> why the ionic radius of the S <sup>2-</sup> ion is smaller than that of the P <sup>3-</sup> ion.	[2]
c) <b>Define</b> 1st ionization energy	[2]
minimum energy regulated to remove an electron from the vale	ence
minimum energy regusted to remove an electron from the vale shell of a gaslow atom to form a positive ion	••••••
d) $\underline{\textbf{Describe}}$ the chemical bonding in group 1 elements, $\underline{\textbf{state}}$ and $\underline{\textbf{explain}}$ the trend i	n melting point. [3]
the bonding in Group 1 is metalic bonding. Down the group the melting point of group elements decrease. Down the group they have the same ENC but greater radius, less ionization energy, so there is less electrostastic attraction between nucleus and valances so there goes decreasing melting point.	ip 1 İst
	[2]
) Write down the equation for 1 <sup>st</sup> ionization of Cl <b>including state symbols</b> .  Cl (g)-> Cl+(g)+e-  Explain why solid potassium can conduct electricity but solid potassium bromide conducted.	[2]
e) Write down the equation for 1 <sup>st</sup> ionization of Cl <b>including state symbols</b> .  Cl (g)-> Cl+(g)+e-  Begin in the equation of 1 <sup>st</sup> ionization of Cl <b>including state symbols</b> .  Cl (g)-> Cl+(g)+e-  Begin in the equation of Cl including state symbols.	[2]
e) Write down the equation for 1 <sup>st</sup> ionization of Cl <b>including state symbols</b> .  Cl (g) -> Cl + (g) +e-  Delocolized electron, K+ and Cl- held tightly in solid KCl crystal structure	[2]

3. A  $0.6125~g\pm0.0001g~$  sample of potassium iodate is dissolved in distilled water and made up to 250.00~ cm $^3\pm0.05~$  cm $^3$  in a volumetric flask. A  $25.00\pm0.05$  cm $^3$  portion of the solution is transferred by a cylinder added to an excess of acidified potassium iodide solution. The iodine formed requires  $24.50\pm0.05~$  cm $^3$  of sodium thiosulfate solution for titration.

IO<sub>3</sub> - (aq) + 5I - (aq) + 6H+(aq) 
$$\rightarrow$$
 3l<sub>2</sub> + 3H<sub>2</sub>O(l)  
2S<sub>2</sub>O<sub>3</sub><sup>2-</sup>(aq) + I<sub>2</sub>(aq)  $\rightarrow$  2I<sup>-</sup>(aq) + S<sub>4</sub>O<sub>6</sub><sup>2-</sup> (aq)

Determine the concentration of the sodium thiosulfate solution with absolute uncertainty.

4. Students A and B are doing the same investigation about titration to determine the concentration of  $25 \, \text{cm}^3 \, \text{NaOH(aq)}$  titrated with 0.100M HCl solution.

Students A and B uses a 25 cm $^3$  pipette with uncertainty of  $\pm\,0.05$ cm $^3$ .

The data collected is on the table:

Experiment	Titration with 0.100M HCl solution using burette, $\pm$ 0.05cm $^3$				
Student A	Initial reading	0.00	0.00	0.00	0.00
	Final reading	11.30	10.80	11.00	10.50
Student B	Initial reading	0.00	10.30	0.00	10.20
	Final reading	10.20	20.40	10.20	20.50

a) Discuss which set of data is more precise.

B has more accurate concentration of NaOH solution, if the literature value for concentration solution is 0.0410 M.	
c) Student A found out that the burette was not rinsed with hydrochloric acid prior to filling explain how this would affect the calculated concentration of base.	State and [2]
d) Student B used the wet conical flask prior to transferring base into it; State and explain he affect the calculated concentration of base. [2]	ow this would
[to	otal 9 marks]
5. This question is about compounds of sodium.	
(a) Sodium can react with oxygen to form sodium oxide. Describe the structure and bonding sodium oxide. [2	in solid 2 marks]
(b)Compare and explain the electrical conductivity of molten sodium oxide. [1 marks]	

(c). Sodium can react with water.	
(i)Write down the balanced equation to present the reaction between sodium and wastate symbol [1 mark]	ater including
(ii) Calculate the volume of gas produced if 3.20g sodium is reacted with excess of war [2 marks]	nter at STP.
(iii) If the student collected 0.0432mol of gas in this experiment, calculate the percent [1 mark]	tage yield.
(iv) Calculate the percentage error in part (iii).	[1 mark]
6. Draw the Lewis structures of a. nitrite b. silicon tetrachloride c. carbon disulfide. [3 marks]	[total 8 marks]