

# Making Do with Less: An Introduction to Compressed Sensing 2

Kurt Bryan

July 5, 2023

## The Marble Problem Again

Recall the marble problem. In the 10-marble version we need to find the sparsest possible solution to the system of 5 linear equations

$$x_1 + x_2 + x_3 + x_5 + x_7 + x_9 = 0.14$$

$$x_5 + x_6 + x_7 + x_9 = 0.44$$

$$x_1 + x_3 + x_4 + x_5 + x_7 + x_8 = -0.30$$

$$x_3 + x_6 + x_8 + x_9 = 0.14$$

$$x_3 + x_5 + x_6 + x_7 + x_8 + x_{10} = -0.30$$

in 10 unknowns. There are many infinitely many solutions—but the sparsest is  $x_3 = -0.3$ ,  $x_9 = 0.44$ , and all other  $x_i = 0$  (but we don't know this).

## The Marble Problem Again

Recall the marble problem. In the 10-marble version we need to find the sparsest possible solution to the system of 5 linear equations

$$x_1 + x_2 + x_3 + x_5 + x_7 + x_9 = 0.14$$

$$x_5 + x_6 + x_7 + x_9 = 0.44$$

$$x_1 + x_3 + x_4 + x_5 + x_7 + x_8 = -0.30$$

$$x_3 + x_6 + x_8 + x_9 = 0.14$$

$$x_3 + x_5 + x_6 + x_7 + x_8 + x_{10} = -0.30$$

in 10 unknowns. There are many infinitely many solutions—but the sparsest is  $x_3 = -0.3$ ,  $x_9 = 0.44$ , and all other  $x_i = 0$  (but we don't know this). We want to find this sparse solution.

# Matrix Formulation

This linear system can be expressed as  $\mathbf{Ax} = \mathbf{b}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 0.14 \\ 0.44 \\ -0.30 \\ 0.14 \\ -0.30 \end{bmatrix}.$$

# Matrix Formulation

This linear system can be expressed as  $\mathbf{Ax} = \mathbf{b}$  where

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 0 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 0.14 \\ 0.44 \\ -0.30 \\ 0.14 \\ -0.30 \end{bmatrix}.$$

The matrix  $\mathbf{A}$  is called the *sensing matrix* or the *measurement matrix*. The vector  $\mathbf{b}$  stems from collected data (weight measurements).

# Matrix Formulation

Our central goal today is to understand some conditions under which an underdetermined linear system  $\mathbf{Ax} = \mathbf{b}$  possesses a unique sparse solution (if it has one at all).

# Matrix Formulation

Our central goal today is to understand some conditions under which an underdetermined linear system  $\mathbf{Ax} = \mathbf{b}$  possesses a unique sparse solution (if it has one at all).

But the matrix  $\mathbf{A}$  does not have to be a 0 – 1 matrix. It could (in principle) be anything. However, it is usually a random matrix—entries may be integers, real numbers, or complex numbers.

# Matrix Formulation

Our central goal today is to understand some conditions under which an underdetermined linear system  $\mathbf{Ax} = \mathbf{b}$  possesses a unique sparse solution (if it has one at all).

But the matrix  $\mathbf{A}$  does not have to be a 0 – 1 matrix. It could (in principle) be anything. However, it is usually a random matrix—entries may be integers, real numbers, or complex numbers.

Let's look at a few concrete examples for intuition.



# Uniqueness

Consider the underdetermined linear system

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

# Uniqueness

Consider the underdetermined linear system

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

If  $\mathbf{b} = \langle 0, 0 \rangle$  then  $\mathbf{x} = \langle 0, 0, 0, 0 \rangle$  is obviously the unique sparsest solution (clearly true in all cases if  $\mathbf{b}$  is the zero vector).

# Uniqueness

Consider the underdetermined linear system

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

If  $\mathbf{b} = \langle 0, 0 \rangle$  then  $\mathbf{x} = \langle 0, 0, 0, 0 \rangle$  is obviously the unique sparsest solution (clearly true in all cases if  $\mathbf{b}$  is the zero vector).

For any nonzero  $\mathbf{b}$  there are always multiple (but finitely many) 2-sparse solutions to this system (e.g., set  $x_3 = x_4 = 0$ , solve for  $x_1$  and  $x_2$ ).

# Uniqueness

Consider the underdetermined linear system

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

If  $\mathbf{b} = \langle 0, 0 \rangle$  then  $\mathbf{x} = \langle 0, 0, 0, 0 \rangle$  is obviously the unique sparsest solution (clearly true in all cases if  $\mathbf{b}$  is the zero vector).

For any nonzero  $\mathbf{b}$  there are always multiple (but finitely many) 2-sparse solutions to this system (e.g., set  $x_3 = x_4 = 0$ , solve for  $x_1$  and  $x_2$ ).

For any  $\mathbf{b}$  there are infinitely many 3 and 4-sparse solutions.

# Uniqueness

What about 1-sparse solutions to

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}?$$

# Uniqueness

What about 1-sparse solutions to

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}?$$

For a typical  $\mathbf{b}$  there may not be a 1-sparse solution, but it's easy to see that if a 1-sparse solution does exist, the solution is unique.

# Uniqueness

To see this write

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

# Uniqueness

To see this write

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

Then the linear system  $\mathbf{Ax} = \mathbf{b}$  can be expressed

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$



# Uniqueness

To see this write

$$\begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

Then the linear system  $\mathbf{Ax} = \mathbf{b}$  can be expressed

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

If a 1-sparse solution exists,  $\mathbf{b}$  is a multiple of one of the columns of  $\mathbf{A}$ , and none of the columns are multiples of each other.

# Uniqueness

More generally, if the sensing matrix  $\mathbf{A}$  has columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , that is,

$$\mathbf{A} = [\mathbf{a}_1 \mid \mathbf{a}_2 \mid \cdots \mid \mathbf{a}_n]$$

then  $\mathbf{Ax} = \mathbf{b}$  can be written

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{b}.$$

# Uniqueness

More generally, if the sensing matrix  $\mathbf{A}$  has columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , that is,

$$\mathbf{A} = [\mathbf{a}_1 \mid \mathbf{a}_2 \mid \cdots \mid \mathbf{a}_n]$$

then  $\mathbf{Ax} = \mathbf{b}$  can be written

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{b}.$$

If none of the columns  $\mathbf{a}_i$  are multiples of each other (that is, no two are parallel vectors) then any 1-sparse solution will be unique.

# Uniqueness

On the other hand, if two columns are multiples of each other, e.g.,

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

then 1-sparse solutions may not be unique!

# Uniqueness

On the other hand, if two columns are multiples of each other, e.g.,

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

then 1-sparse solutions may not be unique!

# Uniqueness

On the other hand, if two columns are multiples of each other, e.g.,

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

then 1-sparse solutions may not be unique! Consider

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}.$$

# Uniqueness

On the other hand, if two columns are multiples of each other, e.g.,

$$\begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & -1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

then 1-sparse solutions may not be unique! Consider

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ -1 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + x_4 \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 5 \end{bmatrix}.$$

Both  $x_3$  and  $x_4$  are candidates for a 1-sparse solution.

# Coherence

For two vectors  $\mathbf{v}$  and  $\mathbf{w}$  we define the *coherence*  $\mu(\mathbf{v}, \mathbf{w})$  of these two vectors as

$$\mu(\mathbf{v}, \mathbf{w}) = \frac{|\mathbf{v} \cdot \mathbf{w}|}{\|\mathbf{v}\| \|\mathbf{w}\|}$$

where  $\mathbf{v} \cdot \mathbf{w}$  is the usual dot product and  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$  is the usual Euclidean length of  $\mathbf{v}$ .



# Coherence

For two vectors  $\mathbf{v}$  and  $\mathbf{w}$  we define the *coherence*  $\mu(\mathbf{v}, \mathbf{w})$  of these two vectors as

$$\mu(\mathbf{v}, \mathbf{w}) = \frac{|\mathbf{v} \cdot \mathbf{w}|}{\|\mathbf{v}\| \|\mathbf{w}\|}$$

where  $\mathbf{v} \cdot \mathbf{w}$  is the usual dot product and  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$  is the usual Euclidean length of  $\mathbf{v}$ .

Recall from basic vector calculus that  $\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos(\theta)$  where  $\theta$  is the angle between  $\mathbf{v}$  and  $\mathbf{w}$ .

# Coherence

For two vectors  $\mathbf{v}$  and  $\mathbf{w}$  we define the *coherence*  $\mu(\mathbf{v}, \mathbf{w})$  of these two vectors as

$$\mu(\mathbf{v}, \mathbf{w}) = \frac{|\mathbf{v} \cdot \mathbf{w}|}{\|\mathbf{v}\| \|\mathbf{w}\|}$$

where  $\mathbf{v} \cdot \mathbf{w}$  is the usual dot product and  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$  is the usual Euclidean length of  $\mathbf{v}$ .

Recall from basic vector calculus that  $\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos(\theta)$  where  $\theta$  is the angle between  $\mathbf{v}$  and  $\mathbf{w}$ .

Then  $0 \leq \mu(\mathbf{v}, \mathbf{w}) \leq 1$  always. We have  $\mu = 1$  exactly when  $\mathbf{v}$  and  $\mathbf{w}$  are parallel,  $\mu = 0$  when they are orthogonal.

# Coherence

If a matrix  $\mathbf{A}$  has columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$  then the coherence  $\mu(\mathbf{A})$  of  $\mathbf{A}$  is defined as

$$\mu(\mathbf{A}) = \max_{i \neq j} \mu(\mathbf{a}_i, \mathbf{a}_j).$$

It's clear that  $0 \leq \mu(\mathbf{A}) \leq 1$  always.

# Coherence

If a matrix  $\mathbf{A}$  has columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$  then the coherence  $\mu(\mathbf{A})$  of  $\mathbf{A}$  is defined as

$$\mu(\mathbf{A}) = \max_{i \neq j} \mu(\mathbf{a}_i, \mathbf{a}_j).$$

It's clear that  $0 \leq \mu(\mathbf{A}) \leq 1$  always.

We have  $\mu(\mathbf{A}) = 0$  exactly when all columns of  $\mathbf{A}$  are orthogonal to each other.

# Coherence

If a matrix  $\mathbf{A}$  has columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$  then the coherence  $\mu(\mathbf{A})$  of  $\mathbf{A}$  is defined as

$$\mu(\mathbf{A}) = \max_{i \neq j} \mu(\mathbf{a}_i, \mathbf{a}_j).$$

It's clear that  $0 \leq \mu(\mathbf{A}) \leq 1$  always.

We have  $\mu(\mathbf{A}) = 0$  exactly when all columns of  $\mathbf{A}$  are orthogonal to each other.

We have  $\mu(\mathbf{A}) = 1$  if any two columns of  $\mathbf{A}$  are multiples of each other.

## Coherence Example 1

Let

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Then  $\mu(\mathbf{a}_1, \mathbf{a}_2) = 0$  (these columns are orthogonal). Also

$$\mu(\mathbf{a}_1, \mathbf{a}_3) = \frac{(1)(1) + (0)(1)}{(1)(\sqrt{2})} = 1/\sqrt{2}.$$

## Coherence Example 1

Let

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Then  $\mu(\mathbf{a}_1, \mathbf{a}_2) = 0$  (these columns are orthogonal). Also

$$\mu(\mathbf{a}_1, \mathbf{a}_3) = \frac{(1)(1) + (0)(1)}{(1)(\sqrt{2})} = 1/\sqrt{2}.$$

Similar computations give pairwise coherence values  $1/\sqrt{5}$ ,  $1/\sqrt{2}$ ,  $2/\sqrt{5}$  and  $3/\sqrt{10}$ . Then  $\mu(\mathbf{A}) = 3/\sqrt{10} \approx 0.948$ .

## Coherence Example 2

Let

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Then  $\mu(\mathbf{a}_3, \mathbf{a}_4) = 1$  (these columns are parallel) and so  $\mu(\mathbf{A}) = 1$ .

It's easy to see that, for example, if  $\mathbf{b} = \langle b, b \rangle$  then there are many 1-sparse solutions to  $\mathbf{Ax} = \mathbf{b}$ .



## An Easy Theorem

A bit of thought shows we've proven that if  $\mu(\mathbf{A}) < 1$  (no columns of  $\mathbf{A}$  are parallel or multiples of each other) then any 1-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique (if such a solution exists).

## An Easy Theorem

A bit of thought shows we've proven that if  $\mu(\mathbf{A}) < 1$  (no columns of  $\mathbf{A}$  are parallel or multiples of each other) then any 1-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique (if such a solution exists).

But if  $\mu(\mathbf{A}) = 1$ , a 1-sparse solution may not be unique (depends on  $\mathbf{b}$ )).

## An Easy Theorem

A bit of thought shows we've proven that if  $\mu(\mathbf{A}) < 1$  (no columns of  $\mathbf{A}$  are parallel or multiples of each other) then any 1-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique (if such a solution exists).

But if  $\mu(\mathbf{A}) = 1$ , a 1-sparse solution may not be unique (depends on  $\mathbf{b}$ )).

Can we relate  $\mu(\mathbf{A})$  to uniqueness for solutions of higher sparsity? Lower values of  $\mu(\mathbf{A})$  let us assert that solutions of greater sparsity are unique.

## Coherence Example 3: Orthogonal Matrices

Consider a matrix  $\mathbf{A}$  with  $\mu(\mathbf{A}) = 0$ . That is, each column of  $\mathbf{A}$  is orthogonal to every other column of  $\mathbf{A}$ , so  $\mathbf{a}_i \cdot \mathbf{a}_j = 0$  if  $i \neq j$ .

## Coherence Example 3: Orthogonal Matrices

Consider a matrix  $\mathbf{A}$  with  $\mu(\mathbf{A}) = 0$ . That is, each column of  $\mathbf{A}$  is orthogonal to every other column of  $\mathbf{A}$ , so  $\mathbf{a}_i \cdot \mathbf{a}_j = 0$  if  $i \neq j$ .

In this case, solutions of ANY sparsity are unique.

## Coherence Example 3: Orthogonal Matrices

Consider a matrix  $\mathbf{A}$  with  $\mu(\mathbf{A}) = 0$ . That is, each column of  $\mathbf{A}$  is orthogonal to every other column of  $\mathbf{A}$ , so  $\mathbf{a}_i \cdot \mathbf{a}_j = 0$  if  $i \neq j$ .

In this case, solutions of ANY sparsity are unique. To see this, write  $\mathbf{Ax} = \mathbf{b}$  as

$$x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + \cdots + x_n \mathbf{a}_n = \mathbf{b}.$$

## Coherence Example 3: Orthogonal Matrices

Consider a matrix  $\mathbf{A}$  with  $\mu(\mathbf{A}) = 0$ . That is, each column of  $\mathbf{A}$  is orthogonal to every other column of  $\mathbf{A}$ , so  $\mathbf{a}_i \cdot \mathbf{a}_j = 0$  if  $i \neq j$ .

In this case, solutions of ANY sparsity are unique. To see this, write  $\mathbf{Ax} = \mathbf{b}$  as

$$x_1 \mathbf{a}_1 + x_2 \mathbf{a}_2 + \cdots + x_n \mathbf{a}_n = \mathbf{b}.$$

Take the dot product of both sides with  $\mathbf{a}_i$  and use column-orthogonality to find  $x_i \|\mathbf{a}_i\|^2 = \mathbf{a}_i \cdot \mathbf{b}$ , so  $x_i = (\mathbf{a}_i \cdot \mathbf{b}) / \|\mathbf{a}_i\|^2$  is uniquely determined.

## Low Coherence Matrices

So  $\mu(\mathbf{A}) = 0$  would be ideal. Except that if  $\mathbf{A}$  is an  $m \times n$  matrix ( $m$  equations in  $n$  unknowns) then the columns can be orthogonal ONLY when  $m \geq n$ .



## Low Coherence Matrices

So  $\mu(\mathbf{A}) = 0$  would be ideal. Except that if  $\mathbf{A}$  is an  $m \times n$  matrix ( $m$  equations in  $n$  unknowns) then the columns can be orthogonal ONLY when  $m \geq n$ .

That is the over (or critically) determined case, but we're interested in the underdetermined case  $m < n$ , probably  $m \ll n$ .

## Low Coherence Matrices

So  $\mu(\mathbf{A}) = 0$  would be ideal. Except that if  $\mathbf{A}$  is an  $m \times n$  matrix ( $m$  equations in  $n$  unknowns) then the columns can be orthogonal ONLY when  $m \geq n$ .

That is the over (or critically) determined case, but we're interested in the underdetermined case  $m < n$ , probably  $m \ll n$ .

So  $\mu(\mathbf{A}) = 0$  is off the table. But for a given  $m \times n$  matrix  $\mathbf{A}$ , the lower the value of  $\mu(\mathbf{A})$ , the better things go for compressed sensing.

## Low Coherence Matrices

So  $\mu(\mathbf{A}) = 0$  would be ideal. Except that if  $\mathbf{A}$  is an  $m \times n$  matrix ( $m$  equations in  $n$  unknowns) then the columns can be orthogonal ONLY when  $m \geq n$ .

That is the over (or critically) determined case, but we're interested in the underdetermined case  $m < n$ , probably  $m \ll n$ .

So  $\mu(\mathbf{A}) = 0$  is off the table. But for a given  $m \times n$  matrix  $\mathbf{A}$ , the lower the value of  $\mu(\mathbf{A})$ , the better things go for compressed sensing.

One lower bound for the coherence of an  $m \times n$  matrix is

$$\mu(\mathbf{A}) \geq \sqrt{\frac{n-m}{m(n-1)}}.$$

# A Theorem

The statement that if  $\mu(\mathbf{A}) < 1$  then any 1-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique can be generalized to

**Theorem:** If

$$\mu(\mathbf{A}) < 1/(2k - 1)$$

then any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique.

# A Theorem

The statement that if  $\mu(\mathbf{A}) < 1$  then any 1-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique can be generalized to

**Theorem:** If

$$\mu(\mathbf{A}) < 1/(2k - 1)$$

then any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique.

Alternatively, this inequality can be turned around to read that if

$$k < \frac{\mu(\mathbf{A}) + 1}{2\mu(\mathbf{A})}$$

then any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique.

# Coherence in Reality

## Coherence in Reality

- Coherence works well for determining when 1-sparse solutions are unique, for it makes it easy to determine when no two columns of the sensing matrix are parallel.

## Coherence in Reality

- Coherence works well for determining when 1-sparse solutions are unique, for it makes it easy to determine when no two columns of the sensing matrix are parallel.
- Coherence isn't as sharp a tool for  $k$ -sparse solutions if  $k \geq 2$ ; the condition  $\mu(\mathbf{A}) < 1/(2k - 1)$  is too stringent.



## Coherence in Reality

- Coherence works well for determining when 1-sparse solutions are unique, for it makes it easy to determine when no two columns of the sensing matrix are parallel.
- Coherence isn't as sharp a tool for  $k$ -sparse solutions if  $k \geq 2$ ; the condition  $\mu(\mathbf{A}) < 1/(2k - 1)$  is too stringent.
- There is a generalization of coherence called the “restricted isometry property (RIP) of order  $k$ ” we can compute for  $\mathbf{A}$  that gives slightly better insight into the uniqueness of sparse solutions.

## Coherence in Reality

- Coherence works well for determining when 1-sparse solutions are unique, for it makes it easy to determine when no two columns of the sensing matrix are parallel.
- Coherence isn't as sharp a tool for  $k$ -sparse solutions if  $k \geq 2$ ; the condition  $\mu(\mathbf{A}) < 1/(2k - 1)$  is too stringent.
- There is a generalization of coherence called the “restricted isometry property (RIP) of order  $k$ ” we can compute for  $\mathbf{A}$  that gives slightly better insight into the uniqueness of sparse solutions.
- Unfortunately computing the RIP for any specific matrix  $\mathbf{A}$  is itself a combinatorially bad computation, and impractical.

# The Nullspace Property

For a given  $\mathbf{A}$ , how can we prove that any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique?

# The Nullspace Property

For a given  $\mathbf{A}$ , how can we prove that any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique?

Suppose that  $\mathbf{x} = \mathbf{v}$  and  $\mathbf{x} = \mathbf{w}$  are distinct  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$ ; let  $\mathbf{u} = \mathbf{v} - \mathbf{w}$ .

# The Nullspace Property

For a given  $\mathbf{A}$ , how can we prove that any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique?

Suppose that  $\mathbf{x} = \mathbf{v}$  and  $\mathbf{x} = \mathbf{w}$  are distinct  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$ ; let  $\mathbf{u} = \mathbf{v} - \mathbf{w}$ . Note that  $\mathbf{u}$  is  $2k$ -sparse and is not the zero vector, and

$$\mathbf{Au} = \mathbf{0}.$$

# The Nullspace Property

For a given  $\mathbf{A}$ , how can we prove that any  $k$ -sparse solution to  $\mathbf{Ax} = \mathbf{b}$  is unique?

Suppose that  $\mathbf{x} = \mathbf{v}$  and  $\mathbf{x} = \mathbf{w}$  are distinct  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$ ; let  $\mathbf{u} = \mathbf{v} - \mathbf{w}$ . Note that  $\mathbf{u}$  is  $2k$ -sparse and is not the zero vector, and

$$\mathbf{Au} = \mathbf{0}.$$

**Punchline:** if there are two distinct  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  then the nullspace of  $\mathbf{A}$  must contain a nontrivial  $2k$ -sparse vector.

# The Nullspace Property

So we can thus show  $k$ -sparse solutions are unique by showing that the nullspace of  $\mathbf{A}$  (the set of all solutions to  $\mathbf{Ax} = 0$ ) contains no  $2k$ -sparse vectors.

## The Nullspace Property

So we can thus show  $k$ -sparse solutions are unique by showing that the nullspace of  $\mathbf{A}$  (the set of all solutions to  $\mathbf{Ax} = 0$ ) contains no  $2k$ -sparse vectors.

One approach is brute force. For example, let

$$\mathbf{A} = \begin{bmatrix} -3 & -3 & -1 & 3 & -2 & 4 \\ 5 & -3 & 3 & 5 & 4 & -4 \\ 1 & 2 & 2 & -2 & -2 & 0 \\ -2 & 3 & 5 & -2 & -4 & -4 \end{bmatrix}$$



## The Nullspace Property

So we can thus show  $k$ -sparse solutions are unique by showing that the nullspace of  $\mathbf{A}$  (the set of all solutions to  $\mathbf{Ax} = 0$ ) contains no  $2k$ -sparse vectors.

One approach is brute force. For example, let

$$\mathbf{A} = \begin{bmatrix} -3 & -3 & -1 & 3 & -2 & 4 \\ 5 & -3 & 3 & 5 & 4 & -4 \\ 1 & 2 & 2 & -2 & -2 & 0 \\ -2 & 3 & 5 & -2 & -4 & -4 \end{bmatrix}$$

Must a 2-sparse solution to  $\mathbf{Ax} = \mathbf{b}$  be unique? Let's check the nullspace of  $\mathbf{A}$  for 4-sparse vectors!

# The Nullspace Property

Suppose  $\mathbf{x} = \langle x_1, x_2, x_3, x_4, 0, 0 \rangle$  is a 4-sparse solution (support  $\{1, 2, 3, 4\}$ ) to  $\mathbf{Ax} = \mathbf{0}$ .

## The Nullspace Property

Suppose  $\mathbf{x} = \langle x_1, x_2, x_3, x_4, 0, 0 \rangle$  is a 4-sparse solution (support  $\{1, 2, 3, 4\}$ ) to  $\mathbf{Ax} = \mathbf{0}$ . Then

$$\begin{bmatrix} -3 & -3 & -1 & 3 \\ 5 & -3 & 3 & 5 \\ 1 & 2 & 2 & -2 \\ -2 & 3 & 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \mathbf{0}.$$

You can check that the only solution is  $x_1 = x_2 = x_3 = x_4 = 0$ .

## The Nullspace Property

Suppose  $\mathbf{x} = \langle x_1, x_2, x_3, x_4, 0, 0 \rangle$  is a 4-sparse solution (support  $\{1, 2, 3, 4\}$ ) to  $\mathbf{Ax} = \mathbf{0}$ . Then

$$\begin{bmatrix} -3 & -3 & -1 & 3 \\ 5 & -3 & 3 & 5 \\ 1 & 2 & 2 & -2 \\ -2 & 3 & 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \mathbf{0}.$$

You can check that the only solution is  $x_1 = x_2 = x_3 = x_4 = 0$ .

Now repeat assuming  $\mathbf{x}$  has support  $\{1, 2, 3, 5\}$ , etc., all  $\binom{6}{4}$  index support subset possibilities.

# The Nullspace Property

This approach for proving  $k$ -sparse uniqueness on an  $m \times n$  system requires solving  $\binom{n}{2k}$  linear systems of size  $m \times 2k$ —way too much computation!

## The Nullspace Property

This approach for proving  $k$ -sparse uniqueness on an  $m \times n$  system requires solving  $\binom{n}{2k}$  linear systems of size  $m \times 2k$ —way too much computation!

But here's an alternate take on the matter that will involve randomness. Let  $\mathbf{u}$  be a vector in  $\mathbb{R}^n$  (might as well be a unit vector).

## The Nullspace Property

This approach for proving  $k$ -sparse uniqueness on an  $m \times n$  system requires solving  $\binom{n}{2k}$  linear systems of size  $m \times 2k$ —way too much computation!

But here's an alternate take on the matter that will involve randomness. Let  $\mathbf{u}$  be a vector in  $\mathbb{R}^n$  (might as well be a unit vector).

If  $\mathbf{A}\mathbf{u} = \mathbf{0}$  this is equivalent to saying that

$$\|\mathbf{A}\mathbf{u}\|^2 = 0.$$

# The Restricted Isometry Property



# The Restricted Isometry Property

We could prove that there are no  $2k$ -sparse vectors in the nullspace of  $\mathbf{A}$  by showing that

$$0 < c_1 \leq \|\mathbf{A}\mathbf{u}\|^2$$

for all  $2k$ -sparse vectors  $\mathbf{u}$ .

# The Restricted Isometry Property

We could prove that there are no  $2k$ -sparse vectors in the nullspace of  $\mathbf{A}$  by showing that

$$0 < c_1 \leq \|\mathbf{A}\mathbf{u}\|^2$$

for all  $2k$ -sparse vectors  $\mathbf{u}$ .

A matrix  $\mathbf{A}$  is said to satisfy the restricted isometry property (RIP) of order  $q$  if

$$0 < c_1 \leq \|\mathbf{A}\mathbf{u}\|^2 \leq c_2$$

for some constants  $c_1, c_2$ , and all  $q$ -sparse unit vectors  $\mathbf{u}$ .

# The Restricted Isometry Property

Suppose  $\mathbf{A}$  is  $m \times n$  and has entries that are random variables (e.g., standard normal/Gaussian).

# The Restricted Isometry Property

Suppose  $\mathbf{A}$  is  $m \times n$  and has entries that are random variables (e.g., standard normal/Gaussian).

In this case, for any fixed unit vector  $\mathbf{u}$ , the quantity  $\|\mathbf{A}\mathbf{u}\|^2$  is itself a random variable.

# The Restricted Isometry Property

Suppose  $\mathbf{A}$  is  $m \times n$  and has entries that are random variables (e.g., standard normal/Gaussian).

In this case, for any fixed unit vector  $\mathbf{u}$ , the quantity  $\|\mathbf{A}\mathbf{u}\|^2$  is itself a random variable. Its distribution may be impossible to work out, but we can use tools from probability to make estimates.

# The Restricted Isometry Property

Suppose  $\mathbf{A}$  is  $m \times n$  and has entries that are random variables (e.g., standard normal/Gaussian).

In this case, for any fixed unit vector  $\mathbf{u}$ , the quantity  $\|\mathbf{A}\mathbf{u}\|^2$  is itself a random variable. Its distribution may be impossible to work out, but we can use tools from probability to make estimates.

In particular, a bit of detailed analysis shows that for any  $\epsilon \in (0, 1)$  and  $\delta > 0$ , if  $m$ ,  $n$ , and  $k$  stand in the proper relation then

$$P(\|\mathbf{A}\mathbf{u}\|^2 > \delta) > 1 - \epsilon$$

for all  $2k$ -sparse unit vectors  $\mathbf{u}$ .

# The Restricted Isometry Property

**Crude Summary:** For any given  $n$  (number of variables) and  $k$  (solution sparsity), if  $m$  (number of measurements) is large enough and  $\mathbf{A}$  is an  $m \times n$  matrix with suitable random entries, then the RIP of order  $2k$  (so  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique) almost certainly holds.

# The Restricted Isometry Property

**Crude Summary:** For any given  $n$  (number of variables) and  $k$  (solution sparsity), if  $m$  (number of measurements) is large enough and  $\mathbf{A}$  is an  $m \times n$  matrix with suitable random entries, then the RIP of order  $2k$  (so  $k$ -sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique) almost certainly holds.

This, despite the fact that verifying the RIP of order  $2k$  for any sizable matrix is almost impossible.



## An Easy Case: Bernoulli Matrices

Let  $\mathbf{A}$  be an  $m \times n$  matrix with entries that are independent signed-Bernoulli ( $\pm 1$ ), equal probability.

## An Easy Case: Bernoulli Matrices

Let  $\mathbf{A}$  be an  $m \times n$  matrix with entries that are independent signed-Bernoulli ( $\pm 1$ ), equal probability.

Under what conditions on  $m$  and  $n$  can we be confident that 1-sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique? This requires no two columns are parallel.

## An Easy Case: Bernoulli Matrices

Let  $\mathbf{A}$  be an  $m \times n$  matrix with entries that are independent signed-Bernoulli ( $\pm 1$ ), equal probability.

Under what conditions on  $m$  and  $n$  can we be confident that 1-sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique? This requires no two columns are parallel.

A typical column  $\mathbf{a}_i$  of  $\mathbf{A}$  has one of  $2^m$  realizations ( $\pm 1$  patterns).

## An Easy Case: Bernoulli Matrices

Let  $\mathbf{A}$  be an  $m \times n$  matrix with entries that are independent signed-Bernoulli ( $\pm 1$ ), equal probability.

Under what conditions on  $m$  and  $n$  can we be confident that 1-sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique? This requires no two columns are parallel.

A typical column  $\mathbf{a}_i$  of  $\mathbf{A}$  has one of  $2^m$  realizations ( $\pm 1$  patterns).

The probability of another column  $\mathbf{a}_j$  being parallel to  $\mathbf{a}_i$  is  $2/2^m$  or  $1/2^{m-1}$  (because this requires  $\mathbf{a}_j = \pm \mathbf{a}_i$ , two possibilities).

## An Easy Case: Bernoulli Matrices

Let  $\mathbf{A}$  be an  $m \times n$  matrix with entries that are independent signed-Bernoulli ( $\pm 1$ ), equal probability.

Under what conditions on  $m$  and  $n$  can we be confident that 1-sparse solutions to  $\mathbf{Ax} = \mathbf{b}$  are unique? This requires no two columns are parallel.

A typical column  $\mathbf{a}_i$  of  $\mathbf{A}$  has one of  $2^m$  realizations ( $\pm 1$  patterns).

The probability of another column  $\mathbf{a}_j$  being parallel to  $\mathbf{a}_i$  is  $2/2^m$  or  $1/2^{m-1}$  (because this requires  $\mathbf{a}_j = \pm \mathbf{a}_i$ , two possibilities).

There are  $n$  columns, so  $n(n-1)/2$  column pairs to check.

## Boole's Inequality

Boole's inequality states that if  $E_1, \dots, E_N$  are events in some probability space then

$$P(E_1 \cup E_2 \cup \dots \cup E_N) \leq P(E_1) + \dots + P(E_N)$$

where  $P(E)$  is the probability of the event  $E$ .

## Boole's Inequality

Boole's inequality states that if  $E_1, \dots, E_N$  are events in some probability space then

$$P(E_1 \cup E_2 \cup \dots \cup E_N) \leq P(E_1) + \dots + P(E_N)$$

where  $P(E)$  is the probability of the event  $E$ . It's not hard to convince yourself with a Venn diagram.

## Boole's Inequality

Boole's inequality states that if  $E_1, \dots, E_N$  are events in some probability space then

$$P(E_1 \cup E_2 \cup \dots \cup E_N) \leq P(E_1) + \dots + P(E_N)$$

where  $P(E)$  is the probability of the event  $E$ . It's not hard to convince yourself with a Venn diagram.

In plain English, the probability of at least one of the  $E_k$  occurring is no larger than the sum on the right above.



## Coherence for Bernoulli Matrices

For our Bernoulli matrices, the probability that at least one pair of columns is parallel is no larger than

$$\sum_{j=1}^{n(n-1)/2} \frac{1}{2^{m-1}} = \left( \frac{n(n-1)}{2} \right) \left( \frac{1}{2^{m-1}} \right) = \frac{n(n-1)}{2^m}.$$

## Coherence for Bernoulli Matrices

For our Bernoulli matrices, the probability that at least one pair of columns is parallel is no larger than

$$\sum_{j=1}^{n(n-1)/2} \frac{1}{2^{m-1}} = \left( \frac{n(n-1)}{2} \right) \left( \frac{1}{2^{m-1}} \right) = \frac{n(n-1)}{2^m}.$$

Or for simplicity, use  $n^2 - n < n^2$  to see this occurs with probability less than  $n^2/2^m$ .

## Coherence for Bernoulli Matrices

For our Bernoulli matrices, the probability that at least one pair of columns is parallel is no larger than

$$\sum_{j=1}^{n(n-1)/2} \frac{1}{2^{m-1}} = \left( \frac{n(n-1)}{2} \right) \left( \frac{1}{2^{m-1}} \right) = \frac{n(n-1)}{2^m}.$$

Or for simplicity, use  $n^2 - n < n^2$  to see this occurs with probability less than  $n^2/2^m$ .

Equivalently, the probability that no pair of columns is parallel is larger than  $1 - n^2/2^m$ . That is

$$P(\mu(\mathbf{A}) < 1) > 1 - n^2/2^m.$$

# Coherence for Bernoulli Matrices

For example, based on

$$P(\mu(\mathbf{A}) < 1) > 1 - n^2/2^m$$

we can see that if  $n = 100$  and  $m = 20$ , 1-sparse solutions are unique with probability at least  $1 - 10^4/2^{20} \approx 0.99$ .

## Coherence for Bernoulli Matrices

For example, based on

$$P(\mu(\mathbf{A}) < 1) > 1 - n^2/2^m$$

we can see that if  $n = 100$  and  $m = 20$ , 1-sparse solutions are unique with probability at least  $1 - 10^4/2^{20} \approx 0.99$ .

If  $n = 10000$  and  $m = 50$ , 1-sparse solutions are unique with probability at least  $1 - 10^8/2^{50} \approx 0.99999991$ .

## Conclusion

Let  $\mathbf{A}$  be an  $m \times n$  matrix.

## Conclusion

Let  $\mathbf{A}$  be an  $m \times n$  matrix. If the entries of  $\mathbf{A}$  are certain types of random variables then it can be shown that if  $m \geq O(k \ln(n/k))$  then the RIP condition of order  $2k$  (uniqueness of  $k$ -sparse solutions) is met with high probability.

## Conclusion

Let  $\mathbf{A}$  be an  $m \times n$  matrix. If the entries of  $\mathbf{A}$  are certain types of random variables then it can be shown that if  $m \geq O(k \ln(n/k))$  then the RIP condition of order  $2k$  (uniqueness of  $k$ -sparse solutions) is met with high probability.

In many cases this becomes (asymptotically) roughly  $m \geq 2k \ln(n/k)$ .



## Conclusion

Let  $\mathbf{A}$  be an  $m \times n$  matrix. If the entries of  $\mathbf{A}$  are certain types of random variables then it can be shown that if  $m \geq O(k \ln(n/k))$  then the RIP condition of order  $2k$  (uniqueness of  $k$ -sparse solutions) is met with high probability.

In many cases this becomes (asymptotically) roughly  $m \geq 2k \ln(n/k)$ .

So verifying that a specific matrix  $\mathbf{A}$  works is hard, but we can prove that “most” random matrices will work, under suitable conditions.

## Conclusion

Let  $\mathbf{A}$  be an  $m \times n$  matrix. If the entries of  $\mathbf{A}$  are certain types of random variables then it can be shown that if  $m \geq O(k \ln(n/k))$  then the RIP condition of order  $2k$  (uniqueness of  $k$ -sparse solutions) is met with high probability.

In many cases this becomes (asymptotically) roughly  $m \geq 2k \ln(n/k)$ .

So verifying that a specific matrix  $\mathbf{A}$  works is hard, but we can prove that “most” random matrices will work, under suitable conditions.

One rule of thumb people have formulated is that  $m > 4k$  is often sufficient.