

Emerging Role of Precision Medicine in Cardiovascular Disease

Jane A. Leopold, Joseph Loscalzo

Abstract: Precision medicine is an integrative approach to cardiovascular disease prevention and treatment that considers an individual's genetics, lifestyle, and exposures as determinants of their cardiovascular health and disease phenotypes. This focus overcomes the limitations of reductionism in medicine, which presumes that all patients with the same signs of disease share a common pathophenotype and, therefore, should be treated similarly. Precision medicine incorporates standard clinical and health record data with advanced panomics (ie, transcriptomics, epigenomics, proteomics, metabolomics, and microbiomics) for deep phenotyping. These phenotypic data can then be analyzed within the framework of molecular interaction (interactome) networks to uncover previously unrecognized disease phenotypes and relationships between diseases, and to select pharmacotherapeutics or identify potential protein–drug or drug–drug interactions. In this review, we discuss the current spectrum of cardiovascular health and disease, population averages and the response of extreme phenotypes to interventions, and population-based versus high-risk treatment strategies as a pretext to understanding a precision medicine approach to cardiovascular disease prevention and therapeutic interventions. We also consider the search for resilience and Mendelian disease genes and argue against the theory of a single causal gene/gene product as a mediator of the cardiovascular disease phenotype, as well as an Erlichian magic bullet to solve cardiovascular disease. Finally, we detail the importance of deep phenotyping and interactome networks and the use of this information for rational polypharmacy. These topics highlight the urgent need for precise phenotyping to advance precision medicine as a strategy to improve cardiovascular health and prevent disease. (*Circ Res.* 2018;122:1302–1315. DOI: 10.1161/CIRCRESAHA.117.310782.)

Key Words: genomics ■ polypharmacy ■ precision medicine ■ proteomics ■ systems biology

It is because we are not exact that we fail

—Paul Ehrlich, 1896.¹

Over the past 50 years, progress toward the eradication of cardiovascular disease has been achieved through the adoption of lifestyle modifications, including dietary, tobacco, and exercise interventions, as well as evidence-based therapies that aim to modify a recognizable and commonly shared cardiovascular or at-risk phenotype. Despite the success of this approach, the related issues of disease prevention and cure have been elusive, presumably because of imprecise deep phenotyping of individuals needed to characterize subgroups of disease. The magnitude of the problem remains considerable. At present, there are 92.1 million adults (>1 in 3) in the United States who have been diagnosed with cardiovascular disease, with a projection that by 2030, at least 44% of the adult population will have this diagnosis.² Although death rates attributable to cardiovascular disease have declined ≈25% from 2004 to 2014 in the United States, cardiovascular disease remains the leading cause of death worldwide accounting for ≈32% of all global deaths, with the expectation that this number will rise to >23.6 million deaths annually by 2030.^{3,4} The contribution of attendant comorbidities and traditional risk factors (eg, hyperlipidemia, hypertension, diabetes mellitus,

metabolic syndrome, and chronic kidney disease), health behaviors (smoking and tobacco use, physical inactivity, nutrition, overweight, and obesity), and other immutable factors (family history) to morbidity and mortality is understood and likely underlies the sex-, age-, race-, ethnic-, regional-, and economic-based differences in disease burden (reviewed in ⁴). A recent meta-analysis stressed the importance of these comorbidities to disease risk. This study, which included data from 9 prospective cohort studies that followed 12878 individuals, found that working toward optimal cardiovascular health metrics to achieve the greatest possible benefit was associated with a decrease in the risk for major adverse cardiac events, including cardiovascular mortality (relative risk [RR], 0.25; 95% confidence interval: 0.10–0.63).⁵ Taken together, there is broad heterogeneity in the clinical profiles and outcomes of individuals with cardiovascular disease, thereby highlighting the urgent need for precision phenotyping.

The current approach to reducing cardiovascular morbidity and mortality in at-risk individuals or those with established disease is based on a traditional reductionist approach and uses a multitiered system with input from patients, the physician and the medical system, and evidence-based medicine at large. The first layer involves patient identification of symptoms that represent a deviation from normal or baseline.

From the Brigham and Women's Hospital and Harvard Medical School, Boston, MA.

Correspondence to Joseph Loscalzo, MD, PhD, Brigham and Women's Hospital, 75 Francis St, Boston, MA 02115. E-mail jloscalzo@bwh.harvard.edu

© 2018 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.117.310782

Nonstandard Abbreviations and Acronyms

GWAS	genome wide association study
SNP	single nucleotide polymorphism

This report initiates entry into the medical system where contact with a physician results in a personalized assessment of the physical signs of disease and implementation of evidence-based therapies whose efficacy is supported by results from clinical trials. Although this paradigm of care can ameliorate symptoms and affect disease progression, this outcome is not always certain, especially when targeting a chronic complex illness like atherothrombotic cardiovascular disease that does not have a single root cause. To address this issue, it is necessary to understand the totality of cardiovascular disease at a granular and integrative molecular level.

Although all patients are fundamentally unique, reductionism in medicine presupposes that patients with common signs and symptoms share the same disease pathophenotype and, therefore, will respond similarly to medical, procedural, and behavioral interventions tested in aggregates of like individuals.⁶⁻⁸ It also places the focus squarely on treatment of established cardiovascular disease without addressing health, prevention, timing of disease inception, or cure and eradication.⁹ Owing to advances in panomics (genomics, transcriptomics, epigenomics, metabolomics, proteomics, and microbiomics), that is, technologies and data analysis that provide in-depth clinical, biological, and molecular phenotyping, there is growing awareness that this conventional approach may be an overly simplistic view of the multiple contributors to and complexity of an individual's cardiovascular disease phenotype. Similarly, it is now understood that other factors, such as exposure to the natural, personal, and social environments, contribute to an individual's highly personalized disease phenotype (reviewed in ¹⁰). Integration of this large body of data points lends itself toward a more exacting (patho)phenotype, one that is amenable to precision medicine.

Why precision medicine? Precision medicine represents a new strategy in the approach to care by targeting prevention and treatment while considering individual differences in genetics, exposures, and lifestyle and health factors that are determinants of a person's disease phenotype (Figure 1). The goal of precision medicine is to identify optimal care for an individual based on a unique personal profile rather than that of the average population. The power of precision medicine lies in the data and requires the synthesis of rapidly changing data sets, ranging from standard clinical, imaging, and laboratory testing to next-generating sequencing, metabolomics, and proteomic studies to historical health record data. Data derived from these sources may also be analyzed using advanced systems biology and network analytical methods to uncover new and unbiased relationships between health and disease factors. The optimal system will also evolve rapidly, learn, and be facile enough to be useful for prevention, diagnosis, and treatment across a broad range of cardiovascular health factors, risk factors, and diseases. The evolution of precision medicine and its application to cardiovascular disease hold the promise

of improving health as well as revolutionizing prevention and treatment options similar to what has occurred in the field of oncology.

Population Averages and Probability Density Function

To move precision medicine into the cardiovascular arena, it is necessary to identify and adopt parameters that reflect what is considered the normal or ideal range (Figure 2A). This is especially important given the diversity in population age and ethnicity where normal standards may change over time. Within the general population, there is variability with some individuals who have ideal health, others who have recognizable disease risk factors, and those who have established disease. Within groups, the transition between these health states tends to be less clear with there generally being a continuum with overlap between categories.⁹ Examples of this point are blood pressure and blood cholesterol levels. In each case, there are individuals with normal levels who are considered healthy, those with increased levels in the absence of cardiovascular disease, those with high levels and disease, as well as those who do not fit into any clear category, and are at an overlap of states.⁹

In clinical trials, this continuum of disease is often simplified with the assumption that all participants have the same phenotype, which can be represented using a typical bell-shaped curve with a discernable median measurement. Individuals entered in the clinical trial are considered based on the variable of interest, such as their cholesterol level, blood pressure, or history of myocardial infarction, and not their complex phenotype. This oversimplification allows results to be described in terms of an average finding on a per group basis. Although this paradigm has been the norm, the hazard of assuming similarity is that only relatively large differences may be detected as significantly different, and the actual range of responses may be undetected.⁹ This concept was illustrated when patients with chronic heart failure were rephenotyped using cluster analysis. This analysis revealed significant heterogeneity between patients with individuals segregating into groups that would not have been predicted based on their initial clinical phenotype.¹¹ In precision medicine, similarity in phenotypes is not assumed for the purpose of studying a population behavior or response. Instead, the goal is to deviate from population averages as the metric and focus on the individual's unique phenotype with an eye toward identifying a response to treatment as one that moves the individual from disease to health and those at health to ideal health.⁹

Responses of the Tails of the Distribution and Implications

The importance of phenotyping and identifying an ideal level of health within community-based populations has implications for defining what constitutes a high-risk phenotype or those individuals who fall in the tail(s) of the distribution curve. For example, individuals with blood pressure or blood cholesterol levels at the population extreme are considered high risk. There have been 2 strategies considered when deciding how to reduce adverse events: a high-risk strategy and a population-based strategy. The high-risk strategy targets only those individuals who fall within




Figure 1. A precision medicine approach to phenotyping. Individuals may have a similar endophenotype but be biologically distinct and have different disease profiles. Using a precision medicine approach, individuals undergo deep phenotyping with data analysis performed using network analysis. This analytic strategy clusters individuals into groups that are different from those based on endophenotype alone. This methodology can be used to optimize medication and behavioral changes to improve health, predict and prognosticate disease, identify biomarkers for disease, enrich clinical trial enrollment, and optimally tailor exposures.

the tail of the distribution curve and have the greatest risk for disease in an attempt to lower levels to what is considered normal or ideal. However, the majority of cases of cardiovascular disease occur in individuals who fall in the average risk group. A population-based strategy targets this latter group, focuses on treating more individuals in the population, and aims for a smaller population-based risk factor reduction. This approach follows the argument put forth by Geoffrey Rose, viz, that shifting the risk distribution curve by only a small amount within a population will have greater societal benefit in reducing or preventing cardiovascular morbidity and mortality than treating only high-risk patients.¹² Others have also argued that significant reduction in the population burden of cardiovascular disease can only occur from a population approach shifting the entire population distribution to lower levels.¹³ Despite the epidemiological cogency of

this argument, adopting a population-based strategy that shifts the population mean also shifts the distribution of individuals who fall within the tails and, therefore, cross thresholds to initiate or stop treatment (Figure 2B). The net outcome can be either beneficial or harmful. Consider differences in the threshold for treatment of cholesterol using guidelines established by the American College of Cardiology/American Heart Association (ACC/AHA) as compared with the US Preventive Services Task Force. Differences in the guidelines suggest that shifting to the US Preventive Services Task Force recommendations from the ACC/AHA guidelines resulted in $\approx 9\%$ of individuals who met ACC/AHA guideline criteria for statin use for primary prevention no longer being recommended for treatment; 55% of this group are younger adults with high mean long-term risk for cardiovascular disease.¹⁴




Figure 2. Population distribution and effect of changing the population mean. **A**, Within a population, the majority of people express a phenotype that centers around a mean with a minority of individuals who exhibit an extreme or high-risk phenotype. **B**, A change in the population average affects the number of individuals who are considered to be at the extremes. This may also influence the number of individuals who reach a predefined threshold for initiating treatment.

Using data from the National Health and Nutrition Examination Survey III study, investigators performed a head-to-head comparison of population-based and high risk-based treatment strategies using a low- or moderate-intensity intervention to determine which had the greatest benefit for reducing cardiovascular events and cardiovascular mortality. The study compared a population-based strategy that treated all, a high-risk strategy that selected individuals within the top 25% low-density lipoprotein cholesterol level, and a high-risk strategy based on individuals within the top 25% for cardiovascular risk as determined by a risk prediction tool. In this case, the investigators found that the high-risk strategy focused on individuals within the top 25% of cardiovascular risk was the most efficient method for preventing cardiovascular events over the long-term but was comparable to a population-based strategy with a more moderate interventional goal. This finding led to the conclusion that while the high-risk treatment strategy was the most efficacious, the population-based prevention strategy was a reasonable option if the intervention had no or minimal side effects.¹³ In contrast, in developing low- and middle-income countries, the high-risk strategy should be used, especially when there are few resources available.¹⁵

Search for Single Pathogenic Genes/Gene Products and Its Flaws

The standard view that all or most cardiovascular diseases have a heritable component has fueled the search for single pathogenic genes for specific disorders. Although cardiovascular disease is significantly broad and encompasses diseases related to blood vessels, the myocardium, heart valves, the conduction system, and developmental abnormalities, there are only a few cardiovascular disorders that can be attributed to a single pathogenic gene. Nonetheless, there are notable examples of monogenic disorders that cause cardiovascular disease, such as a mutation in the low-density lipoprotein receptor (*LDLR*) gene that causes familial hypercholesterolemia,^{16,17} in the β -myosin heavy chain and other sarcomeric proteins that are causal for hypertrophic cardiomyopathy,^{18,19} and a mutation in the fibrillin (*FBN1*) gene that causes Marfan syndrome among others.²⁰ In contrast, atherosclerosis and myocardial infarction are multifaceted diseases with a complex inheritance, cannot be explained by a single pathogenic gene, and are more likely because of perturbations of large, and possibly phenotypically related, genes and other environmental factors.²¹

There are several inherent limitations associated with the search for a single pathogenic gene in complex cardiovascular diseases, and these are exemplified in known monogenic disorders. The first is that the genotype–phenotype relationship can be difficult to ascertain. For example, mutations in the α subunit of the type V voltage-gated sodium channel (*SCN5A*) was initially described as a single causal gene for inherited long-QT syndrome.²² Since that time, other phenotypes have been associated with the mutation, including Brugada syndrome and dilated cardiomyopathy.^{23,24} The second consideration involves penetrance and expressivity with examples of incomplete penetrance in a pedigree with familial hypercholesterolemia and an *LDLR* mutation, and differences in disease expression in families with Marfan syndrome and the same heritable mutation in *FBN1*.^{25,26} Incomplete penetrance of these classic

Mendelian disorders simply suggests other interacting genetic variants and environmental exposures on their pathogenesis.

The limitation of genome-wide association studies (GWAS) as a method designed to implicate a single pathogenic gene, or a limited number of pathogenic genes, is that performing genetic mapping by association only identifies genomic regions that may contribute to the disease process. Although coverage of the genome is improving, the singular information provided by these data are that a pathogenic gene resides nearby with distance determined by the sequencing platform. Results from GWAS support the concept that complex disorders possess genetic heterogeneity of variants (common, low-frequency, or rare), and this is likely the most frequent expression pattern in these types of diseases. This point was exemplified in a GWAS meta-analysis of individuals with myocardial infarction and control subjects. In this study of 60 801 individuals with a myocardial infarction and 123 504 controls, there were common and rare variants identified accounting for genetic heterogeneity.²⁷

Another argument against a single pathogenic gene for atherothrombotic vascular disease and myocardial infarction is illustrated by the relationship between chromosome 9p21 and coronary artery disease or myocardial infarction. Single nucleotide polymorphisms (SNP) in 9p21 were identified by several independent GWAS, and each risk allele was associated with a 29% increased risk of cardiovascular disease.^{28–30} The findings that the SNPs were in noncoding regions, the nearest genes were >100 Kb away, and causality between the nearest genes (*CDKN2B*, *CDKN2A*) and susceptibility to atherosclerosis have not yet been ascertained all argue against the single pathogenic gene explanation.^{28–30} Considering the history of GWAS and the observation that these studies have not identified a single pathogenic gene or SNP as the mechanism underlying coronary artery disease or myocardial infarction indicates further that searching for a single element to explain a complex phenotype is not sufficient. Instead, it is more likely that results from GWAS and next-generation sequencing will contribute to a mechanistic exploration of molecular networks (see below) and to the need for concomitant deep phenotyping multifaceted cardiovascular diseases, such as coronary artery disease and myocardial infarction.

Resilience and Mendelian Disease Genes

Advances in next-generation sequencing have expanded our understanding of the genetic basis of cardiovascular diseases in particular and of all human disease in general. There are >150 000 disease-related genetic variants mapped to >6000 Mendelian disorders (Online Mendelian Inheritance in Man) that have been catalogued in the Human Gene Mutation Database.^{31,32} Although this information is readily available, there are relatively few therapies developed that effectively treat or cure these Mendelian diseases.³³ For common complex diseases, such as atherothrombotic cardiovascular disease, few gene variants have translated directly into predictors of disease risk or severity, thereby limiting their potential to be developed into a diagnostic or therapeutic.

More recently, attention has turned to examining the genetic and environmental factors that predispose to resilience, the ability to adapt successfully to stress. In the context of

Mendelian disease, resilience implies the ability to withstand changes to fitness in the presence of disease-associated mutations.³⁴ The role of second site mutations or environmental modifications that promote resilience to disease has also been established in preclinical models.^{35,36} In addition, clinical studies that have found highly penetrant disease-causing mutations in individuals who do not manifest the disease phenotype, suggesting that there are either genetic or environmental resilience factors that protect individuals from manifesting disease.^{37,38} A healthy lifestyle, defined as the absence of obesity, no current tobacco use, a healthy diet, and regular physical activity, has been shown to serve as an environmental resilience factor and modify genetic risk of cardiovascular disease. In a study of 55 685 individuals who were stratified according to a polygenic risk score comprised 50 SNPs associated with coronary artery disease, individuals with a high genetic risk with a healthy lifestyle had a 46% reduction in the relative risk of coronary events.³⁹

Although many genetic studies have attempted to identify mutations linked to disease, an alternative is to search for genetic factors that prevent disease, or wellness factors, in the presence of Mendelian disease-causing mutations. Although attempts to find resilience factors have been limited, some secondary cardiovascular disease modulators have been identified, including a loss-of-function mutation in pro-protein convertase subtilisin/kexin type 9 and mutations in zinc transporter 8 that protect obese individuals from diabetes mellitus.^{35,40–42}

A more recent study used a different tactic to find resilience factors and studied the genomes of 589 306 individuals who were apparently healthy. Using this approach, they identified 13 adults with what had been viewed as completely penetrant Mendelian disease mutations for 8 different severe childhood diseases, including cystic fibrosis, atelosteogenesis, and Smith–Lemli–Opitz syndrome, but who were without any clinical manifestations of disease. This finding suggested that these individuals have some form of genetic protection or resilience against even these classical Mendelian disorders; however, it was not possible to isolate the resilience factor(s) because of the data available and study design.⁴³

Search for the Ehrlichian Magic Bullet and Its Flaws

The complexities of the cardiovascular disease phenotype also highlight the fact that there is no single therapy, or Ehrlichian magic bullet, that will cure a disease. In 1900, the German Nobel laureate Paul Erlich, a pioneer of chemical biology and chemotherapy, put forth the concept of the magic bullet. He hypothesized that there would be a way to target specifically disease-causing microbes in the same way a bullet fired from a gun hits its target. He referred to this hypothetical agent as *Zauberkugel* or magic bullet. His continued work to cure syphilis using a magic bullet led to the discovery of Salvarsan (an arsenical compound) on August 31, 1909, the 606th compound he tested that was efficacious and free of important side effects.⁴⁴

The magic bullet concept was soon popularized, and its meaning expanded to include a perfect drug to cure a specific disease that acted without side effects. The goal would be achieved by identifying a compound that targets a specific

disease-related molecule and does so in a highly specific manner with no off-target effects. This idea has become an enduring part of cardiovascular (and all of) medicine even though there are numerous examples of therapies or interventions for cardiovascular disease that have been hailed as a magic bullet yet have failed to provide a cure, including stem cells and statins. In fact, this is not surprising because cardiovascular diseases are complex pathophenotypes with significant biological diversity owing to numerous genetic, metabolic, and environmental mediators.

Thus, the idea of a single target in the original, strictest definition of a magic bullet is unlikely to have broad applicability in cardiovascular diseases. In other words, the use of a mono-specific drug(s) to target a multifactorial disease will not be successful. This has led to the idea of targeting multiple targets at once (eg, the polypill). Although this strategy may improve the pharmacological profile, there is also the increased risk of adverse effects. Biologicals such as monoclonal antibodies are less likely to have off-target effects but may show variations in efficacy and unknown off-target effects or on-target toxicity (eg, abciximab and severe bleeding or immune checkpoint inhibitors and autoimmune diseases). This approach has evolved further to fall within the rubric of precision medicine via network-targeted therapy that uses a combination therapy strategy targeting multiple steps in a signaling pathway or network identified by deep phenotyping. This more complex approach could be considered a modern interpretation of Ehrlich's magic bullet theory in the contemporary era.^{44,45}

Importance of Molecular Interaction (Interactome) Networks

Using data from genomic and other panomic profiling, data can be analyzed using molecular interaction or interactome networks to discover disease associations and possible therapeutic targets (Figure 3). Interactome networks play an increasingly important role in the discovery of novel relationships between genes or proteins that may have broad applicability across a spectrum of human diseases. The interactome defines a comprehensive, unbiased set of biologically relevant molecular (ie, gene, protein, metabolite) interactions in a cell, organ, or person. The advantage of examining the interactome to understand disease pathobiology is that the interactome allows for the simultaneous consideration of (ideally) all or a large number of relevant genes, metabolites, and proteins, as well as their interactions as depicted through network representation. These networks and subnetworks allow for discovery of new interactions, ascertainment of pathways governing phenotype, identification of critical regulatory checkpoints in a biological system, and prediction of the consequences of disease in network associations or pathway functions on phenotype. The interactome eliminates the bias introduced by analyzing a data set using a reductionist approach, which examines a limited number of interactions in the data set to find a differentially expressed gene or protein associated with a phenotype without consideration of the network context within which it operates.^{46,47} The current human protein–protein interactome contains 13 460 proteins that have 141 296 physical interactions (protein–protein, metabolic pathways, and kinase




Figure 3. Molecular interaction networks (interactome). The protein–protein interactome describes interactions between proteins in a scale-free (ie, interactions are not random but clustered) manner. Within the interactome, groups of proteins reside in disease neighborhoods (colored areas) that may overlap indicating common mediators (**left, center**). Within the neighborhood, there are several disease modules or groups of protein interactions that are related. Analysis of the interactome may uncover new or previously unrecognized relationships. Gray circles, nodes (ie, proteins in a protein–protein interaction network); black lines, edges (ie, interactions).

substrate).⁴⁸ The interactome network is scale free, indicating that interactions are not random, and the proteins within it exhibit emergent behavior such that detailed information about 1 protein does not predict its functional response within the context of the interactome or of the interactome itself.⁴⁹ Although the human interactome is incomplete owing to the fact that the majority (80%) of pairwise protein interactions have not been analyzed using currently available high-throughput methodology, it remains a valuable tool for discovery of disease-related phenomena.⁴⁸

The human interactome has been shown to contain subnetworks for specific diseases, so-called disease modules. Certain disease modules overlap with one another, and these diseases typically have similar symptoms, patterns of coexpression, and comorbidities while diseases that reside in separate neighborhoods in the network do not share these characteristics and are clinically distinct. Analysis of disease relationships has shown that 7% of disease pairs have overlapping disease neighborhoods in the interactome with unexpected disease relationships, such as that between asthma and celiac disease, which may share a molecular phenotype despite differences in their pathobiology. Interrogation of the shared factors between asthma and celiac disease revealed pathways for immunoglobulin A production as a common intermediary.³³ Interactome-based network analysis has been used to inform GWAS data. For example, the great majority of genes associated with type 2 diabetes mellitus have small effect size and limited statistical significance when analyzed in isolation; however, mapping these alleles to the interactome leads to the identification of novel interactions in pathway clusters that provide unique, highly statistically significant information on mechanism not appreciated by conventional GWAS (Figure 4). Another capability of the interactome is to improve comprehension of the molecular actions of drugs with a goal of identifying drug targets, the potential for adverse reactions, and the possibility of repurposing drugs based on newly identified relevant molecular targets. This concept was investigated for myocardial infarction using a network that incorporated myocardial infarction disease genes, drugs, drug targets, and drug target

interactors or binding partners in the interactome to reveal that many myocardial infarction–related drug targets and disease proteins have close relationships within the disease module in the interactome, and some of the drug target intermediaries are themselves targets for other drugs suggesting potential drug repurposing strategies.⁵⁰ Studies of the interactome have also been used to identify tissue specificity of modules associated with human disease. This has shown that an entire functional subnetwork or pathway incorporating disease gene products, or a disease module, must be expressed within a particular tissue for a disease to be manifest in that tissue, not simply expression of the disease gene product itself. This type of analysis also leads to the creation of a disease–tissue network and allows for the discovery of unexpected disease–tissue associations.⁵¹

Importance of Phenotype

A core principle of precision medicine is establishing the precise phenotype for any given disorder (Figure 5). The ability to understand and recognize the relationship between the elements that comprise health phenotypes and disease (patho) phenotypes is required for precision medicine to improve individual and population outcomes.^{52,53} The foray into precision phenotyping has faced some obstacles, including the unknown function of many genes or metabolites, the incomplete descriptors used to categorize diversity within disease phenotypes, and inadequate data collection or extraction from medical records.^{52,54} These shortcomings have led to the characterization of most phenotype descriptions as imprecise and highlight the necessity of a unified descriptive ontology that is widely adopted. This widely recognized need has led to the proposal that the Human Phenotype Ontology be used as a standard to characterize disease phenotypes using a specific vocabulary; however, it remains to be seen if this system will achieve widespread use.⁵⁵

The importance of obtaining a more detailed, granular phenotype for cardiovascular diseases is underscored by the fact that phenotypic profiling is lagging with respect to panomics technologies, and, that phenotypic information is necessary




Figure 4. Identifying genome-wide association studies (GWAS) genes that have biological relevance. Analysis of GWAS data reveals that there are a large number of genes that, while having genome-wide significance, have modest effect sizes. **A**, Using data from patients with type 2 diabetes mellitus as an example and starting with a module that includes 6 genes of greatest statistical significance, GWAS genes are added to the module in decreasing order of their P value (top). The statistical significance of the growing network as compared with a random network is shown (middle) as well as the size of the resulting module (bottom). **B** and **C**, Of 77 significant GWAS genes, only 5 connect to the interactome cluster initially; however, the addition of the interactor gene (product), *CALM2*, joins disconnected parts of the network and increases both its size and significance. Another disconnected cluster is then joined to the module through consideration of 200 GWAS genes, including the *KCNJ11* (GWAS significant gene) and *ABCC8*. Reproduced from Menche et al⁴⁸ with permission. Copyright ©2015, American Association for the Advancement of Science. OMIM indicates Online Mendelian Inheritance in Man.




Figure 5. The importance of phenotype and its molecular network underpinning. Individuals who share a common endophenotype, such as hypertension or hypercholesterolemia, may have a different phenotype when examined at a molecular level. Using molecular phenotyping, such as genome sequencing, and network analysis, individuals may cluster into distinct phenotypes. These phenotypes have important implications for disease risk, prognosis, or response to medication. These important differences are not discoverable when relying on endophenotype alone (blue nodes, normal or major alleles; red nodes, variant or minor alleles in protein–protein interaction network).

to deconvolute the results of genomic, transcriptomic, metabolic, and proteomic studies. Only recently has the approach to clinical phenotyping moved beyond historical metrics, such as blood pressure, to improve stratification of cardiovascular diseases. Current phenotypic assessments are beginning to incorporate standard clinical traits, such as blood pressure, as well as other complex characteristics, like social, environmental, and personal exposures.^{10,56} Phenotypes are also being assessed continually using personal sensors and wearable devices, illustrating the importance of time trajectories in monitoring early identifiers of disease progression.^{57,58}

There are several recent examples of the use of advanced phenotyping methodologies to discover differences within individuals with a clinical syndrome that has underlying heterogeneity. Clinical phenomapping of patients with heart failure and preserved ejection fraction using unbiased hierarchical clustering analysis revealed 3 distinct groups of patients, with 1 phenogroup having an increased risk of heart failure hospitalization (hazard ratio: 4.2; 95% confidence interval: 2.0–9.1; $P < 0.001$).⁵⁸ Similarly, cluster analysis of 1619 patients with chronic heart failure identified heterogeneity within the group as a whole and 4 phenotypic clusters of patients with differences in their risk of all-cause mortality or all-cause hospitalization.¹¹

The concept of an unbiased or undirected phenome-wide assessment of an individual is also entering the realm of possibility and has been conducted in a single person who was extensively phenotyped during the course of 18 months. In the MyConnectome project, brain function over time was assessed with imaging, functional studies, health surveys, and genomic and metabolomic profiling with data from the latter

examined using network analysis. Using this platform, gene and metabolite expressions were related to dynamic changes in brain connectivity over a timescale of days to months. This pioneering study illustrates the feasibility of this deep temporal phenotyping approach and provides a strategy for translating these types of studies into the cardiovascular space.⁵⁹

Endophenotype Diversity and Pre-Emptive/Preventive Strategies

The endophenotype, or intermediate phenotype, is a biological trait that is quantitative and reflects the function of a biological system. It is heritable and is more closely related to the root cause of a disease than the broad clinical phenotype.^{60,61} GWAS that examine conventional phenotypes can be improved upon by focusing on individual endophenotypes.⁶² Endophenotypes are themselves thought to be subject to natural selection leading to the concept that there are evolutionarily established molecular mechanisms, which regulate endophenotypes, and that these factors translate into genetic predisposition for the correlate downstream clinical phenotypes.^{63–65} This view leads to the expectation that complex molecular mechanisms link genes responsible for both endophenotypes and phenotypes.⁶⁶ Thus, genetic effects on endophenotype and phenotype do not necessarily have to be the same in different populations, and genetic predisposition to an endophenotype does not necessarily translate into predisposition for a downstream phenotype.

Several GWAS studies have illustrated the diverse relationships between endophenotypes and phenotypes. For example, 1 study examined the association of the rs693 and rs562338 polymorphisms in the apolipoprotein B locus with total cholesterol and high-density lipoprotein cholesterol (endophenotype) with myocardial infarction and survival (phenotypes) across 4 population-based studies (MESA [Multi-Ethnic Study of Atherosclerosis], ARIC [Atherosclerosis Risk in Communities], FHS [Framingham Heart Study], CHS [Cardiovascular Health Study]). The endophenotypes were investigated to determine whether they could mediate the associations of the polymorphisms with myocardial infarction or mortality. In this analysis, the SNPs rs693 and rs562338 were robustly associated with the endophenotype (total cholesterol) but had a trivial association with the phenotype (myocardial infarction and survival) despite the well-known relationship between lipids and myocardial infarction.⁶⁷

Network analysis has been used to examine endophenotypes and their overlapping relationships to different diseases. Inflammation, thrombosis, and fibrosis are endophenotypes that are linked pathologically and common to virtually all disease states, both in terms of response to injury and to repair.^{68–71} To explore biological and topological crosstalk between these endophenotypes, network analysis of the inflammasome, thrombosome, and fibrosome was undertaken. Mapping these networks to the human interactome identified inflammation, thrombosis, and fibrosis subregions that were highly overlapping and contained disease genes associated with complex diseases and cardiovascular risk factors. These subregions were found to be integral to the structure of the basic interactome network, thereby highlighting the key role of these processes in (health and) disease.⁷²

With the advent of panomics profiling, endophenotypes and their association with complex disease phenotypes will be refined further. To ensure the integrity of these relationships, however, there is an implicit need for standardization of methodology and platforms, as well as adoption of CLIA (Clinical Laboratory Improvement Amendments)-grade laboratory testing with strict adherence to quality control. Analytical methods to integrate endophenotypic data should favor unbiased approaches, such as network analysis, and account for issues related to multiple testing, which is an inherent limitation of large data set analysis.

Genetic Diversity and Rational Polypharmacy

Although randomized clinical trials have determined the efficacy of many new therapeutics across large populations of individuals with similar clinical phenotypes, heterogeneity in the response to drugs remains a relevant and challenging issue. Aggregating patients with similar disease characteristics into a cohort deemed to be putative therapeutic responders with equal effect obviously ignores this heterogeneity for clinical trial design and therapeutic simplicity even though there is no phenotypic basis to support such an oversimplified categorization (Figure 6A).⁵⁶ Pharmacogenetic variation because of single and cumulative drug-gene interactions can lead to adverse drug interactions^{73,74}; these types of reactions increase with polypharmacy. Approximately 66% of adults age ≥ 65 years have at least ≥ 1 prescription medications that require daily use.^{75–77} It is estimated that $\approx 35\%$ of geriatric patients have adverse drug events, at least half of which are

deemed preventable; and 10% to 17% of hospitalizations in this population are related to drug reactions leading to 51% of adverse drug-related deaths.^{77–79} The application of precision medicine to this problem has revealed that cardiovascular pharmacogenetics can be used to identify genetic diversity markers associated with poor response or adverse reactions to drugs. In fact, drug interactions related to predisposing genetic factors lead to $\approx 47\%$ of warnings of potential interactions associated with side effects.⁷⁹

There are numerous examples of genetic diversity and DNA variants as determinants of response to a drug. In cardiovascular disease, platelet aggregation promotes atherothrombotic vascular disease and stent thrombosis after percutaneous coronary intervention. The P2Y₁₂ inhibitor clopidogrel, which is part of a typical dual antiplatelet regimen, has a range of interindividual variability. Some individuals are known to be clopidogrel nonresponders, a phenotype that has been associated with genetic determinants and an increase in ischemic events.^{80–83} Loss-of-function alleles in CYP2C19 (CYP2C19*2 and CYP2C19*3) have been associated with poor drug responsiveness while the gain-of-function allele CYP2C19*17 is associated with increased bleeding risk.^{84–86} Attempts to use this information for tailored therapy in patients with drug-eluting stents and high on-treatment platelet reactivity, however, did not affect cardiovascular death, myocardial infarction, or stent thrombosis.⁸⁷

It has been suggested that $\approx 50\%$ of patients taking statins stop taking their medications owing to side effects or adverse events.⁸⁸ Of the related genes, only the gene for the soluble




Figure 6. Precision medicine approach to rational polypharmacy. **A**, The current approach to selecting pharmacological therapy involves identifying a group with a common endophenotype, using genomic profiling in select cases to identify potential drug (non) responders, and then testing a medication based on results from clinical trials. **B**, Using a precision medicine approach, individuals with a common endophenotype undergo deep phenotyping, including panomics, exposures, and clinical assessments, and the data are analyzed using network analysis to identify more precise phenotypes and their molecular determinants in the interactome (gray nodes, disease-determining module on network; red, blue, and green nodes, drug targets by phenotype). This information is then used to select agents that target key pathways or proteins.

carrier organic anion transporter 1B1 (*SLCO1B1*), which regulates statin influx and metabolism in the liver, has been associated consistently with myopathy.⁸⁸ Patients who are either homozygous or heterozygous for the rs4263657 polymorphism have an increased risk for rhabdomyolysis with statin use, and it has been suggested that individuals with this SNP avoid statin drugs.⁸⁹

Warfarin, one of the most commonly prescribed anticoagulants, has a narrow therapeutic window but wide interindividual variation.⁹⁰ Warfarin is metabolized primarily in the liver by oxidation by cytochrome P450 2C9 (*CYP2C9*) and inhibits the protein vitamin K epoxide reductase complex subunit 1 (*VKORC1*).^{91,92} Studies have attributed the ~10% to 50% variability in dose requirements to genotypes, notably SNPs in *CYP2C9* (*CYP2C9*2*, *CYP2C9*3*) and *VKORC1* (rs9923231).^{93–95} The Food and Drug Administration has recognized the importance of these genetic variants and updated the drug packaging to include information on dosing based on *CYP2C9* and *VKORC1* genotypes.^{96,97} Two randomized clinical trials, the COAG (Clarification of Optimal Anticoagulation through Genetics) trial and the EU-PACT (European Pharmacogenetics and Anticoagulant Therapy-Warfarin) study, evaluated genotype-guided warfarin dosing.^{98,99} These studies incorporated *CYP2C9* and *VKORC1* genotyping into the clinical algorithm for selecting the warfarin dose. The 1015 patient COAG trial reported that genotyping had no effect on the percent of time in the therapeutic international normalized ratio range while the 455 patient EU-PACT study found that genotyping was useful, resulted in fewer episodes of excess anticoagulation (international normalized ratio levels ≥ 4.0), and led to more rapid achievement of the therapeutic range (21 versus 29 days) than those who were not guided by genotype.^{98,99} The differences observed between these trials has been attributed to trial design, prevalence of the SNPs in the study population, and the demographics of the study populations.¹⁰⁰

Genetic variants have also been identified that modify the response to other relevant cardiovascular drugs, including β -blockers (*ADRB1*, *ADRB2*, *GRK5*, *GRK4*); angiotensin-converting enzyme inhibitors (*ACE*, *AGTRI*); diuretics (*ADD1*, *NPPA*, *NEDD4L*); and calcium channel blockers (*CACNB2*, *CACNA1C*; reviewed in ¹⁰¹). To date, however, genetic testing has not been used routinely to guide selection of these drugs, and many genetic variants require confirmation in larger studies.

Other pilot studies have examined the efficacy of implementing pharmacogenetic profiling coupled with a clinical decision support tool in polypharmacy patients to reduce adverse drug reactions and emergency department visits. In a small study of 110 patients randomized to genetic profiling of CYP 450 genes and prescribed relevant medications, the use of pharmacogenetic profiling was associated with a reduction in the number of emergency department visits (RR, 0.58; 95% confidence interval: 0.34–0.99; $P=0.045$) and hospitalizations at 60 days (RR, 0.48; 95% confidence interval: 0.27–0.82; $P=0.0007$). Interestingly, of 124 recommendations on drug therapy based on pharmacogenetic profiling that were passed on to the patient's clinician, only 96 (77%) were followed. Although exploratory, this study highlights the benefits of

identifying genetic diversity to achieve rational polypharmacy and clinician attitudes toward this approach.¹⁰²

Pharmacogenetic profiling has also been used to determine whether individuals treated with polypharmacy tend to express more rare variants than the general population and, therefore, are less likely to respond to individual medications. Using a deidentified electronic medical record and corresponding samples from a biorepository, investigators identified 326 frequently medicated individuals (defined as prescribed clopidogrel or warfarin in addition to >5 medications from select classes with at least 1 drug known to have a pharmacogenetic interaction). Analysis found that most of the marker allele frequencies in polypharmacy patients did not differ from what was reported for individuals of European descent in the 1000 Genomes Pilot with a single exception in *CYP2D6* (rs1080985).¹⁰³ The relevance of this finding was deemed unclear because it was not certain whether it was the result of sequencing errors in the 1000 Genomes Pilot or a true difference.¹⁰⁴

Precision medicine will advance this field by using enhanced phenotyping for disease stratification to identify groups of patients that differ in drug responsiveness, will benefit from existing pharmacotherapies, or will benefit from rational polypharmacy (ie, the use of multiple drugs chosen based on knowledge of their interactome-based network location and effects on key pathways in the interactome that govern pathophenotype [Figure 6B]).¹⁰⁵ This strategy can also be used to identify biomarkers to guide drug dosing, continue therapy, or limit side effects and adverse drug reactions.¹⁰¹ When and how pharmacogenetic profiling will become a mainstay of clinical practice to guide polypharmacy and limit adverse drug reactions remains to be determined.¹⁰⁶ In part, the slow uptake of clinically applied pharmacogenetics is a reflection of statistical uncertainty, the abundance of genomic variants (even those previously thought to be causative), and the typical association analysis rather than network analysis of the functional consequences of the variants.¹⁰⁷

Precision Versus Personalized Medicine

Although the terms precision medicine and personalized medicine have been used interchangeably, they are not identical but, rather, related and overlapping disciplines. Precision medicine identifies the unique aspects of an individual related to health and disease to select appropriate and effective therapy. Personalized medicine may refer to implementation of these data into a person-specific treatment plan or, simply, the creation of a treatment plan for an individual based on a known biomarker(s). Personalized medicine is not, however, meant to suggest that results from detailed phenotyping will be used to create new drug or other therapies.

Precision medicine is increasingly considered as the art and science of creating individualized assessments of health and disease that are derived from established clinical and pathological indices integrated with state-of-the-art panomic (ie, genomic, transcriptomic, metabolomic, and proteomic) profiling.^{108,109} Thus, precision medicine aims to drive the field away from the implementation of the same pharmacotherapy, lifestyle intervention, or behavioral modification to a group of individuals presumed to share the same phenotype and toward




Figure 7. Challenges in precision medicine. Precision medicine relies on integration and dynamic adaptability of panomic analysis, overall data analysis, and data management; and acceptance by the public, medical, research, big pharma, and policy-maker stakeholders.

personalizing treatments that are aimed at prevention, promotion of health, and amelioration of disease. Although this concept has been embraced with success in the field of oncology, the issue of patient adherence with a personalized therapy identified using precision medicine remains a concern.¹⁰⁹ Estimates indicate that approximately half of all medications are not taken as prescribed and that patients with chronic conditions, such as cardiovascular disease, only take 50% to 60% of the medications prescribed.¹¹⁰ Although this is a valid point and developments in precision and personalized medicine must be matched by patient education and acceptance of these strategies, it is also plausible that poor adherence rates are the result of imprecise medication use leading to limited efficacy and increased side effect risk in a given individual. This issue will clearly require further study.

Future Considerations

Precision medicine is poised to become the next great revolution in the practice of medicine, as well as in the maintenance of cardiovascular health and the prevention and cure of cardiovascular disease. Precision medicine disrupts standard practice and draws from clinical testing, electronic health records, panomics profiling, big data sets, and novel analytical methods, such as systems biology and network science, to create a person-specific phenotype that can then be used to identify an optimal intervention with minimal risk. The obvious benefits of this approach to patients, clinicians, and researchers are numerous and include individual phenotype specificity, identification of individuals with a similar molecular phenotype, selection of best drugs or therapies with maximal efficacy and no or limited adverse reactions, efficient selection and enrichment of clinical trial participants, potential to improve adherence and reduce costs, and creating a paradigm shift in how cardiovascular care is delivered. To accomplish this laudable goal, the medical community at large and other stakeholders will need to overcome barriers to implementation that range from technical to sociopolitical (Figure 7).¹¹¹ At the least, precision medicine will need to demonstrate that

phenotype-based person-specific interventions are superior to the current standard of care and, ultimately, have a population effect by moving the mean on the disease spectrum toward health.¹¹¹ Other barriers are related to large data set collection and focuses on methods to ensure data accuracy, computational power, security and privacy of data sets, renewal of accruing data, and continuous development and refinement of analytical methods. Finally, education, affordability, and public acceptance of the strategy all play key roles in its ultimate implementation.¹¹¹

Despite a clear path forward toward mainstream application of precision medicine, there continues to be debate about whether a precision medicine approach will have a global impact on cardiovascular disease prevention and treatment or will only serve a small group of patients and be relegated to a highly selected niche role. This skepticism has arisen, in part, based on progress in the field to date and perceived challenges in maintaining the physician–patient relationship.^{112,113} Although these concerns are valid points, they are not insurmountable and are currently being addressed conceptually as how best to implement precision medicine in the practice of cardiovascular medicine. The nascent field of precision medicine fully embraces Paul Ehrlich’s concept that imprecision leads to failure and harnesses our most precise technologies and methodologies to improve cardiovascular health and treat disease.

Sources of Funding

This work was supported, in part, by the American Heart Association and National Institutes of Health grants U01 125215 (J.A. Leopold) and HL61795, HG007690, and GM107618 (J. Loscalzo).

Disclosures

None.

References

1. de Kruif P. *Microbe Hunters*. New York, NY: Houghton Mifflin Harcourt; 1954.

2. Heidenreich PA, Trogdon JG, Khavjou OA, et al; American Heart Association Advocacy Coordinating Committee; Stroke Council; Council on Cardiovascular Radiology and Intervention; Council on Clinical Cardiology; Council on Epidemiology and Prevention; Council on Arteriosclerosis; Thrombosis and Vascular Biology; Council on Cardiopulmonary; Critical Care; Perioperative and Resuscitation; Council on Cardiovascular Nursing; Council on the Kidney in Cardiovascular Disease; Council on Cardiovascular Surgery and Anesthesia, and Interdisciplinary Council on Quality of Care and Outcomes Research. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation*. 2011;123:933–944. doi: 10.1161/CIR.0b013e31820a55f5.
3. Roth GA, Forouzanfar MH, Moran AE, Barber R, Nguyen G, Feigin VL, Naghavi M, Mensah GA, Murray CJ. Demographic and epidemiologic drivers of global cardiovascular mortality. *N Engl J Med*. 2015;372:1333–1341. doi: 10.1056/NEJMoa1406656.
4. Benjamin EJ, Blaha MJ, Chiuve SE, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146–e603. doi: 10.1161/CIR.0000000000000485.
5. Fang N, Jiang M, Fan Y. Ideal cardiovascular health metrics and risk of cardiovascular disease or mortality: a meta-analysis. *Int J Cardiol*. 2016;214:279–283. doi: 10.1016/j.ijcard.2016.03.210.
6. Snyderman R, Meade C, Drake C. Value of personalized medicine. *JAMA*. 2016;315:613. doi: 10.1001/jama.2015.17136.
7. Snyderman R, Dreke CD. Personalized health care: unlocking the potential of genomic and precision medicine. *J Precision Med*. 2015;1:38–41.
8. Naylor S. What's in a name? The evolution of p-medicine. *J Precision Med*. 2015;1:15–29.
9. Antman EM, Loscalzo J. Precision medicine in cardiology. *Nat Rev Cardiol*. 2016;13:591–602. doi: 10.1038/nrcardio.2016.101.
10. Bhatnagar A. Environmental determinants of cardiovascular disease. *Circ Res*. 2017;121:162–180. doi: 10.1161/CIRCRESAHA.117.306458.
11. Ahmad T, Pencina MJ, Schulte PJ, O'Brien E, Whellan DJ, Piña IL, Kitzman DW, Lee KL, O'Connor CM, Felker GM. Clinical implications of chronic heart failure phenotypes defined by cluster analysis. *J Am Coll Cardiol*. 2014;64:1765–1774. doi: 10.1016/j.jacc.2014.07.979.
12. Rose G. Sick individuals and sick populations. *Int J Epidemiol*. 1985;14:32–38.
13. Zulman DM, Vijan S, Omenn GS, Hayward RA. The relative merits of population-based and targeted prevention strategies. *Milbank Q*. 2008;86:557–580. doi: 10.1111/j.1468-0009.2008.00534.x.
14. Pagidipati NJ, Navar AM, Mulder H, Sniderman AD, Peterson ED, Pencina MJ. Comparison of recommended eligibility for primary prevention statin therapy based on the us preventive services task force recommendations vs the ACC/AHA guidelines. *JAMA*. 2017;317:1563–1567.
15. Babu RB, Alam M, Helis E, Fodor JG. Population-based versus high-risk strategies for the prevention of cardiovascular diseases in low- and middle-income countries. *Indian Heart J*. 2012;64:439–443. doi: 10.1016/j.ihj.2012.08.001.
16. Lehrman MA, Goldstein JL, Brown MS, Russell DW, Schneider WJ. Internalization-defective LDL receptors produced by genes with nonsense and frameshift mutations that truncate the cytoplasmic domain. *Cell*. 1985;41:735–743.
17. Lehrman MA, Schneider WJ, Südhof TC, Brown MS, Goldstein JL, Russell DW. Mutation in LDL receptor: Alu-Alu recombination deletes exons encoding transmembrane and cytoplasmic domains. *Science*. 1985;227:140–146.
18. Jarcho JA, McKenna W, Pare JA, Solomon SD, Holcombe RF, Dickie S, Levi T, Donis-Keller H, Seidman JG, Seidman CE. Mapping a gene for familial hypertrophic cardiomyopathy to chromosome 14q1. *N Engl J Med*. 1989;321:1372–1378. doi: 10.1056/NEJM198911163212005.
19. Geisterfer-Lowrance AA, Kass S, Tanigawa G, Vosberg HP, McKenna W, Seidman CE, Seidman JG. A molecular basis for familial hypertrophic cardiomyopathy: a beta cardiac myosin heavy chain gene missense mutation. *Cell*. 1990;62:999–1006.
20. Dietz HC, Cutting GR, Pyeritz RE, Maslen CL, Sakai LY, Corson GM, Puffenberger EG, Hamosh A, Nanthakumar EJ, Currstin SM. Marfan syndrome caused by a recurrent de novo missense mutation in the fibrillin gene. *Nature*. 1991;352:337–339. doi: 10.1038/352337a0.
21. Kathiresan S, Srivastava D. Genetics of human cardiovascular disease. *Cell*. 2012;148:1242–1257. doi: 10.1016/j.cell.2012.03.001.
22. Wang Q, Shen J, Splawski I, Atkinson D, Li Z, Robinson JL, Moss AJ, Towbin JA, Keating MT. SCN5A mutations associated with an inherited cardiac arrhythmia, long QT syndrome. *Cell*. 1995;80:805–811.
23. Chen Q, Kirsch GE, Zhang D, et al. Genetic basis and molecular mechanism for idiopathic ventricular fibrillation. *Nature*. 1998;392:293–296. doi: 10.1038/32675.
24. McNair WP, Ku L, Taylor MR, Fain PR, Dao D, Wolfel E, Mestroni L; Familial Cardiomyopathy Registry Research Group. SCN5A mutation associated with dilated cardiomyopathy, conduction disorder, and arrhythmia. *Circulation*. 2004;110:2163–2167. doi: 10.1161/01.CIR.0000144458.58660.BB.
25. Hobbs HH, Leitersdorf E, Leffert CC, Cryer DR, Brown MS, Goldstein JL. Evidence for a dominant gene that suppresses hypercholesterolemia in a family with defective low density lipoprotein receptors. *J Clin Invest*. 1989;84:656–664. doi: 10.1172/JCI114212.
26. Faivre L, Collod-Beroud G, Loeys BL, et al. Effect of mutation type and location on clinical outcome in 1,013 probands with Marfan syndrome or related phenotypes and FBN1 mutations: an international study. *Am J Hum Genet*. 2007;81:454–466. doi: 10.1086/520125.
27. Nikpay M, Goel A, Won HH, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet*. 2015;47:1121–1130. doi: 10.1038/ng.3396.
28. Helgadottir A, Thorleifsson G, Manolescu A, et al. A common variant on chromosome 9p21 affects the risk of myocardial infarction. *Science*. 2007;316:1491–1493. doi: 10.1126/science.1142842.
29. McPherson R, Pertsemlidis A, Kavaslar N, Stewart A, Roberts R, Cox DR, Hinds DA, Pennacchio LA, Tybjaerg-Hansen A, Folsom AR, Boerwinkle E, Hobbs HH, Cohen JC. A common allele on chromosome 9 associated with coronary heart disease. *Science*. 2007;316:1488–1491. doi: 10.1126/science.1142447.
30. Samani NJ, Erdmann J, Hall AS, et al; WTCCC and the Cardiogenics Consortium. Genomewide association analysis of coronary artery disease. *N Engl J Med*. 2007;357:443–453. doi: 10.1056/NEJMoa072366.
31. McKusick VA. Mendelian Inheritance in Man and its online version, OMIM. *Am J Hum Genet*. 2007;80:588–604. doi: 10.1086/514346.
32. Stenson PD, Mort M, Ball EV, Shaw K, Phillips A, Cooper DN. The Human Gene Mutation Database: building a comprehensive mutation repository for clinical and molecular genetics, diagnostic testing and personalized genomic medicine. *Hum Genet*. 2014;133:1–9. doi: 10.1007/s00439-013-1358-4.
33. Dietz HC. New therapeutic approaches to mendelian disorders. *N Engl J Med*. 2010;363:852–863. doi: 10.1056/NEJMra0907180.
34. Soule T. Resilient individuals improve evolutionary search. *Artif Life*. 2006;12:17–34. doi: 10.1162/106454606775186437.
35. Hartman JL 4th. Buffering of deoxyribonucleotide pool homeostasis by threonine metabolism. *Proc Natl Acad Sci USA*. 2007;104:11700–11705.
36. Friend SH, Schadt EE. Translational genomics. Clues from the resilient. *Science*. 2014;344:970–972. doi: 10.1126/science.1255648.
37. Cooper DN, Krawczak M, Polychronakos C, Tyler-Smith C, Kehrer-Sawatzki H. Where genotype is not predictive of phenotype: towards an understanding of the molecular basis of reduced penetrance in human inherited disease. *Hum Genet*. 2013;132:1077–1130. doi: 10.1007/s00439-013-1331-2.
38. Sosnay PR, Siklosi KR, Van Goor F, et al. Defining the disease liability of variants in the cystic fibrosis transmembrane conductance regulator gene. *Nat Genet*. 2013;45:1160–1167. doi: 10.1038/ng.2745.
39. Khera AV, Emdin CA, Drake I, et al. Genetic risk, adherence to a healthy lifestyle, and coronary disease. *N Engl J Med*. 2016;375:2349–2358. doi: 10.1056/NEJMoa1605086.
40. Louie RJ, Guo J, Rodgers JW, White R, Shah N, Pagant S, Kim P, Livstone M, Dolinski K, McKinney BA, Hong J, Sorscher EJ, Bryan J, Miller EA, Hartman JL 4th. A yeast phenomic model for the gene interaction network modulating CFTR-ΔF508 protein biogenesis. *Genome Med*. 2012;4:103. doi: 10.1186/gm404.
41. Cohen J, Pertsemlidis A, Kotowski IK, Graham R, Garcia CK, Hobbs HH. Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. *Nat Genet*. 2005;37:161–165. doi: 10.1038/ng1509.
42. Flannick J, Thorleifsson G, Beer NL, et al; Go-T2D Consortium; T2D-GENES Consortium. Loss-of-function mutations in SLC30A8 protect against type 2 diabetes. *Nat Genet*. 2014;46:357–363. doi: 10.1038/ng.2915.
43. Chen R, Shi L, Hakenberg J, et al. Analysis of 589,306 genomes identifies individuals resilient to severe Mendelian childhood diseases. *Nat Biotechnol*. 2016;34:531–538. doi: 10.1038/nbt.3514.
44. Strebhardt K, Ullrich A. Paul Ehrlich's magic bullet concept: 100 years of progress. *Nat Rev Cancer*. 2008;8:473–480. doi: 10.1038/nrc2394.

45. Bosch F, Rosich L. The contributions of Paul Ehrlich to pharmacology: a tribute on the occasion of the centenary of his Nobel Prize. *Pharmacology*. 2008;82:171–179. doi: 10.1159/000149583.
46. Barabási AL, Gulbahce N, Loscalzo J. Network medicine: a network-based approach to human disease. *Nat Rev Genet*. 2011;12:56–68. doi: 10.1038/nrg2918.
47. Vidal M, Cusick ME, Barabási AL. Interactome networks and human disease. *Cell*. 2011;144:986–998. doi: 10.1016/j.cell.2011.02.016.
48. Menche J, Sharma A, Kitsak M, Ghiassian SD, Vidal M, Loscalzo J, Barabási AL. Disease networks. Uncovering disease-disease relationships through the incomplete interactome. *Science*. 2015;347:1257601. doi: 10.1126/science.1257601.
49. Chan SY, Loscalzo J. The emerging paradigm of network medicine in the study of human disease. *Circ Res*. 2012;111:359–374. doi: 10.1161/CIRCRESAHA.111.258541.
50. Wang RS, Loscalzo J. Illuminating drug action by network integration of disease genes: a case study of myocardial infarction. *Mol Biosyst*. 2016;12:1653–1666. doi: 10.1039/c6mb00052e.
51. Kitsak M, Sharma A, Menche J, Guney E, Ghiassian SD, Loscalzo J, Barabási AL. Tissue specificity of human disease module. *Sci Rep*. 2016;6:35241. doi: 10.1038/srep35241.
52. Delude CM. Deep phenotyping: the details of disease. *Nature*. 2015;527:S14–S15. doi: 10.1038/527S14a.
53. Robinson PN, Mungall CJ, Haendel M. Capturing phenotypes for precision medicine. *Cold Spring Harb Mol Case Stud*. 2015;1:a000372. doi: 10.1101/mcs.a000372.
54. Wei WQ, Denny JC. Extracting research-quality phenotypes from electronic health records to support precision medicine. *Genome Med*. 2015;7:41. doi: 10.1186/s13073-015-0166-y.
55. Köhler S, Vasilevsky NA, Engelstad M, et al. The human phenotype ontology in 2017. *Nucleic Acids Res*. 2017;45:D865–D876. doi: 10.1093/nar/gkw1039.
56. MacRae CA, Vasan RS. The future of genetics and genomics: closing the phenotype gap in precision medicine. *Circulation*. 2016;133:2634–2639. doi: 10.1161/CIRCULATIONAHA.116.022547.
57. McConnell MV, Shcherbina A, Pavlovic A, et al. Feasibility of obtaining measures of lifestyle from a smartphone app: the MyHeart Counts Cardiovascular Health Study. *JAMA Cardiol*. 2017;2:67–76. doi: 10.1001/jamacardio.2016.4395.
58. Shah SJ, Katz DH, Selvaraj S, Burke MA, Yancy CW, Gheorghiade M, Bonow RO, Huang CC, Deo RC. Phenomapping for novel classification of heart failure with preserved ejection fraction. *Circulation*. 2015;131:269–279. doi: 10.1161/CIRCULATIONAHA.114.010637.
59. Poldrack RA, Laumann TO, Koyejo O, et al. Long-term neural and physiological phenotyping of a single human. *Nat Commun*. 2015;6:8885. doi: 10.1038/ncomms9885.
60. Gottesman II, Gould TD. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry*. 2003;160:636–645. doi: 10.1176/appi.ajp.160.4.636.
61. Cannon TD, Keller MC. Endophenotypes in the genetic analyses of mental disorders. *Annu Rev Clin Psychol*. 2006;2:267–290. doi: 10.1146/annurev.clinpsy.2.022305.095232.
62. Willer CJ, Schmidt EM, Sengupta S, et al; Global Lipids Genetics Consortium. Discovery and refinement of loci associated with lipid levels. *Nat Genet*. 2013;45:1274–1283. doi: 10.1038/ng.2797.
63. Willer CJ, Sanna S, Jackson AU, et al. Newly identified loci that influence lipid concentrations and risk of coronary artery disease. *Nat Genet*. 2008;40:161–169. doi: 10.1038/ng.76.
64. Teslovich TM, Musunuru K, Smith AV, et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature*. 2010;466:707–713. doi: 10.1038/nature09270.
65. Nesse RM, Ganten D, Gregory TR, Omenn GS. Evolutionary molecular medicine. *J Mol Med (Berl)*. 2012;90:509–522. doi: 10.1007/s00109-012-0889-9.
66. Button KS, Ioannidis JP, Mokrysz C, Nosek BA, Flint J, Robinson ES, Munafò MR. Empirical evidence for low reproducibility indicates low pre-study odds. *Nat Rev Neurosci*. 2013;14:877. doi: 10.1038/nrn3475-c6.
67. Kulminkski AM, Kermogitski Y, Culminkskaya I, Loika Y, Arbeev KG, Bagley O, Duan M, Arbeeva L, Ukraintseva SV, Wu D, Stallard E, Yashin AI. Uncoupling associations of risk alleles with endophenotypes and phenotypes: insights from the ApoB locus and heart-related traits. *Aging Cell*. 2017;16:61–72. doi: 10.1111/acel.12526.
68. Loscalzo J, Kohane I, Barabási AL. Human disease classification in the postgenomic era: a complex systems approach to human pathobiology. *Mol Syst Biol*. 2007;3:124. doi: 10.1038/msb4100163.
69. Fox EA, Kahn SR. The relationship between inflammation and venous thrombosis. A systematic review of clinical studies. *Thromb Haemost*. 2005;94:362–365. doi: 10.1160/TH05-04-0266.
70. Wakefield TW, Strieter RM, Prince MR, Downing LJ, Greenfield LJ. Pathogenesis of venous thrombosis: a new insight. *Cardiovasc Surg*. 1997;5:6–15.
71. Libby P, Simon DI. Inflammation and thrombosis: the clot thickens. *Circulation*. 2001;103:1718–1720.
72. Ghiassian SD, Menche J, Chasman DI, et al. Endophenotype network models: common core of complex diseases. *Sci Rep*. 2016;6:27414. doi: 10.1038/srep27414.
73. Wilkinson GR. Drug metabolism and variability among patients in drug response. *N Engl J Med*. 2005;352:2211–2221. doi: 10.1056/NEJMr032424.
74. Cardelli M, Marchegiani F, Corsonello A, Lattanzio F, Provinciali M. A review of pharmacogenetics of adverse drug reactions in elderly people. *Drug Saf*. 2012;35(suppl 1):3–20. doi: 10.1007/BF03319099.
75. Qato DM, Alexander GC, Conti RM, Johnson M, Schumm P, Lindau ST. Use of prescription and over-the-counter medications and dietary supplements among older adults in the United States. *JAMA*. 2008;300:2867–2878. doi: 10.1001/jama.2008.892.
76. Hajjar ER, Cafiero AC, Hanlon JT. Polypharmacy in elderly patients. *Am J Geriatr Pharmacother*. 2007;5:345–351. doi: 10.1016/j.amjopharm.2007.12.002.
77. Safran DG, Neuman P, Schoen C, Kitchman MS, Wilson IB, Cooper B, Li A, Chang H, Rogers WH. Prescription drug coverage and seniors: findings from a 2003 national survey. *Health Aff (Millwood)*. 2005;Suppl Web Exclusives:W5-152–W155-166.
78. Budnitz DS, Lovegrove MC, Shehab N, Richards CL. Emergency hospitalizations for adverse drug events in older Americans. *N Engl J Med*. 2011;365:2002–2012. doi: 10.1056/NEJMs1103053.
79. Lazarou J, Pomeranz BH, Corey PN. Incidence of adverse drug reactions in hospitalized patients: a meta-analysis of prospective studies. *JAMA*. 1998;279:1200–1205.
80. Oqueli E, Hiscok M, Dick R. Clopidogrel resistance. *Heart Lung Circ*. 2007;16(suppl 3):S17–S28. doi: 10.1016/j.hlc.2007.03.012.
81. Ahmad T, Voora D, Becker RC. The pharmacogenetics of antiplatelet agents: towards personalized therapy? *Nat Rev Cardiol*. 2011;8:560–571. doi: 10.1038/nrcardio.2011.111.
82. Collet JP, Hulot JS, Pena A, Villard E, Esteve JB, Silvain J, Payot L, Brugier D, Cayla G, Beygui F, Bensimon G, Funck-Brentano C, Montalescot G. Cytochrome P450 2C19 polymorphism in young patients treated with clopidogrel after myocardial infarction: a cohort study. *Lancet*. 2009;373:309–317. doi: 10.1016/S0140-6736(08)61845-0.
83. Mega JL, Close SL, Wiviott SD, Shen L, Hockett RD, Brandt JT, Walker JR, Antman EM, Macias W, Braunwald E, Sabatine MS. Cytochrome p-450 polymorphisms and response to clopidogrel. *N Engl J Med*. 2009;360:354–362. doi: 10.1056/NEJMoa0809171.
84. Mega JL, Hochholzer W, Frelinger AL 3rd, Kluk MJ, Angiolillo DJ, Kereiakes DJ, Isserman S, Rogers WJ, Ruff CT, Contant C, Pencina MJ, Scirica BM, Longtine JA, Michelson AD, Sabatine MS. Dosing clopidogrel based on CYP2C19 genotype and the effect on platelet reactivity in patients with stable cardiovascular disease. *JAMA*. 2011;306:2221–2228. doi: 10.1001/jama.2011.1703.
85. Geisler T, Schaeffeler E, Dippon J, Winter S, Buse V, Bischofs C, Zuern C, Moerike K, Gawaz M, Schwab M. CYP2C19 and nongenetic factors predict poor responsiveness to clopidogrel loading dose after coronary stent implantation. *Pharmacogenomics*. 2008;9:1251–1259. doi: 10.2217/14622416.9.9.1251.
86. Sibbing D, Koch W, Gebhard D, Schuster T, Braun S, Stegherr J, Morath T, Schömig A, von Beckerath N, Kastrati A. Cytochrome 2C19*17 allelic variant, platelet aggregation, bleeding events, and stent thrombosis in clopidogrel-treated patients with coronary stent placement. *Circulation*. 2010;121:512–518. doi: 10.1161/CIRCULATIONAHA.109.885194.
87. Price MJ, Berger PB, Teirstein PS, et al; GRAVITAS Investigators. Standard- vs high-dose clopidogrel based on platelet function testing after percutaneous coronary intervention: the GRAVITAS randomized trial. *JAMA*. 2011;305:1097–1105. doi: 10.1001/jama.2011.290.
88. Wei MY, Ito MK, Cohen JD, Brinton EA, Jacobson TA. Predictors of statin adherence, switching, and discontinuation in the USAGE survey: understanding the use of statins in America and gaps in patient education. *J Clin Lipidol*. 2013;7:472–483. doi: 10.1016/j.jacl.2013.03.001.
89. Link E, Parish S, Armitage J, Bowman L, Heath S, Matsuda F, Gut I, Lathrop M, Collins R; SEARCH Collaborative Group. SLCO1B1 variants and statin-induced myopathy—a genome-wide study. *N Engl J Med*. 2008;359:789–799. doi: 10.1056/NEJMoa0801936.

90. Klein TE, Altman RB, Eriksson N, Gage BF, Kimmel SE, Lee MT, Limdi NA, Page D, Roden DM, Wagner MJ, Caldwell MD, Johnson JA; International Warfarin Pharmacogenetics Consortium. Estimation of the warfarin dose with clinical and pharmacogenetic data. *N Engl J Med*. 2009;360:753–764. doi: 10.1056/NEJMoa0809329.
91. Zabalza M, Subirana I, Sala J, Lluís-Ganella C, Lucas G, Tomás M, Masiá R, Marrugat J, Brugada R, Elosua R. Meta-analyses of the association between cytochrome CYP2C19 loss- and gain-of-function polymorphisms and cardiovascular outcomes in patients with coronary artery disease treated with clopidogrel. *Heart*. 2012;98:100–108. doi: 10.1136/hrt.2011.227652.
92. Johnson JA, Cavallari LH. Warfarin pharmacogenetics. *Trends Cardiovasc Med*. 2015;25:33–41. doi: 10.1016/j.tcm.2014.09.001.
93. Jorgensen AL, FitzGerald RJ, Oyee J, Pirmohamed M, Williamson PR. Influence of CYP2C9 and VKORC1 on patient response to warfarin: a systematic review and meta-analysis. *PLoS One*. 2012;7:e44064. doi: 10.1371/journal.pone.0044064.
94. Cooper GM, Johnson JA, Langae TY, Feng H, Stanaway IB, Schwarz UI, Ritchie MD, Stein CM, Roden DM, Smith JD, Veenstra DL, Rettie AE, Rieder MJ. A genome-wide scan for common genetic variants with a large influence on warfarin maintenance dose. *Blood*. 2008;112:1022–1027. doi: 10.1182/blood-2008-01-134247.
95. Takeuchi F, McGinnis R, Bourgeois S, Barnes C, Eriksson N, Soranzo N, Whittaker P, Ranganath V, Kumanduri V, McLaren W, Holm L, Lindh J, Rane A, Wadelius M, Deloukas P. A genome-wide association study confirms VKORC1, CYP2C9, and CYP4F2 as principal genetic determinants of warfarin dose. *PLoS Genet*. 2009;5:e1000433. doi: 10.1371/journal.pgen.1000433.
96. Finkelman BS, Gage BF, Johnson JA, Brensinger CM, Kimmel SE. Genetic warfarin dosing: tables versus algorithms. *J Am Coll Cardiol*. 2011;57:612–618. doi: 10.1016/j.jacc.2010.08.643.
97. Gage BF, Eby C, Johnson JA, et al. Use of pharmacogenetic and clinical factors to predict the therapeutic dose of warfarin. *Clin Pharmacol Ther*. 2008;84:326–331. doi: 10.1038/clpt.2008.10.
98. Kimmel SE, French B, Kasner SE, et al; COAG Investigators. A pharmacogenetic versus a clinical algorithm for warfarin dosing. *N Engl J Med*. 2013;369:2283–2293. doi: 10.1056/NEJMoa1310669.
99. Pirmohamed M, Burnside G, Eriksson N, et al; EU-PACT Group. A randomized trial of genotype-guided dosing of warfarin. *N Engl J Med*. 2013;369:2294–2303. doi: 10.1056/NEJMoa1311386.
100. Ross S, Nejat S, Paré G. Use of genetic data to guide therapy in arterial disease. *J Thromb Haemost*. 2015;13(suppl 1):S281–S289. doi: 10.1111/jth.12924.
101. Zaiou M, El Amri H. Cardiovascular pharmacogenetics: a promise for genomically-guided therapy and personalized medicine. *Clin Genet*. 2017;91:355–370. doi: 10.1111/cge.12881.
102. Elliott LS, Henderson JC, Neradilek MB, Moyer NA, Ashcraft KC, Thirumaran RK. Clinical impact of pharmacogenetic profiling with a clinical decision support tool in polypharmacy home health patients: A prospective pilot randomized controlled trial. *PLoS One*. 2017;12:e0170905. doi: 10.1371/journal.pone.0170905.
103. Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, Gibbs RA, Hurles ME, McVean GA; 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature*. 2010;467:1061–1073. doi: 10.1038/nature09534.
104. Oetjens MT, Denny JC, Ritchie MD, Gillani NB, Richardson DM, Restrepo NA, Pulley JM, Dilks HH, Basford MA, Bowton E, Masys DR, Wilke RA, Roden DM, Crawford DC. Assessment of a pharmacogenomic marker panel in a polypharmacy population identified from electronic medical records. *Pharmacogenomics*. 2013;14:735–744. doi: 10.2217/pgs.13.64.
105. Garmaroudi FS, Handy DE, Liu YY, Loscalzo J. Systems pharmacology and rational polypharmacy: nitric oxide-cyclic GMP signaling pathway as an illustrative example and derivation of the general case. *PLoS Comput Biol*. 2016;12:e1004822. doi: 10.1371/journal.pcbi.1004822.
106. Stingl Kirchheiner JC, Brockmüller J. Why, when, and how should pharmacogenetics be applied in clinical studies?: current and future approaches to study designs. *Clin Pharmacol Ther*. 2011;89:198–209. doi: 10.1038/clpt.2010.274.
107. Manrai AK, Funke BH, Rehm HL, Olesen MS, Maron BA, Szolovits P, Margulies DM, Loscalzo J, Kohane IS. Genetic misdiagnoses and the potential for health disparities. *N Engl J Med*. 2016;375:655–665. doi: 10.1056/NEJMs1507092.
108. Mirnezami R, Nicholson J, Darzi A. Preparing for precision medicine. *N Engl J Med*. 2012;366:489–491. doi: 10.1056/NEJMp1114866.
109. Fuster V. A first dilemma in cardiovascular medicine: adherence versus personalized therapy. *J Am Coll Cardiol*. 2014;64:1059–1060. doi: 10.1016/j.jacc.2014.07.936.
110. Bosworth HB, Granger BB, Mendys P, et al. Medication adherence: a call for action. *Am Heart J*. 2011;162:412–424. doi: 10.1016/j.ahj.2011.06.007.
111. Kohane IS. Health care policy. Ten things we have to do to achieve precision medicine. *Science*. 2015;349:37–38. doi: 10.1126/science.aab1328.
112. Joyner MJ. Precision medicine, cardiovascular disease and hunting elephants. *Prog Cardiovasc Dis*. 2016;58:651–660. doi: 10.1016/j.pcad.2016.02.004.
113. Fuster V. A second dilemma in cardiovascular medicine: personalized medicine versus personal interaction with the patient. *J Am Coll Cardiol*. 2014;64:1292–1293. doi: 10.1016/j.jacc.2014.08.006.