

Arsen - Microbiome Analysis

Jan Waelchli

2022-08-25

Contents

Experimental Setup	2
Description all data	2
Sequencing Depth	2
Normalization	4
Sample Control	5
Taxonomy	6
Phyla abundance plot	6
Effect of all factors on phyla abundances	8
Alpha diversity	9
Method	9
Genotype*Arsen Effect	9
Beta diversity	11
Method	11
Genotype*Arsen Effect	11
Taxa Response	14
Method	14
Genotype*Arsen Effect	14

Experimental Setup


WT maize plants and bx1 mutants with a W22 background were grown. Half of them were treated with 0 ppm arsen (CTRL) and the other half with 100 ppm arsen. We analyse the shift of the bacterial and fungal communities in the plant rhizosphere.

Description all data

Sequencing Depth

Figure 1 | Reads tracking

We plot the amount of reads during each pipeline step. This allows us to see where we loses reads and if the samples from the different groups behave similar.




Conclusion: We lose the expected amount of reads. Samples from different group behave very similar.

Number of sequences

We show the sum, range and median of sequecnese over all samples.

Taxa	removed_samples	sum	min	max	median
Bacteria	0	2484130	37055	77593	49508
Fungi	1	146569	2172	5112	3151

Figure 2 | Sequencing depth



Normalization

Asymptotic Kruskal-Wallis Test & Normalization

To decide on how to normalize the data we follow the recommendation of Weiss et al. (2017, Microbiome Journal) and inspect whether there are differences in sequencing depths between the different arsen-treatments and genotypes by using the non-parametric Kruskal-Wallis Test.

```
## [1] "Bacteria"

##
## Kruskal-Wallis rank sum test
##
## data:  sample_depth by group
## Kruskal-Wallis chi-squared = 4.9192, df = 3, p-value = 0.1778

## [1] "Fungi"

##
## Kruskal-Wallis rank sum test
##
## data:  sample_depth by group
## Kruskal-Wallis chi-squared = 3.798, df = 3, p-value = 0.2841
```

Conclusion: We don't find significant differences between the groups in bacteria or fungi. We follow the recommendation of Weiss et al. (2017) to use TSS normalization for samples with small sequencing-depth differences.

Outlier Detection

We use the method CLOUD developed by Montassier et al. 2018, which is a non-parametric detection test for outliers. We perform the test with Bray-Curtis distances from the normalized data for each substrate and each plastic treatment individually. We set the number of nearest neighbors to 60% of the samples size and chose an empirical outlier percentile of 0.1. We remove all outliers from our data.

Table 2: Number of outliers

Species	Arsen.CTRL	Arsen.arsen
Bacteria	2	2
Fungi	2	2

Sample Control

Sample Size

We end up with the following number of samples per treatment for the analysis.

Table 3: Bacteria: Sample profile


	CTRL	arsen
WT	10	10
bx1	12	12

Table 4: Fungi: Sample profile

	CTRL	arsen
WT	10	10
bx1	12	11


Figure 3 | Rarefaction plot

We plot a rarefaction plot with the remaining samples to check if the sequence depth is enough to capture the microbial diversity.



Conclusion: All samples were sequenced deep enough.

Figure 4.2 | Fungi: Phylum level taxonomy



Effect of all factors on phyla abundances

We test if there are any difference between the phyla abundances between genotypes, arsen-treatments or their interaction by performing a PERMANOVA (permutations = 999).

Bacteria

Table 5: Bacteria: PERMANOVA

	Df	SumOfSqs	R2	F	Pr(>F)
arsen	1	0.003888	0.04373	1.904	0.097
genotype	1	0.002971	0.03341	1.455	0.188
arsen:genotype	1	0.000367	0.004128	0.1797	0.957
Residual	40	0.08169	0.9187	NA	NA
Total	43	0.08892	1	NA	NA

Fungi

Table 6: Fungi: PERMANOVA

	Df	SumOfSqs	R2	F	Pr(>F)
arsen	1	0.00398	0.01002	0.4147	0.729
genotype	1	0.008926	0.02247	0.9301	0.427
arsen:genotype	1	0.01005	0.0253	1.047	0.368
Residual	39	0.3743	0.9422	NA	NA
Total	42	0.3972	1	NA	NA

Conclusion: No differences found.

Alpha diversity

We answer the following questions for the alpha diversity in each substrate:

- **Q1: Has arsen changed the beta diversity?**
- **Q2: Is beta diversity different between the genotypes?**
- **Q3: Are there differences in beta diversity between the different genotypes after treating plants with arsen?**

Method

We calculate the Shannon diversity for each sample with the normalized data.

Genotype*Arsen Effect

We investigate the effect on alpha diversity by the factors of genotype, arsen and the interaction between them. We model the alpha diversity against these factors in an aov-model and perform a F-Test.


Table 7: Bacteria: F test

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
genotype	1	0.02808	0.02808	1.244	0.2713
arsen	1	0.001565	0.001565	0.06932	0.7937
genotype:arsen	1	0.03245	0.03245	1.438	0.2376
Residuals	40	0.9029	0.02257	NA	NA

Table 8: Fungi: F test

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
genotype	1	0.04784	0.04784	1.999	0.1653
arsen	1	0.004894	0.004894	0.2045	0.6536
genotype:arsen	1	0.01819	0.01819	0.7601	0.3886
Residuals	39	0.9331	0.02393	NA	NA

Figure 5 | Genotype*Arsen effect on alpha diversity



Conclusion: No effect has been found.

Beta diversity

We answered the following question for the bacterial and fungal beta diversity in each compartment:

- **Q1: Has arsen changed the beta diversity?**
- **Q2: Is beta diversity different between the genotypes?**
- **Q3: Are there differences in beta diversity between the different genotypes after treating plants with arsen?**

Method

First we use the function ‘adonis()’ (package vegan) to analyze the beta diversity with a PERMANOVA (permutations = 999). Then, we graphically represent the beta diversity with a PCoA (unconstrained ordination) and a CAP plot (constrained ordination).

Genotype*Arsen Effect

We investigate the full model to see which factors alters the beta diversity.

Table 9: Bacteria: PERMANOVA

	Df	SumOfSqs	R2	F	Pr(>F)
genotype	1	0.04415	0.03271	1.469	0.034
arsen	1	0.0492	0.03646	1.637	0.015
genotype:arsen	1	0.05384	0.0399	1.791	0.005
Residual	40	1.202	0.8909	NA	NA
Total	43	1.35	1	NA	NA

Table 10: Fungi: PERMANOVA

	Df	SumOfSqs	R2	F	Pr(>F)
genotype	1	0.1526	0.0328	1.448	0.049
arsen	1	0.2483	0.05335	2.356	0.002
genotype:arsen	1	0.1431	0.03076	1.358	0.079
Residual	39	4.109	0.8831	NA	NA
Total	42	4.653	1	NA	NA

Figure 6.1 | PCoA - genotype:arsen effect on beta diversity





Figure 6.2 | CAP - genotype:arsen effect on beta diversity



Conclusion: There are differences in the bacterial and fungal communities due to the arsenic treatment, the genotypes and their interactions. We can explain about 4% of bacterial and 3% of fungal variety due to the arsen:genotype interaction effect.

Taxa Response

Is there a core of sensitive microbial taxa? We searched sensitive ASVs – ASVs being differential abundant between WT and bx1. We answer the following question in non-arsenic and arsenic conditions:

Q1: Are there sensitive ASVs between control and WT and bx1 samples in non-arsenic and arsenic soil?

Method

We answered the question by using four different tools to measure differential abundances - aldex2, acomb, maaslin2 and metagenomeSeq - and predict ASVs to be different if they were detected by 2 or more tools.

Genotype*Arsen Effect

We check for each ASV if it is sensitive or not. Then, we show how many ASVs has been changed between the genotypes and how much of the relative abundance belongs to those sensitive ASVs.

Table 11: Bacteria: genotype effect (continued below)

taxa	arsen	lower in WT	unchanged	higher in WT
bacteria	CTRL	0	1236	0
bacteria	arsen	0	1284	0

rel. abu. of sens. ASVs
0%
0%

0% in non-arsenic and 0% in arsenic conditions of the bacterial community was changed in abundance due to genotype.

Table 13: Fungi: gneotype effect

taxa	arsen	lower in WT	unchanged	higher in WT	rel. abu. of sens. ASVs
fungi	CTRL	0	167	1	0.316%
fungi	arsen	0	176	0	0%

0.32% in non-arsenic and 0% in arsenic conditions of the fungal community was changed in abundance due to genotype.

Conclusion: Most ASVs are insensitive.