# **Problem Statement - Part II**

## **Question 1:**

What is the optimal value of alpha for ridge and lasso regression? What will be the changes in the model if you choose double the value of alpha for both ridge and lasso? What will be the most important predictor variables after the change is implemented?

## Answer1:

Optimal value of alpha for ridge: 1

Optimal value of alpha for lasso: 10

After make the double alpha for ridge and lasso i.e. 2 and 20

<u>For Ridge:</u> Coeff values are increasing as alpha will increase.r2\_score of train data is also drop from .807 to 0.45

<u>For Lasso:</u> As alpha value increased more feautrues removed from model. But r2score is also dropped by 1% in both test and train data

*Top Features:* Neighborhood\_NoRidge, Neighborhood\_NridgHt, OverallQual, overallQual Neighborhood\_Veenkar

### **Question 2:**

You have determined the optimal value of lambda for ridge and lasso regression during the assignment. Now, which one will you choose to apply and why?

### Answer 2:

The r2\_score of lasso is slightly higher than ridge for the test dataset so we will choose lasso regression to solve this problem

model without affecting the model accuracy. Which makes are model generalized and simple and accurate.

### **Question 3**

After building the model, you realised that the five most important predictor variables in the lasso model are not available in the incoming data. You will now have to create another model excluding the five most important predictor variables. Which are the five most important predictor variables now?

#### Answer 3:

Fmive most important predictor variables

- 11stFlrSF-----First Floor square feet
- GrLivArea-----Above grade (ground) living area square feet
- Street\_Pave-----Pave road access to property
- RoofMatl\_Metal-----Roof material\_Metal
- RoofStyle\_Shed-----Type of roof(Shed)

## **Question 4:**

How can you make sure that a model is robust and generalisable? What are the implications of the same for the accuracy of the model and why?

## Answer 4:

The model should be generalized so that the test accuracy is not lesser than the training score. The model should be accurate for datasets other than the ones which were used during training. Too much importance should not given to the outliers so that the accuracy predicted by the model is high. To ensure that this is not the case, the outliers analysis needs to be done and only those which are relevant to the dataset need to be retained. Those outliers which it does not make sense to keep must be removed from the dataset. If the model is not robust, It cannot be trusted for predictive analysis.