

My neat title here

Figures




Figure 1: World map showing the spatial distribution of the study sites.




Figure 2: Results of the linear regressions between $a_{\text{CDOM}}(350)$ and $a_{\text{CDOM}}(\lambda)$. (A) Determination coefficient (R^2), (B) slope and (C) intercept of the linear regressions. Panels contain the results of 251 linear models, each based on 2321 data points. Note that at $\lambda = 350$ nm, $R^2 = 1$, slope = 1 and intercept = 0.




Figure 3: Boxplots showing the distribution of **(A)** absorption coefficients at 350 nm ($a_{CDOM}(350)$), **(B)** dissolved organic carbon (DOC) and **(C)** the so-called a^* . Y-axis are log-transformed given the wide ranges spanned by the data.




Figure 4: (A) Global relationship between absorption at 350 nm $a_{\text{CDOM}}(350)$ and dissolved organic carbon. The blue line is the fitted values of a linear model $y = \log(x)$, $R^2 = 0.93$, $p < 0.00001$, $n = 11431$. (B) Barplot showing the determination coefficient (R^2) of the linear relationships between $a_{\text{CDOM}}(350)$ and DOC by ecosystems.




Figure 5: Segmentation analysis performed on the linear relationship between SUVA₂₅₄ and salinity ($R^2 = 0.74, p < 0.00001, n = xxx$). Dashed vertical lines represent the identified breakpoints at salinity 8.66 and 26.84.