


# Figures


Note that all figures are scaled by a factor of 2 for display.




**Figure 1:** Location of the ice camp located on the Qikiqtarjuaq Island in the Baffin Bay. Projection used: EPSG-4326.




**Figure 2:** Temporal evolution of the snow and sea-ice thickness for both ice camp missions. The dashed horizontal line represents the snow/ice interface.




**Figure 3:** Temporal evolution of the salinity for both ice camp missions. Note that for visualization, salinity below  $31.5 \text{ g kg}^{-1}$  have been squished at  $31.5 \text{ g kg}^{-1}$ .




**Figure 4:** Temporal evolution of daily photosynthetically available radiation (PAR) at the sea-ice/water interface (1.3 m depth) for both ice camp missions. The horizontal dashed line show the 0.415 mol photons m<sup>-2</sup> d<sup>-1</sup> threshold often used in the literature as the minimum light requirement for primary production.




**Figure 5:** Beam attenuation coefficients ( $c$ ,  $\text{m}^{-1}$ ) measured in 2016 using an ACS before and during the phytoplankton bloom. Note that the colors of the lines correspond to wavelength frequencies.



**Figure 6:** Temporal evolution of the nitrates in the first 60 m of the water column for both ice camp missions.




**Figure 7:** Temporal evolution of chlorophyll a in ice and water (depth-integrated) for both ice camp missions. Note that the water chlorophyll a have been integrated over the first 100 m of the water column whereas the ice chlorophyll a was measured on the bottom 0-10 cm of the ice cores.




**Figure 8:** Taxo. TODO: I asked P.-L. to produce a better composite image.





**Figure 9:** Temporal evolution of  $F_v/F_m$  for ice (last cm) and water underneath the ice (depths 1.5 m, 10 m, 40 m) samples for the ice camp 2016 between May 6<sup>th</sup> and July 8<sup>th</sup>.  $F_v/F_m$  monitoring on ice samples stopped on Day 172-June 20<sup>th</sup> because the Chl a fluorescence signal was not reliable anymore.  $F_v/F_m$  monitoring on 40 m and 60 m depth samples was limited between May 13<sup>th</sup> and June 24<sup>th</sup> and between June 29<sup>th</sup>-July 08<sup>th</sup>, respectively. The gray shaded area represents the range at which the algae are optimally growing.



**Figure 10:** Time series of the abundance of the zooplankton community (ind m<sup>-1</sup>) measured over the first 100 m and 350 m of the water column in 2016 using the zooscan.