

Green Edge ice camp campaigns: understanding the processes controlling the under-ice Arctic phytoplankton spring bloom

Philippe Massicotte¹, Rémi Amiraux^{1,2}, Marie-Pier Amyot¹, Philippe Archambault^{1,3}, Mathieu Ardyna^{4,5}, Laurent Arnaud⁶, Lise Artigue⁷, Cyril Aubry¹, Pierre Ayotte^{8,9,10}, Guislain Bécu¹, Simon Bélanger¹¹, Ronald Benner¹², Henry C. Bittig^{4,13}, Annick Bricaud⁴, Éric Brossier¹⁴, Flavienne Bruyant¹, Laurent Chauvaud², Debra Christiansen-Stowe¹⁵, Hervé Claustre⁴, Véronique Cornet-Barthaux¹⁶, Pierre Coupel¹, Christine Cox¹⁴, Aurelie Delaforge¹⁷, Thibaud Dezutter¹, Céline Dimier¹⁸, Florent Dominé¹, Francis Dufour¹, Christiane Dufresne^{19,20,3}, Dany Dumont^{19,20,3}, Jens Ehn¹⁷, Brent Else²¹, Joannie Ferland¹, Marie-Hélène Forget¹, Louis Fortier¹, Martí Galí¹, Virginie Galindo^{19,20,3}, Morgane Gallinari², Nicole Garcia¹⁶, Catherine Gérikas-Ribeiro^{22,23}, Margaux Gourdal¹, Priscilla Gourvil²⁴, Clemence Goyens²⁵, Pierre-Luc Grondin¹, Pascal Guillot²⁶, Caroline Guilmette¹, Marie-Noëlle Houssais²⁷, Fabien Joux²⁸, Léo Lacour¹, Thomas Lacour²⁹, Augustin Lafond¹⁶, José Lagunas¹, Catherine Lalande¹, Julien Laliberté¹, Simon Lambert-Girard¹, Jade Larivière¹, Johann Lavaud¹, Anita LeBaron¹, Karine Leblanc¹⁶, Florence Le Gall²², Justine Legras¹⁶, Mélanie Lemire^{8,30,10}, Maurice Levasseur^{1,3}, Edouard Leymarie⁴, Aude Leynaert², Adriana Lopes dos Santos³¹, Antonio Lourenço³², David Mah³¹, Claudie Marec^{1,33}, Dominique Marie³⁴, Nicolas Martin³², Constance Marty¹⁴, Sabine Marty³⁵, Guillaume Massé¹, Atsushi Matsuoka¹, Lisa Matthes¹⁷, Brivaela Moriceau², Pierre-Emmanuel Muller¹⁴, Christopher-John Mundy¹⁷, Griet Neukermans¹⁴, Laurent Oziel^{1,4}, Christos Panagiotopoulos¹⁶, Jean-Jacques Pangazi¹⁴, Ghislain Picard³⁶, Marc Picheral⁴, France Pinczon du Sel¹⁴, Nicole Pogorzelec¹⁷, Ian Probert²⁴, Bernard Queguiner¹⁶, Patrick Raimbault¹⁶, Joséphine Ras⁴, Eric Rehm¹, Erin Reimer¹, Jean-François Rontani¹⁶, Søren Rysgaard¹⁷, Blanche Saint-Béat¹, Makoto Sampei³⁷, Julie Sansoulet¹, Sabine Schmidt³⁸, Richard Sempéré¹⁶, Caroline Sévigny³⁹, Yuan Shen^{40,41}, Margot Tragin³⁴, Jean-Éric Tremblay¹, Daniel Vaulot^{34,31}, Gauthier Verin¹, Frédéric Vivier³², Anda Vladoiu^{42,43}, Jeremy Whitehead²¹, and Marcel Babin¹

¹UMI Takuvik, CNRS/Université Laval, Québec, QC Canada

²Univ Brest, CNRS, IRD, Ifremer, LEMAR, F-29280 Plouzane, France

³Québec-Océan

⁴Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, F-06230 Villefranche-sur-Mer, France

⁵Department of Earth System Science, Stanford University, Stanford, CA, 94305, USA

⁶UMR 5001, IGE, CNRS, Grenoble, France

⁷LEGOS, University of Toulouse, CNRS, CNES, IRD, UPS, 31400 Toulouse, France

⁸Axe Santé des populations et pratiques optimales en santé, Centre de recherche du CHU de Québec - Université Laval

⁹Centre de toxicologie du Québec, INSPQ

¹⁰Département de médecine sociale et préventive, Université Laval, Québec QC Canada

¹¹Département de Biologie, Chimie et Géographie (groupes BORÉAS et Québec-Océan), Université du Québec à Rimouski, 300 allé des Ursulines, Rimouski, QC, G5L 3A1

- ¹²University of South Carolina, Department of Biological sciences, Columbia, SC 29208 USA
¹³Leibniz Institute for Baltic Sea Research Warnemünde, IOW, Rostock-Warnemünde, Germany
¹⁴Independent collaborator
¹⁵Institut nordique du Québec, Université Laval, Québec, QC Canada
¹⁶Aix-Marseille Univ., Université de Toulon, CNRS, IRD, MIO, UM110, Marseille, 13288, France
¹⁷Centre for Earth Observation Science, University of Manitoba, Winnipeg, Manitoba, Canada
¹⁸FR3761, Institut de la Mer de Villefranche, CNRS, 06230 Villefranche-sur-mer, France
¹⁹Institut des sciences de la mer de Rimouski
²⁰Université du Québec à Rimouski
²¹Department of Geography, University of Calgary, Calgary, Alberta, Canada
²²CNRS, Sorbonne Université, UMR7144, Team ECOMAP, Station Biologique de Roscoff, Roscoff, France
²³GEMA Center for Genomics, Ecology & Environment, Universidad Mayor, Camino La Pirámide, 5750, Huechuraba, Santiago, Chile
²⁴Sorbonne Université, CNRS, FR2424, Centre de Ressources Biologiques Marines, Station Biologique de Roscoff, France
²⁵Royal Belgian Institute of Natural Sciences (RBINS), Operational Directorate Natural Environment, 29 Rue Vautierstraat, 1000 Brussels, Belgium
²⁶Québec Océan & Amundsen Science, Université Laval, Québec, QC Canada
²⁷LOCEAN, CNRS/Sorbonne Université/IRD/MNHN, 4 place Jussieu, F-75005 Paris, France
²⁸Sorbonne Université, CNRS, Laboratoire d'Océanographie Microbienne (LOMIC), Observatoire Océanologique de Banyuls, 66650 Banyuls/mer, France
²⁹IFREMER, Physiology and Biotechnology of Algae Laboratory, rue de l'Ile d'Yeu, 44311, Nantes, France
³⁰Institut de biologie intégrative et des systèmes
³¹Asian School of the Environment, Nanyang Technological University, 50 Nanyang Avenue, Singapore 639798
³²LOCEAN-IPSL, CNRS, Sorbonne Université, Paris, France
³³Univ Brest, CNRS, IUEM, UMS3113, F-29280 Plouzane, France
³⁴UMR 7144, Sorbonne Université and CNRS, Station Biologique, 29680 Roscoff, France
³⁵Norwegian institute for water research, Gaustadalleen 21, 0349 Oslo, Norway
³⁶Institut des Géosciences de l'Environnement 54, rue Molière 38402 - Saint Martin d'Hères, France
³⁷Faculty of fisheries sciences, Hokkaido University, Hakodate, Japan
³⁸UMR CNRS 5805 EPOC - OASU, Université de Bordeaux, 33615 PESSAC CEDEX, FRANCE
³⁹Environnement et changement climatique Canada
⁴⁰School of the Earth, Ocean and Environment, University of South Carolina, Columbia, South Carolina, 29208, USA
⁴¹Present address: Ocean Sciences Department, University of California, Santa Cruz, California, 95064, USA
⁴²LOCEAN-IPSL, Sorbonne Université, Paris, France
⁴³Applied Physics Laboratory, University of Washington, Seattle, Washington, USA

Correspondence: Philippe Massicotte (philippe.massicotte@takuvik.ulaval.ca)

Abstract. The Green Edge initiative was developed to investigate the processes controlling the primary productivity and the fate of organic matter produced during the Arctic phytoplankton spring bloom (PSB) and to determine its role in the ecosystem. Two field campaigns were conducted in 2015 and 2016 at an ice camp located on landfast sea ice southeast of Qikiqtarjuaq Island in Baffin Bay (67.4797N, 63.7895W). During both expeditions, a large suite of physical, chemical and biological variables was measured beneath a consolidated sea ice cover from the surface to the bottom at 360 m depth to better understand the factors driving the PSB. Key variables such as temperature, salinity, radiance, irradiance, nutrient concentrations, chlorophyll-a concentration, bacteria, phytoplankton

and zooplankton abundance and taxonomy, carbon stocks and fluxes were routinely measured at the ice camp. Here, we present the results of a joint effort to tidy and standardize the collected data sets that will facilitate their
10 reuse in other Arctic studies. The dataset is available at <https://www.seanoe.org/data/00487/59892/> (Massicotte et al., 2019a).

1 Introduction

In the Arctic Ocean, the phytoplankton spring bloom (PSB) initiates the period of highest biomass primary production of the year (Sakshaug, 2004; Perrette et al., 2011; Ardyna et al., 2013). Although it was discovered that the PSB
15 may occur more extensively and more frequently beneath a consolidated ice-pack (Arrigo et al., 2012, 2014; Assmy et al., 2017), only a small number of research initiatives (e.g., Fortier et al., 2002; Galindo et al., 2014; Mundy et al., 2009, 2014; Wassmann et al., 1999; Gosselin et al., 1997) have been investigating the processes controlling the Arctic PSB in the ice-covered water column. Additionally, ice algal communities play an important role within the Arctic food web and for the carbon export to the benthos during the winter-spring transition (Leu et al., 2015). However,
20 primary production within the Arctic ice-pack is still poorly understood. The Green Edge project was conceived in an effort to better understand the Arctic PSB from the level of fundamental physical, chemical and biological processes to that of their interactions within the ecosystem, and at spatial scales ranging from local to pan-Arctic. Besides studying each major component of the processes controlling Arctic PSB, another objective of Green Edge
25 was to investigate its impact on the nutrient and carbon dynamics within the ecosystem. A total of three Green Edge campaigns were conducted: two ice camp campaigns on landfast sea ice in 2015 and 2016, and an oceanographic cruise aboard the *CCGS Amundsen* in Baffin Bay in 2016. In this article, we present an overview of an extensive and comprehensive data set acquired during two surveys conducted at the Green Edge ice camp.

2 Study area, environmental conditions and sampling strategy

The field campaigns were conducted on landfast sea ice southeast of the Qikiqtarjuaq Island in Baffin Bay (67.4797N,
30 63.7895W, Fig. 1) in 2015 (April 24 - July 17) and in 2016 (April 20 to July 27). These periods were chosen in order to capture the dynamics of the sea-ice algae and phytoplankton spring blooms, from bloom initiation to termination. The field operations took place at a location (the "ice camp") south of the Qikiqtarjuaq Island where the water depth is 360 m. Continuous records of wind speed and air temperature were made with a meteorological station (Automated Meteo Mat equipped with temperature (HC2S3) and wind (05305-L) sensors (Campbell Scientific) positioned near (< 100 m) the tent (Polarhaven, Weatherhaven) in which water sampling was carried out. During the sampling periods, the study site experienced changes in snow cover and ice thickness (Fig. 2). In 2015, the snow and ice thickness varied between 2-40 cm (mean = 21 cm) and 103-136 cm (mean = 121) respectively. In 2016, the snow and ice thickness varied between 0.3-49 cm (mean = 19 cm) and 106-149 cm (mean = 128 cm) respectively. For both years, snowmelt began at the beginning of June and lasted for approximately two to three weeks (Oziel
35

40 et al., 2019). Water sampling was usually carried out every two days through a 1×1 m hole in the ice pack shielded by the tent. For the analysis of nutrient concentration, photosynthetic parameters, primary production, chlorophyll a (chl a), phytoplankton taxonomy and carbon stocks such as dissolved organic carbon (DOC), particulate organic carbon (POC), water samples were collected at 1.5, 5, 10, 20, 40 and 60 m using 10 or 20-L Niskin bottles. Details about specific measurements such as zooplankton and bacteria abundances are provided in the following sections.



Figure 1. Location of the ice camp located near the Qikiqtarjuaq Island in the Baffin Bay. Projection used: EPSG-4326.




Figure 2. Temporal evolution of the snow and sea-ice thickness for both ice camp missions. The dashed horizontal line represents the snow/ice interface.

45 3 Data quality control and data processing

Different quality control procedures were adopted to ensure the integrity of the data. First, the raw data were visually screened to eliminate errors originating from the measurement devices, including sensors (systematic or random) and errors inherent from measurement procedures and methods. Statistical summaries such as average, standard deviation and range were computed to detect and remove anomalous values in the data. Then, data were
 50 checked for duplicates and remaining outliers. Once raw measurements were cleaned, data were structured and regrouped into plain text comma-separated (CSV) files. Each of these files was constructed to gather variables of the same nature (ex.: nutrients). In each of these files, a minimum number of variables (columns) were always included so the different data sets can be easily merged together (Table 1). More than 120 different variables have been measured during the Green Edge landfast-ice expeditions. The complete list of variables is presented in Table
 55 2 and detailed metadata information can be found on the LEFE-CYBER online repository <http://www.obs-vlfr.fr/proof/php/GREENEDGE/greenedge.php>. The processed and tidied version of the data is hosted at SEANOE (SEA scieNtific Open data Edition) under the CC-BY license (<https://www.seanoe.org/data/00487/59892/>, Massicotte et al. (2019a)). In the following sections, we present a subset of these variables along with the methods used to collect

and measure them. For each of these variables, time series or vertical profiles are used to describe the data. Data
60 cleaning and visualization were performed with R 3.6.1 (R Core Team, 2019). The code used to produce the figures and the analysis presented in this paper is available under the GNU GPLv3 licence <https://github.com/PMassicotte/greenedge-icecamp-data-paper>. The code used to process and tidy the data provided by each researcher is also publicly available <https://gitlab.com/Takuvik/greenedge-database> under the GNU GPLv3 licence.

4 Data description: an overview

65 4.1 Physical data

Conductivity, temperature and depth (CTD) vertical profiles were measured using a Sea-Bird SBE19plusV2 CTD system (factory calibrated prior to the expedition) deployed from inside the Polarhaven tent between the surface and a 350 m depth. The data were post-processed according to the standard procedures recommended by the manufacturer and averaged into 1-m vertical bins. During the sampling periods, salinity was generally greater than 31.5
70 g kg⁻¹ (range: 4–34.4 g kg⁻¹). Flushes of freshwater at the ocean surface due to snow/ice melt started slowly at the beginning of June with the largest peaks/pulses taking place late June when salinity decreased to approximately 4 g kg⁻¹ (Fig. 3).




Figure 3. Temporal evolution of the salinity in the first 100 meters of the water column for both campaigns. Note that for visualization, salinity below 31.5 g kg^{-1} have been binned to 31.5 g kg^{-1} . Note that salinity as low as 4 g kg^{-1} was observed during flushes of freshwater at the ocean surface due to snow/ice melt (dark blue color in the figure).

Ocean current profiles in the water column were measured using a downward-looking 300 kHz Sentinel Workhorse Acoustic Doppler Current Profiler (ADCP, RDI Teledyne) mounted directly beneath the sea ice bottom. The study site
75 was dominated by seawater originating from the Arctic Ocean modulated by spring-neap tidal cycles (14 days) and semidiurnal M2 periods (≈ 12.4 hours). Vertical profiles of water column turbulence were measured on June 23 of 2016 during a spring tidal cycle (≈ 12.4) using a self-contained autonomous microprofiler (SCAMP, Precision Measurement Engineering, California, U.S.A.). The turbulence profile (i.e. a median profile of the rate of dissipation of turbulent kinetic energy, ϵ) showed a mixing layer depth of about 20–25 m characterized by an elevated dissipation
80 rate with values above $10^{-8} \text{ W kg}^{-1}$. The reader is referred to the paper by Oziel et al. (2019) for detailed methods, visualization and discussion of the CTD, SCAMP and ADCP data.

Vertical profiles (surface to 200 m) of CTD and bio-optical properties were measured every hour during a M2 tidal cycle measured on June 9, 2016 (an example of modelled surface tidal height versus time is shown in supplementary Fig. A1). These observations (Fig. 4) illustrate that internal tidal waves caused large vertical isopycnal displacements
85 (20–30 m) of all observed physical and biogeochemical properties below 50 m depth across the semi-diurnal M2 period. Hence, as vertical profiles of physical and bio-optical variables were measured at approximately the same time

each day, properties (assuming they follow a conservative mixing behaviour) will appear to be vertically displaced. Therefore, when comparing properties from vertical profiles taken at the ice camp, we suggest that comparisons of profile variables should be made on isopycnal (constant density) coordinates, rather than depth coordinates (Fig. 90 4).




Figure 4. Temporal evolution of physical (temperature) and bio-optical (CDOM fluorescence) variables with superimposed lines of potential density anomaly (σ_θ , kg m^{-3}) during a 13-h tidal cycle. Surface tidal height versus time at Qikiqtarjuaq is shown in blue. (A-B) Plotted versus pressure coordinates (equivalent to depth in meters). (C-D) The same data plotted versus potential density anomaly σ_θ coordinates (kg m^{-3}). The tidal survey was performed on 2015-06-09.

4.2 Underwater bio-optical data

4.2.1 Radiance and irradiance measurements with ICE-Pro

A total of 173 and 89 vertical light profiles were measured in 2015 and 2016, respectively, using a factory-calibrated ICE-Pro (an ice-floe version of the C-OPS, or Compact-Optical Profiling System, from Biospherical Instruments Inc.).

The ICE-Pro was equipped with radiometers for both downward plane irradiance (E_d , $\text{W m}^{-2} \text{nm}^{-1}$) and either upward irradiance (E_u , $\text{W m}^{-2} \text{nm}^{-1}$) in 2015 or upward radiance (L_u , $\text{W m}^{-2} \text{sr}^{-1} \text{nm}^{-1}$) in 2016. The profiles were taken at two sites, separated by approximately 40 m. In order to perform the profiles, the ICE-Pro was deployed through

auger holes that had been drilled at distances of 82 and 113 m from the tent and cleaned of ice chunks. Once
 100 the ICE-Pro was underneath the ice layer, fresh clean snow was shovelled back into the hole to avoid, as much as
 possible, having a bright spot above the sensors (see supplementary Fig. B1 and Table B1). The frame was then
 manually lowered at a rate of approximately 0.3 m s^{-1} . The above-surface reference sensor was fixed on a steady
 tripod installed approximately 2 m above the ice surface and above all neighbouring camp features. Data process-
 ing and validation were performed using a protocol inspired by that of Smith1984, which is now used by several
 105 space agencies. Measurements were taken between 380 and 875 nm at 19 discrete spectral wavebands. Vertical
 profiles were usually measured in duplicates or triplicates. Time series of daily photosynthetically active radiation
 (PAR) at the sea-ice/water interface (1.3 m depth) are shown in Fig. 5. In 2016, PAR started to increase rapidly in the
 second week of May, compared to early June in 2015. Overall, PAR at 1.3 m in the water column was also greater
 110 in 2016 than in 2015 and reached the threshold of $0.415 \text{ mol photons m}^{-2} \text{ d}^{-1}$, above which light is sufficient for
 net growth (Letelier et al., 2004), a few days earlier. Further information about in situ underwater irradiance and
 radiance measurements can be found in Massicotte et al. (2018).




Figure 5. Temporal evolution of daily photosynthetically available radiation (PAR) at the sea-ice/water interface (1.3 m depth) for both ice camp missions. The horizontal dashed line shows the $0.415 \text{ mol photons m}^{-2} \text{ d}^{-1}$ threshold often used in the literature as the minimum light requirement for primary production.

4.2.2 Underwater photos and videos of ice bottom

Several vertical profiles to 30 m were performed using a GoPro Hero 4 camera mounted on the ICE-Pro and pointing up, towards the ice bottom (see Fig. B1 and Table B1). Still images were captured every five seconds during descent, as well as a video was taken of the complete descent. These photos and videos were used for a qualitative
 115 assessment of the pronounced spatial and temporal heterogeneity of the under-ice environment and the associated water column nekton community between the two profiling locations.

4.2.3 Irradiance measurements with TriOS

To quantify the impact of the heterogeneous radiation field under sea ice on irradiance measurements, replicated spectral irradiance profiles were collected beneath landfast sea ice from 5 May to 8 June 2015 and from 14 June to
120 4 July 2016. The replicates were made on each sampling day, under different surface conditions. In 2015, measurements were performed prior to melt onset, under different snow depths. In 2016, measurements began after the onset of snowmelt and were performed beneath sea ice with a wet snow cover, shallow melt ponds and white ice. The deployed sensor array consisted of a surface reference radiometer, which recorded incident downwelling planar irradiance, $E_d(0, \phi)$, and three radiometers attached to a custom-built double-hinged aluminum pole (under-ice
125 L-arm) to measure downwelling planar irradiance, $E_d(z, \phi)$, downwelling scalar irradiance, $\dot{E}_d(z, \phi)$, and upwelling scalar irradiance, $\dot{E}_u(z, \phi)$. These four hyperspectral radiometers (two planar RAMSES-ACC and two scalars RAMSES-ASC, TriOS GmbH, Germany) measured pressure and tilt internally and recorded irradiance spectra in the wavelength range from 320 to 950 nm at a resolution of 3.3 nm (190 channels). Transmitted irradiance was recorded along with vertical profiles by lowering the L-arm manually through a 20-inches auger hole with a winch and 1.5-
130 m aluminum poles extensions. In 2015, 17 vertical profiles were collected in 0.4 - 0.5-m depth steps from the ice bottom to a water depth of 18 m. In 2016, 11 profiles were recorded to a depth of 20 m under different sea ice surface conditions. Differences between planar and scalar PAR measurements were used to derive the downwelling average cosine, μ_d , an index of the angular structure of the downwelling under-ice radiation field which, in practice, can be used to convert between downwelling scalar, \dot{E}_d , and planar, E_d , irradiance. The average cosine was smaller
135 prior to snowmelt in 2015 compared to after snowmelt (≈ 0.6 vs. 0.7), when melt ponds covered the ice surface in 2016 (Fig. 6). Further details about the sampling procedure, data processing and results can be found in Matthes et al. (2019).




Figure 6. **(A)** Under-ice vertical profiles of downwelling planar and scalar irradiance at 442 nm, 532 nm and for PAR. Note the log scale for the irradiance measurements. **(B)** Calculated downwelling average cosine (unitless) was measured beneath snow-covered sea ice on 16 May 2015, beneath bare ice on 20 June 2016 and beneath a melt pond on 4 July 2016.

4.2.4 Inherent optical properties (IOP)

IOPs measurements were made using an optical frame equipped with the physical and bio-optical sensors that
140 were factory calibrated before each field campaign. A Seabird SBE-9 CTD measured temperature, salinity, and pressure. A WetLabs AC-S was used for spectral beam attenuation (c , m^{-1}) and total absorption (a , m^{-1}) between 405 and 740 nm, and a BB9 (WetLabs) and a BB3 (WetLabs) were utilized for backscattering coefficients (bb , m^{-1}) between 440 and 870 nm. During both campaigns, pure water calibration was performed for the AC-S sensor on each sampling day and linear regression as a function of time was computed for each wavelength of absorption and attenuation
145 signals. Then, the offset applied during the data processing was taken on this linear regression at the exact date of the measurement. Figure 7 shows two vertical profiles of attenuation coefficients at different wavelengths acquired during pre-bloom and bloom conditions in 2016. One can see that during the bloom, attenuation increased markedly in the 0-50 m surface layer due to higher phytoplankton biomass.




Figure 7. Beam attenuation coefficients (c , m^{-1}) measured in 2016 using an ACS before and during the phytoplankton bloom. Note that the colors of the lines correspond to wavelength frequencies.

4.2.5 Other optical measurements

150 Other optical variables measured during both field campaigns included absorbance of particulate matter, absorbance of dissolved organic matter, snow and sea-ice transmittance, snow/ice hyperspectral and hyperangular hemispherical-directional-reflectance (Goyens et al., 2018) and surface spectral albedo (Table 2). Downwelling spectral irradiance above the surface ($1^\circ \times 1^\circ$) spatial resolution, daily temporal resolution, interpolated hourly) was also computed based on the radiative transfer model SBDART (Ricchiazzi et al., 1998) as described in Laliberté et al.
155 (2016) and Randelhoff et al. (2019).

4.3 Nutrients

Nitrate, nitrite, phosphate and silicate concentrations were measured from water filtered through $0.7\text{ }\mu\text{m}$ Whatman GF/F filters and through $0.2\text{ }\mu\text{m}$ cellulose acetate membranes. Filtrates were collected into sterile 20 mL polyethylene vials, poisoned with $100\text{ }\mu\text{L}$ of mercuric chloride (60 mg L^{-1}) and subsequently stored in the dark prior to analysis. Nutrient concentrations were determined using an automated colorimetric procedure described in Aminot and Kérouel (2007). Figure 8 shows an overview of the dynamics of nitrate which is often the limiting nutrient for phytoplankton growth in the ocean (Tremblay and Gagnon, 2009). It can be seen that the depletion of the nitrates started approximately mid-June for both years, coinciding with the initiation of the phytoplankton bloom. However, the depletion was observed deeper in the water column in 2016 compared to 2015 due to stronger currents and
160 a longer sampling period in 2016 (Oziel et al., 2019). Other nutrients such as dissolved organic and inorganic carbon (DOC/DIC), particulate organic and inorganic carbon (POC/PIC), total organic carbon (TOC), phosphate (PO₄), orthosilicic acid (Si(OH)₄), and ammonium (NH₄), were also measured during both campaigns (Table 2). Detailed information about analytical procedures can be found in the LEFE-CYBER online repository. A comprehensive discussion about nutrient dynamics during the Green Edge missions can be found in Grondin et al. (2019).




Figure 8. Temporal evolution of the nitrates in the first 60 m of the water column for both ice camp missions.

170 4.4 Bacteria and Phytoplankton

4.4.1 Flow cytometry

The abundances of pico-phytoplankton, nano-phytoplankton and bacteria were measured by flow cytometry. Samples (1.5 mL) were preserved with a mix of glutaraldehyde and Pluronic (Marie et al., 2014) and frozen at -80°C. Samples were analyzed on a FACS Canto flow cytometer (Becton Dickinson) in the laboratory at the Station Biologique de Roscoff. The abundance (cells mL^{-1}) of phytoplankton populations was determined on unstained samples and cells were discriminated by their red chlorophyll autofluorescence. Bacterial abundance was determined based on the fluorescence of SYBR Green-stained DNA (Marie et al., 1997). In both 2015 and 2016, bacterial concentrations were initially low, of the order of 100 000 cells mL^{-1} , and quite uniform throughout the water column. During the bloom, bacterial abundance increased continuously, reaching values of one million cells mL^{-1} (Fig. 9). Simultaneously, the distribution of highest abundance became stratified with a higher concentration found near the surface in early July before it moved down to the subsurface (between 10 and 20 m) later in July (Fig. 9). In 2015, the sampling period did not extend long enough to capture the full progression of bacterial community development.




Figure 9. Concentration of bacteria in the water column at the ice camp in 2015 and 2016.

4.5 Phytoplankton

4.5.1 Chlorophyll a

185 Chl a and accessory pigments concentrations were determined by high-performance liquid chromatography (HPLC) following Ras2008. Concentrations were measured using volumes between 0.1 and 1 L of melted ice and volumes
 between 1 and 2.5 L of seawater. Water was filtered onto Whatman GF/F 25 mm filters and stored at -80°C until
 analysis. Filters were extracted in 100% methanol, disrupted by sonication and clarified by filtration. Pigments were
 analyzed using an Agilent Technologies 1200 Series system with a narrow reversed-phase C8 Zorbax Eclipse XDB
 190 column (150 × 3 mm, 3.5 µm particle size) which was maintained at 60°C. Figure 10 shows the temporal evolution of
 surface integrated chl a in the bottom 10 cm of the ice cover and the water column for both years. At the beginning
 of the sampling periods in 2015 and 2016, total chl a concentrations in the bottom of the ice and the water column
 were of approximately the same magnitude ($\approx 5 \text{ mg m}^{-2}$). Later in the season, when the snowpack and the ice sheet
 started to melt (between June and July), and at the onset of the PSB, chl a in the water column increased rapidly to
 195 reach concentrations of 145 mg m^{-2} in 2015 and 113 mg m^{-2} in 2016. At the same time, or slightly before, chl a in
 the ice bottom started to decrease rapidly to concentrations varying between 0.1 and 0.3 mg m^{-2} .




Figure 10. Temporal evolution of chlorophyll a in ice and water (depth-integrated) for both ice camp missions. Note that the water chlorophyll a have been integrated over the first 100 m of the water column whereas the ice chlorophyll a was measured on the bottom 0-10 cm of the ice cores. The details of the calculations to determine the approximate dates of phytoplankton bloom initiation can be found in Oziel et al. (2019).

Primary production during the phytoplankton bloom was incompletely sampled in 2015, while in 2016 it was monitored from the onset under melting sea ice in May to its termination in July (Fig. 11). During the ice-covered period in 2015, primary production, as well as nitrate assimilation (rNO_3), occurred at very low but detectable rates 200 reaching 8 and 0.4 $\text{mmol m}^{-2} \text{ d}^{-1}$, respectively. Phytoplankton production rates were higher in the ice than in the water column, representing approximately 80% and 40% for primary production and rNO_3 , respectively. Estimated assimilated concentrations of total carbon and nitrate within the ice cover were 30-96 and 1.4-4.6 mmol m^{-2} during 205 this period. The break-up of the sea ice cover was characterized by a rapid increase in primary production and rNO_3 . During this period of high light transmission through the melting ice cover (day 169 to 190), concentrations of assimilated total carbon and rNO_3 reached 60 and 8 mmol m^{-2} , respectively, leading to a complete nitrate depletion. The quantities of total carbon and nitrate assimilated during the 2016 PSB in the water column were 562 and 97 mmol m^{-2} , respectively.




Figure 11. Temporal evolution of primary production a in ice and water (depth-integrated) for both ice camp missions.

4.5.2 Phytoplankton taxonomy

The phytoplankton community species composition was determined using an Imaging FlowCytobot (IFCB, Woods Hole Oceanographic Institute, Sosik and Olson (2007), Olson and Sosik (2007)). The size range targeted was between 1 and 150 μm , while the image resolution of approximately 3.4 pixels μm^{-1} limited the identification of cell < 10 μm to broad functional groups. A 150 μm Nitex mesh was used to avoid clogging of the fluidics system by large particles, although this might have induced a bias in the results by preventing large cells to be sampled. For each melted ice and seawater sample, 5 mL were analyzed and Milli-Q water was run between samples with high biomass in order to prevent contamination between samples. Image acquisition was triggered by chl a in vivo fluorescence, with excitation and emission wavelengths of 635 and 680 nm, respectively. Grayscale images were processed to extract regions of interest (ROIs) and their associated features (e.g.: geometry, shape, symmetry, texture, etc.), using a custom made MATLAB (2013b) code (Sosik and Olson (2007), Olson and Sosik (2007); processing codes are available at <https://github.com/hsosik/ifcb-analysis>). A total of 231 features (see the full list and description at <https://github.com/hsosik/ifcb-analysis/wiki/feature-file-documentation>) were derived on the resulting ROIs and were used for automatic classification using random forest algorithms with the EcoTaxa application (Picheral et al., 2017). A learning set was manually prepared for each year, with ca. 20 000 images annotated and used for

automatic prediction. Each automatically annotated image was further validated by visual examination and corrected when necessary. The final 2015 and 2016 datasets consist of 124 247 and 57 397 annotated images and
225 their associated features in 39 and 35 taxonomic categories, respectively (Fig. 12). As it was impossible to count the number of cells in each image, we assumed one cell per image. To account for potential underestimations of cell abundance when colonies or chains were imaged, the biovolume of each living protist on images was computed during image processing according to Moberg and Sosik (2012). Using carbon to volume ratios from Menden-Deuer and Lessard (2000), biovolume was converted into carbon estimates, as described in Laney and Sosik (2014). De-
230 tailed information about sea ice algae and phytoplankton community composition can be found in Grondin et al. (2019).




Figure 12. Images of protists sampled with the IFCB. Scale bar on images is 10 µm. Note that images are not to scale. (A) *Anabaena* sp. (B) *Nitzschia frigida* (C) *Polarella glacialis* (D) Flagellate (E) *Euglena* (F) *Pseudo-nitzschia* sp. (G) *Ceratium* sp. (H) *Thalassiosira nordenskioeldii* with *Attheya septentrionalis* (I) *Peridiniella catenata* (J) *Navicula pelagica* (K) *Phaeocystis* sp. colony (L) *Chaetoceros* sp. (M) *Entomoneis* sp. (N) *Synedropsis hyperborea* (O) Ciliate (P) Pennate diatom (Q) *Eucampia* sp. (R) *Melosira* sp.

4.5.3 Physiology of the phytoplankton community

The photosynthetic potential of microalgae was assessed by measuring Fv/Fm , namely the maximum photochemical efficiency of Photosystem II (PSII), via dynamic chl a fluorescence:

$$235 \quad \frac{Fv}{Fm} = \frac{(Fm - F0)}{Fm} \quad (1)$$

where Fm and $F0$ are the maximum and minimum PSII chl a fluorescence yields, respectively. Chl a fluorescence was recorded with a Water-PAM fluorometer (Walz, Germany) on melted sea-ice (last centimeter of the cores) and

water samples collected at different depths (i.e. 1.5 m, 10 m, 40 m, 60 m). Measurements were performed after storing samples in 50 mL dark Falcon tubes (Corning Life Sciences, USA) on ice for at least 1 h. For further technical details, see Galindo et al. (2017). Fv/Fm is often used as an index for evaluating the physiological condition of microalgal communities. For algae that are growing optimally, the Fv/Fm ratio ranges between 0.50 and 0.75 in the absence of cyanobacteria. Below 0.50, algal growth is considered to be limited by nutrient availability and/or light stress (Suggett et al., 2010). Figure 13 shows the temporal evolution of Fv/Fm for ice algae and phytoplankton for the ice camp in 2016. At the beginning of the sampling period, all samples showed Fv/Fm above 0.55. While in ice Fv/Fm ranged between 0.60 and 0.75 until the beginning of June, it decreased to ca. 0.20-0.35 in water. This decrease of Fv/Fm (Fig. 13A) is coincident with a sharp increase in PAR under the ice sheet (Fig. 5), which may have induced light stress in phytoplankton and ice algae communities. After approximately 1 month, phytoplankton became acclimated to this new light environment and Fv/Fm increased back to 0.60-0.75 by the beginning of June. From that time on (corresponding to higher irradiance transmittance through ice, see Fig. 5), Fv/Fm in ice decreased dramatically to an approximate value of 0.20 while Fv/Fm in the water column generally remained between 0.60 and 0.75 for depths between 10 and 60 m (note however the large decrease at 40 m on June 13). In contrast, Fv/Fm at 1.5 m was lower and noisier with values varying between 0.45 and 0.60.




Figure 13. (A) Temporal evolution of F_v/F_m for ice (last cm) and water underneath the ice (depths 1.5 m, 10 m, 40 m) samples for the ice camp 2016 between May 6th and July 8th. F_v/F_m monitoring on ice samples stopped on June 20th because the chl a fluorescence signal was not reliable anymore. F_v/F_m monitoring on 40 m and 60 m depth samples was limited between May 13th and June 24th and between June 29th and July 08th, respectively. The gray shaded area represents the range at which the algae are optimally growing. **(B)** The light saturation parameter, E_k , an index of photoadaptation of the phytoplankton community measured at 1.5 m, 5 m and 10 m depth. Note de log scale on the y axis.

In addition to the photosynthetic potential of microalgae, photosynthetic parameters were measured from seawater incubated at different irradiance levels in the presence of ^{14}C labelled sodium bicarbonate. The light saturation parameter, E_k , is an indication of the physiological state of the phytoplankton community. Figure 13B shows the increase of E_k as the phytoplankton community grows between May and July of 2016 at 1.5 m, 5 m and 10 m depth. Between 1.5 m and 10 m depth, E_k varied between 15 and 194 $\mu\text{mol m}^{-2} \text{s}^{-1}$ ($61 \pm 37 \mu\text{mol m}^{-2} \text{s}^{-1}$, $n = 69$) which fall in range within values reported in other marine studies conducted at high-latitudes (Bouman et al., 2018; Massicotte et al., 2019b). The observed increase in E_k over the growing season suggests that the phytoplankton community became more photo-adapted to increasing available irradiance (Fig. 5).

4.6 Zooplankton

Zooplankton was collected from a ring net deployed under the ice at the ice camp between April 22 and June 10 in 2015 and between May 16 and July 18 in 2016. This sampler, composed of a 1 m diameter circular frame mounted with a 4 m long 200 µm mesh size conical plankton net was lowered cod-end first to avoid filtration
265 during the descent, using an electric winch. An additional 50 µm net with an aperture of 10 cm in diameter was attached to the side of the metal ring to sample eggs and small zooplankton larvae while the main net collected the mesozooplankton fraction. This sampling device was hauled vertically from a depth of 100 m (2015 and 2016) or 350 m (only in 2016), 10 m above the seafloor to the surface at a speed of about 30m min⁻¹. The filtered volume
270 was estimated by a KC Denmark flowmeter placed in the mouth of the 200 µm mesh net. Samples were preserved in 10% buffered formalin seawater solution for further taxonomic analyses. Classification and count of the 200 µm mesh net samples from both campaigns were performed using the zooscan by the PIQv team at l'Observatoire Océanographique de Villefranche-sur-Mer, France, following their protocol. Figure 14 shows the time series of the abundance of copepods (the dominant group of zooplankton in the Arctic) for the first 100 m and 350 m of the water column in 2016.




Figure 14. Time series of the abundance of the copepods (ind m⁻³) measured over the first 100 m and 350 m of the water column in 2016 using the zooscan. For visualization, only the six most abundant groups are presented in decreasing order of importance. Note the different *y* axes in both panels.

275 Highest copepod abundance was observed in late May and early June in both the top 100 m and over 350 m hauling depths. At the beginning of the sampling period, abundance was approximately 10 times higher in the first 100 m of the water column than over 350 m, suggesting that copepods were agglomerating near the surface to exploit the ice algae production before the start of phytoplankton production. Abundance started to decrease during the first week of June. The family of Oithonidae and the order of Calanoida were the two most abundant groups 280 over the 2 sampling depths. Oithonidae was more abundant over the top 100m layer as this group is probably mainly composed of small epipelagic *Oithona similis* one of the most numerous copepods in the Arctic. Calanoida, the most common copepod order, which includes the families Calanidae (including species such as *Calanus spp.*) and Acartiidae, was the dominant group over the 350m depth haul.

4.7 Other data

285 An exhaustive list of all measured variables is presented in Table 2 along with contact information of principal investigators associated with each measured parameter.

5 Recommendations and lessons learned

As with any Arctic surveys, a large number of measurements were acquired during the Green Edge project. Although initial recommendations on good practices about collection, processing and storage of collected data were communicated to all scientists, extensive efforts, such as data standardization, had to be performed to assemble the data. It is important for reducing possible errors, that a uniformized data management plan should be prepared and distributed prior to each mission. Furthermore, dedicated data management specialists should be involved from the beginning of the project to ensure the data are adequately collected, tidied, stored, backed up and archived.

6 Conclusions

295 The comprehensive data set assembled during both Green Edge ice-camp campaigns allowed us to study the fundamental physical, chemical and biological processes controlling the Arctic PSB. In this paper, only a handful of variables have been presented. The reader can find the complete list of measured variables in Table 2, all of which are also fully available in the data repository. Furthermore, a collection of scientific research papers is currently being submitted to a special issue of the Elementa journal entitled *Green Edge -The phytoplankton spring bloom in*
300 *the Arctic Ocean: past, present and future response to climate variations, and impact on carbon fluxes and the marine food web*. The uniqueness and comprehensiveness of this data set offer more opportunities to reuse it for other applications.

7 Code and data availability

The raw data provided by all the researchers, as well as metadata, are available on the LEFE-CYBER repository
305 (<http://www.obs-vlfr.fr/proof/php/GREENEDGE/greenedge.php>). The data presented in this paper and in Table 2 are hosted at SEANOE (SEA scieNtific Open data Edition) under the CC-BY license (<https://www.seanoe.org/data/00487/59892/>, Massicotte et al. (2019a)). Detailed metadata are associated with each file including the principal investigator's contact information. For specific questions, please contact the principal investigator associated with the data (see Table 2).

Tables

Table 1: Descriptions of the minimal variables included in each data set (i.e. in each CSV file).

Variable	Description
date	Sampling date (UTC)
latitude	Latitude of the sampling location (degree decimals).
longitude	Longitude of the sampling location (degree decimals).
sample_type	Origin of the water ("water", "ice", "meltpond").
sample_source	Source of the water ("niskin", "underice" "0-1 cm", "0-3 cm", "3-10 cm", "rosette").
depth_m	Depth at which measurement was made.
snow_thickness	Qualitative value describing the snow cover under which measurement was made ("thin_snow", "thick_snow").
mission	Mission identifier ("ice_camp_2015", "ice_camp_2016")
pi	Name(s) of the principal investigator(s) responsible of the measured variable.

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year.

Year	Parameter	Sampling method	Principal investigators	Processed
2015	Absorption coefficient	In-water profiler	Becu G. / Babin M.	Available
2015	Absorption (particulate)	Camp ice sample	Ehn J. / Cox C.	Available
2015	Absorption (particulate)	Camp water sample	Ehn J. / Cox C.	Available
2015	Absorption (particulate)	Camp ice sample	Matsuoka A. / Bricaud A. / Ferland J.	Available
2015	Absorption (particulate)	Camp water sample	Matsuoka A. / Bricaud A. / Ferland J.	Available
2015	ADCP (Mooring)	Mooring	Marec C.	Available
2015	Aerosol optical depth	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Aerosol relative humidity	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Air Relative Humidity	Meteorological Tower	Massé G.	Available
2015	Air Temperature	Meteorological Tower	Massé G.	Available
2015	Alkalinity total (TA)	Camp water sample	Else B. / Whitehead J.	Available
2015	Ammonium (NH_4^+)	Camp water sample	Raimbault P.	Data not available yet
2015	Ammonium (NH_4^+ , assimilation)	Camp water sample	Raimbault P.	Available
2015	Ammonium (NH_4^+ , regeneration)	Camp water sample	Raimbault P.	Available
2015	Angstrom coefficient	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Attenuation coefficient	In-water profiler	Becu G. / Babin M.	Available
2015	Backscattering coefficient	In-water profiler	Becu G. / Babin M.	Available
2015	Bacterial sequencing	Air filtration	Amiraux R.	Available
2015	Bacterial sequencing	Camp water sample	Amiraux R.	Available
2015	Bacterial sequencing	Ice core	Amiraux R.	Available
2015	Bacterial sequencing	Sediment trap	Amiraux R.	Available
2015	Brine salinity and volume	Sea ice core	Galindo V. / Rysgaard S.	Available
2015	Chlorophyll a	In-water profiler	Becu G. / Babin M.	Available
2015	Chlorophyll a	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Chlorophyll a and Phaeopigments (concentration)	Camp water sample	Babin M. / Ferland J.	Available
2015	Chlorophyll a and phaeopigments (concentration)	Camp water sample	Raimbault P.	Data not available yet
2015	Chromophoric dissolved organic matter absorption	In-water profiler	Becu G. / Babin M.	Available
2015	Chromophoric dissolved organic matter absorption	Camp water sample	Matsuoka A. / Ferland J. / Babin M.	Available
2015	Conductivity, temperature, and depth (CTD)	In-water profiler	Becu G. / Babin M.	Available
2015	Conductivity, temperature, and depth (CTD)	In-water profiler	Guillot P. / Babin M. / Marec C.	Available
2015	Cryptophytes (abundance)	Camp water sample	Vaulot D. / Marie D.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (continued)

Year	Parameter	Sampling method	Principal investigators	Processed
2015	Diffuse attenuation coefficient (Kd)	Profile mode	Becu G. / Babin M.	Available
2015	Dimethyl sulfide (DMS)	Camp water sample	Levasseur M.	Available
2015	Dimethyl sulfide (DMS)	Melt pond water sample	Levasseur M.	Available
2015	Dimethyl sulfide (DMS)	Sea ice core	Levasseur M.	Available
2015	Dimethylsulfoniopropionate (DMSP)	Camp water sample	Levasseur M.	Available
2015	Dimethylsulfoniopropionate (DMSP)	Melt pond water sample	Levasseur M.	Available
2015	Dimethylsulfoniopropionate (DMSP)	Sea ice core	Levasseur M.	Available
2015	Dissolved inorganic Carbon (DIC)	Camp water sample	Else B. / Whitehead J.	Available
2015	Dissolved organic matter (sugars)	Rosette	Sempéré R. / Panagiotopoulos C.	Available
2015	Dissolved organic nitrogen (release)	Camp water sample	Raimbault P.	Available
2015	Downwelling irradiance	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Downwelling Irradiance above the surface ($E_d(0^+)$)	Surface mode	Babin M. / Galí M.	Available
2015	Downwelling Irradiance above the surface ($E_d(0^+)$)	Profile mode	Becu G. / Babin M.	Available
2015	Downwelling Irradiance ($E_d(z)$)	Profile mode	Becu G. / Babin M.	Available
2015	$E_d(0^+)$ spectra from SBDART radiative transfer simulations	Surface mode	Babin M. / Galí M.	Available
2015	Faecal pellets flux	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Fluorescence Variable (phytoplankton)	Camp water sample	Galindo V. / Rysgaard S.	Data not available yet
2015	Fluorescence Variable (phytoplankton)	Sediment Trap	Galindo V. / Rysgaard S.	Available
2015	Fluorescence Variable (phytoplankton)	Surface mode	Galindo V. / Rysgaard S.	Data not available yet
2015	Hemispherical directional reflectance function	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Hemispherical Directional Reflectance Factor	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Heterotrophic bacteria (abundance)	Camp water sample	Vaulot D. / Marie D.	Available
2015	Heterotrophic nanoflagellates	Camp water sample	Joux F.	Available
2015	Ice and snow temperature	Meteorological Tower	Massé G.	Available
2015	Ice thickness	Camp ice sample	Galindo V. / Rysgaard S.	Available
2015	Irradiance (downwelling, upwelling)	Surface-, Under-water profile-mode	Matthes L. / Ehni J. / Lambert-Girard S. / Mundy C.J.	Available
2015	Isoprenoid lipids	Camp water sample	Massé G. / Guilmette C.	Data not available yet
2015	Isoprenoid lipids	Sea ice core	Massé G. / Guilmette C.	Data not available yet
2015	Net radiation	Surface mode	Else B.	Available
2015	Nitrate (NO_3^-)	Camp water sample	Raimbault P.	Available
2015	Nitrate (NO_3^-)	Sea ice core	Raimbault P.	Available
2015	Nitrate (NO_3^- , assimilation)	Camp water sample	Raimbault P.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (*continued*)

Year	Parameter	Sampling method	Principal investigators	Processed
2015	Nitrification	Camp water sample	Raimbault P.	Available
2015	Nitrite (NO_2^-)	Camp water sample	Raimbault P.	Available
2015	Nitrite (NO_2^-)	Sea ice core	Raimbault P.	Available
2015	PAR from SBDART radiative transfer simulations	Surface mode	Babin M. / Galí M.	Available
2015	Particle Size Distribution	In-water profiler	Becu G. / Babin M.	Available
2015	Particles size	Underwater Vision Profiler (UVP)	Marec C. / Picheral M.	Available
2015	Particulate Carbon (PC)	Camp water sample	Babin M. / Ferland J.	Available
2015	Particulate mass	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Particulate Nitrogen (PN)	Camp water sample	Babin M. / Ferland J.	Available
2015	Particulate nitrogen (PN)	Sediment Trap	Fortier L. / Lalande C.	Data not available yet
2015	Particulate organic carbon (POC)	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Particulate organic carbon (POC)	Camp water sample	Raimbault P.	Available
2015	Particulate organic nitrogen (PON)	Camp water sample	Raimbault P.	Available
2015	Particulate Organic Phosphorus (POP)	Camp water sample	Raimbault P.	Data not available yet
2015	PDMPO uptake	Camp water sample	Leynaert A.	Data not available yet
2015	PDMPO uptake per species	Camp water sample	Leynaert A.	Data not available yet
2015	Phosphate ($(\text{PO}_4)^{3-}$)	Camp water sample	Raimbault P.	Available
2015	Phosphate ($(\text{PO}_4)^{3-}$)	Sea ice core	Raimbault P.	Available
2015	Photosynthetically available radiation (PAR)	Surface mode	Babin M. / Galí M.	Available
2015	Photosynthetically available radiation (PAR)	Profile mode	Becu G. / Babin M.	Available
2015	Photosynthetic nanoeukaryotes (abundance)	Camp water sample	Vaulot D. / Marie D.	Available
2015	Photosynthetic parameters	Camp water sample	Ferland J. / Babin M.	Available
2015	Photosynthetic picoeukaryotes (abundance)	Camp water sample	Vaulot D. / Marie D.	Available
2015	Phytoplankton	Camp water sample	Ferland J. / Grondin P.L. / Babin M. / Marec C.	Available
2015	Phytoplankton (taxonomy)	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Pigments	Sea ice core	Galindo V. / Rysgaard S.	Available
2015	Pigments	Camp water sample	Ras J. / Claustré H.	Available
2015	Primary production	Camp water sample	Raimbault P.	Available
2015	Rrs (I^+)	Profile mode	Becu G. / Babin M.	Available
2015	Salinity	Sea ice core	Galindo V. / Rysgaard S.	Available
2015	Salinity-induced bacterial biomarker	Ice core	Amiraux R. / Rontani J-F.	Available
2015	Salinity-induced bacterial biomarker	Sediment trap	Amiraux R. / Rontani J-F.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (continued)

Year	Parameter	Sampling method	Principal investigators	Processed
2015	Sea ice concentration	Surface mode	Massicotte P.	Available
2015	Silica Biogenic (BSi)	Camp water sample	Leynaert A.	Available
2015	Silica Biogenic (BSi) dissolution rate	Camp water sample	Leynaert A.	Available
2015	Silicate $Si(OH)_4$ - absorption kinetics	Camp water sample	Leynaert A.	Available
2015	Silica (uptake rate)	Camp water sample	Leynaert A.	Available
2015	$Si(OH)_4$	Camp water sample	Leynaert A.	Available
2015	$Si(OH)_4$	Camp water sample	Raimbault P.	Available
2015	$Si(OH)_4$	Sea ice core	Raimbault P.	Available
2015	Snow depth	Camp snow sample	Galindo V.J. Rysgaard S.	Available
2015	Snow depth	Meteorological Tower	Massé G.	Available
2015	Sugars	Sediment Trap	Sempéré R. / Panagiotopoulos C.	Available
2015	Surface Albedo	Surface mode	Verin G.	Available
2015	Suspended particulate material (SPM)	Camp water sample	Babin M. / Ferland J.	Available
2015	Swimmers	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Synechococcus (abundance)	Camp water sample	Vaulot D. / Marie D.	Available
2015	Temperature	Sea ice core	Galindo V. / Rysgaard S.	Available
2015	Total organic carbon (TOC)	Rosette	Sempéré R. / Panagiotopoulos C.	Available
2015	Total organic carbon (TOC)	Camp water sample	Raimbault P.	Available
2015	Total organic nitrogen (TON)	Camp water sample	Raimbault P.	Available
2015	Total organic phosphorus (TOP)	Camp water sample	Raimbault P.	Available
2015	Transmittance through ice	Surface mode	Verin G.	Available
2015	Under-ice export fluxes of biogenic matter (fresh)	Sediment Trap	Fortier L. / Lalande C.	Available
2015	Under-ice photos and video	GoPro Hero 4 on radiometer profiler	Rehm E.	Available
2015	Upwelling Irradiance ($E_u(z)$)	Profile mode	Becu G. / Babin M.	Available
2015	Upwelling radiance ($L_u(z)$)	Surface mode	Belanger S. / Goyens C. / Leymarie E.	Available
2015	Upwelling radiance ($L_u(z)$)	Profile mode	Becu G. / Babin M.	Available
2015	Vertical profile of snow density	Surface mode	Verin G.	Available
2015	Vertical profile of Specific Surface Area	Surface mode	Verin G.	Available
2015	Virus (abundance)	Camp water sample	Joux F.	Available
2015	Wind Direction	Meteorological Tower	Massé G.	Available
2015	Wind Speed	Meteorological Tower	Massé G.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (continued)

Year	Parameter	Sampling method	Principal investigators	Processed
2015	Zooplankton (Abundances)	Plankton net	Fortier L. / Aubry C	Available
2015	Zooplankton (Abundances)	Plankton net (LOKI)	Fortier L. / Aubry C	Available
2015	Zooplankton (Taxonomy)	Plankton Net	Fortier L. / Aubry C	Available
2015	Zooplankton (Taxonomy)	Plankton net (LOKI)	Fortier L. / Aubry C	Available
2015	Zooplankton vertical distribution	Underwater Vision Profiler (UVP)	Marec C. / Sophie R. / Picheral M.	Available
2016	234Th (dissolved)	Rosette	Schmidt S.	Data not available yet
2016	234Th (particulate)	Rosette	Schmidt S.	Data not available yet
2016	Absorption coefficient	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Absorption (particulate)	Camp ice sample	Matsuoka A. / Bricaud A. / Ferland J.	Available
2016	Absorption (particulate)	Camp water sample	Matsuoka A. / Bricaud A. / Ferland J.	Available
2016	ADCP (Mooring)	Mooring	Oziel L. / Houssais M.-N. / Babin M. / Lagunas J.	Available
2016	Air Relative Humidity	Meteorological Tower	Massé G.	Available
2016	Air Temperature	Meteorological Tower	Massé G.	Available
2016	Ammonium (NH_4^+)	Camp water sample	Raimbault P.	Data not available yet
2016	Ammonium (NH_4^+ , assimilation)	Camp water sample	Raimbault P.	Available
2016	Ammonium (NH_4^+ , regeneration)	Camp water sample	Raimbault P.	Available
2016	Attenuation coefficient	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Backscattering coefficient	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Bacterial cultures	Camp water sample	Joux F.	Available
2016	Bacterial cultures	Sea ice core	Joux F.	Available
2016	Bacterial production	Sea ice core	Joux F. / Galindo V.	Available
2016	Bacterial production	Camp water sample	Joux F. / Galindo V.	Available
2016	Brine salinity and volume	Sea ice core	Galindo V. / Rysgaard S.	Available
2016	Chlorophyll a	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Chlorophyll a	Sediment Trap	Fortier L. / Lalande C.	Available
2016	Chlorophyll a and phaeopigments (concentration)	Camp water sample	Babin M. / Ferland J.	Available
2016	Chromophoric dissolved organic matter absorption	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Chromophoric dissolved organic matter absorption	Camp water sample	Matsuoka A. / Ferland J. / Babin M.	Available
2016	Chromophoric dissolved organic matter fluorescence	Camp water sample	Matsuoka A. / Ferland J.	Available
2016	Conductivity, temperature, and depth (CTD)	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Conductivity, temperature, and depth (CTD)	In-water profiler	Guillot P. / Lagunas J.	Available
2016	Cryptophytes (abundance)	Camp water sample	Vaulot D.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (continued)

Year	Parameter	Sampling method	Principal investigators	Processed
2016	Cultures of sorted populations	Camp water sample	Vaulot D.	Available
2016	Diatoms (bacilliarophyta) abundance	Camp water sample	Leblanc K. / Queguiner B. / Lafond.A	Available
2016	Diatoms (bacilliarophyta) taxonomy	Camp water sample	Leblanc K. / Queguiner B. / Lafond.A	Available
2016	Diffuse attenuation coefficient (Kd)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	Dissolved organic carbon (HTCO)	Rosette	Matsuoka A. / Benner R. / Ferland J.	Available
2016	Dissolved organic matter (Amino acids)	Rosette	Matsuoka A. / Benner R. / Ferland J.	Available
2016	Dissolved organic matter (sugars)	Rosette	Panagiotopoulos C./ R Sempéré	Available
2016	Dissolved organic nitrogen (release)	Camp water sample	Raimbault P.	Available
2016	Downwelling irradiance	Surface mode	Belanger S. / Goyens C. / Lambert Girard S.	Available
2016	Downwelling irradiance	Surface mode	Lambert-Girard S. / Leymarie E.	Available
2016	Downwelling Irradiance above the surface ($E_d(0^+)$)	Surface mode	Babin M. / Galí M.	Available
2016	Downwelling Irradiance above the surface ($E_d(0^+)$)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	Downwelling Irradiance ($E_d(z)$)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	$E_d(0^+)$ spectra from SBDART radiative transfer simulations	Surface mode	Babin M. / Galí M.	Available
2016	Faecal pellets flux	Sediment Trap	Fortier L. / Lalande C.	Available
2016	Hemispherical directional reflectance distribution function	Surface mode	Belanger S. / Goyens C. / Lambert-Girard S.	Data not available yet
2016	Heterotrophic bacteria (abundance)	Camp water sample	Vaulot D.	Available
2016	Hydro SCAMP (temperature, salinity, chlorophyll, turbidity, etc.)	In-water profiler	Vladoiu A. / Dumont D. / Sévigny C.	Available
2016	Ice and snow temperature	Meteorological Tower	Massé G.	Data not available yet
2016	Ice thickness	Camp ice sample	Galindo V. / Rysgaard S.	Available
2016	Irradiance (downwelling)	Surface-, Ice Bottom-mode	Matthes L. / Ehn J. / Lambert-Girard S./ Mundy C.J.	Available
2016	Irradiance (downwelling)	Under-ice irradiance transects, ROV	Matthes L. /Lambert-Girard S./ Ehn J./Mundy C.J	Available
2016	Irradiance (downwelling, upwelling)	Surface-, Under-water profile-mode	Matthes L. / Ehn J. / Lambert-Girard S./ Mundy C.J.	Available
2016	Isoprenoid lipids	Camp water sample	Massé G. / Guilmette C.	Data not available yet
2016	Isoprenoid lipids	Sea ice core	Massé G. / Guilmette C.	Data not available yet
2016	Lipid biomarkers	Collected organisms	Dufour F. / Massé G. / Ayotte P. / Lemire M.	Data not available yet
2016	Lipid tracers of bacteria stress	Camp water sample	Rontani J.-F. / Amiraux R.	Data not available yet
2016	Lipid tracers of bacteria stress	Sea ice core	Rontani J.-F. / Amiraux R.	Data not available yet
2016	Nitrate (NO_3^-)	Sediment Trap	Rontani J.-F. / Amiraux R.	Data not available yet
2016	Nitrate (NO_3^-)	Camp water sample	Raimbault P.	Available
2016	Nitrate (NO_3^- , assimilation)	Sea ice core	Raimbault P.	Available
2016	Nitrate (NO_3^- , assimilation)	Camp water sample	Raimbault P.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (*continued*)

Year	Parameter	Sampling method	Principal investigators	Processed
2016	Nitrification	Camp water sample	Raimbault P.	Available
2016	Nitrite (NO_2^-)	Camp water sample	Raimbault P.	Available
2016	Nitrite (NO_2^-)	Sea ice core	Raimbault P.	Available
2016	Nutrients bioassay	Experiment	Delaforge A / Mundy Cj	Data not available yet
2016	Nutrients bioassay	Experiment	Galindo V. / Rysgaard S.	Data not available yet
2016	PAR from SBDART radiative transfer simulations	Surface mode	Babin M. / Galí M.	Available
2016	Particle Size Distribution	In-water IOP profiler	Becu G. / Babin M.	Data not available yet
2016	Particle Size Distribution	In-water profiler	L. Stemmann / Lagunas J.	Data not available yet
2016	Particles size	Underwater Vision Profiler (UVP)	Lagunas J. / Picheral M.	Available
2016	Particulate Carbon (PC)	Camp water sample	Babin M. / Ferland J.	Available
2016	Particulate mass	Sediment Trap	Fortier L. / Lalonde C.	Available
2016	Particulate Nitrogen (PN)	Camp water sample	Babin M. / Ferland J.	Available
2016	Particulate nitrogen (PN)	Sediment Trap	Fortier L. / Lalonde C.	Available
2016	Particulate organic carbon (POC)	Sediment Trap	Fortier L. / Lalonde C.	Available
2016	Particulate organic carbon (POC)	Camp water sample	Raimbault P.	Available
2016	Particulate organic nitrogen (PON)	Camp water sample	Raimbault P.	Available
2016	Particulate Organic Phosphorus (POP)	Camp water sample	Raimbault P.	Data not available yet
2016	PDMPO uptake	Camp water sample	Leblanc K. / Queguiner B.	Data not available yet
2016	PDMPO uptake per species	Camp water sample	Leblanc K. / Queguiner B.	Data not available yet
2016	Phosphate ($(\text{PO}_4)^{3-}$)	Camp water sample	Raimbault P.	Available
2016	Phosphate ($(\text{PO}_4)^{3-}$)	Sea ice core	Raimbault P.	Available
2016	Photosynthetically available radiation (PAR)	Surface mode	Babin M. / Galí M.	Available
2016	Photosynthetically available radiation (PAR)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	Photosynthetic eukaryotes (morphology)	Camp water sample	Vaulot D.	Available
2016	Photosynthetic nanoeukaryotes (abundance)	Camp water sample	Vaulot D.	Available
2016	Photosynthetic parameters	Camp water sample	Ferland J. / Babin M.	Available
2016	Photosynthetic parameters (variable fluorescence)	Camp water sample	Lavaud J. / Galindo V. / Rysgaard S.	Data not available yet
2016	Photosynthetic parameters (variable fluorescence)	Sediment Trap	Lavaud J. / Galindo V. / Rysgaard S.	Data not available yet
2016	Photosynthetic parameters (variable fluorescence)	Sea ice core	Lavaud J. / Galindo V. / Rysgaard S.	Data not available yet
2016	Photosynthetic picoeukaryotes (abundance)	Camp water sample	Vaulot D.	Available
2016	Phytoplankton	Camp water sample	Ferland J. / Grondin P.L. / Babin M.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (continued)

Year	Parameter	Sampling method	Principal investigators	Processed
2016	Phytoplankton (taxonomy)	Sediment Trap	Fortier L. / Lalande C.	Available
2016	Pigments	Camp water sample	Ras J. / Claustre H./Galindo V. / Rysgaard S.	Available
2016	Primary production	Camp water sample	Raimbault P.	Available
2016	Prokaryotic diversity	Camp water sample	Joux F.	Data not available yet
2016	Prokaryotic diversity	Sea ice core	Joux F.	Data not available yet
2016	Rrs (0°)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	Salinity	Sea ice core	Galindo V. / Rysgaard S.	Available
2016	Scattering Coefficient	In-water IOP profiler	Becu G. / Babin M.	Available
2016	Sea ice concentration	Surface mode	Massicotte P.	Available
2016	Selenium	Collected organisms	Dufour F., Massé G., Ayotte P., Lemire M.	Available
2016	Silica Biogenic (BSi)	Camp water sample	Leynaert A./Moriceau B./ Leblanc K./Queguiner B.	Available
2016	Silica Biogenic (BSi) dissolution rate	Camp water sample	Moriceau B.	Available
2016	Silica Lithogenic (LSi)	Camp water sample	Leynaert A./Moriceau B./ Leblanc K./Queguiner B.	Data not available yet
2016	Silicate $Si(OH)_4$ - absorption kinetics	Camp water sample	Leynaert A.	Available
2016	Silica (uptake rate)	Camp water sample	Leynaert A.	Available
2016	$Si(OH)_4$	Camp water sample	Leynaert A. / Moriceau B.	Available
2016	$Si(OH)_4$	Camp water sample	Raimbault P.	Available
2016	$Si(OH)_4$	Sea ice core	Raimbault P.	Available
2016	Snow depth	Camp snow sample	Galindo V./Rysgaard S.	Available
2016	Spectral downwelling radiance angular distribution	Under-water sensor	Lambert-Girard S. / Leymarie E.	Available
2016	Spectral transmittance through ice	Surface mode	Verin G./Picard. G.	Available
2016	Surface spectral albedo	Surface mode	Verin G./Picard. G.	Available
2016	Suspended particulate material (SPM)	Camp water sample	Babin M. / Ferland J.	Available
2016	Swimmers	Sediment Trap	Fortier L. / Lalande C.	Available
2016	Synechococcus (abundance)	Camp water sample	Vaulot D.	Available
2016	Temperature	Sea ice core	Galindo V. / Rysgaard S.	Available
2016	Total organic carbon (TOC)	Camp water sample	Raimbault P.	Available
2016	Total organic carbon (TOC) and dissolved organic carbon (DOC)	Rosette	Panagiotopoulos C./ Sempéré R.	Available
2016	Total organic nitrogen (TON)	Camp water sample	Raimbault P.	Available
2016	Total organic phosphorus (TOP)	Camp water sample	Raimbault P.	Available
2016	Under-ice export fluxes of biogenic matter (fresh)	Sediment Trap	Fortier L. / Lalande C.	Available
2016	Upwelling Irradiance ($E_u(z)$)	Optical radiometers profiling system	Becu G. / Babin M.	Available

Table 2: Parameters measured during the Green Edge ice camp surveys. Parameters are ordered by alphabetical order and sampling year. (*continued*)

Year	Parameter	Sampling method	Principal investigators	Processed
2016	Upwelling radiance ($L_u(z)$)	Surface mode	Belanger S. / Goyens C. / Lambert-Girard S.	Data not available yet
2016	Upwelling radiance ($L_u(z)$)	Optical radiometers profiling system	Becu G. / Babin M.	Available
2016	Vertical profile of snow density	Surface mode	Verin G./Picard. G.	Available
2016	Vertical profile of Specific Surface Area	Surface mode	Verin G./Picard. G.	Available
2016	Virus (abundance)	Camp water sample	Joux F.	Available
2016	Wind Direction	Meteorological Tower	Massé G.	Available
2016	Wind Speed	Meteorological Tower	Massé G.	Available
2016	Zooplankton (Abundances)	Plankton net	Fortier L. / Aubry C	Available
2016	Zooplankton fecal pellet production rate	Plankton net	Fortier L. / Sampei M	Available
2016	Zooplankton grazing rate	Plankton net	Fortier L. / Sampei M	Available
2016	Zooplankton (Taxonomy)	Plankton Net	Fortier L. / Aubry C	Available
2016	Zooplankton vertical distribution	Underwater Vision Profiler (UVP)	Lagunas J. / Picheral M.	Available




Figure A1. Surface tidal height versus time at Qikiqtarjuaq measured on 2015-06-09.

Appendix B: GoPro Hero 4 photos



Figure B1. Video frame (00:58) from GoPro Hero 4 recording of C-OPS descent from 0 to 30 m, 18 May 2015 at the "low snow" hole. Note the streaks of nekton swimming across the upper left quadrant of the frame. Many planktons were seen in this profile, indicating an active under-ice community. A profile of the "high snow" hole on the same day, just 40 m away, showed no such plankton activity.

Table B1. Examples of GoPro Hero 4 photos at the low and high snow holes in 2015 demonstrating the spatial variability of the ice bottom across time and space.

	18 May 2015	31 May 2015	12 June 2015
Low snow			
High snow			

325 *Author contributions.* Ghislain Picard designed the snow optical measurements and participated in the 2015 campaign along with Gauthier Verin who sampled the 2015 and 2016 snow-related measurements. Anda Vladouli, Caroline Sevigny and Dany Dumont deployed and Marie-Noëlle Houssais added her contribution to the analysis of the Self-Contained Autonomous MicroProfiler (SCAMP) on 23 June 2016 and quality-controlled, processed, analyzed and interpreted the data. Guislain Becu, Claudie Marec performed the setup and deployment of the CTD inside the tent in 2015. CTD setup and deployment was performed by José 330 Lagunas, Christiane Dufresne, in 2016. Guislain Becu, Griet Neukermans, Eric Rehm, Simon Lambert-Girard and Laurent Oziel, Jade Larivière, Joannie Ferland, Julien Laliberté, performed the setup, calibration, and deployments of the ICE-Pro optical profiler outside the tent and the IOP frame inside the tent. Eric Rehm performed the 13-h tidal cycle measurements in 2015. Griet Neukermans and Eric Rehm deployed the GoPro Hero on the ICE-Pro. Claudie Marec performed the setup and installation of IFCB in the lab in 2015. Joannie Ferland performed the setup and installation of the IFCB in the lab in 2016. Joannie Ferland, 335 Erin Reimer, Atsushi Matsuoka, Marie-Hélène Forget and Pierre-Luc Grondin performed the measurements. Pierre-Luc Grondin analyzed the data. Claudie Marec and José Lagunas performed the setup and deployment of an In-water profiler for particle size distribution and zooplankton vertical distribution (UVP Underwater Visio Profiler). Claudie Marec and José Lagunas performed setup and water sampling in both 2015 and 2016 campaigns. Claudie Marec was involved in the design and deployment of the ADCP in 2015, José Lagunas deployed the instrument in 2016. Atsushi Matsuoka coordinated the sampling strategy of discrete 340 waters in terms of examining the linkages between optical and organic matter properties. Atsushi Matsuoka and Annick Bricaud wrote the protocols for both CDOM and particulate absorptions. For aCDOM, Atsushi Matsuoka, Joannie Ferland, Marie-Hélène Forget, Erin Reimer, and Pierre-Luc Grondin contributed to the measurements. For ap, Atsushi Matsuoka, Céline Dimier, Léo Lacour, Joséphine Ras, Mathieu Ardyna, Henry Bittig, Blanche St-Béat and Thomas Lacour contributed to the measurements. In 2015, particulate spectral absorption was also done by Lisa Matthes, Christine Quiring and Jens Ehn. Nicole Pogorzelec (who 345 also did snow and ice salinity and overall chl-a filtrations in the field lab). Marie-Pier Amyot worked on tidying and uniformizing the data. Martí Galí ran the radiative transfer calculations and compared them to irradiance measurements taken on the ice camps. Lisa Matthes, Simon Lambert-Girard, Bob Hodgson, Jens Ehn, Nicole Pogorzelec and CJ Mundy designed and/or carried out the TriOS and ROV under-ice irradiance measurements Christos Panagiotopoulos and Richard Sempéré coordinated the sampling strategy for sugars/DOC and the analyses. Remi Amiraux collected the samples. Between October 2014 and July 2016, 350 Éric Brossier and France Pinczon du Sel conducted measurements, collected clams, maintained equipment, kept a time-lapse photography record and represented the Greenedge team in Qikiqtarjuaq outside of the sampling season. Debra Christiansen Stowe coordinated logistics in Qikiqtarjuaq, in support of the 2016 ice camp. Makoto Sampei designed and curried copepods incubations to collect fecal pellets out at the ice camp in 2016. Makoto Sampei made microscopic observations on the collected fecal pellets in the laboratory. Sea ice and snow hemispherical directional reflectance were measured on the ice camp in 2015 355 by Sabine Marty and Clémence Goyens. The set-up was designed by Sabine Marty, Edouard Leymarie, Simon Bélanger and Clémence Goyens. They also processed and analyzed the data.

Competing interests. The authos declar no competing interests.

Acknowledgements. The GreenEdge project is funded by the following French and Canadian programs and agencies: ANR (Contract #111112), CNES (project #131425), IPEV (project #1164), CSA, Fondation Total, ArcticNet, LEFE and the French Arctic Initiative 360 (GreenEdge project). This project would not have been possible without the support of the Hamlet of Qikiqtarjuaq and the members of the community as well as the Inuksuit School and its Principal Jacqueline Arsenault. The project was conducted under the scientific coordination of the Canada Excellence Research Chair in Remote Sensing of Canada's new Arctic frontier and the CNRS & Université Laval Takuviq Joint International laboratory (UMI3376). The field campaign was successful thanks to the contribution of A. Wells, M. Benoît-Gagné, and E. Devred from the Takuviq laboratory as well as R. Hodgson from the University of 365 Manitoba. Pascale Bouruet-Aubertot and Yannis Cuypers who provided the SCAMP and contributed to the processing, quality control, analysis and interpretation of the data. We also thank Michel Gosselin, Québec-Océan, the CCGS Amundsen and the Polar Continental Shelf Program for their in-kind contribution to the logistic and scientific equipment. Thanks to Etienne Ouellet for IT support and data infrastructure management.

References

- 370 Aminot, A. and Kérouel, R.: Dosage automatique des nutriments dans les eaux marines : méthodes en flux continu., 2007.
- Ardyna, M., Babin, M., Gosselin, M., Devred, E., Bélanger, S., Matsuoka, A., and Tremblay, J.-É.: Parameterization of vertical chlorophyll <i>a</i> in the Arctic Ocean: impact of the subsurface chlorophyll maximum on regional, seasonal, and annual primary production estimates, *Biogeosciences*, 10, 4383–4404, <https://doi.org/10.5194/bg-10-4383-2013>, <https://www.biogeosciences.net/10/4383/2013/>, 2013.
- 375 Arrigo, K. R., Perovich, D. K., Pickart, R. S., Brown, Z. W., van Dijken, G. L., Lowry, K. E., Mills, M. M., Palmer, M. A., Balch, W. M., Bahr, F., Bates, N. R., Benitez-Nelson, C., Bowler, B., Brownlee, E., Ehn, J. K., Frey, K. E., Garley, R., Laney, S. R., Lubelczyk, L., Mathis, J., Matsuoka, A., Mitchell, B. G., Moore, G. W. K., Ortega-Retuerta, E., Pal, S., Polashenski, C. M., Reynolds, R. A., Schieber, B., Sosik, H. M., Stephens, M., and Swift, J. H.: Massive Phytoplankton Blooms Under Arctic Sea Ice, *Science* (80-), 336, 1408–1408, <https://doi.org/10.1126/science.1215065>, <http://www.sciencemag.org/cgi/doi/10.1126/science.1215065>, 2012.
- 380 Arrigo, K. R., Perovich, D. K., Pickart, R. S., Brown, Z. W., van Dijken, G. L., Lowry, K. E., Mills, M. M., Palmer, M. A., Balch, W. M., Bates, N. R., Benitez-Nelson, C. R., Brownlee, E., Frey, K. E., Laney, S. R., Mathis, J., Matsuoka, A., Greg Mitchell, B., Moore, G., Reynolds, R. A., Sosik, H. M., and Swift, J. H.: Phytoplankton blooms beneath the sea ice in the Chukchi sea, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 105, 1–16, <https://doi.org/10.1016/j.dsr2.2014.03.018>, <http://dx.doi.org/10.1016/j.dsr2.2014.03.018><http://linkinghub.elsevier.com/retrieve/pii/S0967064514000836><https://linkinghub.elsevier.com/retrieve/pii/S0967064514000836>, 2014.
- 385 Assmy, P., Fernández-Méndez, M., Duarte, P., Meyer, A., Randelhoff, A., Mundy, C. J., Olsen, L. M., Kauko, H. M., Bailey, A., Chierici, M., Cohen, L., Doulgeris, A. P., Ehn, J. K., Fransson, A., Gerland, S., Hop, H., Hudson, S. R., Hughes, N., Itkin, P., Johnsen, G., King, J. A., Koch, B. P., Koenig, Z., Kwasniewski, S., Laney, S. R., Nicolaus, M., Pavlov, A. K., Polashenski, C. M., Provost, C., Rösel, A., Sandbu, M., Spreen, G., Smedsrød, L. H., Sundfjord, A., Taskjelle, T., Tatarek, A., Wiktor, J., Wagner, P. M., Wold, A., Steen, H., and Granskog, M. A.: Leads in Arctic pack ice enable early phytoplankton blooms below snow-covered sea ice, *Sci. Rep.*, 7, 40 850, <https://doi.org/10.1038/srep40850>, <http://www.nature.com/articles/srep40850>, 2017.
- Bouman, H. A., Platt, T., Doblin, M., Figueiras, F. G., Gudmundsson, K., Gudfinnsson, H. G., Huang, B., Hickman, A., Hiscock, M., Jackson, T., Lutz, V. A., Mélin, F., Rey, F., Pepin, P., Segura, V., Tilstone, G. H., van Dongen-Vogels, V., and Sathyendranath, S.: Photosynthesis-irradiance parameters of marine phytoplankton: synthesis of a global data set, *Earth Syst. Sci. Data*, 10, 251–266, <https://doi.org/10.5194/essd-10-251-2018>, <https://www.earth-syst-sci-data.net/10/251/2018/>, 2018.
- 395 Fortier, M., Fortier, L., Michel, C., and Legendre, L.: Climatic and biological forcing of the vertical flux of biogenic particles under seasonal Arctic sea ice, *Mar. Ecol. Prog. Ser.*, 225, 1–16, <https://doi.org/10.3354/meps225001>, <http://www.int-res.com/abstracts/meps/v225/p1-16/>, 2002.
- Galindo, V., Levasseur, M., Mundy, C. J., Gosselin, M., Tremblay, J.-É., Scarratt, M., Gratton, Y., Papakiriakou, T., Poulin, M., and Lizotte, M.: Biological and physical processes influencing sea ice, under-ice algae, and dimethylsulfoniopropionate during spring in the Canadian Arctic Archipelago, *J. Geophys. Res. Ocean.*, 119, 3746–3766, <https://doi.org/10.1002/2013JC009497>, <http://doi.wiley.com/10.1002/2013JC009497>, 2014.
- 400 Galindo, V., Gosselin, M., Lavaud, J., Mundy, C., Else, B., Ehn, J., Babin, M., and Rysgaard, S.: Pigment composition and photoprotection of Arctic sea ice algae during spring, *Mar. Ecol. Prog. Ser.*, 585, 49–69, <https://doi.org/10.3354/meps12398>, <http://www.int-res.com/abstracts/meps/v585/p49-69/>, 2017.

- Gosselin, M., Levasseur, M., Wheeler, P. A., Horner, R. A., and Booth, B. C.: New measurements of phytoplankton and ice algal production in the Arctic Ocean, Deep Sea Res. Part II Top. Stud. Oceanogr., 44, 1623–1644, [https://doi.org/10.1016/S0967-0645\(97\)00054-4](https://doi.org/10.1016/S0967-0645(97)00054-4), <http://linkinghub.elsevier.com/retrieve/pii/S0967064597000544>, 1997.
- Goyens, C., Marty, S., Leymarie, E., Antoine, D., Babin, M., and Bélanger, S.: High Angular Resolution Measurements of the Anisotropy of Reflectance of Sea Ice and Snow, Earth Sp. Sci., 5, 30–47, <https://doi.org/10.1002/2017EA000332>, <http://doi.wiley.com/10.1002/2017EA000332>, 2018.
- Grondin, P.-L., Ferland, J., Massicotte, P., Galindo, V., Ras, J., Claustre, H., Garcia, N., Raimbault, P., Coupel, P., Tremblay, J.-É., Forget, M.-H., Karp-Boss, L., and Babin, M.: Protists succession from ice algae to under-ice phytoplankton spring blooms in Baffin Bay, in prep, 2019.
- Laliberté, J., Bélanger, S., and Frouin, R.: Evaluation of satellite-based algorithms to estimate photosynthetically available radiation (PAR) reaching the ocean surface at high northern latitudes, Remote Sens. Environ., 184, 199–211, <https://doi.org/10.1016/j.rse.2016.06.014>, <http://linkinghub.elsevier.com/retrieve/pii/S0034425716302528>, <http://dx.doi.org/10.1016/j.rse.2016.06.014>, 2016.
- Laney, S. R. and Sosik, H. M.: Phytoplankton assemblage structure in and around a massive under-ice bloom in the Chukchi Sea, Deep Sea Res. Part II Top. Stud. Oceanogr., 105, 30–41, <https://doi.org/10.1016/j.dsr2.2014.03.012>, <http://linkinghub.elsevier.com/retrieve/pii/S0967064514000770>, 2014.
- Letelier, R. M., Karl, D. M., Abbott, M. R., and Bidigare, R. R.: Light driven seasonal patterns of chlorophyll and nitrate in the lower euphotic zone of the North Pacific Subtropical Gyre, Limnol. Oceanogr., 49, 508–519, <https://doi.org/10.4319/lo.2004.49.2.0508>, <http://doi.wiley.com/10.4319/lo.2004.49.2.0508>, 2004.
- Leu, E., Mundy, C., Assmy, P., Campbell, K., Gabrielsen, T., Gosselin, M., Juul-Pedersen, T., and Gradinger, R.: Arctic spring awakening – Steering principles behind the phenology of vernal ice algal blooms, Prog. Oceanogr., 139, 151–170, <https://doi.org/10.1016/j.pocean.2015.07.012>, <http://linkinghub.elsevier.com/retrieve/pii/S0079661115001640>, 2015.
- Marie, D., Partensky, F., Jacquet, S., and Vaulot, D.: Enumeration and Cell Cycle Analysis of Natural Populations of Marine Picoplankton by Flow Cytometry Using the Nucleic Acid Stain SYBR Green I., Appl. Environ. Microbiol., 63, 186–193, <http://www.ncbi.nlm.nih.gov/pubmed/16535483>, http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_type=Abstract&term=PMC1389098, 1997.
- Marie, D., Rigaut-Jalabert, F., and Vaulot, D.: An improved protocol for flow cytometry analysis of phytoplankton cultures and natural samples, Cytom. Part A, 85, 962–968, <https://doi.org/10.1002/cyto.a.22517>, <http://doi.wiley.com/10.1002/cyto.a.22517>, 2014.
- Massicotte, P., Bécu, G., Lambert-Girard, S., Leymarie, E., and Babin, M.: Estimating Underwater Light Regime under Spatially Heterogeneous Sea Ice in the Arctic, Appl. Sci., 8, 2693, <https://doi.org/10.3390/app8122693>, <http://www.mdpi.com/2076-3417/8/12/2693>, 2018.
- Massicotte, P., Amiraux, R., Amyot, M.-P., Archambault, P., Ardyna, M., Arnaud, L., Artigue, L., Aubry, C., Ayotte, P., Bécu, G., Bélanger, S., Benner, R., Bittig, H. C., Bricaud, A., Brossier, É., Bruylants, F., Chauvaud, L., Christiansen-Stowe, D., Claustre, H., Cornet-Barthaux, V., Coupel, P., Cox, C., Delaforge, A., Dezutter, T., Dimier, C., Dominé, F., Dufour, F., Dufresne, C., Dumont, D., Ehn, J., Else, B., Ferland, J., Forget, M.-H., Fortier, L., Galí, M., Galindo, V., Gallinari, M., Garcia, N., Gérakas-Ribeiro, C., Gourdal, M., Gourvil, P., Goyens, C., Grondin, P.-L., Guillot, P., Guillette, C., Houssais, M.-N., Joux, F., Lacour, L., Lacour, T., Lafond, A., Lagunas, J., Lalande, C., Laliberté, J., Lambert-Girard, S., Larivière, J., Lavaud, J., Lebaron, A., Leblanc, K., Le Gall, F., Legras, J., Lemire, M., Levasseur, M., Leymarie, E., Leynaert, A., Lopes Dos Santos, A., Lourenço, A., Mah, D., Marec, C.,

- 445 Marie, D., Martin, N., Marty, C., Marty, S., Massé, G., Matsuoka, A., Matthes, L., Moriceau, B., Muller, P.-E., Mundy, C. J., Neukermans, G., Oziel, L., Panagiotopoulos, C., Pangazi, J.-J., Picard, G., Picheral, M., Pinczon Du Sel, F., Pogorzelec, N., Probert, I., Queguiner, B., Raimbault, P., Ras, J., Rehm, E., Reimer, E., Rontani, J.-F., Rysgaard, S., Saint-Béat, B., Sampei, M., Sansoulet, J., Schmidt, S., Sempéré, R., Sévigny, C., Shen, Y., Tragin, M., Tremblay, J.-É., Vaulot, D., Verin, G., Vivier, F., Vladoiu, A., Whitehead, J., and Babin, M.: The Green Edge initiative: understanding the processes controlling the under-ice Arctic phytoplankton spring bloom, <https://doi.org/10.17882/59892>, <https://www.seanoe.org/data/00487/59892/>, 2019a.
- 450 Massicotte, P., Peeken, I., Katlein, C., Flores, H., Huot, Y., Castellani, G., Arndt, S., Lange, B. A., Tremblay, J.-É., and Babin, M.: Sensitivity of Phytoplankton Primary Production Estimates to Available Irradiance Under Heterogeneous Sea Ice Conditions, *J. Geophys. Res. Ocean.*, p. 2019JC015007, <https://doi.org/10.1029/2019JC015007>, <http://doi.wiley.com/10.1029/2019JC015007><https://onlinelibrary.wiley.com/doi/abs/10.1029/2019JC015007>, 2019b.
- 455 Matthes, L. C., Ehn, J. K., L.-Girard, S., Pogorzelec, N. M., Babin, M., and Mundy, C. J.: Average cosine coefficient and spectral distribution of the light field under sea ice: Implications for primary production, *Elem Sci Anth*, 7, 25, <https://doi.org/10.1525/elementa.363>, <https://www.elementascience.org/article/10.1525/elementa.363/>, 2019.
- Menden-Deuer, S. and Lessard, E. J.: Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton, *Limnol. Oceanogr.*, 45, 569–579, <https://doi.org/10.4319/lo.2000.45.3.0569>, <http://doi.wiley.com/10.4319/lo.2000.45.3.0569>, 2000.
- 460 Moberg, E. A. and Sosik, H. M.: Distance maps to estimate cell volume from two-dimensional plankton images, *Limnol. Oceanogr. Methods*, 10, 278–288, <https://doi.org/10.4319/lom.2012.10.278>, <http://doi.wiley.com/10.4319/lom.2012.10.278>, 2012.
- Mundy, C., Gosselin, M., Gratton, Y., Brown, K., Galindo, V., Campbell, K., Levasseur, M., Barber, D., Papakyriakou, T., and Bélanger, S.: Role of environmental factors on phytoplankton bloom initiation under landfast sea ice in Resolute Passage, Canada, *Mar. Ecol. Prog. Ser.*, 497, 39–49, <https://doi.org/10.3354/meps10587>, <http://www.int-res.com/abstracts/meps/v497/p39-49/>, 2014.
- 465 Mundy, C. J., Gosselin, M., Ehn, J., Gratton, Y., Rossnagel, A., Barber, D. G., Martin, J., Tremblay, J.-É., Palmer, M., Arrigo, K. R., Darnis, G., Fortier, L., Else, B., and Papakyriakou, T.: Contribution of under-ice primary production to an ice-edge upwelling phytoplankton bloom in the Canadian Beaufort Sea, *Geophys. Res. Lett.*, 36, L17 601, <https://doi.org/10.1029/2009GL038837>, <http://doi.wiley.com/10.1029/2009GL038837>, 2009.
- Olson, R. J. and Sosik, H. M.: A submersible imaging-in-flow instrument to analyze nano-and microplankton: Imaging FlowCytobot, *Limnol. Oceanogr. Methods*, 5, 195–203, <https://doi.org/10.4319/lom.2007.5.195>, <http://doi.wiley.com/10.4319/lom.2007.5.195>, 2007.
- 470 Oziel, L., Massicotte, P., Randelhoff, A., Ferland, J., Vladoiu, A., Lacour, L., Galindo, V., Lambert-Girard, S., Dumont, D., Cuypers, Y., Bouruet-Aubertot, P., Mundy, C.-J., Ehn, J., Béchu, G., Marec, C., Forget, M.-H., Garcia, N., Coupel, P., Raimbault, P., Houssais, M.-N., and Babin, M.: Environmental factors influencing the seasonal dynamics of spring algal blooms in and beneath sea ice in western Baffin Bay, *Elem Sci Anth*, 7, 34, <https://doi.org/10.1525/elementa.372>, <https://www.elementascience.org/article/10.1525/elementa.372/>, 2019.
- Perrette, M., Yool, A., Quartly, G. D., and Popova, E. E.: Near-ubiquity of ice-edge blooms in the Arctic, *Biogeosciences*, 8, 515–524, <https://doi.org/10.5194/bg-8-515-2011>, <http://www.biogeosciences.net/8/515/2011/>, 2011.
- Picheral, M., Colin, S., and Irisson, J.-O.: EcoTaxa, a tool for the taxonomic classification of images., <http://ecotaxa.obs-vlfr.fr>, 2017.
- 480 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.r-project.org/>, 2019.

Randelhoff, A., Oziel, L., Massicotte, P., Bécu, G., Galí, M., Lacour, L., Dumont, D., Vladou, A., Marec, C., Bruyant, F., Houssais, M.-N., Tremblay, J.-É., Deslongchamps, G., and Babin, M.: The evolution of light and vertical mixing across a phytoplankton ice-edge bloom, *Elem Sci Anth*, 7, 20, <https://doi.org/10.1525/elementa.357>, <https://www.elementascience.org/article/10.1525/elementa.357/>, 2019.

485 Ricchiazzi, P., Yang, S. R., Gautier, C., and Sowle, D.: SBDART: A research and teaching software tool for plane-parallel radiative transfer in the Earth's atmosphere, *Bull. Am. Meteorol. Soc.*, 79, 2101–2114 ST – SBDART: A research and teaching so, 1998.

Sakshaug, E.: Primary and Secondary Production in the Arctic Seas, in: Org. Carbon Cycle Arct. Ocean, pp. 57–81, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-18912-8_3, http://link.springer.com/10.1007/978-3-642-18912-8_3, 2004.

490 Sosik, H. M. and Olson, R. J.: Automated taxonomic classification of phytoplankton sampled with imaging-in-flow cytometry, *Limnol. Oceanogr. Methods*, 5, 204–216, <https://doi.org/10.4319/lom.2007.5.204>, <http://doi.wiley.com/10.4319/lom.2007.5.204>, 2007.

495 Suggett, D. J., Moore, C. M., Geider, R. J., Perkins, R. G., Kromkamp, J. C., Serôdio, J., Lavaud, J., Jesus, B., Mouget, J. L., Lefebvre, S., Forster, R. M., Suggett, D. J., Moore, C. M., Geider, R. J., Perkins, R. G., Kromkamp, J. C., Serôdio, J., Lavaud, J., Jesus, B., Mouget, J. L., Lefebvre, S., and Forster, R. M.: Chlorophyll a Fluorescence in Aquatic Sciences: Methods and Applications, Springer Netherlands, Dordrecht, <https://doi.org/10.1007/978-90-481-9268-7>, <http://link.springer.com/10.1007/978-90-481-9268-7>, 2010.

500 Tremblay, J.-É. and Gagnon, J.: The effects of irradiance and nutrient supply on the productivity of Arctic waters: a perspective on climate change, in: *Influ. Clim. Chang. Chang. Arct. Sub-Arctic Cond.*, pp. 73–93, Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-1-4020-9460-6_7, http://link.springer.com/10.1007/978-1-4020-9460-6_7, 2009.

Wassmann, P., Ratkova, T., Andreassen, I., Vernet, M., Pedersen, G., and Rey, F.: Spring Bloom Development in the Marginal Ice Zone and the Central Barents Sea, *Mar. Ecol.*, 20, 321–346, <https://doi.org/10.1046/j.1439-0485.1999.2034081.x>, <http://doi.wiley.com/10.1046/j.1439-0485.1999.2034081.x>, 1999.