

Conformal Inference Assignment 2

(Deadline: October 18)

Wen, Zehai

Temps de compilation : 2026-10-10 19:24:46-04:00

Remark All references are to our course textbook "Theoretical Foundations of Conformal Prediction" by Angelopoulos et al. (2026).

Exercise 1

Consider split conformal prediction with $\mathcal{Y} = \mathbb{R}$ and a given model $\hat{f} : \mathcal{X} \rightarrow \mathbb{R}$. Construct a score function $s(x, y)$ that would yield intervals of the form $\mathcal{C}(x) = [\hat{f}(x) - \gamma, \hat{f}(x) + 2\gamma]$ for some $\gamma > 0$.

Solution Put

$$s(x, y) = \begin{cases} 2|\hat{f}(x) - y| & y < \hat{f}(x) \\ |\hat{f}(x) - y| & y \geq \hat{f}(x). \end{cases}$$

Then, with $\hat{q}$ being a quantile of the scores on the validation set and $\mathcal{C}(x) = \{y \in \mathcal{Y} \mid s(x, y) \leq \hat{q}\}$, we will have the desired interval with $\gamma = \frac{\hat{q}}{2}$. ///

Exercise 2

Consider split conformal prediction with $\mathcal{Y} = \mathbb{R}^d$ and a given model $\hat{f} : \mathcal{X} \rightarrow \mathbb{R}^d$ whose coordinates are $\hat{f}(x) = (\hat{f}_1(x), \dots, \hat{f}_d(x))$. Construct a score function $s(x, y)$ that would yield intervals of the form

$$\mathcal{C}(x) = \prod_{j=1}^d [\hat{f}_j(x) - c, \hat{f}_j(x) + c]$$

for a $c > 0$.

Solution Put

$$s(x, y) = \max_{j=1, \dots, d} |\hat{f}_j(x) - y_j|.$$

Then, with $\hat{q}$ being a quantile of the scores on the validation set and $\mathcal{C}(x) = \{y \in \mathcal{Y} \mid s(x, y) \leq \hat{q}\}$, we will have the desired interval with $c = \hat{q}$. ///

Exercise 3

Consider split conformal prediction with $\mathcal{Y} = \mathbb{R}$, two given models $\hat{f}_i : \mathcal{X} \rightarrow \mathbb{R}$ for $i = 1, 2$, and the score function

$$s(x, y) = \min_{i=1,2} |y - \hat{f}_i(x)|.$$

Derive an explicit expression for $\mathcal{C}(x)$ in terms of $\hat{f}_1(x)$, $\hat{f}_2(x)$, and the quantile $\hat{q}$ of the scores on the validation set. What is the shape of the prediction when $\hat{q}$ is small or when it is large? Briefly explain why this might be attractive if the conditional distribution of Y given $X = x$ is bimodal.

Solution By definition, it is immediately clear that

$$\mathcal{C}(x) = \{y \in \mathcal{Y} \mid s(x, y) \leq \hat{q}\} = [\hat{f}_1(x) - \hat{q}, \hat{f}_1(x) + \hat{q}] \cup [\hat{f}_2(x) - \hat{q}, \hat{f}_2(x) + \hat{q}].$$

If $\hat{f}_1(x) = \hat{f}_2(x)$, then $\mathcal{C}(x)$ is an interval of length $2\hat{q}$ centered at $\hat{f}_1(x) = \hat{f}_2(x)$. If we assume that the two models are different at x , that is, $\hat{f}_1(x) \neq \hat{f}_2(x)$, then $\mathcal{C}(x)$ is the union of two intervals when $\hat{q}$ is small enough, and it is a single interval when $\hat{q}$ is large enough. If the conditional distribution of Y given $X = x$ is bimodal, then the score might be of interest because $\hat{f}_1(x)$ and $\hat{f}_2(x)$ might capture the two modes of the distribution and the predicted interval will be centred at both modes. ///

Exercise 4

Fix $n > 2$. Construct a random vector $(X_1, \dots, X_n)$ defined on some probability space such that (X_i, X_j) is exchangeable for all $i \neq j$ but $(X_1, \dots, X_n)$ is not exchangeable.

Solution Let (X, Y) be a jointly normal random vector. It is exchangeable if and only if $\mathbb{E}[X] = \mathbb{E}[Y]$ and $\text{Var}(X) = \text{Var}(Y)$. With this in mind, let $(X_1, \dots, X_n)$ be a jointly normal random vector such that $\mathbb{E}[X_i] = \mathbb{E}[X_j]$ and $\text{Var}(X_i) = \text{Var}(X_j)$ for all $i \neq j$ but $\text{Cov}(X_1, X_2) \neq \text{Cov}(X_1, X_3)$. Then (X_i, X_j) is exchangeable for all $i \neq j$ but $(X_1, \dots, X_n)$ is not exchangeable. For a concrete example, say $\mathbb{E}[X_i] = 0$ and $\text{Var}(X_i) = 1$ for all i , $\text{Cov}(X_1, X_2) = -1$, and all other covariances are zero. ///