

# Can Behavior Change Explain Increases in the Proportion of Genital Ulcers Attributable to Herpes in Sub-Saharan Africa?

## A Simulation Modeling Study

ELINE L. KORENROMP, PhD,\* ROEL BAKKER, PhD,\* SAKE J. DE VLAS, PhD,\* N. JAMIE ROBINSON, PhD,<sup>†</sup> RICHARD HAYES, DSc,<sup>‡</sup> AND J. DIK F. HABBEMA, PhD\*

**Background:** The proportion of cases of genital ulcer disease attributable to herpes simplex virus type 2 (HSV-2) appears to be increasing in sub-Saharan Africa.

**Goal:** To assess the contributions of HIV disease and behavioral response to the HIV epidemic to the increasing proportion of genital ulcer disease (GUD) attributable to HSV-2 in sub-Saharan Africa.

**Study Design:** Simulations of the transmission dynamics of ulcerative sexually transmitted diseases (STDs) and HIV with use of the model STDSIM.

**Results:** In simulations, 28% of GUD was caused by HSV-2 before a severe HIV epidemic. If HIV disease was assumed to double the duration and frequency of HSV-2 recurrences, this proportion rose to 35% by year 2000. If stronger effects of HIV were assumed, this proportion rose further, but because of increased HSV-2 transmission this would shift the peak in HSV-2 seroprevalence to an unrealistically young age. A simulated 25% reduction in partner-change rates increased the proportion of GUD caused by HSV-2 to 56%, following relatively large decreases in chancroid and syphilis.

**Conclusion:** Behavioral change may make an important contribution to relative increases in genital herpes.

THE DISTRIBUTION OF CAUSES of genital ulcer disease (GUD) differs widely between countries within Africa. The proportion of GUD attributable to herpes simplex virus type 2 (HSV-2) appears to be higher or increasing in countries with severe HIV epidemics.<sup>1–3</sup> For example, the proportion of GUD cases that were herpes-culture-positive increased from 11% in 1986–1988 to 21% in 1990–1992 among HIV-positive cases in Rwanda<sup>4</sup>; from 3% before

From the \*Department of Public Health, Faculty of Medicine, Erasmus University, Rotterdam, Rotterdam, The Netherlands; <sup>†</sup>GlaxoSmithKline, Greenford, United Kingdom; and <sup>‡</sup>London School of Hygiene and Tropical Medicine, London, United Kingdom

1981 to 14% in 1994 among gold miners in Johannesburg, South Africa<sup>5</sup>; and from 7% in 1984 to 40% in 1998 in Durban, South Africa.<sup>2</sup> Increases have also been reported in the occurrence of genital herpes as a proportion of new diagnoses of sexually transmitted diseases (STDs) in Durban (from 7% in 1989 to 11% in 1997) and Harare, Zimbabwe (from 6% in 1982 to 10% in 1997). Moreover, GUD appears to constitute a larger proportion of STD diagnoses in the countries of sub-Saharan Africa (SSA) with severe HIV epidemics than in West Africa.<sup>6</sup>

Several explanations have been proposed for the apparent increase in HSV-2 as a cause of GUD in SSA. Immunosuppression during advanced HIV disease can increase the duration, severity, and incidence of herpetic recurrences, leading to an increased herpes ulcer load.<sup>7,8</sup> This effect may be enhanced if the increased herpes ulcer load in HIV-infected patients increases the transmission of HSV-2. Decreases in the prevalence of bacterial causes of GUD, in particular chancroid, may induce a relative increase in HSV-2 as a cause of GUD. Bacterial GUD can decrease because of selective HIV-attributable mortality among high-risk groups<sup>9</sup> and behavior change as a result of HIV control programs promoting safer sex.<sup>10–13</sup> Improved management of bacterial STDs may also contribute to decreases in bacterial GUD.<sup>2,14–16</sup> Finally, apparent increases in genital herpes may reflect changes in detection rather than true shifts in GUD etiology.<sup>2,17</sup> Improved HSV-2 detection could result from increased awareness among clinicians and patients of herpes as a cause of GUD, as well as improved diagnostic choices such as the use of polymerase chain reaction as a complement to viral culture and clinical appearance.<sup>1</sup>

The authors thank Nico Nagelkerke and Kate Orroth for useful comments on the manuscript. We are also grateful to Frances Cowan and Richard White for advice on the HSV-2 model representation.

The study was financially supported by Glaxo Wellcome Research and Development.

Correspondence: Eline L. Korenromp, Department of Public Health, Faculty of Medicine, Erasmus University, Rotterdam, P.O. Box 1738, 3000 DR Rotterdam, The Netherlands. E-mail: korenromp@mgz.fgg.eur.nl

Received for publication June 5, 2001, revised August 14, 2001, and accepted August 17, 2001.

The data obtained to date cannot directly enable epidemiologists to distinguish between these explanations. However, their implications about the burden of disease and the control of ulcerative STD and possibly HIV in SSA may differ. This article presents simulations of the influence of these possible causes on the epidemiology of ulcerative STD. We used the stochastic microsimulation model STDSIM<sup>9,18,19</sup> to simulate the spread of ulcerative STD and HIV for a typical SSA population with a severe HIV epidemic. Comparing simulation outcomes with empirical data, we discuss the likelihood and importance of the suggested explanations.

## Methods

### Microsimulation Model STDSIM

STDSIM simulates the natural history and transmission of multiple STDs and HIV in a population of individuals with assigned characteristics that change over time. The formation and dissolution of heterosexual partnerships and transmission of STDs during contacts between sexual partners are modeled as stochastic events.<sup>9,18,19</sup> Recently the model was adapted to include the simulation of genital infection with HSV-2 and effects of HIV infection on the natural history and transmission of STDs, including HSV-2, allowing us to study the epidemiology of HSV-2 in relation to that of other ulcerative STDs and HIV disease.

### Simulated Population

Assumptions about demography, sexual behavior, and health care were chosen to reflect conditions in a typical SSA city with a severe and advanced HIV epidemic. For this, we adapted a previous model representation of a population in rural Tanzania,<sup>9</sup> by specifying a higher frequency of prostitute visits by men (25% more than in the Tanzania simulation), a higher frequency of client contacts per prostitute (an average of two per week), and an earlier time of introduction of HIV (1980). To adequately reproduce HSV-2 seroprevalence patterns observed in SSA cities,<sup>20,21</sup> we further adapted the representation of sexual behavior by specifying (1) larger age differences between sexual partners (with males being an average of 3.3 years older than their spouses) and (2) an increase in the mean age of sexual debut to 18 years for men and a corresponding decrease to 15 years for women.

### Representation of Herpes Simplex Virus Type 2

Figure 1 shows the model representation of HSV-2 infection; the corresponding parameter values are given in Table 1. Infection starts with a primary ulcer episode of an average 3 weeks' duration.<sup>22–26</sup> The incubation period is ignored because it is short<sup>23,24</sup> and unlikely to influence HSV-2 transmission dynamics.




Fig. 1. Diagrammatic representation of the natural history of genital herpes simplex virus type 2 infection in the STDSIM model. Drawing not to scale. Corresponding parameter values are given in Table 1.

After the primary episode, patients progress to a long *early latent* phase, during which ulcers can recur. In the remainder of this article, the terms *ulcer* and *recurrence* denote both recognized and unrecognized genital lesions. The frequency of recurrences is known to decrease over the first years after primary infection.<sup>25,27–33</sup> Since no studies involving long-term follow-up have been conducted, it is however unknown whether recurrences stop after a certain period or continue at a much lower frequency. Comparisons of age-profile data for clinical presentations of herpetic recurrences and GUD (which peak between ages 30 and 40 years<sup>29–32,34–38</sup>), HSV-2 seroprevalence (which peaks at an older age<sup>39,40</sup>), and HSV-2 incidence (which peaks between ages 20 and 30 years<sup>22,41</sup>) suggest that the majority of symptomatic recurrences occur within 15 to 20 years after infection.

Among clinic patients, frequencies of five to eight recurrences annually are reported as typical,<sup>29,31,32,36,38,42</sup> but studies likely oversampled individuals with relatively severe and frequent recurrences.<sup>38,43</sup> Follow-up studies of HSV-2-infected persons sampled from general populations showed recurrence frequencies of two to four annually.<sup>38,44</sup> In line with these observations, we estimated that the early latent phase during which ulcers recur lasts for an average of 15 years, during which ulcers recur at a mean interval of 6 months. To reflect the large heterogeneity between individuals,<sup>32</sup> we assumed that the duration of the early latent phase and the interval between recurrences varies randomly between episodes within individuals, according to exponential distributions. After the early latent phase, simulated patients progress into a *late latent phase*, during which they remain HSV-2 seropositive but no longer have recurrences.

Recurrences were specified to last for an average of 1 week. Because this duration appears to be relatively homo-

TABLE 1. STDSIM Representation of Natural History and Transmission of Ulcerative Sexually Transmitted Disease

Disease, Stage	Mean Duration*	Transmission Probability <sup>†</sup>		Cofactor Effect on HIV Transmission <sup>†‡</sup>	% of Ulcers That Are Recognized	
		F	F→M		M	F
HSV-2						
Primary ulcer	3 weeks	0.30	0.15	25×	30	30
Early latent	15 years <sup>§  </sup>	0.005 <sup>  </sup>	0.0025 <sup>  </sup>	NA <sup>  </sup>	NA <sup>  </sup>	NA <sup>  </sup>
	1 week <sup>§</sup> ; interval between ulcers: 6 months <sup>§</sup>					
Recurrent ulcer		0.20	0.10	10×	15	7.5
Late latent	Lifelong	0	0	NA	NA	NA
Syphilis						
Infectious	6 months	0.30	0.20	10×	80	50
Latent	15 years	0	0	NA	NA	NA
Chancroid	10 weeks	0.20	0.15	25×	90	70
HIV infection						
Primary	10 weeks	0.045	0.015	NA	NA	NA
Asymptomatic	3 or 5 years <sup>  </sup>	0.00225	0.00075	NA	NA	NA
Symptomatic	4 or 2 years <sup>  </sup>	0.00225	0.00075	NA	NA	NA
AIDS	40 weeks	0.01125	0.00375	NA	NA	NA

\*Individual stage durations were sampled from a Weibull distribution function with shape parameter 2, except for the duration of the early latent stage of HSV-2 and the interval between recurrent HSV-2 ulcers, which used exponential distributions.

<sup>†</sup>Per contact; equal for recognized and unrecognized ulcers.

<sup>‡</sup>For both susceptibility (HIV-negative partner) and infectivity (HIV-positive partner).

<sup>§</sup>In scenarios 1 and 3 (of no effects of HIV on the natural history of HSV-2) and in scenario 2, only for individuals not symptomatic with HIV. In scenario 2a, values of these parameters for symptomatic HIV patients are 30 years, 2 weeks, and 3 months; in scenario 2b, 60 years, 4 weeks, and 1.5 month. See also Methods (Scenarios of Effects of HIV on the Natural History of HSV-2).

<sup>||</sup>Except during recurrent ulcers.

<sup>||</sup>For scenarios 1, 2a, and 3, asymptomatic phase is 5 years and symptomatic phase is 2 years; for scenario 2b, asymptomatic phase is 3 years and symptomatic phase is 4 years (see also Methods section on scenarios, noted above).

F = females; M = males; NA = not applicable.

geneous,<sup>22,23,28,35,38</sup> we assumed a Weibull distribution with shape parameter 2. Patients who have ulcers are considered to be infectious; a proportion of patients recognize symptoms. On the basis of their relative clinical severity,<sup>22,26</sup> we infer that rates of infectiousness and symptom-recognition are higher in association with primary ulcers than with recurrences.

No data are available on per-contact transmission probabilities for HSV-2. Genital shedding, a likely correlate of infectivity, is higher during symptomatic episodes than asymptomatic or in-between episodes and higher during primary episodes than recurrences.<sup>22,23</sup> Because it is not known how shedding levels translate into transmission probabilities, the latter were chosen from a range of estimates for other ulcerative STDs.<sup>45–47</sup> Transmission was assumed to be twice as efficient from males to females than from females to males.<sup>48–50</sup> To account for episodes of viral shedding in the absence of clinical signs,<sup>32,34,51–53</sup> a continuous low level of infectiousness throughout the early latent phase (i.e., in between recurrences) was specified.

Between 10% and 50% of HSV-2-seropositive individuals in Western settings report a history of genital herpes,<sup>54,55</sup> and symptoms are more common in men than in women.<sup>27,56,57</sup> In a Tanzanian population, 28% and 8% of HSV-2-seropositive men and women reported having ulcers in the past, as compared with 3% and 2% of HSV-2-seronegative

men and women; over the previous year, ulcers were reported by 11% and 6% of HSV-2-seropositive men and women, as compared with 1% and 2% of seronegative men and women.<sup>57</sup> On the basis of these data, we estimated that 30% of primary herpetic ulcers and 15% and 7.5% of recurrent herpetic ulcers in men and women, respectively, are recognized in SSA populations. Recognition was assumed to occur randomly and independently over different episodes and individuals, i.e., recognition at one episode did not influence recognition at subsequent episodes.

#### *Scenarios of Effects of HIV on the Natural History of Herpes Simplex Virus Type 2*

HIV disease is associated with threefold to fourfold increases in the number of HSV-2 culture-positive days<sup>58–61</sup> and the incidence of clinical ulcers.<sup>62</sup> The level of HSV-2 shedding during symptomatic ulcers increases with a decreasing CD4 cell count.<sup>58,63</sup> In addition, herpetic recurrences may last longer and be more severe in HIV-infected patients.<sup>3,60,64</sup>

Translating these effects into model parameters is not straightforward. It is unknown how shedding levels and clinical signs quantitatively translate into infectivity, symptom recognition, and cofactor effects. Studies likely oversample HIV-infected patients in the later, symptomatic

stages of disease. No prospective studies of the duration of untreated recurrences in relation to HIV status have been reported. To reflect these uncertainties, we simulated several scenarios. In scenario 1, HIV was assumed not to affect the natural history of HSV-2. In scenario 2a, HIV doubled the duration and the frequency of herpetic recurrences in patients with AIDS and in the last 2 years of the AIDS incubation period. In addition, the (remaining) duration of the early latent phase was doubled from the onset of symptomatic HIV. This ensured that most simulated HIV-infected patients continued to have recurrent ulcers until their deaths associated with AIDS. Thus, HSV-2-infected patients with HIV disease in this simulation suffered herpetic ulcers for an average of 8 weeks per year, as compared with a mean of 2 weeks for patients not infected with HIV. In scenario 2b, stronger biologic effects were assumed: HIV quadrupled the duration and frequency of recurrences and the (remaining) duration of the early latent phase in AIDS and the last 4 years of the AIDS incubation period. This corresponds to an average ulcer load of 32 weeks annually.

#### *Representation of HIV, Syphilis, Chancroid, and STD Treatment*

The STDSIM representations of syphilis, chancroid, and HIV disease (Table 1) were based on the scientific literature. Syphilis was represented by two consecutive stages, with the first, the *infectious stage*, corresponding to primary and secondary syphilis.<sup>65</sup> For computations of ulcer incidence, we assumed that each syphilis infection causes two ulcers during the infectious stage. Chancroid was represented as a continuous ulcerative episode lasting an average of 10 weeks.<sup>66,67</sup> Transmission probabilities were specified at a lower level (15–20%; Table 1) than the single available empirical estimate (43%, for male-to-female transmission<sup>45</sup>). This was done in order to have the model predict a realistic prevalence level and a realistic fraction of ulcers attributable to chancroid (see Results). To allow for the effects of HIV disease on the HSV-2 natural history described above, the AIDS incubation period was divided into three phases: *primary HIV disease*, *asymptomatic pre-AIDS*, and *symptomatic pre-AIDS*.

It was assumed that throughout the simulation period, 5% of symptomatic episodes of chancroid and the first syphilis stage were cured with antibiotic treatment, which reduced the episode duration to 2 weeks.

#### *STD Cofactor Effects on HIV Transmission*

We assumed that the presence of ulcers enhances the infectivity of HIV and the susceptibility to HIV. Because of the lack of data on STD cofactor magnitudes from controlled experimental studies, we estimated these magnitudes on the basis of data from observational studies of prostitutes and their clients in Nairobi (Table 1).<sup>68,69</sup> In line with their

relative clinical severity, herpetic recurrences were attributed a lower cofactor effect (10-fold per contact) than were primary HSV-2 and chancroid (25-fold per contact). For the infectious stage of syphilis, during which several ulcer episodes may occur,<sup>65</sup> an average cofactor effect of 10-fold was applied throughout. Identical cofactor effects were assumed for recognized and unrecognized ulcers.

#### *Scenario of Changing Sexual Behavior*

Several SSA countries have documented recent reductions in risk behavior, including decreases in casual sex and in prostitution and increases in condom use.<sup>10–13,70–73</sup> We assessed the influence of such behavioral changes on the proportion of GUD attributable to herpes in scenario 3, with a 25% reduction in the proportion of men visiting prostitutes, combined with a 25% reduction in relationship-formation rates, after 1990. This type and magnitude of behavior change provided good fit against survey data from Uganda.<sup>10–12,72,74</sup>

#### *Simulation Design*

To reduce random fluctuations associated with stochastic simulations, 100 simulation runs were conducted for each scenario. All outcomes are reported as averages over 100 runs and focus on the general adult population (aged 15–49 years). In univariate sensitivity analyses, simulations were rerun for a set of alternative quantifications in which the values of HSV-2 biomedical parameters were doubled and halved.

## RESULTS

#### *Herpes Simplex Virus Type 2 Epidemiology Before the HIV Epidemic*

The model provided a reasonable fit to data on HSV-2 seroprevalence by age and sex in Ndola, Zambia (Fig. 2), and other SSA populations.<sup>20,21</sup> Seroprevalence among women exceeds that among men, especially in the youngest age group. This reflects that (1) women become sexually active at younger ages than men; (2) women typically have older sex partners (whereas men have younger partners<sup>75,76</sup>) and thus more often meet HSV-2-infected partners; and (3) male-to-female transmission is more efficient than female-to-male transmission.<sup>48–50</sup> This sex difference was somewhat smaller in the model than that evident in the data, perhaps partly because simulated age differences between partners (e.g., a mean of 3.3 years in marriages) may have been less than in reality.<sup>77,78</sup>

The simulated incidence of primary herpetic ulcers was 2.6 and 2.7 per 100 person-years among men and women, respectively (in the total population, including both HSV-2-positive and HSV-2-negative persons), with corresponding peaks in the age groups of 20 to 24 years and 15 to 19 years. For recurrences, including unrecognized episodes,






Fig. 2. Simulated herpes simplex virus type 2 seroprevalence (%) in the general population, by age and sex, in a hypothetical city in sub-Saharan Africa (SSA) in the year 1980. In the absence of empirical data on HSV-2 seroprevalence from general SSA populations in the 1980s, we used data from Ndola, Zambia, in 1997 for comparison.<sup>21</sup>

simulated incidence was 41 and 53 per 100 person-years among men and women. Recurrences peaked at age 25 to 34 years for men and 20 to 29 years for women. The overall simulated incidence of recognized herpetic ulcers was 5.9 per 100 person-years.

#### Ulcer Etiology Before the HIV Epidemic

Based on simulated incidence of herpetic ulcers, syphilis and chancroid, we derived a distribution of aetiologies of incident ulcers. Of all incident ulcers in 1980, 72% were attributable to HSV-2, 24% to chancroid and 4% to syphilis. For comparison with proportions reported in surveys or seen in STD clinics, we also derived the distribution for recognized ulcers only. Among recognized ulcers, 28% were attributable to HSV-2, reflecting the specified relatively poor recognition of herpetic recurrences (Table 1); 64% and 8%, respectively, were caused by chancroid and syphilis.

#### Trends in Ulcer Epidemiology During the HIV Epidemic

The simulated HIV epidemic reached a prevalence of 31% in 2000, allowing a clear illustration of the possible effects of HIV on the epidemiology of HSV-2 and GUD.

When it was assumed that HIV had no enhancing effect on HSV-2 (scenario 1), the simulated incidence of recognized herpetic ulcers was stable between 1980 and 2000, at approximately six per 100 person-years (Fig. 3). The proportion of recognized ulcers attributed to HSV-2 remained roughly stable at 28%. HSV-2 seroprevalence decreased from 48% to 43% because of the higher HIV-related mortality in high-risk groups (Fig. 4). Because of this same

effect, the incidence of chancroid also fell during the HIV epidemic, and the proportion of recognized GUD attributable to chancroid decreased from 64% in 1980 to 59% by 2000. For syphilis, incidence and the contribution to GUD were fairly stable over time.

The moderately strong effects of HIV on HSV-2 (scenario 2a) increased the incidence of recognized herpetic ulcers from 5.9 to 7.4 per 100 person-years between 1980 and 2000. Combined with the relatively large decrease in the incidence of chancroid due to selective AIDS-related mortality, these effects increased the proportion of recognized




Fig. 3. Simulated incidence of recognized ulcers in the general population (aged 15–49 years), by etiology, in 1980 (leftmost bar) and 2000 (right bars): (A) Absolute incidence (per 100 person-years). (B) Proportional contribution (%) to genital ulcer disease (GUD). Scenario 1: no interactions; scenario 2a/moderate HIV: moderately strong effects of HIV on the natural history of herpes simplex virus type 2; scenario 2b/strong HIV: very strong effects of HIV on the natural history of HSV-2; scenario 3: behavior change, starting in 1990; scenario 4: combination of behavior change and moderately strong effects of HIV on HSV-2. For complete descriptions of scenarios, see Methods.




Fig. 4. Simulated prevalence (%) in the general population (15–49 years) over time of (A) herpes simplex virus type 2; (B) chancroid; (C) syphilis (ulcerative and nonulcerative stages combined, corresponding to serological syphilis); and (D) HIV in alternative scenarios. For complete descriptions of scenarios, see Methods.

GUD attributable to HSV-2 from 28% to 35%. In absolute terms, the increase in herpetic ulcer incidence counterbalanced the decrease in GUD incidence due to selective HIV-attributable mortality. As a result, HSV-2 seroprevalence and the overall incidence of GUD (of any etiology) remained stable between 1980 and 2000. Similar but more pronounced shifts in GUD etiology were predicted for scenario 2b, involving very strong effects of HIV. Here, the fraction of GUD attributable to HSV-2 increased to 48%. The absolute incidence of herpetic ulcers increased from six to 13 per 100 person-years, whereas the HSV-2 seroprevalence increased from 48% in 1980 to 58% in 2000.

Behavior change (scenario 3) reduced the incidence and prevalence of chancroid and syphilis considerably and rapidly (Figs. 3 and 4). The rates for chancroid decreased more markedly than those of syphilis because of its lower reproductive number.<sup>47</sup> Both HSV-2 seroprevalence and the incidence of herpetic ulcers decreased relatively little (from 48% to 39% and from 5.9 to 4.8 per 100 person-years, respectively). Since HSV-2 is a life-long infection with a high baseline prevalence and recurrent nature, herpetic ulcerations can be reduced only by reductions in new infections in the youngest age groups. Therefore, it takes a long time for a decrease in HSV-2 transmission to have an impact on the incidence of her-

petic ulcers at a population level. As a result of the large decrease in chancroid relative to the occurrences of herpes and syphilis, the proportion of recognized incident GUD attributable to HSV-2 increased from 28% in 1980 to 56% in 2000, whereas the proportion caused by chancroid decreased from 64% to 28%.

In a fourth scenario, combining behavior change and moderately strong effects of HIV, the incidence of herpetic ulcers remained unchanged over time, whereas that of chancroid and syphilis decreased, increasing the proportion of GUD attributable to HSV-2 to 61% by year 2000.

Figure 5 shows the age- and sex-specific seroprevalence of HSV-2 in year 2000 in the various scenarios. Effects of HIV on herpetic ulcerations increased HSV-2 seroprevalence, especially in the younger age groups. Very strong effects of HIV (scenario 2b) would shift the peak from the oldest group to a plateau from age 25 to 29 years onward for men and age 20 to 24 years for women, resulting in an unrealistic age pattern.<sup>20,21</sup> Behavior change, in contrast, slightly decreased HSV-2 seroprevalence in the younger age groups, who have the highest partnership-change rates. The age pattern in the combined scenario 4 resembled that in scenario 1, of no interactions.




Fig. 5. Simulated herpes simplex virus type 2 seroprevalence (%), by age and sex, in 2000 in alternative scenarios. For complete descriptions of scenarios, see Methods. Empirical data, for comparison, are from Ndola, Zambia, in 1997.<sup>21</sup>

### Sensitivity Analyses

To explore the robustness of the results, we assessed their sensitivity to variations in model assumptions on HSV-2 and its possible interactions with HIV.

Of the parameters tested, the average duration of the early latent phase, the probability distribution function assumed for this duration, and the interval between recurrent ulcers were the most important determinants of HSV-2 seroprevalence and ulcer incidence levels. Although halving durations considerably reduced HSV-2 seroprevalence, because of saturation. In all simulations, the gender difference in HSV-2 seroprevalence was insensitive to the relative efficiencies of male-to-female and female-to-male transmission, indicating that its population distribution is mainly determined by sexual behavior patterns. Across all quantifications, behavior change shifted GUD etiologic distributions much more than moderately strong biologic effects of HIV. Of the different possible effects of HIV, that on the frequency of herpetic recurrences had the greatest impact on GUD etiology. Of the two assumed components of behavioral change, a 25% reduction in relationship-formation rates had more of an impact on GUD etiology than did a 25% reduction in prostitute visits.

## Discussion

### Herpes Simplex Virus Type 2 Transmission Dynamics

In order to simulate both observed HSV-2 seroprevalences among adults in SSA ( $\geq 45\%$ ) and plausible frequencies of herpetic recurrences ( $< 5$  per 100 person-years), a high level of asymptomatic HSV-2 transmission had to be assumed. This finding supports the empirical evidence that

a large proportion of herpetic ulcerations go unrecognized and that these contribute substantially to transmission.<sup>49,54,55</sup>

Simulated HSV-2 epidemiology and GUD etiologic distribution were robust against most uncertainties in the natural history of HSV-2 (sensitivity analysis). This corroborates findings with simpler transmission models, in which interventions that had an impact on HSV-2 infectivity and ulceration had little influence on seroprevalence levels.<sup>79,80</sup> The duration of the early latent phase, during which ulcers recur, and the variability of this duration within the population were critical, however. Better data about these parameters, from long-term follow-up studies of unselected populations, would improve understanding of the transmission dynamics of HSV-2.

### GUD Incidence and Etiology

The model showed that realistic prevalences of HSV-2 and syphilis are consistent with an incidence of recognized GUD as high as 20 per 100 person-years. The simulated fraction of the population with recognized GUD in the last year, in contrast, was only 9% (scenarios 1 and 2), corresponding with proportions reported for SSA populations.<sup>20,37</sup> The difference between the incidence rate and the proportion with ulcers is the result of recurrence of ulcers (such as for HSV-2) and reinfections (such as for chancroid), as well as of clustering of ulcers caused by different STD, because of shared risk factors.

The simulated etiologic distribution of recognized GUD in 1980 (Fig. 3) matched data from the 1980s from SSA cities. Chancroid was the predominant cause of GUD.<sup>2,4,81–86</sup> The observed proportions should be viewed with caution, however, because diagnosis was based on either culture methods with limited sensitivity (likely resulting in underestimation) or clinical appearance (possibly causing overestimation).<sup>82,83,87</sup> The simulated 28% attributable to HSV-2 is higher than values reported for Nairobi (2–16%),<sup>82–85</sup> Kigali (11–19%),<sup>4,86</sup> and Durban and Johannesburg (3–7%).<sup>2,5</sup> These values, however, are from older studies in which culture was used to diagnose herpes.<sup>1</sup> The low simulated proportion attributable to syphilis (8%) matches proportions found in Nairobi (3–10%)<sup>82,85</sup> and Zimbabwe (6%).<sup>81</sup> Studies in Johannesburg and Rwanda showed proportions of 15% and 19%,<sup>4,88</sup> but because these diagnoses were based on serology (which remains positive for years after the disappearance of ulcers),<sup>1,65,89</sup> these may be overestimates.

### Behavioral Change Versus Biologic Effects of HIV

In both scenario 2b and scenario 3, the proportion of GUD attributable to HSV-2 increased considerably during the HIV epidemic (from 28% to 48% and 56%, respectively), resulting in GUD etiologic distributions in year 2000 (Fig. 5) in the range of those observed in advanced HIV

epidemics. In Uganda, 75% to 85% of reported ulcers of which the cause was known were attributable to HSV-2, 6% to 9% were due to chancroid, and 4% to 17% were due to syphilis.<sup>37,64</sup> In Malawi, the three infections accounted for about equal proportions of GUD.<sup>90</sup> In rural Zimbabwe, 32% of diagnosed ulcers were attributable to HSV-2, 47% to chancroid, and 22% to syphilis.<sup>91</sup>

Seemingly very strong biologic effects (scenario 2b) had to be assumed to explain a shift in GUD etiology approaching that caused by apparently realistic magnitudes of behavior change (scenario 3), and only in scenario 2b would HSV-2 seroprevalence and GUD incidence increase in absolute terms. The limited sensitivity of HSV-2 epidemiology in the general population to enhancement of herpetic ulceration in HIV-infected patients is first explained by the fact that not all HSV-2-infected patients are infected with HIV. Even in a severe HIV epidemic, not all HSV-2-infected patients contract HIV, and those who do contract it die earliest, which limits the combined prevalence. Second, the majority of partners of patients infected with HSV-2 or HIV are already infected with HSV-2; thus, increased infectivity does not always increase transmission in the population. Long-term follow-up studies in HIV-positive and HIV-negative populations could help determine whether HIV truly enhances herpetic ulceration as strongly as assumed here.

We may have overestimated the increase in the proportion of GUD attributable to HSV-2 because the simulation did not allow for a possible enhancement by HIV disease of chancroidal ulceration.<sup>7</sup> Because of its low reproductive number,<sup>47</sup> the latter effect could markedly increase the spread of chancroid during HIV epidemics. It should also be noted that the simulated HIV epidemic was, at a prevalence of 31% in 2000, more advanced than in many SSA populations. For less severe epidemics, increases in herpetic ulceration due to HIV would be less. Furthermore, we assumed sexual activity to be unaffected by symptoms. If many patients with ulcers were to temporarily abstain or reduce the frequency of intercourse,<sup>64,92</sup> we may have overestimated the influence of herpetic ulceration on HSV-2 transmission. Conversely, we ignored an effect of HIV on the severity of herpetic ulcers, which might lead HIV-infected patients to more often recognize these. This may have caused our model to underestimate the shift in GUD etiology possible among clinical cases in severe, advanced HIV epidemics.

The simulated shifts in GUD etiology in scenarios 2b and 3 were mediated through different mechanisms, which had opposing effects on HSV-2 seroprevalence and the absolute incidence of herpetic ulcers, chancroid, and syphilis (Figs. 3, 4, and 5). Does comparison of these predicted concomitant effects with empirical data allow inference as to which scenario best reflects reality?

No data are available on HSV-2 seroprevalence over time

periods spanning the course of SSA HIV epidemics.<sup>2</sup> One study showed an increase in seroprevalence in urban and rural Zaire between 1959 and 1985 (from 21% to 60% and from 6% to 32%, respectively).<sup>39</sup> But sampling for these surveys was among young men in 1959<sup>93</sup> and in undefined general populations in 1985, circumstances leaving doubts about their comparability in terms of age and risk profile. Moreover, this putative time trend occurred largely before the HIV epidemic in this region. For herpetic ulcer incidence, longitudinal data allowing assessment of time trends in SSA are also lacking. The contrast between scenarios 2b and 3 in their effect on the absolute incidence of herpetic GUD and HSV-2 seroprevalence therefore does not allow proper validation. Future population-based surveillance on trends in these indicators may help to solve this issue.

Data on time trends in bacterial ulcerative and nonulcerative STDs in several SSA settings are available, and some match the simulated reduction in rates of syphilis, chancroid, and recognized GUD in scenario 3. For example, the number of chancroid diagnoses decreased sevenfold in Harare between 1990 and 1998<sup>2</sup>; prevalences of gonorrhea, chlamydia, and syphilis decreased twofold to threefold among women in Nairobi between 1992 and 1997<sup>94</sup>; in Malawi there were 1.1-fold to twofold decreases in the prevalence of syphilis, trichomoniasis, gonorrhea, and genital ulcers between 1990 and 1996<sup>95</sup>; and factory workers in Mwanza, Tanzania, reported 35% less GUD in 1994 than in 1991.<sup>70</sup>

Comparison between empirical data and simulations on age patterns in HSV-2 seroprevalence suggests that the true effect of HIV on herpetic ulceration is less strong than assumed in scenario 2b. The simulated saturation of HSV-2 seroprevalence at ages 20 to 29 years in this scenario (Fig. 5) is inconsistent with available data from SSA populations. For example, in Ndola, Harare, and Kisumu, HSV-2 seroprevalence peaked above age 45 or 35 years for both sexes.<sup>20,21</sup> This inconsistency was apparent in spite of conservative assumptions about the relation between ulceration and infectivity, since we specified a nonzero infectivity throughout the early latent phase in between recurrences and ignored a possibly higher infectivity with the occurrence of recognized (more severe) ulcers than with unrecognized episodes. If HSV-2 infectivity in SSA populations would correlate more strongly with (recognized) ulcers than in these simulations, the specified strong effects of HIV would result in even more unrealistic age patterns. Behavior change, in contrast, did not produce unrealistic age patterns in HSV-2 seroprevalence, although the limited available epidemiologic data did not allow us to check whether this scenario fit better for populations with recorded behavior change than for populations without. For Ndola, the simulated decrease in HSV-2 seroprevalence among the young after reduction of risk behavior improved the fit for males but worsened it for females (Figs. 2 and 5). For this advanced HIV epidemic, scenario 4 (combining behavior change with moderately strong effects of HIV) fit observed age patterns in



HSV-2 seroprevalence as well as scenario 1 (with no interactions) did.

### *Other Explanations?*

The large influence of behavior change can explain why increases in HSV-2 as a cause of GUD are also pronounced in developing countries without severe HIV epidemics, such as Southeast Asia.<sup>96–98</sup> For example, in Thailand the incidence of clinical syphilis and chancroid decreased threefold and 20-fold between 1987 and 1993,<sup>98</sup> the period of the 100%-condom-use program that successfully constrained the spread of HIV.<sup>99</sup>

The simulated reduction in partner-change rates is not the only form of behavior change that can underlie relative increases in herpes. With increased condom use, the shift in GUD etiology could be even more marked than with partner reduction, because condom efficacy is likely lower against HSV-2—which can cause lesions outside the condom-protected genital area—than against bacterial STDs.

Behavior changes have not been noted in all countries in which the etiology of GUD has changed, not in, e.g., South Africa.<sup>2,14–16,100</sup> Also, the only published reports on absolute increases in genital herpes—which, according to our simulations, could result in advanced HIV epidemics if biologic effects of HIV are strong and there are no behavioral responses—come from Singapore and India.<sup>101,102</sup> Because Singapore and India at the time of these studies did not have advanced HIV epidemics, these absolute increases cannot be explained by HIV-related immunosuppression. This suggests that risk reduction and, in severe HIV epidemics, HIV-related immunosuppression are not the main explanations of (relative) increases in HSV-2 in all countries. Increased detection of genital herpes because of greater awareness may well be another important factor.

Finally, improved antibiotic management of STDs may have contributed to relative increases in herpes, through decreases in syphilis and especially chancroid. Improved care for STDs probably contributed to GUD epidemiologic shifts in more developed (e.g., Southeast Asian) countries.<sup>101</sup> For SSA populations, whose treatment facilities, symptom recognition, and treatment-seeking behavior are much poorer,<sup>103</sup> it is questionable whether STD treatment has so far played an important role. However, improvement of STD treatment, coupled with population-based health education to improve clinic attendance, certainly has the potential to enhance ongoing trends in GUD epidemiology in this region.

### *Conclusion and Implications*

Our results suggest that seeming increases in genital herpes in SSA are mainly relative—or artifacts reflecting changes in detection—rather than absolute. Even in severe HIV epidemics, the overall incidence of GUD and HSV-2 seroprevalence

are unlikely to increase in absolute terms, because the effects of HIV-related enhancement of herpetic ulceration are at a population level easily offset by factors causing GUD to decrease, such as HIV-attributable mortality. Behavioral response to the HIV epidemic is a particularly potent cause of relative increases in herpes among patients with GUD, through rapid and large decreases in chancroid.

The increasing proportion of HSV-2 as a cause of GUD in patients presenting to clinics implies that, as long as antiviral therapy for HSV-2 is not evident in developing countries, education on the prevention of transmission and on avoiding possible consequences of ulcers, including enhanced transmission of HIV (e.g., through temporary abstinence or condom use), becomes an increasingly important component of STD management.

### **References**

1. Morse SA. Etiology of genital ulcer disease and its relationship to HIV infection [editorial]. *Sex Transm Dis* 1999; 26:63–65.
2. O'Farrell N. Increasing prevalence of genital herpes in developing countries: implications for heterosexual HIV transmission and STI control programmes. *Sex Transm Infect* 1999; 75:377–384.
3. Chen CY, Ballard RC, Beck-Sague CM, et al. HIV infection and genital ulcer disease in South Africa: the herpetic connection. *Sex Transm Dis* 2000; 27:21–29.
4. Bogaerts J, Kestens L, van Dyck E, Martinex Tello W, Akingeneye J, Mukantabana V. Genital ulcers in a primary health clinic in Rwanda: impact of HIV infection on diagnosis and ulcer healing (1986–1992). *Int J STD AIDS* 1998; 9:706–710.
5. Htun Y, Dangor Y, Fehler G. Temporal changes in the aetiology of genital ulcer disease among STD clinic attenders in South Africa: the influence of concomitant HIV infection. Presented at the 5th Joint Conference of the STD and Infectious Diseases Societies of South Africa. Capetown, South Africa, 1997.
6. O'Farrell N. High prevalence of genital ulcers in countries worst affected by HIV in Africa [abstract TuPeC340g]. Presented at the 13th International AIDS conference. Durban, South Africa, 2000.
7. Wasserheit JN. Epidemiological synergy: interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. [review]. *Sex Transm Dis* 1992; 19:61–77.
8. Wald A, Corey L, Handsfield HH, Holmes KK. Influence of HIV infection on manifestations and natural history of other STDs. *Annu Rev Publ Health* 1993; 14:19–42.
9. Korenromp EL, Van Vliet C, Grosskurth H, et al. Model-based evaluation of single-round mass STD treatment for HIV control in a rural African population. *AIDS* 2000; 14:573–593.
10. Asiimwe-Okiror G, Opio AA, Musinguzi J, Madraa E, Tembo G, Carael M. Change in sexual behaviour and decline in HIV infection among young pregnant women in urban Uganda. *AIDS* 1997; 11:1757–1763.
11. Kilian AhD, Gregson S, Ndyabangi B, et al. Reductions in risk behaviour provide the most consistent explanation for declining HIV-1 prevalence in Uganda. *AIDS* 1999; 13:391–398.
12. Kamali A, Carpenter LM, Whitworth JA, Pool R, Ruberantwari A, Ojwiya A. Seven-year trends in HIV-1 infection rates, and changes in sexual behaviour, among adults in rural Uganda. *AIDS* 2000; 14:427–434.
13. Gregson S, Zhuwau T, Anderson RM, Chandiwana SK. Is there evidence for behaviour change in response to AIDS in rural Zimbabwe? *Soc Sci Med* 1998; 46:321–330.
14. Dangor Y, Htun H, Radebe F, Fehler G, Tshabalala V, Ballard RC. Epidemiological, clinical and microbiological features of genital ulcer disease in Southern Africa: influence of HIV infection [abstract 218]. Presented at the 13th meeting of the International Society for STD Research. Denver, 1999.

15. Steen R, DeCoito T, Ralepeli S, et al. Rapid control of chancroid in a South African mining company [abstract 337]. Presented at the 13th Meeting of the International Society for STD Research. Denver, 1999.
16. De Coito T, Ralepeli S, Steen R, et al. Sustained reductions of STD in a South African mining community [abstract 485]. Presented at the 13th meeting of the International Society for STD research. Denver, 1999.
17. Cowan FM, Johnson AM, Ashley R, Corey L, Mindel A. Relationship between antibodies to herpes simplex virus (HSV) and symptoms of HSV infection. *J Infect Dis* 1996; 174:470–475.
18. Van der Ploeg CPB, Van Vliet C, De Vlas SJ, et al. STDSIM: a microsimulation model for decision support on STD control. *Interfases* 1998; 28:84–100.
19. Korenromp EL, Van Vliet C, Bakker R, De Vlas SJ, Habbema JDF. HIV spread and partnership reduction for different patterns of sexual behaviour: a study with the microsimulation model STD-SIM. *Math Popul Studies* 2000; 8:135–173.
20. McFarland W, Gwanzura L, Bassett MT, et al. Prevalence and incidence of herpes simplex virus type 2 infection among male Zimbabwean factory workers. *J Infect Dis* 1999; 180:1459–1465.
21. Robinson NJ, Hosseini M. Age-specific herpes simplex virus type 2 seroprevalence in general populations: estimates from around the globe [abstract 118]. Presented at the MSSVD/ASTDA STI at the Millennium Conference. Baltimore, 2000.
22. Rawls WE, Gardner HL, Flanders RW, Lowry SP, Kaufman RH, Melnick JL. Genital herpes in two social groups. *Am J Obstet Gynecol* 1971; 110:682–689.
23. Corey L, Adams HG, Brown ZA, Holmes KK. Genital herpes simplex virus infections: clinical manifestations, course, and complications. *Ann Intern Med* 1983; 98:958–972.
24. Cone RW, Hobson AC, Palmer J, Remington M, Corey L. Extended duration of herpes simplex virus DNA in genital lesions detected by the polymerase chain reaction. *J Infect Dis* 1991; 164:757–760.
25. Koelle DM, Benedetti J, Langenberg A, Corey L. Asymptomatic reactivation of herpes simplex virus in women after the first episode of genital herpes. *Ann Intern Med* 1992; 116:433–437.
26. Corey L, Benedetti J, Critchlow C, et al. Treatment of primary first-episode genital herpes simplex virus infections with acyclovir: results of topical, intravenous and oral therapy. *J Antimicrob Chemother* 1983; 12(suppl B):79–88.
27. Jeansson S, Molin L. On the occurrence of genital herpes simplex virus infection. *Acta Dermatovener (Stockholm)* 1974; 54:479–485.
28. Rattray MC, Corey L, Reeves WC, Vontver LA, Holmes KK. Recurrent genital herpes among women: symptomatic vs. asymptomatic viral shedding. *Br J Vener Dis* 1978; 54:262–265.
29. Knox SR, Corey L, Blough HA, Lerner AM. Historical findings in subjects from a high socioeconomic group who have genital infections with herpes simplex virus. *Sex Transm Dis* 1982; 9:15–20.
30. Mertz GJ, Jones CC, Mills J, et al. Long-term acyclovir suppression of frequently recurring genital herpes simplex virus infection: a multicenter double-blind trial. *JAMA* 1988; 260:201–206.
31. Brock BV, Selke S, Benedetti J, Douglas JM, Corey L. Frequency of asymptomatic shedding of herpes simplex virus in women with genital herpes. *JAMA* 1990; 263:418–420.
32. Wald A, Zeh J, Selke S, Ashley RL, Corey L. Virologic characteristics of subclinical and symptomatic genital herpes infections. *N Engl J Med* 1995; 333:770–775.
33. Corey L, Wald A. Genital herpes. In: Holmes KK, Mardh P-A, Sparling PF, et al, eds. *Sexually Transmitted Diseases*. 3rd ed. New York: McGraw-Hill, 1999; 285–312.
34. Adam E, Kaufman RH, Mirkovic RR, Melnick JL. Persistence of virus shedding in asymptomatic women after recovery from herpes genitalis. *Obstet Gynecol* 1979; 54:171–173.
35. Guinan ME, MacCalman J, Kern ER, Overall JC Jr, Spruance SL. The course of untreated recurrent genital herpes simplex infection in 27 women. *N Engl J Med* 1981; 304:759–763.
36. Mertz GJ, Coombs RW, Ashley R, et al. Transmission of genital herpes in couples with one symptomatic and one asymptomatic partner: a prospective study. *J Infect Dis* 1988; 157:1169–1177.
37. Wawer MJ, Sewankambo NK, Serwadda D, et al. Control of sexually transmitted diseases for AIDS prevention in Uganda: a randomised community trial. *Lancet* 1999; 353:525–535.
38. Wald A, Zeh J, Selke S, et al. Reactivation of genital herpes simplex virus type 2 infection in asymptomatic seropositive persons. *N Engl J Med* 2000; 342:844–850.
39. Nahmias AJ, Lee FK, Beckman-Nahmias S. Sero-epidemiological and -sociological patterns of herpes simplex virus infection in the world. *Scand J Infect Dis Suppl* 1990; 69:19–36.
40. Kamali A, Nunn AJ, Mulder DW, Van Dyck E, Dobbins JG, Whitworth JAG. Seroprevalence and incidence of genital ulcer infections in a rural Ugandan population. *Sex Transm Infect* 1999; 75:98–102.
41. Kinghorn GR. Epidemiology of genital herpes [review]. *J Int Med Res* 1994; 22(suppl 1):14A–23A.
42. Wald A, Zeh J, Barnum G, Davis LG, Corey L. Suppression of subclinical shedding of herpes simplex virus type 2 with acyclovir [see comments]. *Ann Intern Med* 1996; 124(1 Pt 1):8–15.
43. Benedetti J, Corey L, Ashley R. Recurrence rates in genital herpes after symptomatic first-episode infection. *Ann Intern Med* 1994; 121:847–854.
44. Langenberg A, Benedetti J, Jenkins J, Ashley R, Winter C, Corey L. Development of clinically recognizable genital lesions among women previously identified as having “asymptomatic” herpes simplex virus type 2 infection. *Ann Intern Med* 1989; 110:882–887.
45. Plummer FA, D’Costa LJ, Nsanze H, Dylewski J, Karasira P, Ronald AR. Epidemiology of chancroid and *Haemophilus ducreyi* in Nairobi, Kenya. *Lancet* 1983; 2:1293–1295.
46. Csonka GW, Oates JK. Syphilis. In: Csonka GW, Oates JK, eds. *Sexually Transmitted Diseases: A Textbook of Genitourinary Medicine*. London: Bailliere Tindall, 1990:227–276.
47. Brunham RC, Plummer FA. A general model of sexually transmitted disease epidemiology and its implications for control. *Med Clin North Am* 1990; 74:1339–1352.
48. Mertz GJ, Ashley R, Burke RL, et al. Double-blind, placebo-controlled trial of a herpes simplex virus type 2 glycoprotein vaccine in persons at high risk for genital herpes infection. *J Infect Dis* 1990; 161:653–660.
49. Mertz GJ, Benedetti J, Ashley R, Selke SA, Corey L. Risk factors for the sexual transmission of genital herpes. *Ann Intern Med* 1992; 116:197–202.
50. Bryson Y, Dillon M, Bernstein DI, Radolf J, Zakowski P, Garratty E. Risk of acquisition of genital herpes simplex virus type 2 in sex partners of persons with genital herpes: a prospective couple study. *J Infect Dis* 1993; 167:942–946.
51. McCaughy ML, Fleagle GS, Docherty JJ. Inapparent genital herpes simplex virus infection in college women. *J Med Virol* 1982; 10:283–290.
52. Corey L. The current trend in genital herpes: progress in prevention. *Sex Transm Dis* 1994; 21:S38–S44.
53. Wald A, Corey L, Cone R, Hobson A, Davis G, Zeh J. Frequent genital herpes simplex virus 2 shedding in immunocompetent women: effect of acyclovir treatment. *J Clin Invest* 1997; 99:1092–1097.
54. Mertz GJ. Epidemiology of genital herpes infections. *Infect Dis Clin North Am* 1993; 7:825–839.
55. Brugha R, Keersmaekers K, Renton A, Meheus A. Genital herpes infection: a review. *Int J Epidemiol* 1997; 26:698–709.
56. Stavraky KM, Rawls WE, Chiavetta J, Donner AP, Wanklin JM. Sexual and socioeconomic factors affecting the risk of past infections with herpes simplex virus type 2. *Am J Epidemiol* 1983; 118:109–121.
57. Obasi A, Mosha F, Quigley M, et al. Antibody to HSV-2 as a marker of sexual risk behaviour in rural Tanzania. *J Infect Dis* 1999; 179:16–24.
58. Augenbraun M, Feldman J, Chirgwin K, et al. Increased genital shedding of HSV type 2 in HIV-seropositive women. *Ann Intern Med* 1995; 123:845–847.
59. Hitti J, Watts DH, Burchett SK, et al. Herpes simplex virus seropositivity and reactivation at delivery among pregnant women infected with human immunodeficiency virus-1. *Am J Obstet Gynecol* 1997; 177:450–454.

60. Schacker T, Zeh J, Hu HL, Hill E, Corey L. Frequency of symptomatic and asymptomatic herpes simplex virus type 2 reactivations among human immunodeficiency virus-infected men. *J Infect Dis* 1998; 178:1616–1622.
61. Mbopi-Keou F-X, Grésenguet G, Mayaud P, et al. Genital HSV-2 shedding is increased in HIV-infected women in Africa. *AIDS* 1999; 13:536–537.
62. Kaul R, Kimani J, Nagelkerke NJD, et al. Risk factors for genital ulcerations in Kenyan sex workers: the role of HIV-1 infection. *Sex Transm Dis* 1997; 24:387–392.
63. Bagdades EK, Pillay D, Squire SB, O'Neil C, Johnson MA, Griffiths PD. Relationship between herpes simplex virus ulceration and CD4+ cell counts in patients with HIV infection. *AIDS* 1992; 6:1317–1320.
64. Kanya MR, Nsubuga P, Grant RM, Hellman N. The high prevalence of genital herpes among patients with genital ulcer disease in Uganda. *Sex Transm Dis* 1995; 22:351–354.
65. Holmes KK, Mardh P-A, Sparling PF, et al, eds. *Sexually Transmitted Diseases*. 3rd ed. New York: McGraw-Hill, 1999.
66. Rauschkolb JE. Circumcision in treatment of chancroidal lesions of male genitalia. *Arch Dermatol Syph* 1939; 39:319–328.
67. Hanschell HM. Sulphanilamide in the treatment of chancroid. *Lancet* 1938; 1:886–888.
68. Hayes RJ, Schulz KF, Plummer FA. The cofactor effect of genital ulcers on the per-exposure risk of HIV transmission in sub-Saharan Africa [review]. *J Trop Med Hyg* 1995; 98:1–8.
69. Korenromp EL, De Vlas NJ, Nagelkerke NJD, Habbema JDF. Estimating the magnitude of STD cofactor effects on HIV transmission: how well can it be done? *Sex Transm Dis* 2001; 28:613–621.
70. Ng'weshemi JZL, Boerma JT, Pool R, et al. Changes in male sexual behaviour in response to the AIDS epidemic: evidence from a cohort study in urban Tanzania. *AIDS* 1996; 10:1415–1420.
71. Jackson DJ, Rakwar JP, Richardson BA, et al. Decreased incidence of sexually transmitted diseases among trucking company workers in Kenya: results of a behavioural risk-reduction programme. *AIDS* 1997; 11:903–909.
72. Wawer MJ, Serwadda D, Gray RH, et al. Trends in HIV-1 prevalence may not reflect trends in incidence in mature epidemics: data from the Rakai population-based cohort, Uganda. *AIDS* 1997; 11:1023–1030.
73. Meekers D. Going underground and going after women: trends in sexual risk behaviour among gold miners in South Africa. *Int J STD AIDS* 2000; 11:21–26.
74. Konde-Lule JK, Musagara M, Musgrave S. Focus group interviews about AIDS in Rakai district of Uganda. *Soc Sci Med* 1993; 37:679–684.
75. Garnett GP, Anderson RM. Factors controlling the spread of HIV in heterosexual communities in developing countries: patterns of mixing between different age and sexual activity classes. *Philos Trans R Soc Lond B Biol Sci* 1993; 342:137–159.
76. Gregson S, Nyamukapa CA, Garnett GP, et al. Rapid rise in age-specific HIV prevalence in young women in rural Zimbabwe: the contribution of sexual mixing patterns. Presented at the International Union for the Scientific Study of Populations Conference on Partnership Networks and the Spread of HIV and Other Infections. Chang Mai, Thailand, 2000.
77. Munguti K, Grosskurth H, Newell J, et al. Patterns of sexual behaviour in a rural population in north-western Tanzania. *Soc Sci Med* 1997; 44:1553–1561.
78. Glynn J. Sexual behaviour and HIV risk in young adults in Kisumu, Kenya and Ndola, Zambia. Presented at the International Union for the Scientific Study of Populations Conference on Partnership Networks and the Spread of HIV and Other Infections. Chang Mai, Thailand, 2000.
79. Blower SM, Porco TC, Darby G. Predicting and preventing the emergence of antiviral drug resistance in HSV-2. *Nat Med* 1998; 4:673–678.
80. White PJ, Garnett GP. Use of antiviral treatment and prophylaxis is unlikely to have a major impact on the prevalence of herpes simplex virus type 2. *Sex Transm Infect* 1999; 75:49–54.
81. Haines M, Goldsmid JM, Kennedy S. A study of gonorrhoea in the Rhodesian African. *Centr Afr J Med* 1978; 24:140–144.
82. Fast MV, D'Costa LJ, Nsanze H, et al. The clinical diagnosis of genital ulcer disease in men in the tropics. *Sex Transm Dis* 1984; 11:72–76.
83. Hazlett DTG, Bowmer MI, Nsanze H, et al. Herpes simplex virus in clinically suspected chancroid in Nairobi, Kenya. *Afr J Sex Transm Dis* 1984; 1:16–18.
84. Plummer FA, D'Costa LJ, Nsanze H, et al. Clinical and microbiologic studies of genital ulcers in Kenyan women. *Sex Transm Dis* 1985; 12:193–197.
85. Greenblatt RM, Lukehart SA, Plummer FA, et al. Genital ulceration as a risk factor for HIV infection. *AIDS* 1988; 2:47–50.
86. Bogaerts J, Ricart CA, Van Dyck E, Piot P. The etiology of genital ulceration in Rwanda. *Sex Transm Dis* 1989; 16:123–126.
87. Htun Y, Morse SA, Dangor Y, et al. Comparison of clinically directed, disease specific, and syndromic protocols for the management of genital ulcer disease in Lesotho. *Sex Transm Infect* 1998; 74(suppl 1):S23–S28.
88. Duncan MO, Bilgeri YR, Fehler HG. The diagnosis of sexually acquired genital infections in black patients in Johannesburg, South Africa. *South Afr J STD* 1981; 1:20–23.
89. Malonza IM, Tyndall MW, Ndinya-Achola JO, et al. A randomized, double-blind, placebo-controlled trial of single-dose ciprofloxacin versus erythromycin for the treatment of chancroid in Nairobi, Kenya. *J Infect Dis* 1999; 180:1886–1893.
90. Behets FMT, Liomba G, Lule G, et al. Sexually transmitted diseases and HIV control in Malawi: a field study of genital ulcer disease. *J Infect Dis* 1995; 171:451–455.
91. Le Bacq F, Mason PR, Gwanzura L, Robertson VJ, Latif AS. HIV and other sexually transmitted diseases at a rural hospital in Zimbabwe. *Genitourin Med* 1993; 69:352–356.
92. O'Farrell N, Hoosen AA, Coetzee KD, Van den Ende J. Sexual behaviour in Zulu men and women with genital ulcer disease. *Genitourin Med* 1992; 68:245–248.
93. Motulsky AG, Vandepitte J, Fraser GR. Population genetic studies in the Congo. I. Glucose-6-phosphate dehydrogenase deficiency, hemoglobin S, and malaria. *Am J Hum Genet* 1966; 18:514–537.
94. Moses S, Ngugi EN, Costigan A, et al. Declining STD and HIV prevalences among antenatal and family planning clinic attendees in Nairobi, Kenya, from 1992–1999 [abstract ThOrC 727]. Presented at the 13th International AIDS Conference. Durban, South Africa, 2000.
95. Taha TE, Dallabetta GA, Hoover DR, et al. Trends of HIV-1 and STD among pregnant and post-partum women in urban Malawi. *AIDS* 1998; 12:197–203.
96. Chua SH, Cheong WK. Genital ulcer disease in patients attending a public sexually transmitted disease clinic in Singapore: an epidemiologic study. *Ann Acad Med Singapore* 1995; 24:510–514.
97. Taylor DN, Duangmani C, Suvongse C, et al. The role of *Haemophilus ducreyi* in penile ulcers in Bangkok, Thailand. *Sex Transm Dis* 1984; 11:148–151.
98. Beyrer C, Jitwatcharanan K, Natpratan C, et al. Molecular methods for the diagnosis of genital ulcer disease in a sexually transmitted disease clinic population in Northern Thailand: predominance of HSV infection. *J Infect Dis* 1998; 178:243–246.
99. Hanenberg RS, Rojanapithayakorn W, Kunasol P, Sokal DC. Impact of Thailand's HIV-control programme as indicated by the decline of sexually transmitted diseases. *Lancet* 1994; 344:243–245.
100. Low-Beer D, Barnett T, Whiteside A, Stoneburner R. Knowledge diffusion and the personalisation of risk as key indicators of behaviour change in Uganda compared to Southern Africa [abstract ThPeDs 787]. Presented at the 13th International AIDS Conference. Durban, South Africa, 2000.
101. Goh CL. Epidemiology of sexually transmitted diseases in Singapore, 1977 to 1993. *Ann Acad Med Singapore* 1995; 24:495–499.
102. Kumar B, Rajagopalan M. Rising incidence of genital herpes in an STD clinic in north India [letter]. *Genitourin Med* 1991; 67:353–354.
103. Adler MW. Sexually transmitted diseases control in developing countries. *Genitourin Med* 1996; 72:83–88.