

Assignment 2

Prabhath Chellingi - CS20BTECH11038

Download all python codes from

https://github.com/PRABHATH-cs20-11038/AI1103/tree/main/Assignment_2

and latex-tikz codes from

https://github.com/PRABHATH-cs20-11038/AI1103/tree/main/Assignment_2

1 PROBLEM

(GATE – EC – 66) Consider two identical boxes B_1 and B_2 , where the box $B_i (i = 1, 2)$ contains $i + 2$ red and $5 - i - 1$ white balls. A fair die is cast. Let the number of dots shown on the top face of the die be N . If N is even or 5, then two balls are drawn with replacement from the box B_1 , otherwise, two balls are drawn with replacement from the box B_2 . The probability that the two drawn balls are of different colours is

2 SOLUTION

Let $X \in \{1, 2, 3, 4, 5, 6\}$ be the random variables of a die,

$$\Pr(X = N) = \begin{cases} \frac{1}{6} & 1 \leq N \leq 6 \\ 0 & \text{otherwise} \end{cases} \quad (2.0.1)$$

$$\Pr(X = m) \cdot \Pr(X = n) = 0 \quad (2.0.2)$$

$\forall m, n \in \{1, 2, 3, 4, 5, 6\}$ as a single die cannot show more than one outcome at a roll.

Let $Y \in \{0, 1\}$ represent the die where,

$1 \Rightarrow$ the die with outcome $N = \{2, 4, 5, 6\}$,

$0 \Rightarrow N = \{1, 3\}$.

$$\Pr(Y = 1) =$$

$$\Pr((X = 2) + (X = 4) + (X = 5) + (X = 6)) \quad (2.0.3)$$

by using Boolean logic and (2.0.2),

$$\Pr(Y = 1) = \frac{2}{3} \quad (2.0.4)$$

$$\Pr(Y = 0) = 1 - \Pr(Y = 1) = \frac{1}{3} \quad (2.0.5)$$

$$\Rightarrow \Pr(B_1) = \Pr(Y = 1) = \frac{2}{3} \quad (2.0.6)$$

$$\Rightarrow \Pr(B_2) = \Pr(Y = 0) = \frac{1}{3} \quad (2.0.7)$$

Let $C \in \{0, 1\}$ where,

$0 \Rightarrow$ red balls,

$1 \Rightarrow$ white balls.

TABLE 0

TABLE OF NUMBER OF BALLS

Box	No. of red balls ($i + 2$)	No. of white balls ($5 - i - 1$)	Total balls
B_1	$n(C = 0 B_1) = 3$	$n(C = 1 B_1) = 3$	$n(C B_1) = 6$
B_2	$n(C = 0 B_2) = 4$	$n(C = 1 B_2) = 2$	$n(C B_2) = 6$

TABLE 0

TABLE OF PROBABILITY OF TAKING BALLS FROM EACH BOX

Box	Probability of taking red ball	Probability of taking white ball
B_1	$\Pr(C = 0 B_1) = 1/2$	$\Pr(C = 1 B_1) = 1/2$
B_2	$\Pr(C = 0 B_2) = 2/3$	$\Pr(C = 1 B_2) = 1/3$

The probability of picking 2^{nd} ball is not effected by picking 1^{st} ball because the 2^{nd} ball is chose after replacement.

Selecting two balls with replacement is a Bernoulli distribution of 2 trails,

TABLE 0

TABLE OF NO. OF WAYS OF SELECTING TWO DIFFERENT COLOURED BALLS

Cases	Trail 1	Trail 2
$(B_1, C = 0, C = 1)$	$\Pr(C = 0 B_1)$	$\Pr(C = 1 B_1)$
$(B_1, C = 1, C = 0)$	$\Pr(C = 1 B_1)$	$\Pr(C = 0 B_1)$
$(B_2, C = 0, C = 1)$	$\Pr(C = 0 B_2)$	$\Pr(C = 1 B_2)$
$(B_2, C = 1, C = 0)$	$\Pr(C = 1 B_2)$	$\Pr(C = 0 B_2)$

TABLE 0
TABLE OF VARIABLES DESCRIPTION

Variables	Description
$\Pr((C = 0, C = 1) B_1)$	Probability of selecting two different coloured balls from box B_1
$\Pr((C = 0, C = 1) B_2)$	Probability of selecting two different coloured balls from box B_2
$\Pr(T)$	Total probability of selecting two different coloured balls

$$\begin{aligned} \Rightarrow \Pr((C = 0, C = 1)|B_1) = \\ \Pr(C = 0|B_1) \cdot \Pr(C = 1|B_1) \\ + \Pr(C = 1|B_1) \cdot \Pr(C = 0|B_1) \quad (2.0.8) \end{aligned}$$

$$\Pr((C = 0, C = 1)|B_1) = \frac{1}{2} \quad (2.0.9)$$

$$\begin{aligned} \Rightarrow \Pr((C = 0, C = 1)|B_2) = \\ \Pr(C = 0|B_2) \cdot \Pr(C = 1|B_2) \\ + \Pr(C = 1|B_2) \cdot \Pr(C = 0|B_2) \quad (2.0.10) \end{aligned}$$

$$\Pr((C = 0, C = 1)|B_1) = \frac{4}{9} \quad (2.0.11)$$

by using Bayes theorem,

$$\begin{aligned} \Pr(T) = \\ \Pr((C = 0, C = 1)|B_1) \cdot \Pr(B_1) + \\ \Pr((C = 0, C = 1)|B_2) \cdot \Pr(B_2) \quad (2.0.12) \end{aligned}$$

$$\Pr(T) = \left(\frac{1}{2}\right)\left(\frac{2}{3}\right) + \left(\frac{4}{9}\right)\left(\frac{1}{3}\right) \quad (2.0.13)$$

Hence, the probability of selecting two different coloured balls from the boxes is

$$\Pr(T) = \frac{13}{27} \quad (2.0.14)$$

Probability–
simulation: 0.48015,
actual: 0.48148148148148145