

Rajiv Gandhi University of Health Sciences, Karnataka

Curriculum Development Cell

CONFIRMATION FOR REGISTRATION OF SUBJECTS FOR DISSERTATION

Registration No.	: 09_T018_106054
Name of the Candidate	: AMRITA MISHRA
Address	: C-582, Second Floor, lane no.-7, Majlish Park, Delhi-110033
Name of the Institution	: Krupanidhi College of Physiotherapy, Bangalore
Course of Study and Subject	: MPT
Date of Admission to Course	: 06/06/2019
Title of the Topic	: Comparison of Proprioceptive Neuro-Muscular Facilitation and Mulligan Stretching on Hamstring Tightness in Recreational Football Players.
Brief resume of the intended work	: Attached
Signature of the Student	:
Guide Name	: DR. V. KALIDASAN
Remarks of the Guide	: Approved and Forwarded
Signature of the Guide	:
Co-Guide Name	:
Signature of the Co-Guide	:
HOD Name	: DR. MILAN ANAND
Signature of the HOD	:
Principal Name	:
Principal Mobile No.	:
Principal E-mail ID	:
Remarks of the Principal	:
Principal Signature	:

Print above Details

RAJIV GANDHI UNIVERSITY OF HEALTH SCIENCES
BANGALORE KARNATAKA

ANNEXURE II

PROFORMA FOR REGISTRATION OF SUBJECT FOR DISSERTATION

1	NAME OF THE CANDIDATE & ADDRESS	AMRITA MISHRA C-582, Second Floor, lane no.-7,Majlish Park, Delhi-110033
2	NAME OF THE INSTITUTION	KRUPANIDHI COLLEGE OF PHYSIOTHERAPY
3	COURSE OF STUDY AND SUBJECT	MASTER OF PHYSIOTHERAPY IN MUSCULOSKELETAL AND SPORTS
4	DATE OF ADMISSION TO COURSE	06.06.2019
5	TITLE OF TOPIC : COMPARISON OF PROPRIOCEPTIVE NEURO-MUSCULAR FACILITATION AND MULLIGAN STRETCHING ON HAMSTRING TIGHTNESS IN RECREATIONAL FOOTBALL PLAYERS.	

6	Brief Resume of the intended work
---	--

Introduction

Muscles ability to lengthen , allowing a joint to move throughout the range is known as muscle flexibility¹.Flexibility is one of the standard components for the prevention of injuries and for rehabilitation as well².Clinicians and sports medicine specialists routinely survey hamstring tightness (i.e., hamstring flexibility) since short hamstring have been identified with various lower limb injuries².

During the previous years in India, football among numerous youngsters has promoted as a recreational sport. As football is a game that includes extraordinary physical quality, such adolescents, who are not under experts guidance are at the risk of getting exposed to musculoskeletal injuries because of lack of appropriate warm-up and stretching's³.

High intensity sprinting is an important component for scoring a goal in football game⁴. while looking for athletes overall physical fitness, hamstring flexibility is one of the greatest concern which has a significant impact on athletes performance due to hamstring injuries⁵.

There has been a great importance of muscle stretching for prevention of injuries, as well as skill development and physical capacity¹.There are various stretching techniques for the muscle such as static stretching, ballistic stretching, and proprioceptive neuromuscular facilitation (PNF) stretching¹.

PNF techniques are more effective than the traditional stretching techniques for flexibility enhancement⁶.Mulligan traction straight leg raise is a technique for muscle stretching to increase the limited hip flexion range of motion which is limited due to hamstring shortening⁷.

6.1 Need of the study

Prevalence of hamstring tightness is present significantly at higher rates among athletes engaged with contact sports⁸.

Due to the hamstring tightness the performance in the game is affected.

Proprioceptive neuromuscular facilitation is been proved more effective than static stretching in football players¹.

TSLR Mulligan stretch is better than BLR Mulligan stretch in improving the biceps femoris muscle performance, flexibility⁹.

Young football players with great hamstring flexibility will have better performance in tests of sprinting².

Though there are studies available but there is a dearth in studies comparing the effects of PNF and mulligan TSLR stretching among the Amateur football players and that is where the need arises to compare the effects of PNF and mulligan TSLR stretching among the College Level football players.

6.2A. Aim and Objectives of the study Aim

of the study :

To evaluate the effectiveness of proprioceptive neuromuscular facilitation and mulligan stretching on hamstring tightness in recreational football players.

Objectives of the study :

- 1) To evaluate PNF stretching on sprint, agility and flexibility performance on Recreational football players.
- 2) To evaluate Mulligan stretching on sprint, agility and flexibility performance on Recreational football players.
- 3) To compare the effect of PNF and Mulligan stretching on sprint, agility and flexibility among the Recreational football players.

6.2B. Hypothesis

Null Hypothesis (H0):

There is no significant difference among PNF and Mulligan stretching on sprint, agility and flexibility among the Recreational football players. **Alternate Hypothesis (H1):**

There is significant difference between the PNF and Mulligan stretching on sprint, agility and flexibility among the Recreational football players.

6.3. Review of Literature

1) T Burgess , et al.

States that Contract-Relax Agonist-Contract (CRAC), when applied to stretch the hamstring muscles of active males, resulted in a large increase of active knee extension range of motion, without decreasing performance. Therefore, CRAC appears to be a safe and effective method of increasing the length of the hamstrings pre-sport activity and should be utilized by sports physiotherapists if deemed necessary. It was also shown to be beneficial following the initial assessment¹⁰.

2) Laura Deguzman et al.

Concluded the PNF intervention elicited a greater improvement in knee extension angle compared to the foam rolling intervention, but not when compared to the dynamic warmup intervention¹¹.

3) Landon Lempke, et al.

Stated that PNF stretching was not demonstrated to be more effective at increasing hamstring extensibility compared to static stretching. The literature reviewed suggests both are effective methods for increasing hip-flexion ROM¹².

4) Ramachandran. S, et al.

Conducted a study to find the effectiveness of two PNF stretching techniques (Hold Relax and Contract Relax-Antagonist Contract) on improving hamstring flexibility concluded that both PNF hold relax and PNF contract relax antagonist contract stretching is effective in improving hamstring flexibility. However PNF contract relax antagonist contract stretching technique was better than PNF hold relax stretching technique in improving the hamstring muscle flexibility¹³.

5) Erick D. Richman et al.

designed to determine the effect of a 6-minute protocol of self massage known as self-myofascial release (SMR) using a foam rolling device in conjunction with a general warmup and sport-specific dynamic stretching (DS) session on flexibility and explosive athletic performance stated that self-myofascial release in the form of foam rolling after a general warm-up and preceding a dynamic stretching session seems to improve squat jump and countermovement jump with no detriment to flexibility, drop jump, sprint, and agility performance in comparison with light walking and dynamic stretching¹⁴.

6) Wootack Lim

Investigated changes in hamstring flexibility in relation to intensity of proprioceptive neuromuscular facilitation stretching and changes in pain over time, and examine the correlations between pain level and target intensity or flexibility gain concluded that

Repetitive high intensity stretching may cause heavy burden on muscle tissues, and pain

caused by high-intensity stretching can hinder muscle performance. Moderate stretching intensity is recommended and considered conducive to maintaining the effects of stretching while ensuring its safety¹⁵.

7) Robin KV, Dr. YC Louis Raj

Revealed that a single session of PNF stretching on hamstring muscle is an effective and suitable method to improve the flexibility of hamstring muscles¹⁶.

8) Mohamed Arshadh

Concluded that there is no significant difference in Hamstring muscle flexibility following Muscle energy techniques and Proprioceptive neuromuscular facilitation types of stretching. Both Muscle energy techniques and Proprioceptive neuromuscular facilitation types of stretching are equally effective on hamstring flexibility¹⁷.

9) IzzetIslamoglu et al.

Researched the effect of different static stretching time on some physical performance parameters and concluded that the length of static stretching duration increased in flexibility. Also, static stretching decreased vertical jumping performance and did not affect speed and agility¹⁸.

10) Cem Kurt, IlkayFirtin

Compared the acute effects of static and dynamic stretching exercises on flexibility, agility, fatigue index and anaerobic performance in professional football players concluded that static and/or dynamic stretching exercises can be applied in addition to aerobic running to enhance flexibility¹⁹.

11) Yildirim MS et al.

Compare the effects of static stretching, proprioceptive neuromuscular facilitation (PNF) stretching and Mulligan technique on hip flexion range of motion (ROM) in subjects with bilateral hamstring tightness concluded that 4-week stretching intervention is beneficial for increasing hip flexion ROM in bilateral hamstring tightness. However, PNF stretching and Mulligan TSLR technique are superior to typical static stretching. These two interventions can be alternatively used for stretching in hamstring tightness⁷.

12) Dr. PatitapabanMohanty et al.

Compare the effect of dynamic soft tissue mobilization and Mulligan traction straight leg raise in increasing hamstrings flexibility by active knee extension range of motion and passive SLR concluded that Mulligan's traction straight leg raise and Dynamic Soft Tissue Mobilization improved active range of knee and passive straight leg raise range of motion in normal individuals⁵.

13) Bekie Mendes et al.

Purposed study to compare the flexibility and sprint performance values according to age and gender, the features of physical fitness performance of boys and girls who play football concluded that the relationship between speed and flexibility was varied by gender. The significant relation was found between flexibility values and 10 m sprint performance values of male footballers, while the significant relation was found between flexibility values and 20 m sprint performance values of female footballers. In order to improve the sprint performance, which is a specific skill of football, flexibility of the muscles should be trained with a program at an early age²⁰.

14) F. Garcia-Pinillos et al.

Stated that Hamstring flexibility is a key factor for performing football-specific skills, such as sprinting, jumping, agility, and kicking in young football players and muscle flexibility must be specifically trained in football players beginning at early ages².

15) Jesudas Mazumdar et al.

Compared mulligan traction straight leg raise technique vs muscle energy technique on hamstring tightness in asymptomatic male concluded that Mulligan TSLR and MET are effective in reducing the hamstring tightness. When both groups are compared MET was found to be more effective than TSLR²¹.

16) Mandeepkaur et al.

Compared the Effectiveness of Static Stretching Versus PNF Stretching of Hamstring Muscles Following Superficial Heat in Athletes and stated both the methods of stretching were effective but the PNF group shows more improvement than static group, so the PNF stretching is more effective than static stretching¹.

17) Márk Váci1. et al.

Investigate the effects of a short-term in-season plyometric training program on power, agility and knee extensor strength indicate that plyometric training consisting of high impact unilateral and bilateral exercises induced remarkable improvements in lower extremity power and maximal knee extensor strength, and smaller improvements in soccer-specific agility²².

18) Pratishtha, K., Jagga, V.

Stated that Mulligan TSLR stretch is more effective than BLR stretch in improving biceps femoris muscle performance, flexibility and pelvic rotation⁹.

19) Nagarwal A.K. et al.

Conducted a study to determine the effectiveness of two PNF stretching technique for improving hamstring flexibility and to compare the effectiveness of two PNF stretching techniques (Hold-Relax and Contract-Relax) for improving hamstring flexibility concluding that Both the techniques viz. PNF Hold Relax and PNF-CRAC are almost equal

in their clinical effectiveness for improving hamstring flexibility and that either of the techniques may be used in clinical practice for improving hamstring flexibility²³.

20) Anis Chaouachi, et al.

Investigated the effects of static and dynamic stretching alone and in combination on subsequent agility, sprinting, and jump performance concluded that individuals who wish to implement static stretching should include an adequate warm-up and dynamic sport specific activities with at least 5 or more minutes of recovery before their sport activity²⁴.

21) Pooja D Kapadia, et al.

Conducted a comparative study on immediate effects of traction straight leg and bent leg raise on hamstring muscle flexibility in normal individuals and stated that both Mulligan's Traction Straight Leg Raise and Bent leg Raise are effective in improving the hamstring muscle flexibility in normal individuals, but Mulligan's traction straight leg raise technique was more effective in improving hamstring flexibility²⁵.

7	Materials and Methodology
---	----------------------------------

7.1. Source of Data

Krupanidhi football Academy

7.2 Method of Collection of Data

7.2.1 Study Design

Two group Quasi experimental design

7.2.2 Sample Design

Purposive sampling technique

7.2.3 Sample Size

A sample of 82 subjects with hamstring tightness college level football According to the following calculations :

Sample size calculation:

$$SS_{UKP} = \frac{z^2 p (1-p)}{e^2}$$

$$e^2$$

z= 1.96 p= power of confidence

i.e. 68% e= 0.05 type 1 error

$$SS_{UKP} = \frac{(1.96)^2 \times 0.68 (1-0.68)}{(0.05)^2}$$

$$(0.05)^2$$

$$= \frac{3.8416 \times 0.68 \times 0.32}{0.0025}$$

$$0.0025$$

$$= \frac{0.83593216}{0.0025}$$

$$0.0025$$

$$= 334.372$$

$$= 334 \text{ (sample size for unknown population)}$$

$$SS_{KP} = SS_{UKP}$$

	$1 + \frac{SS_{UKP}}{N}$ $= \frac{334}{1 + 334/110}$ $= 334 / 4.03636364$ $= 82 \text{ (Sample size for known population)}$ <p>7.3 Selection criteria</p> <p>1. Inclusion criteria:</p> <p>A. College Level football players, all the participants were non-professional, collegiate and university-level competitive players. These players periodically participating in fitness training and conditioning, sport or other competition, practices or events requiring physical strength, agility, flexibility, range of motion, speed or stamina and who were not affiliated with an amateur, educational or professional athletic organization or any associated that sponsors athletic programs or events in the state²⁶.</p> <p>B. Tightness of hamstring muscle evaluated under AKE^{19, 1}.</p> <p>C. No recent injuries in trunk and lower extremities^{19, 1}.</p> <p>D. Age - 18-23 years^{19, 1}</p> <p>E. Gender - male and female^{19, 1}</p> <p>2. Exclusion criteria:</p> <p>A. Muscle soreness^{19, 1}</p> <p>B. Hamstring spasm or tendinopathy during last 6months^{19, 1}.</p> <p>C. Inflammatory conditions of hip and knee joint^{19, 1}.</p> <p>D. Neurological involvement of lower limb^{19, 1}</p> <p>E. Radiculopathy of lower limb^{19, 1}</p>
--	---

7.4 Outcome Measures

Active Knee Extension Test (AKE) :

Patient will be lying in supine lying position, the pelvis will be stabilized with the help of the stabilization strap. A rectangular frame is placed then the hip and knee is maintained at 90° with the reference to the rectangular frame.

Then the goniometer is placed over the lateral femoral condyle, with one arm aligned along the thigh in direction of the greater trochanter and the other arm aligned over the leg in the lateral

malleolus. The subject will be then instructed to extend the knee and hold this position for 2-3 seconds and the goniometer reading will be recorded⁵.

Illinois Agility Test :

The length of the field will be 10m, while the width (separation between the beginning and the finishing point) will be 5m four cones were put at the focal points of testing zone at the distance of 3.3m from each other. The four cones will be utilized to stamp the beginning, finish, and two turning points.

The test will be performed by the subjects while lying face down, with their hands at the shoulder level. The trail will begin on the “GO” order and accordingly, the subjects ran as quick as could be expected under the circumstances. The trial will be finished when the players will cross the end goal without having thumped over any cones¹⁹.

40 Yards Dash Test :

The subjects will run a full speed 40yards sprint on the field. The time will be recorded for sprint on hand held stop watch and the average time will be recorded to the nearest hundredth of a second for each subjects²⁷.

7.5 Study Duration

1 Year

7.6 Procedure

The ethical clearance will be obtained from the Institutional ethical committee. After explanation of procedure, a written informed consent will be taken from the individual participants. AKE test will be performed on the subjects and scores will be recorded accordingly. Based on the scores subjects will be categorised under hamstring tightness. Then the subjects will be divided into two groups. Randomization will be done for allocating into 2 groups. Group A

PNF stretching will be given.

Group B

Mulligan traction straight leg raise stretching (TSLR) will be.

After dividing the subjects into two groups.

After that the participants pre testing will be done for flexibility using Active knee extension test, sprint using 40yards dash test and agility using Illinois test.

Group – A will be given PNF stretching i.e, Contract relax and Group – B will be given Mulligan TSLR stretching for 3times per week for 6 weeks.

After completion of 6-weeks of intervention the participants will be assessed for flexibility, sprint, and agility.

Methods of application

PNF STERTCHING :contract relax technique will be used. The subject will be lying supine the hip and knee will be maintained at 90° then the subject will be guided to perform the SLR. SLR will be performed until the subject feel the stretch of hamstring muscle. Then maintaining the same position for 15 seconds, the therapist will support the leg of participants at ankle. After that the subject will be instructed to perform the continuous contraction of muscles for 6sec and performing hip extension. The resistance will be applied at the ankle. After this the subject was asked to perform resisted hip flexion. The participant will be maintaining hip flexion for 6seconds. 30 seconds rest will be given at the end of each repetition of the above procedure. Total three repetitions will be given for each leg and continuous instruction were given^{10, 23}.

MULLIGAN TRACTION STRAIGHT LEG RAISE STRETCHING: In supine lying traction will be applied to the leg while lifting the limb through the pain free range of motion. The maximum traction force will be applied along the longitudinal axis of the leg while the knee remains extended. At the same time the limb will be passively moved through the range of motion until the beginning of inconvenience and afterwards it will come back to the resting position. It will be ensured that there will be no pain during the procedure. The direction of the leg raise will be altered if the subject complains of pain. Pain free three repetitions will be given to each subject⁷ with 30seconds rest in between the repetitions.

After 6weeks of intervention data will be collected and analyzed.

Statistical Analysis

The data on response variables Sprint, Agility and Flexibility in both groups is carefully elicited and analyzed through following statistical technique by using statistical software SPSS version 20.0.

- 1) The descriptive measures of mean, range and standard deviation will be used to describe the pre and post score in both the groups.
- 2) T-test will be used to compare pre and post changes in flexibility , sprint and agility, in both the groups.

7.7 Does the study require any investigation or intervention to be conducted on patients or other humans or animals? If so please describe.

Yes, the study to be conducted on human subjects who are college level football players.

7.8 Has ethical clearance been obtained from the subject and the institution?

Yes, ethical clearance has been obtained from the institution.

8	List of References
---	---------------------------

	<ol style="list-style-type: none"> 1) Kaur M, Paul R, Kumar S, Arora R, Arora L. A Randomized Controlled Trial to Compare the Effectiveness of Static Stretching Versus PNF Stretching of Hamstring Muscles Following Superficial Heat in Athletes. <i>International Journal of Scientific and Research Publications</i>. 2014 Jul;4(7). 2) García-Pinillos F, Ruiz-Ariza A, Moreno del Castillo R, Latorre-Román PÁ. Impact of limited hamstring flexibility on vertical jump, kicking speed, sprint, and agility in young football players. <i>Journal of sports sciences</i>. 2015 Jul 21;33(12):1293-7. 3) Bhosale N, Yeole U, Chogle A, Khatri S. Assessment of Lower Extremity Flexibility in Recreational Football Players. 4) Gajdosik R, Lusin G. Hamstring muscle tightness: reliability of an active-knee-extension test. <i>Physical therapy</i>. 1983 Jul 1;63(7):1085-8. 5) Singh, Basant&Pattnaik, Mrs&Mohanty, Dr. (2016). A Comparative Study of Mulligan Traction Straight Leg Raising and Dynamic Soft Tissue Mobilization to Increase Hamstringss Flexibility. <i>IOSR Journal of Nursing and Health Science</i>. 05. 80-88. 10.9790/1959-0504058088. 6) Zakaria A, Melam G, Buragadda S. Efficacy of PNF stretching techniques on hamstring tightness in young male adult population. <i>World J Med Sci</i>. 2012;7(1):23-6. 7) Yıldırım MS, Ozyurek S, Tosun OÇ, Uzer S, Gelecek N. Comparison of effects of static, proprioceptive neuromuscular facilitation and Mulligan stretching on hip flexion range of motion: a randomized controlled trial. <i>Biology of sport</i>. 2016 Mar;33(1):89. 8) Weerasekara I, Kumari HM, Weeraratna N, Withanage C, Wanniarachchi C, Mariyanayagam Y, Vigneshwaran S, Shivaraja P, Suraweera H. The prevalence of hamstring tightness among the male athletes of University of Peradeniya in 2010. <i>Journal of SLSAJ</i>. 2012;12:56-8. 9) Pratihtha K, Jagga V. Effect of mulligan stretching techniques (TSLR AND BLR) on biceps femoris muscle and pelvic rotation by using surface EMG and bubble inclinometer respectively. <i>Journal of Exercise Science and Physiotherapy</i>. 2012 Jun;8(1):39. 10) Lempke L, Wilkinson R, Murray C, Stanek J. The Effectiveness of PNF Versus Static Stretching on Increasing Hip-Flexion Range of Motion. <i>Journal of sport rehabilitation</i>. 2018 May 1;27(3):289-94. 11) Deguzman L, Flanagan SP, Stecyk S, Montgomery MM. The Immediate Effects of Selfadministered Dynamic Warm-up, Proprioceptive Neuromuscular Facilitation, and Foam Rolling on Hamstring Tightness. <i>Athletic Training and Sports Health Care</i>. 2018 May 16;10(3):108-16.
--	---

12)	Lempke L, Wilkinson R, Murray C, Stanek J. The Effectiveness of PNF Versus Static Stretching on Increasing Hip-Flexion Range of Motion. <i>Journal of sport rehabilitation</i> . 2018 May 1;27(3):289-94.
13)	Sivagnanam, Ramachandran & Jibipaul, & N, Senthilkumar & .S, Sudhakar. (2018). Comparative Effect Of Pnf Stretching Technique On Hamstring Flexibility. 4.
14)	Richman ED, Tyo BM, Nicks CR. Combined effects of self-myofascial release and dynamic stretching on range of motion, jump, sprint, and agility performance. <i>The Journal of Strength & Conditioning Research</i> . 2019 Jul 1;33(7):1795-803.
15)	Lim W. Optimal intensity of PNF stretching: maintaining the efficacy of stretching while ensuring its safety. <i>Journal of physical therapy science</i> . 2018;30(8):1108-11.
16)	Robin KV, Raj YL. Impact of PNF stretching on muscular flexibility in football.
17)	Arshadh, Mohamed. (2016). Effect's of different muscle stretching techniques on hamstring muscle flexibility in amateur soccer players..
18)	Islamoglu I, Atan T, Unver S, Cavusoglu G. Effects of different durations of static stretching on flexibility, jumping, speed and agility performance. <i>The Anthropologist</i> . 2016 Mar 1;23(3):454-61.
19)	Kurt C, Firtin İ. Comparison of the acute effects of static and dynamic stretching exercises on flexibility, agility and anaerobic performance in professional football players. <i>Turkish Journal of Physical Medicine & Rehabilitation/TurkiyeFiziksel Tip VeRehabilitasyonDergisi</i> . 2016 Jul 1;62(3).
20)	MENTES B, ERCIN T, Kenan UZ. Examination of flexibility and sprint performance values of adolescent footballers. <i>Turkish Journal of Sport and Exercise</i> . 2015;17(3):16-20.
21)	Mazumdar J, Shriwas JK. A comparison between Mulligan traction straight leg raise technique vs muscle energy technique on hamstring tightness in asymptomatic male. <i>Int J Physiother Res</i> . 2014;2(2):412-17.
22)	Vácz M, Tollár J, Meszler B, Juhász I, Karsai I. Short-term high intensity plyometric training program improves strength, power and agility in male soccer players. <i>Journal of human kinetics</i> . 2013 Mar 1;36(1):17-26.
23)	Nagarwal AK, Zutshi K, Ram CS, Zafar R, Hamdard J. Improvement of hamstring flexibility: A comparison between two PNF stretching techniques. <i>International journal of sports science and engineering</i> . 2010;4(1):25-33.

24)	Chaouachi A, Castagna C, Chtara M, Brughelli M, Turki O, Galy O, Chamari K, Behm DG. Effect of warm-ups involving static or dynamic stretching on agility, sprinting, and jumping performance in trained individuals. <i>The Journal of Strength & Conditioning Research</i> . 2010 Aug 1;24(8):2001-11.
25)	Kapadia PD, Meshram VK. A comparative study on immediate effects of traction straight leg and bent leg raise on hamstring muscle flexibility in normal individuals. <i>IJAR</i> . 2019;5(4):274-8.
26)	Sudhakar S .et.al effect of blood pressure regulating gene polymorphisms on the endurance capacity of athletes: <i>biomedicine</i> : 2018;38(2): 267-273.
27)	Guiser ZJ. Examination of Hamstring Flexibility and Maximal Sprint Speed.
28)	Kirmizigil B, Ozcaldiran B, Colakoglu M. Effects of three different stretching techniques on vertical jumping performance. <i>The Journal of Strength & Conditioning Research</i> . 2014 May 1;28(5):1263-71.
29)	Yu B, Queen RM, Abbey AN, Liu Y, Moorman CT, Garrett WE. Hamstring muscle kinematics and activation during overground sprinting. <i>Journal of biomechanics</i> . 2008 Nov 14;41(15):3121-6.
30)	Craig KL, Goman DM, Shields ML, Stewart JL. The effects of hamstring stretching on vertical jump in healthy young adults.
31)	vanDoormaal MC. The relation between hamstring flexibility and hamstring injuries in male amateur soccer players (Master's thesis).
32)	García-Pinillos F, Ruiz-Ariza A, Moreno del Castillo R, Latorre-Román PÁ. Impact of limited hamstring flexibility on vertical jump, kicking speed, sprint, and agility in young football players. <i>Journal of sports sciences</i> . 2015 Jul 21;33(12):1293-7.
33)	Altmann S, Ringhof S, Neumann R, Woll A, Rumpf MC. Validity and reliability of speed tests used in soccer: A systematic review. <i>PloS one</i> . 2019;14(8).
34)	Hachana Y, Chaabène H, Nabli MA, Attia A, Moualhi J, Farhat N, Elloumi M. Test-retest reliability, criterion-related validity, and minimal detectable change of the Illinois agility test in male team sport athletes. <i>The Journal of Strength & Conditioning Research</i> . 2013 Oct 1;27(10):2752-9.
35)	Mann JB, Ivey PJ, Brechue WF, Mayhew JL. Validity and reliability of hand and electronic timing for 40-yd sprint in college football players. <i>The Journal of Strength & Conditioning Research</i> . 2015 Jun 1;29(6):1509-14.

36)	Jung, Jihye& Choi, Wonjae& Lee, Yonghyuk& Kim, Jiwoo& Kim, Hyunju& Lee, Kyoungho& Lee, Jaewoo& Lee, Seungwon. (2017). Immediate effect of self-myofascial release on hamstring flexibility. <i>Physical Therapy Rehabilitation Science</i> . 6. 45-51. 10.14474/ptrs.2017.6.1.45.
37)	Tsolakis C, Douvis A, Tsigganos G, Zacharogiannis E, Smirniotou A. Acute effects of stretching on flexibility, power and sport specific performance in fencers. <i>Journal of Human Kinetics</i> . 2010 Dec 1;26:105-14.

38)	Pagaduan JC, Pojskić H, Užičanin E, Babajić F. Effect of various warm-up protocols on jump performance in college football players. <i>Journal of human kinetics</i> . 2012 Dec 1;35(1):127-32.
39)	Hasebe K, Okubo Y, Kaneoka K, Takada K, Suzuki D, Sairyo K. The effect of dynamic stretching on hamstrings flexibility with respect to the spino-pelvic rhythm. <i>The Journal of Medical Investigation</i> . 2016;63(1.2):85-90.
40)	Goral, Kemal. (2015). Examination of agility performances of soccer players according to their playing positions. <i>The Sport Journal</i> . 10.17682/sportjournal/2015.004.
41)	Shultz S, Olszewski A, Ramsey O, Schmitz M, Wyatt V, Cook C. A systematic review of outcome tools used to measure lower leg conditions. <i>International journal of sports physical therapy</i> . 2013 Dec;8(6):838.
42)	Makhlouf I, Chaouachi A, Chaouachi M, Ben Othman A, Granacher U, Behm DG. Combination of agility and plyometric training provides similar training benefits as combined balance and plyometric training in young soccer players. <i>Frontiers in physiology</i> . 2018 Nov 13;9:1611.
43)	Barbosa GD, Dantas GA, Silva BR, Pinheiro SM, Santos HH, Vieira WH. Intra-rater and inter-instrument reliability on range of movement of active knee extension. <i>Motriz: Revista de Educação Física</i> . 2017 Mar;23(1):53-9.
44)	van der Made AD, Wieldraaijer T, Kerkhoffs GM, Kleipool RP, Engebretsen L, Van Dijk CN, Golanó P. The hamstring muscle complex. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> . 2015 Jul 1;23(7):2115-22.
45)	Koli BK, Anap DB. Prevalence And Severity Of Hamstring Tightness Among College Student: A Cross Sectional Study. <i>International Journal of Clinical and Biomedical Research</i> . 2018.
46)	Clanton TO, Coupe KJ. Hamstring strains in athletes: diagnosis and treatment. <i>JAAOS Journal of the American Academy of Orthopaedic Surgeons</i> . 1998 Jul 1;6(4):237-48.

	<p>47) Mann JB, Ivey PJ, Brechue WF, Mayhew JL. Validity and reliability of hand and electronic timing for 40-yd sprint in college football players. The Journal of Strength & Conditioning Research. 2015 Jun 1;29(6):1509-14.</p> <p>48) Hollis J. The Effects Of Static Stretching Of The Hamstring Muscles In A Warm-Up On Performance Among Football Players: A Systematic Literature Review.</p> <p>49) Biener E. The effect of different warm-up protocols on speed and agility.</p> <p>50) Chinnavan E, Gopaladhas S, Kaikondan P. Effectiveness of pilates training in improving hamstring flexibility of football players. Bangladesh Journal of Medical Science. 2015 Jun 20;14(3):265-9.</p>
	<p>51) Roopa Desai, TusharPalekar, Amruta Deshpande, Carryover effect of Mulligan Traction straight leg raise in individuals with hamstring tightness, GLOBAL JOURNAL FOR RESEARCH ANALYSIS : VOLUME-6 Issue-5 May-2017</p>

APPENDIX-I

CONSENT FORM

Title: Comparison Of Proprioceptive Neuro-Muscular Facilitation And Mulligan Stretching On Hamstring Tightness In College Level Football Players.

Principal Investigator: Amrita Mishra, MPT – Musculoskeletal and Sports

About this form:

This form gives you important information about a research study. Please read it carefully. One of our staff members will be with you to answer any questions you may have about the study and what you will be asked to do. If you decide to be a participant (called a "subject"), you will have to sign this form. We will give you a copy of it to keep.

Why is this research being done?

The purpose of this research is to find out the effectiveness of proprioceptive neuro-muscular facilitation and mulligan stretching on hamstring tightness in college level football players. We are inviting you to join the study. We would like about 82 people to take part in this study.

How long will I take part in this study?

You will spend 6 weeks from the beginning to the end of the study. During this time, you will come to this clinic for treatment and for testing.

What will happen in this research study?

If you agree to be in this study, you will be assigned randomly into groups to receive one of two treatment techniques. One of these is called Proprioceptive neuromuscular facilitation stretching.

The other group will receive mulligan traction straight leg raise technique. Before your program begins, you will perform a sprint, agility and flexibility test.

If you are in the proprioceptive neuromuscular facilitation group, you will come to the clinic and will take treatment. Thrice a week, you will visit clinic to take treatment.

If you are in the Mulligan Traction Straight leg raise group, you will come to the clinic thrice a week and will receive treatment. your daily activities with one of us.

You will return to the clinic to repeat the tests after 6 week of intervention.

What are the risks and possible discomforts from being in this research study?

During the intervention and during your activities program, it is possible that you could feel pain. You may feel tired at the end of the intervention or activities period. If this happens, a short rest period would be a good idea.

What are the possible benefits from being in this study?

You may feel more active and may be more comfortable doing your daily household and outdoor activities.

Every institution that has an IRB will have several additional "boilerplate" sections.

These will include:

- 1) Description of alternative treatments (if they exist)
- 2) Assurance that nonparticipation will not affect routine medical care
- 3) Financial compensation (if relevant)
- 4) Injury statement, indicating the patient's responsibility for cost of care if needed.
- 5) Contact individual including phone number.
- 6) Privacy and HIPAA authorization, explaining what information will be taken and how it will be used.
- 7) Right to withdraw from the study at any time without prejudice or bias

Statement of Subject or Person Giving Consent/ Assent

I have read this consent form. This research study has been explained to me, including risks and possible benefits (if any), other options for treatments or procedures, and other important things about the study. I have had the opportunity to ask questions.

If you understand the information we have given you, and would like to take part in this research study, and also agree to allow your health information to be used and shared as described above, then please sign below.

Name of the Subject:

Signature of the subject:

Date:

Name of the Witness (if required):

Signature of the Witness:

Date:

Statement of Individual Obtaining Consent:

I have explained the research to the study subject, and I have answered all questions about this research study to the best of my ability.

Investigator or Person Obtaining Consent form:

Date:

The question format of this sample form is taken from the standard Informed Consent Form of Partners Human Research Committee, Partners HealthCare System, Boston, 2007.

9	SIGNATURE OF CANDIDATE	
10	REMARKS OF GUIDE	Approved and Forwarded
11.1	NAME AND DESIGNATION	Dr. V. Kalidasan Professor Krupanidhi College Of Physiotherapy
11.2	SIGNATURE	
11.3	CO GUIDE (IF ANY)	
11.4	SIGNATURE	
11.5	HEAD OF DEPARTMENT	Dr. Milan Anand Principal Krupanidhi College of Physiotherapy
11.6	SIGNATURE	
12.1	REMARKS OF CHARIMAN AND PRINCIPAL	Approved and Forwarded
12.2	SIGNATURE	