

Two-Pass Method A: Authentic Native MISESERI Error Indicator Across Donor Deformation Stages
(a) Candidate B: $u_2 = 0.00537$ mm
(Step 2, Frame 50, Pre-Peak Inception)
Notch Tip Share: 62.2%
Secondary Lig. Share: 1.5%
Peak Error: 1657 MPa (Notch Tip)
(b) Candidate C: $u_2 = 0.00711$ mm
(Step 2, Frame 150, Primary Ligament Deflection)
Notch Tip Share: 56.9%
Secondary Lig. Share: 2.0%
Peak Error: 1601 MPa (Notch Tip)
(c) Candidate D: $u_2 = 0.00776$ mm
(Step 2, Frame 200, Hole Cavity Penetration)
Notch Tip Share: 3.0%
Secondary Lig. Share: 10.0%
Peak Error: 411 MPa (Pinned Corner)
(d) Terminal Baseline: $u_2 = 0.00959$ mm
(Step 2, Frame 337, Fully Softened Post-Severance)
Notch Tip Share: 0.3%
Secondary Lig. Share: 94.4%
Peak Error: 1437 MPa (Sec. Lig.)
x [mm]
y [mm]
Native MISESERI Stress Error η_{Mises} [MPa] (Log Scale)
10^0
10^1
10^2
10^3