

# Recurrence of substitutive Sturmian words

Pablo Rotondo

LIGM, Université Paris-Est Marne-la-Vallée

Joint work with  
Brigitte Vallée


**Journée de l'axe AlgoComb, Normastic,**  
**Caen**, 28 May, 2019.

# Context

## ► Probabilistic analysis

Object/experiment/execution?

⇒ Models, averages, distribution?



# Context

## ► Probabilistic analysis


Object/experiment/execution?

⇒ Models, averages, distribution?

## ► Word Combinatorics

Study of *words*

⇒ subwords (factors), frequencies



**Thue-Morse**

$\sigma: 0 \mapsto 01, 1 \mapsto 10$


01101001...

# Context

## ► Probabilistic analysis

Object/experiment/execution?

⇒ Models, averages, distribution?



## ► Word Combinatorics

Study of *words*

⇒ subwords (factors), frequencies

## ► Sturmian Words


simplest not eventually periodic.

⇒ recurrence: worst case, **average?**

**Thue-Morse**

$\sigma: 0 \mapsto 01, 1 \mapsto 10$

01101001...



# Plan of the talk

## 1. Sturmian words

- General Sturmian words
- Substitutive words

## 2. Recurrence function

- Definition and classical results
- Our models and results

## 3. Quadratic Irrational slope

- Model
- Main result

## 4. Toolbox for the proofs

## 5. Conclusion

# Complexity and Sturmian words

## Definition

**Complexity function** of an infinite word  $u \in \mathcal{A}^{\mathbb{N}}$

$$p_u: \mathbb{N} \rightarrow \mathbb{N}, \quad p_u(n) = \#\{\text{factors of length } n \text{ in } u\}.$$

# Complexity and Sturmian words

## Definition

**Complexity function** of an infinite word  $u \in \mathcal{A}^{\mathbb{N}}$

$$p_u: \mathbb{N} \rightarrow \mathbb{N}, \quad p_u(n) = \#\{\text{factors of length } n \text{ in } u\}.$$

## Important property

$u \in \mathcal{A}^{\mathbb{N}}$  is **not** eventually periodic

$$\iff p_u(n+1) > p_u(n) \text{ for all } n \in \mathbb{N}$$

.

# Complexity and Sturmian words

## Definition

**Complexity function** of an infinite word  $u \in \mathcal{A}^{\mathbb{N}}$

$$p_u: \mathbb{N} \rightarrow \mathbb{N}, \quad p_u(n) = \#\{\text{factors of length } n \text{ in } u\}.$$

## Important property

$u \in \mathcal{A}^{\mathbb{N}}$  is **not** eventually periodic

$$\iff p_u(n+1) > p_u(n) \text{ for all } n \in \mathbb{N}$$

$$\implies p_u(n) \geq n + 1.$$

Sturmian words are the “**simplest**” that are **not** eventually periodic.



# Complexity and Sturmian words

## Definition

**Complexity function** of an infinite word  $u \in \mathcal{A}^{\mathbb{N}}$

$$p_u: \mathbb{N} \rightarrow \mathbb{N}, \quad p_u(n) = \#\{\text{factors of length } n \text{ in } u\}.$$

## Important property

$u \in \mathcal{A}^{\mathbb{N}}$  is **not** eventually periodic

$$\iff p_u(n+1) > p_u(n) \text{ for all } n \in \mathbb{N}$$

$$\implies p_u(n) \geq n + 1.$$

Sturmian words are the “**simplest**” that are **not** eventually periodic.

## Definition

$u \in \{0, 1\}^{\mathbb{N}}$  is Sturmian  $\iff p_u(n) = n + 1$  for each  $n \geq 0$ .

# Explicit construction

Given  $\alpha, \beta \in [0, 1)$  we define

$$\underline{\mathfrak{S}}_{\alpha, \beta}(n) = \lfloor (n+1)\alpha + \beta \rfloor - \lfloor n\alpha + \beta \rfloor ,$$

$$\overline{\mathfrak{S}}_{\alpha, \beta}(n) = \lceil (n+1)\alpha + \beta \rceil - \lceil n\alpha + \beta \rceil ,$$

for  $n \geq 0$ .




Figure : Sequences  $\underline{\mathfrak{S}}_{\alpha, \beta}$  and  $\overline{\mathfrak{S}}_{\alpha, \beta}$  are discrete codings of  $y = \alpha x + \beta$ .

# Explicit construction

Given  $\alpha, \beta \in [0, 1)$  we define

$$\underline{\mathfrak{S}}_{\alpha, \beta}(n) = \lfloor (n+1)\alpha + \beta \rfloor - \lfloor n\alpha + \beta \rfloor ,$$

$$\overline{\mathfrak{S}}_{\alpha, \beta}(n) = \lceil (n+1)\alpha + \beta \rceil - \lceil n\alpha + \beta \rceil ,$$

for  $n \geq 0$ .




Figure : Sequences  $\underline{\mathfrak{S}}_{\alpha, \beta}$  and  $\overline{\mathfrak{S}}_{\alpha, \beta}$  are discrete codings of  $y = \alpha x + \beta$ .

## Theorem [Morse & Hedlund '40]

►  $u$  is Sturmian  $\iff$  there are  $\alpha, \beta \in [0, 1)$ ,  $\alpha$  irrational, such that

$$u_i = \underline{\mathfrak{S}}_{\alpha, \beta}(i), \quad \text{for all } i \geq 0, \quad \text{or } u_i = \overline{\mathfrak{S}}_{\alpha, \beta}(i), \quad \text{for all } i \geq 0.$$

# Substitutive words

## Definition (Substitutive word)

A word  $u$  is substitutive iff  $\sigma(u) = u$  for a primitive morphism  $\sigma$ .

**Primitivity:**  $\sigma$  is primitive iff the associated *matrix*  $M_\sigma$  is primitive

$$M_\sigma = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} |\sigma(0)|_0 & |\sigma(0)|_1 \\ |\sigma(1)|_0 & |\sigma(1)|_1 \end{bmatrix} \end{matrix}$$

# Substitutive words

## Definition (Substitutive word)

A word  $u$  is substitutive iff  $\sigma(u) = u$  for a primitive morphism  $\sigma$ .

**Primitivity:**  $\sigma$  is primitive iff the associated *matrix*  $M_\sigma$  is primitive

$$M_\sigma = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} |\sigma(0)|_0 & |\sigma(0)|_1 \\ |\sigma(1)|_0 & |\sigma(1)|_1 \end{bmatrix} \end{matrix}$$

## Example (Fibonacci word)

Consider the **primitive** morphism  $\sigma: 0 \mapsto 01, 1 \mapsto 0$

$$M_\sigma = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$

# Substitutive words

## Definition (Substitutive word)

A word  $u$  is substitutive iff  $\sigma(u) = u$  for a primitive morphism  $\sigma$ .

**Primitivity:**  $\sigma$  is primitive iff the associated *matrix*  $M_\sigma$  is primitive

$$M_\sigma = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} |\sigma(0)|_0 & |\sigma(0)|_1 \\ |\sigma(1)|_0 & |\sigma(1)|_1 \end{bmatrix} \end{matrix}$$

## Example (Fibonacci word)

Consider the **primitive** morphism  $\sigma: 0 \mapsto 01, 1 \mapsto 0$

$$M_\sigma = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$

its *fixed point*  $f_\infty$  can be constructed by iteration

$f_0 = 0, f_1 = 01, f_2 = 010, f_3 = 01001, \dots f_\infty = 0100101001001\dots$

## Substitutive *Sturmian* words

Slope  $\alpha$  of a *substitutive* Sturmian word is a quadratic irrational,

(1) Solution of a quadratic equation over  $\mathbb{Z}$  determined by  $M_\sigma$ .

## Substitutive *Sturmian* words

Slope  $\alpha$  of a *substitutive* Sturmian word is a **quadratic irrational**,

- (1) Solution of a quadratic equation over  $\mathbb{Z}$  determined by  $M_\sigma$ .
- (2) Eventually periodic **continued fraction expansion (CFE)**.



## Substitutive *Sturmian* words

Slope  $\alpha$  of a *substitutive* Sturmian word is a **quadratic irrational**,

- (1) Solution of a quadratic equation over  $\mathbb{Z}$  determined by  $M_\sigma$ .
- (2) Eventually periodic **continued fraction expansion (CFE)**.

**Reminder** for CFEs

$$\alpha = [m_1, m_2, \dots] := \frac{1}{m_1 + \frac{1}{m_2 + \ddots}}$$

where  $m_1, m_2, \dots \in \mathbb{Z}_{>0}$  are called the **quotients**.

# Substitutive *Sturmian* words

Slope  $\alpha$  of a *substitutive* Sturmian word is a **quadratic irrational**,

- (1) Solution of a quadratic equation over  $\mathbb{Z}$  determined by  $M_\sigma$ .
- (2) Eventually periodic **continued fraction expansion (CFE)**.

**Reminder** for CFEs

$$\alpha = [m_1, m_2, \dots] := \frac{1}{m_1 + \frac{1}{m_2 + \dots}}$$

where  $m_1, m_2, \dots \in \mathbb{Z}_{>0}$  are called the **quotients**.


## Theorem (Characterization by continued fractions)

The Sturmian word

$\underline{S}(\alpha, \alpha)$  is **substitutive**



$\alpha$  is **qi** and preperiod is of form given here.



# Recurrence

## Definition (Recurrence function)

Consider an infinite word  $u$ . Its recurrence function is:

$$R_u(n) = \inf \{m \in \mathbb{N} : \text{every factor of length } m \\ \text{contains all the factors of length } n\}.$$

# Recurrence

## Definition (Recurrence function)

Consider an infinite word  $\mathbf{u}$ . Its recurrence function is:

$$R_{\mathbf{u}}(n) = \inf \{m \in \mathbb{N} : \text{every factor of length } m \\ \text{contains all the factors of length } n\}.$$

- **Cost** we have to pay to **discover** the **factors** if we start from an arbitrary point in  $\mathbf{u} = u_1 u_2 \dots$

# Recurrence

## Definition (Recurrence function)

Consider an infinite word  $\mathbf{u}$ . Its recurrence function is:

$$R_{\mathbf{u}}(n) = \inf \{m \in \mathbb{N} : \text{every factor of length } m \\ \text{contains all the factors of length } n\}.$$

- ▶ **Cost** we have to pay to **discover** the **factors** if we start from an arbitrary point in  $\mathbf{u} = u_1u_2\dots$
- ▶ Inequality **relating** the functions,

$$R_{\mathbf{u}}(n) \geq n + p_{\mathbf{u}}(n) - 1.$$

# Recurrence

## Definition (Recurrence function)

Consider an infinite word  $u$ . Its recurrence function is:

$$R_u(n) = \inf \{m \in \mathbb{N} : \text{every factor of length } m \\ \text{contains all the factors of length } n\}.$$

- **Cost** we have to pay to **discover** the **factors** if we start from an arbitrary point in  $u = u_1u_2 \dots$
- Inequality **relating** the functions,

$$R_u(n) \geq \underbrace{n}_{\text{first factor}} + \underbrace{p_u(n) - 1}_{\substack{\text{count } +1 \\ \text{for every other factor}}}.$$

# Recurrence of Sturmian words: a link to arithmetic

Theorem (Morse, Hedlund, 1940)

*The recurrence function is piecewise affine and satisfies*

$$R_\alpha(n) = n - 1 + q_{k-1}(\alpha) + q_k(\alpha), \quad \text{for } q_{k-1}(\alpha) \leq n < q_k(\alpha).$$

Truncating the expansion at depth  $k$  we get a convergent

$$\frac{p_k(\alpha)}{q_k(\alpha)} = \frac{1}{m_1 + \frac{1}{m_2 + \cdots + \frac{1}{m_k}}}.$$

The denominators  $q_k(\alpha)$  are called the **continuants** of  $\alpha$ .


## Recurrence quotient

$$S(\alpha, n) := \frac{R_\alpha(n) + 1}{n} = 1 + \frac{q_{k-1}(\alpha) + q_k(\alpha)}{n}, \quad q_{k-1}(\alpha) \leq n < q_k(\alpha).$$




# Recurrence quotient

$$S(\alpha, n) := \frac{R_\alpha(n) + 1}{n} = 1 + \frac{q_{k-1}(\alpha) + q_k(\alpha)}{n}, \quad q_{k-1}(\alpha) \leq n < q_k(\alpha).$$



Recurrence quotient  $\alpha = e^{-1}$ .



Recurrence quotient  $\alpha = \phi^{-2}$ .

# Studies of the recurrence function

- Usual studies of  $R_{\alpha}(n)$  give
  - ▶ information about **extreme** cases.
  - ▶ results for **almost all**  $\alpha$ .

# Studies of the recurrence function

- Usual studies of  $R_\alpha(n)$  give
  - ▶ information about **extreme** cases.
  - ▶ results for **almost all**  $\alpha$ .

**Thm** (Morse & Hedlund '40)  $\forall \epsilon > 0$ , for almost every  $\alpha$

$$\limsup_{n \rightarrow \infty} \frac{S(\alpha, n)}{\log n} = \infty, \quad \lim_{n \rightarrow \infty} \frac{S(\alpha, n)}{(\log n)^{1+\epsilon}} = 0.$$

# Studies of the recurrence function

- Usual studies of  $R_\alpha(n)$  give
  - ▶ information about **extreme** cases.
  - ▶ results for **almost all**  $\alpha$ .

**Thm** (Morse & Hedlund '40)  $\forall \epsilon > 0$ , for almost every  $\alpha$

$$\limsup_{n \rightarrow \infty} \frac{S(\alpha, n)}{\log n} = \infty, \quad \lim_{n \rightarrow \infty} \frac{S(\alpha, n)}{(\log n)^{1+\epsilon}} = 0.$$

- In our **probabilistic** setting we
  - ▶ fix an integer  $n$  (we want  $n \rightarrow \infty \dots$ )
  - ▶ pick an irrational  $\alpha$  **uniformly at random** from
    - (1) the **“generic” reals** from  $[0, 1]$
    - (2) **quadratic irrationals** of **“size”**  $\leq D$  and let  $D \rightarrow \infty$ .

# Studies of the recurrence function

- Usual studies of  $R_\alpha(n)$  give
  - ▶ information about **extreme** cases.
  - ▶ results for **almost all**  $\alpha$ .

**Thm** (Morse & Hedlund '40)  $\forall \epsilon > 0$ , for almost every  $\alpha$

$$\limsup_{n \rightarrow \infty} \frac{S(\alpha, n)}{\log n} = \infty, \quad \lim_{n \rightarrow \infty} \frac{S(\alpha, n)}{(\log n)^{1+\epsilon}} = 0.$$

- In our **probabilistic** setting we
  - ▶ fix an integer  $n$  (we want  $n \rightarrow \infty \dots$ )
  - ▶ pick an irrational  $\alpha$  **uniformly at random** from
    - (1) the **“generic” reals** from  $[0, 1]$
    - (2) **quadratic irrationals** of **“size”**  $\leq D$  and let  $D \rightarrow \infty$ .
  - ▶ study **expectations**  $\mathbb{E}_\alpha[S(\alpha, n)]$ , **distributions**  $\mathbb{P}_\alpha(S(\alpha, n) \leq \lambda)$

## Theorem (uniform $\alpha \in (0, 1)$ , [R.,Vallée,17])

The random variable  $\alpha \mapsto S(\alpha, n)$  admits a limiting distribution

$$\lim_{n \rightarrow \infty} \mathbb{P}(\alpha : S(\alpha, n) \leq \lambda) = \int_{[2, \lambda]} g(y) dy ,$$

for  $\lambda \geq 2$  (and 0 otherwise), where the density  $g$  equals

$$g(\lambda) = \begin{cases} \frac{12}{\pi^2} \frac{1}{\lambda-1} \log(1 + \frac{\lambda-2}{1}) & \text{if } \lambda \in [2, 3] \\ \frac{12}{\pi^2} \frac{1}{\lambda-1} \log(1 + \frac{1}{\lambda-2}) & \text{if } \lambda \in [3, \infty) \end{cases} .$$




Figure : Histogram with  $\epsilon = 1/n$ .




Figure : Limit density.

For  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ ,

$$\begin{aligned} S(\alpha, n) &= 1 + \frac{q_{k-1}(\alpha) + q_k(\alpha)}{n} = 1 + \frac{q_k(\alpha)}{n} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)} + 1 \right) \\ &= f \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right), \end{aligned}$$

with

$$f(x, y) = 1 + y(1 + x), \quad (x, y) \in \mathcal{D} := \{(x, y) \in \mathbb{R}_{\geq 0} : xy \leq 1 < y\}.$$

For  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ ,

$$\begin{aligned} S(\alpha, n) &= 1 + \frac{q_{k-1}(\alpha) + q_k(\alpha)}{n} = 1 + \frac{q_k(\alpha)}{n} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)} + 1 \right) \\ &= f \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right), \end{aligned}$$

with


$$f(x, y) = 1 + y(1 + x), \quad (x, y) \in \mathcal{D} := \{(x, y) \in \mathbb{R}_{\geq 0} : xy \leq 1 < y\}.$$

Theorem (uniform  $\alpha \in (0, 1)$ , [R., Vallée, 17])


Limit distribution for  $\alpha \mapsto S(\alpha, n)$  (+ more general class) given by

$$\lim_{n \rightarrow \infty} \mathbb{P}(\alpha : S(\alpha, n) \leq \lambda) = \frac{6}{\pi^2} \iint_{\mathcal{D}_\lambda} \omega(x, y) dx dy,$$

$$\mathcal{D}_\lambda = \{(x, y) \in \mathcal{D} : f(x, y) \leq \lambda\}, \quad \omega(x, y) = \frac{1}{y(1+x)}.$$



The domain  $\mathcal{D}_\lambda$   
 $\lambda = 2.5, \quad \lambda = 3.5.$





# The index of tours $\ell(\alpha, n)$

## **Simplifying assumptions for the talk.**

- ▶ Slopes  $\alpha$  that are reduced quadratic irrationals, i.e., corresponding to purely periodic expansions.
- ▶ Periods may be *primitive* or not. Here we omit this detail.

# The index of tours $\ell(\alpha, n)$

## Simplifying assumptions for the talk.

- ▶ Slopes  $\alpha$  that are reduced quadratic irrationals, i.e., corresponding to purely periodic expansions.
- ▶ Periods may be *primitive* or not. Here we omit this detail.

Thus fix  $\alpha = \left[ \overline{m_1, \dots, m_p} \right]$ , i.e., *period*  $(m_1, \dots, m_p)$ .

# The index of tours $\ell(\alpha, n)$

## Simplifying assumptions for the talk.

- ▶ Slopes  $\alpha$  that are reduced quadratic irrationals, i.e., corresponding to purely periodic expansions.
- ▶ Periods may be *primitive* or not. Here we omit this detail.

Thus fix  $\alpha = \left[ \overline{m_1, \dots, m_p} \right]$ , i.e., **period**  $(m_1, \dots, m_p)$ .

## Definition ( $\ell$ -th tour)

The interval

$$I_\ell(\alpha) := \left( q_{\ell p}(\alpha), q_{(\ell+1)p}(\alpha) \right]$$

is called the  $\ell$ -th tour of  $\alpha$ , and

$$q_{\ell p+1}(\alpha), \dots, q_{(\ell+1)p}(\alpha)$$

are said to be the continuants of the  $\ell$ -th tour.

## Theorem

Fix  $\alpha = [m_1, \dots, m_p]$ , i.e., *period*  $(m_1, \dots, m_p)$ .

Then for every fixed  $r$  the following limit exists

$$Q_r(\alpha) := \lim_{\ell \rightarrow \infty} \frac{q_{\ell p + r}(\alpha)}{q_{\ell p}(\alpha)},$$

furthermore convergence is exponential in  $\ell$ .

## Theorem

Fix  $\alpha = [m_1, \dots, m_p]$ , i.e., *period*  $(m_1, \dots, m_p)$ .

Then for every fixed  $r$  the following limit exists

$$Q_r(\alpha) := \lim_{\ell \rightarrow \infty} \frac{q_{\ell p + r}(\alpha)}{q_{\ell p}(\alpha)},$$

furthermore convergence is exponential in  $\ell$ .

$\implies$  Define the size of  $\alpha$  by  $\varrho(\alpha) := Q_p(\alpha)$ .

## Theorem

Fix  $\alpha = [m_1, \dots, m_p]$ , i.e., *period*  $(m_1, \dots, m_p)$ .

Then for every fixed  $r$  the following limit exists

$$Q_r(\alpha) := \lim_{\ell \rightarrow \infty} \frac{q_{\ell p + r}(\alpha)}{q_{\ell p}(\alpha)},$$

furthermore convergence is exponential in  $\ell$ .

$\implies$  Define the size of  $\alpha$  by  $\varrho(\alpha) := Q_p(\alpha)$ .




Figure : Logarithmic plot of the recurrence quotient  $S(\alpha, n)$  for  $\alpha = [3, 3, 3, 1, 1] = \frac{5\sqrt{317}-63}{86}$

# Model for quadratic irrationals

Quadratic irrationals present two *striking features*

- ▶ **Countable** and dense subset of  $[0, 1]$ .
- ▶ **Periodic structure** (after re-scaling) with respect to tours.

# Model for quadratic irrationals

Quadratic irrationals present two *striking features*

- ▶ Countable and dense subset of  $[0, 1]$ .
- ▶ Periodic structure (after re-scaling) with respect to tours.

Model takes these into account

- ▶ Pick  $\alpha$  uniformly at random from the finite

$$\mathcal{S}_D := \left\{ \alpha \text{ quadratic irrational} : \varrho(\alpha) \leq D \right\},$$



# Model for quadratic irrationals

Quadratic irrationals present two *striking features*

- ▶ **Countable** and dense subset of  $[0, 1]$ .
- ▶ **Periodic structure** (after re-scaling) with respect to tours.

**Model** takes these into account

- ▶ Pick  $\alpha$  **uniformly at random** from the **finite**

$$\mathcal{S}_D := \left\{ \alpha \text{ quadratic irrational} : \varrho(\alpha) \leq D \right\},$$

- ▶ **Restriction** to  $\ell$ -th tour  $\Gamma_\ell(\alpha)$

$$S_\ell(\alpha, n) = \llbracket n \in \Gamma_\ell(\alpha) \rrbracket S(\alpha, n).$$

# Main result for substitutive Sturmian words

For **quadratic irrationals**

probabilities are **discrete** and defined from

$$R_D(\ell, \lambda) := \left\{ (\alpha, n) : \alpha \in \mathcal{S}_D, n \in \Gamma_\ell(\alpha), S(\alpha, n) \leq \lambda \right\},$$

Main result (?) (R., Vallée, 19)

**Limit distribution** for  $\alpha \mapsto S(\alpha, n)$  over **quadratic irrationals**

$$\lim_{D, u, \ell \rightarrow \infty} \frac{\left| R_D(\ell, \lambda) \cap \left\{ \frac{n}{q_{\ell p}} \in u \cdot (1, \theta) \right\} \right|}{|\mathcal{S}_D| \cdot u \cdot (\theta - 1)} = \frac{6}{\pi^2} \iint_{\mathcal{D}_\lambda} \omega(x, y) dx dy,$$

$$\mathcal{D}_\lambda = \{(x, y) \in \mathcal{D} : f(x, y) \leq \lambda\}, \quad \omega(x, y) = \frac{1}{y(1+x)}.$$

## An analytic “dictionary”

A **prefix**  $(m_1, \dots, m_k)$  of the **CFE** defines an homography  $g \in \mathcal{H}^k$

$$g(x) := \frac{1}{m_1 + \frac{1}{\ddots + \frac{1}{m_k + x}}}$$

associated with an **operator**, its **generating function**,

$$\mathbf{H}_{[g],s}[f](x) := |g'(x)|^{s/2} f(g(x)) .$$

# An analytic “dictionary”

A **prefix**  $(m_1, \dots, m_k)$  of the **CFE** defines an homography  $g \in \mathcal{H}^k$

$$g(x) := \frac{1}{m_1 + \frac{1}{\ddots + \frac{1}{m_k + x}}}$$

associated with an **operator**, its **generating function**,

$$\mathbf{H}_{[g],s}[f](x) := |g'(x)|^{s/2} f(g(x)) .$$

**Generating functions.**

- $\mathbf{H}_s := \sum_{g \in \mathcal{H}} \mathbf{H}_{[g],s}$  describes all prefixes of depth 1.

# An analytic “dictionary”

A **prefix**  $(m_1, \dots, m_k)$  of the **CFE** defines an homography  $g \in \mathcal{H}^k$

$$g(x) := \frac{1}{m_1 + \frac{1}{\ddots + \frac{1}{m_k + x}}}$$

associated with an **operator**, its **generating function**,

$$\mathbf{H}_{[g],s}[f](x) := |g'(x)|^{s/2} f(g(x)).$$

## Generating functions.

- ▶  $\mathbf{H}_s := \sum_{g \in \mathcal{H}} \mathbf{H}_{[g],s}$  describes all prefixes of depth 1.
- ▶  $\mathbf{H}_s^k = \mathbf{H}_s \circ \dots \circ \mathbf{H}_s$  describes all prefixes of depth  $k$ .

# An analytic “dictionary”

A **prefix**  $(m_1, \dots, m_k)$  of the **CFE** defines an homography  $g \in \mathcal{H}^k$

$$g(x) := \frac{1}{m_1 + \frac{1}{\ddots + \frac{1}{m_k + x}}}$$

associated with an **operator**, its **generating function**,

$$\mathbf{H}_{[g],s}[f](x) := |g'(x)|^{s/2} f(g(x)).$$

## Generating functions.

- ▶  $\mathbf{H}_s := \sum_{g \in \mathcal{H}} \mathbf{H}_{[g],s}$  describes all prefixes of depth 1.
- ▶  $\mathbf{H}_s^k = \mathbf{H}_s \circ \dots \circ \mathbf{H}_s$  describes all prefixes of depth  $k$ .
- ▶ and  $(\mathbf{I} - \mathbf{H}_s)^{-1} = \mathbf{I} + \mathbf{H}_s + \mathbf{H}_s^2 + \dots$  describes **all prefixes**.

## Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

# Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

**Key properties:**

- **Singularities** of Mellin transform **transfer** into limits.



# Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

**Key properties:**

- **Singularities** of Mellin transform **transfer** into limits.

$$f(u) \sim \operatorname{Res}[f^*(\rho); s = \sigma] \cdot u^{-\sigma} (1 + O(u^a)) , \quad u \rightarrow 0^+ .$$

# Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

**Key properties:**

- **Singularities** of Mellin transform **transfer** into limits.

$$f(u) \sim \operatorname{Res}[f^*(\rho); s = \sigma] \cdot u^{-\sigma} (1 + O(u^a)) , \quad u \rightarrow 0^+ .$$

- **Harmonic sums** produce **generating functions** of *Dirichlet* type

# Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

**Key properties:**

- **Singularities** of Mellin transform **transfer** into limits.

$$f(u) \sim \operatorname{Res}[f^*(\rho); s = \sigma] \cdot u^{-\sigma} (1 + O(u^a)) , \quad u \rightarrow 0^+ .$$

- **Harmonic sums** produce **generating functions** of *Dirichlet* type

$$G(x) := \sum \lambda_k g(\mu_k x) \implies G^*(\rho) = \left( \sum \lambda_k \mu_k^{-\rho} \right) \cdot g^*(\rho) .$$

# Principles of the proofs

**Mellin transform** of  $f: [0, \infty) \rightarrow \mathbb{C}$  is defined by

$$f^*(\rho) := \int_0^\infty f(u) u^{\rho-1} du.$$

**Key properties:**

- **Singularities** of Mellin transform **transfer** into limits.

$$f(u) \sim \text{Res}[f^*(\rho); s = \sigma] \cdot u^{-\sigma} (1 + O(u^a)) , \quad u \rightarrow 0^+ .$$

- **Harmonic sums** produce **generating functions** of *Dirichlet* type

$$G(x) := \sum \lambda_k g(\mu_k x) \implies G^*(\rho) = \left( \sum \lambda_k \mu_k^{-\rho} \right) \cdot g^*(\rho) .$$

$\Rightarrow$  For **generic**  $\alpha$ :

Mellin transform of distribution yields  $(\mathbf{I} - \mathbf{H}_{\rho/2+1})^{-1} [G_\rho](0)$ .

# Principles of the proofs

For *quadratic irrational*  $\alpha$ , additional steps:

- ▶  $\alpha$  is **fixed point**  $x_g^* \in (0, 1)$

$$x_g^* = g(x_g^*) = \frac{1}{m_1 + \frac{1}{\ddots + \frac{1}{m_k + x_g^*}}}$$

of some  $g \in \mathcal{H}^*$ .

- ▶ generating function related to *trace* of operators

$$f \mapsto (\mathbf{I} - \mathbf{H}_{s/2})^{-1} [L_{\lambda, \rho} \cdot (\mathbf{I} - \mathbf{H}_{(s+\rho)/2})^{-1} [f]] .$$

## Final remarks and conclusion

- ▶ Strong parallels between the respective results and methods.

## Final remarks and conclusion

- ▶ Strong parallels between the respective results and methods.
- ▶ Results hold more generally for **Q-functions**, i.e.,

$$\Lambda(\alpha, n) = f_{\Lambda} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right)$$

for  $k = k(\alpha, n)$  such that  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ .

## Final remarks and conclusion

- ▶ Strong parallels between the respective results and methods.
- ▶ Results hold more generally for **Q-functions**, i.e.,

$$\Lambda(\alpha, n) = f_{\Lambda} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right)$$

for  $k = k(\alpha, n)$  such that  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ .

- ▶ For generic  $\alpha$  model: the density on  $\alpha$  can be more general.



## Final remarks and conclusion

- ▶ Strong parallels between the respective results and methods.
- ▶ Results hold more generally for **Q-functions**, i.e.,

$$\Lambda(\alpha, n) = f_{\Lambda} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right)$$

for  $k = k(\alpha, n)$  such that  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ .

- ▶ For generic  $\alpha$  model: the density on  $\alpha$  can be more general.  
 $\Rightarrow$  **Asymptotic independence** between  $p_k/q_k$  and  $q_{k-1}/q_k$ .

## Final remarks and conclusion

- ▶ Strong parallels between the respective results and methods.
- ▶ Results hold more generally for **Q-functions**, i.e.,

$$\Lambda(\alpha, n) = f_{\Lambda} \left( \frac{q_{k-1}(\alpha)}{q_k(\alpha)}, \frac{q_k(\alpha)}{n} \right)$$

for  $k = k(\alpha, n)$  such that  $q_{k-1}(\alpha) \leq n < q_k(\alpha)$ .

- ▶ For generic  $\alpha$  model: the density on  $\alpha$  can be more general.  
 $\Rightarrow$  **Asymptotic independence** between  $p_k/q_k$  and  $q_{k-1}/q_k$ .
- ▶ Multidimensional analogs?

Thank you!