

EXPLAINING EXTREME EVENTS OF 2016

From A Climate Perspective



Special Supplement to the
Bulletin of the American Meteorological Society
Vol. 99, No. 1, January 2018

EXPLAINING EXTREME EVENTS OF 2016 FROM A CLIMATE PERSPECTIVE

Editors

Stephanie C. Herring, Nikolaos Christidis, Andrew Hoell, James P. Kossin,
Carl J. Schreck III, and Peter A. Stott

Special Supplement to the

Bulletin of the American Meteorological Society

Vol. 99, No. 1, January 2018

AMERICAN METEOROLOGICAL SOCIETY

CORRESPONDING EDITOR:

Stephanie C. Herring, PhD
NOAA National Centers for Environmental Information
325 Broadway, E/CC23, Rm 1B-131
Boulder, CO 80305-3328
E-mail: stephanie.herring@noaa.gov

COVER CREDIT:

©The Ocean Agency / XL Catlin Seaview Survey / Chrisophe Bailhache—A panoramic image of coral bleaching at Lizard Island on the Great Barrier Reef, captured by The Ocean Agency / XL Catlin Seaview Survey / Christophe Bailhache in March 2016.

HOW TO CITE THIS DOCUMENT

Citing the complete report:

Herring, S. C., N. Christidis, A. Hoell, J. P. Kossin, C. J. Schreck III, and P. A. Stott, Eds., 2018: Explaining Extreme Events of 2016 from a Climate Perspective. *Bull. Amer. Meteor. Soc.*, **99** (I), SI–SI57.

Citing a section (example):

Quan, X.W., M. Hoerling, L. Smith, J. Perlwitz, T. Zhang, A. Hoell, K. Wolter, and J. Eischeid, 2018: Extreme California Rains During Winter 2015/16: A Change in El Niño Teleconnection? [in “Explaining Extreme Events of 2016 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **99** (I), S54–S59, doi:10.1175/BAMS-D-17-0118.1.

EDITORIAL AND PRODUCTION TEAM

Riddle, Deborah B., Lead Graphics Production, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Love-Brotak, S. Elizabeth, Graphics Support, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Veasey, Sara W., Visual Communications Team Lead, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Fulford, Jennifer, Editorial Support, Telesolv Consulting LLC, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Griffin, Jessica, Graphics Support, Cooperative Institute for Climate and Satellites-NC, North Carolina State University, Asheville, NC

Misch, Deborah J., Graphics Support, Telesolv Consulting LLC, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Osborne, Susan, Editorial Support, Telesolv Consulting LLC, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Sprain, Mara, Editorial Support, LAC Group, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

Young, Teresa, Graphics Support, Telesolv Consulting LLC, NOAA/NESDIS National Centers for Environmental Information, Asheville, NC

TABLE OF CONTENTS

Abstract.....	ii
1. Introduction to Explaining Extreme Events of 2016 from a Climate Perspective	1
2. Explaining Extreme Ocean Conditions Impacting Living Marine Resources	7
3. CMIP5 Model-based Assessment of Anthropogenic Influence on Record Global Warmth During 2016.....	11
4. The Extreme 2015/16 El Niño, in the Context of Historical Climate Variability and Change	16
5. Ecological Impacts of the 2015/16 El Niño in the Central Equatorial Pacific	21
6. Forcing of Multiyear Extreme Ocean Temperatures that Impacted California Current Living Marine Resources in 2016	27
7. CMIP5 Model-based Assessment of Anthropogenic Influence on Highly Anomalous Arctic Warmth During November–December 2016.....	34
8. The High Latitude Marine Heat Wave of 2016 and Its Impacts on Alaska.....	39
9. Anthropogenic and Natural Influences on Record 2016 Marine Heat waves.....	44
10. Extreme California Rains During Winter 2015/16: A Change in El Niño Teleconnection?....	49
11. Was the January 2016 Mid-Atlantic Snowstorm "Jonas" Symptomatic of Climate Change?...54	
12. Anthropogenic Forcings and Associated Changes in Fire Risk in Western North America and Australia During 2015/16.....	60
13. A Multimethod Attribution Analysis of the Prolonged Northeast Brazil Hydrometeorological Drought (2012–16).....	65
14. Attribution of Wintertime Anticyclonic Stagnation Contributing to Air Pollution in Western Europe.....	70
15. Analysis of the Exceptionally Warm December 2015 in France Using Flow Analogues.....	76
16. Warm Winter, Wet Spring, and an Extreme Response in Ecosystem Functioning on the Iberian Peninsula	80
17. Anthropogenic Intensification of Southern African Flash Droughts as Exemplified by the 2015/16 Season	86
18. Anthropogenic Enhancement of Moderate-to-Strong El Niño Events Likely Contributed to Drought and Poor Harvests in Southern Africa During 2016	91
19. Climate Change Increased the Likelihood of the 2016 Heat Extremes in Asia	97
20. Extreme Rainfall (R20mm, RX5day) in Yangtze–Huai, China, in June–July 2016: The Role of ENSO and Anthropogenic Climate Change.....	102
21. Attribution of the July 2016 Extreme Precipitation Event Over China's Wuhang	107
22. Do Climate Change and El Niño Increase Likelihood of Yangtze River Extreme Rainfall?....113	
23. Human Influence on the Record-breaking Cold Event in January of 2016 in Eastern China.....	118
24. Anthropogenic Influence on the Eastern China 2016 Super Cold Surge.....	123
25. The Hot and Dry April of 2016 in Thailand.....	128
26. The Effect of Increasing CO ₂ on the Extreme September 2016 Rainfall Across Southeastern Australia.....	133
27. Natural Variability Not Climate Change Drove the Record Wet Winter in Southeast Australia	139
28. A Multifactor Risk Analysis of the Record 2016 Great Barrier Reef Bleaching	144
29. Severe Frosts in Western Australia in September 2016.....	150
30. Future Challenges in Event Attribution Methodologies.....	155

ABSTRACT—Stephanie C. Herring, Nikolaos Christidi, Andrew Hoell, James P. Kossin, Carl J. Schreck III, and Peter A. Stott

This sixth edition of explaining extreme events of the previous year (2016) from a climate perspective is the first of these reports to find that some extreme events were not possible in a preindustrial climate. The events were the 2016 record global heat, the heat across Asia, as well as a marine heat wave off the coast of Alaska. While these results are novel, they were not unexpected. Climate attribution scientists have been predicting that eventually the influence of human-caused climate change would become sufficiently strong as to push events beyond the bounds of natural variability alone. It was also predicted that we would first observe this phenomenon for heat events where the climate change influence is most pronounced. Additional retrospective analysis will reveal if, in fact, these are the first events of their kind or were simply some of the first to be discovered.

Last year, the editors emphasized the need for additional papers in the area of “impacts attribution” that investigate whether climate change’s influence on the extreme event can subsequently be directly tied to a change in risk of the socio-economic or environmental impacts. Several papers in this year’s report address this challenge, including Great Barrier Reef bleaching, living marine resources in the Pacific, and ecosystem productivity on the Iberian Peninsula. This is an increase over the number of impact attribution papers than in the past, and are hopefully a sign that research in this area will continue to expand in the future.

Other extreme weather event types in this year’s edition include ocean heat waves, forest fires, snow storms, and frost, as well as heavy precipitation, drought, and extreme heat and cold events over land. There were

a number of marine heat waves examined in this year’s report, and all but one found a role for climate change in increasing the severity of the events. While human-caused climate change caused China’s cold winter to be less likely, it did not influence U.S. storm Jonas which hit the mid-Atlantic in winter 2016.

As in past years, the papers submitted to this report are selected prior to knowing the final results of whether human-caused climate change influenced the event. The editors have and will continue to support the publication of papers that find no role for human-caused climate change because of their scientific value in both assessing attribution methodologies and in enhancing our understanding of how climate change is, and is not, impacting extremes. In this report, twenty-one of the twenty-seven papers in this edition identified climate change as a significant driver of an event, while six did not. Of the 131 papers now examined in this report over the last six years, approximately 65% have identified a role for climate change, while about 35% have not found an appreciable effect.

Looking ahead, we hope to continue to see improvements in how we assess the influence of human-induced climate change on extremes and the continued inclusion of stakeholder needs to inform the growth of the field and how the results can be applied in decision making. While it represents a considerable challenge to provide robust results that are clearly communicated for stakeholders to use as part of their decision-making processes, these annual reports are increasingly showing their potential to help meet such growing needs.

6. FORCING OF MULTIYEAR EXTREME OCEAN TEMPERATURES THAT IMPACTED CALIFORNIA CURRENT LIVING MARINE RESOURCES IN 2016

MICHAEL G. JACOX, MICHAEL A. ALEXANDER, NATHAN J. MANTUA, JAMES D. SCOTT, GAELLE HERVIEUX,
ROBERT S. WEBB, AND FRANCISCO E. WERNER

Significant impacts on California Current living marine resources in 2016 resulted from sustained extremely high ocean temperatures forced by a confluence of natural drivers and likely exacerbated by anthropogenic warming.

Introduction. Recent record high sea surface temperature anomalies (SSTa) in the California Current Large Marine Ecosystem (CCLME; Fig. 6.1a) produced dramatic impacts on marine life (Cavole et al. 2016; Peterson et al. 2016; Welch 2016). While effects on many species and fisheries may have been short-lived, salmon fisheries, for example, were heavily impacted in 2016 due to multiyear persistence of unfavorable conditions. Negative impacts on CCLME salmon fisheries are likely to persist until at least 2019, as poor stream and 2014–16 ocean conditions directly influence the 2016–19 Chinook salmon abundance. U.S. West Coast Chinook salmon catches in 2016 were approximately 52% of the average catch since 2006, quotas for Chinook salmon fisheries were not met, and spawning escapements to the Klamath and Sacramento River basins were very low (PFMC 2017a). For 2017, the Klamath River Chinook salmon abundance forecast is the lowest on record, and salmon fishing has been sharply restricted from southern Oregon to southern California (PFMC 2017b).

Our analysis focuses on the climatic drivers of the 2014–16 CCLME warm period and its extremity in the context of the past century. This study is motivated by an important question from a fisheries management

perspective: to what extent were the 2014–16 extremes due to natural variability versus anthropogenic climate change?

Temperature impacts on salmon. Salmon are a subarctic species that thrive in marine habitats featuring lipid-rich food-webs with cool-water plankton and fish communities. Warm periods in the CCLME are characterized by sharp reductions in cool, nutrient-rich, highly productive upwelled and subarctic water (Chavez et al. 2002; Checkley and Barth 2009), a shift from lipid-rich to lipid-poor zooplankton (Peterson and Schwings 2003), and an influx of predators to the nearshore areas critical for salmon early marine survival (e.g., Pearcy 1992; Wells et al. 2017). These shifts in the prey base and predator distributions favor reduced growth and survival rates for CCLME salmon (e.g., Daly et al. 2017), and anomalously warm CCLME SSTs are associated with low post-release survival rates for hatchery-origin coho and Chinook salmon from southeast Alaska to California (Sharma et al. 2012; Kilduff et al. 2015). While links between salmon abundance (or catch) and SST are not easily evaluated with time series correlations (see online supplement material), a strong link between record-warm 2014–16 CCLME SSTs and negative impacts on the West Coast salmon fishery in 2016 is evidenced by a shift to subtropical species and widespread negative impacts (increased mortality rates, reduced reproductive success and/or abundance) on top predators like sea birds, salmon, and marine mammals that typically thrive under neutral or cool SST conditions (Cavole et al. 2016; Peterson et al. 2016; Welch 2016).

Data and methods. For 1920–2016 CCLME SST observations, we used the 1° Hadley Centre Sea Ice and Sea

AFFILIATIONS: JACOX—Institute of Marine Sciences, University of California, Santa Cruz, and NOAA Southwest Fisheries Science Center, Environmental Research Division, Monterey, California; ALEXANDER AND WEBB—NOAA Earth System Research Laboratory, Physical Sciences Division, Boulder, Colorado; MANTUA—NOAA Southwest Fisheries Science Center, Fisheries Ecology Division, Santa Cruz, California; SCOTT AND HERVIEUX—NOAA Earth System Research Laboratory, Physical Sciences Division, and Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado; WERNER—NOAA Fisheries, Silver Spring, Maryland

DOI:10.1175/BAMS-D-17-0119.1

A supplement to this article is available online (10.1175/BAMS-D-17-0119.2)




FIG. 6.1. (a) Observed (HadISST) 2014–16 mean northeast Pacific SSTa ($^{\circ}$ C) relative to the 1920–2016 mean. Black line outlines the CCLME. (b),(c) CCLME annual mean SSTa ($^{\circ}$ C) from HadISST (black line), model ensemble mean (red line), and range of individual ensemble members in percentiles (gray shading): 25%–75% (dark), 10%–90% (medium) and 0–100% (light). (d),(e) Smoothed histograms of CCLME annual mean SSTa ($^{\circ}$ C) for 1920–2016 observations (black) and from all model ensemble members during 1920–50 (green), 1920–2016 (blue) and 2000–30 (red). Histograms were calculated using a SSTa bin width of 0.2° C. Model values are from (b),(d) CESM-LENS and (c),(e) CMIP5. Observed annual mean CCLME SSTa in $^{\circ}$ C (and standardized units) were 1.15° C (2.2σ), 1.71° C (3.3σ), and 0.95° C (1.8σ) in 2014, 2015, and 2016, respectively.

Surface Temperature (HadISST; Rayner et al. 2003) dataset. For spatial SST_a correlation analyses we used the 1982–present, higher resolution (0.25°) NOAA Optimum Interpolation Sea Surface Temperature, version 2 (Banzon et al. 2016; Reynolds et al. 2007).

Anthropogenic forcing contributions to extreme warming were assessed using SST_a distributions from “historical” (1920–50) and “present” (2000–30) periods in the Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor et al. 2012) ensemble and the Community Earth System Model Large Ensemble Project (CESM-LENS; Kay et al. 2015). We used 26 and 30 members from the CMIP5 and CESM-LENS ensembles, respectively. For each ensemble, historical external forcing was applied until 2005, after which representative concentration pathway 8.5 (RCP8.5) external forcing was applied (Lamarque et al. 2010, 2011) to provide continuous simulations of the twentieth and twenty-first centuries. The change in risk of an extreme event due to anthropogenic forcing is estimated using the fraction of attributable risk, FAR = $1 - (P_0/P_1)$, where P_0 is the probability of an event in the historical period and P_1 is the probability of the same event in the present period (Stott et al. 2004).

Forcing of CCLME SST_a is explored using first-order auto-regressive [AR(1)] models of the form

$$\text{SST}_{\text{t}} = a^* \text{SST}_{\text{t}-1} + b_i^* F_i + \varepsilon_{\text{t}}$$

where a is the lag-1 autoregression coefficient such that $a^* \text{SST}_{\text{t}-1}$ represents damped persistence, F_i are forcing functions and their regression coefficients, respectively, and ε is a residual error term (noise).

CCLME SST anomalies in the context of variability and change. The 1920–2016 distribution of observed annual mean CCLME SST_a is positively skewed, with more extreme warm anomalies than cold, and the 2014–16 values in the tail of the distribution (Fig. 6.1d). The CESM-LENS and CMIP5 distributions are nearly Gaussian and generally match the observed histogram (Figs. 6.1d,e), with the observed record 2015 SST_a near the upper bound of both ensembles (Figs. 6.1b,c). From the historical period (1920–50) to the present period (2000–30), increases in the ensemble mean (standard deviation) of SST_a are 0.47°C (0.10°C) in CESM-LENS and 0.65°C (0.06°C) in CMIP5. There is a long-term tendency for warmer SST_a to occur later in the observed record, although it is unclear if the increase is linear (Johnstone and Mantua 2014). Indeed, simulated CCLME SST_a exhibit little trend from 1920 to ~2000, after which they increase rapidly (Figs. 6.1b,c), similar to nonlinear changes that emerge for

coastal upwelling in the CCLME (Brady et al. 2017).

In 2015, the observed annual mean CCLME SST_a was 1.7°C , or 3.3 standard deviations (σ) above the mean, the highest value in the 1920–2016 record (Fig. 6.1b). The persistence of this heat wave was also remarkable; 2014–16 was the warmest 3-year period on record, with mean SST_a of 1.3°C , 3.1σ above the mean of all 3-year periods from 1920–2016 (Fig. ES6.1). The annual and three-year mean SST_a observed in 2015 and 2014–16 are never reached in the historical period for either CMIP5 or CESM-LENS (~1700 total simulated years under 1920–50 external forcing). In the “present” period, mean 2015 SST_a and 2014–16 SST_a occur approximately 2%–4% and 7%–9% of the time, respectively (Table ES6.1). Therefore, for these events, FAR = 1. However, one must take care when interpreting FAR as over shorter periods it can be influenced by natural variability; we discuss this variability in the next section.

Forcing of SST anomalies in the CCLME. Bond et al. (2015) showed that record SST_a in the Gulf of Alaska (GOA) in 2014 were caused by a persistent ridge of high sea level pressure anomalies that reduced surface wind speeds and weakened normal cooling processes over the 2013/14 winter. In 2015, northeast Pacific SST extremes expanded to include an area encompassing Alaska to Baja California (Gentemann et al. 2017). Di Lorenzo and Mantua (2016; hereafter DM2016) showed this persistent marine heatwave was a consequence of two atmospheric forcing/ocean response patterns, the 2014 GOA pattern and the 2015 northeast Pacific Arc pattern, linked with ENSO via teleconnections.

We examined the forcing of CCLME SST anomalies using AR(1) models in which observed SST_a derive from damped persistence of pre-existing anomalies plus some forcing. As the CCLME is a coastal upwelling system, the alongshore wind is a dominant forcing via mechanisms that include coastal and offshore upwelling, horizontal advection, and surface heat fluxes (Johnstone and Mantua 2014). The SST_a tendency has maximum correlations with meridional wind stress off the coasts of California and Baja California, from $\sim 130^{\circ}$ to 140°W (Fig. 6.2a). An AR(1) model forced by this index of local atmospheric forcing reproduces much of the observed SST_a variance (Fig. 6.2b). However, it fails to reproduce the extreme 2014–16 warming.

A lag-correlation analysis of residuals from the AR(1) model suggests an important influence of GOA SST_a at lead times of ~6 months (Fig. 6.2c), and




FIG. 6.2. Forcing of SST anomalies ($^{\circ}\text{C}$) in the CCLME. (a) Correlation of meridional wind stress and SST_a tendency. Black contour outlines the CCLME, white contour bounds the region of highest correlation ($r > 0.6$), which was used to force the AR(1) model shown in (b). **(c)** Residual of the AR(1) model in (b) correlated with SST_a 6 months prior suggests GOA (gray box) SST_a as a precursor to CCLME SST_a (white contour bounds $r > 0.5$). **(d)** AR(1) model with 6-month lead SST_a in the GOA added as a second forcing term. **(e)** Residual of the AR(1) model in (d) correlated with basin-wide SST_a suggests a role for ENSO variability. **(f)** AR(1) model with the EqSOI added as a third forcing term.

inclusion of 6-month lead GOA SSTa dramatically improves the AR(1) model, particularly in 2014. While a mechanistic evaluation of the GOA influence on the CCLME is beyond the scope of this paper, this finding is consistent with a tendency found in the historical record for warm GOA SSTa to evolve into an Arc pattern warming the following year (DM2016).

Correlating residuals from our second AR(1) model with basin-wide SSTa produces a spatial pattern that implicates ENSO variability (Fig. 6.2e). While one pathway for ENSO forcing is through the alongshore wind (Alexander et al. 2002; Jacox et al. 2015), which is already in our model, ENSO may impart additional variance through coastal trapped waves or anomalous poleward advection. Inclusion of the NOAA/CPC Equatorial Southern Oscillation Index (EqSOI) only modestly improves the overall performance of the AR(1) model, but improvements are visible for calendar years impacted by strong El Niños (Fig. 6.2f). The influence of the 2015/16 El Niño is visible in our AR(1) model, though its timing was earlier (a larger ENSO influence in 2015 than 2016; Frishknecht et al. 2017) and its impact on the CCLME weaker (Jacox et al. 2016) than common ENSO indices (e.g., Niño3.4) suggest.

Discussion. While a FAR calculation of the 2014–16 CCLME SSTa suggests an important role for anthropogenic warming, over shorter timescales the FAR is also influenced by natural internal variability, especially in the CCLME (Weller et al. 2015). The analysis outlined in Fig. 6.2 suggests roles for multiple drivers of SSTa in the CCLME, specifically atmospheric variability off the North American west coast, a lagged response to GOA SSTa, and ENSO teleconnections impacting the CCS. While these forcing mechanisms share some variance ($r = 0.4\text{--}0.6$), the 2014–16 period had notably strong and sustained forcing from all three (Fig. ES6.2). The superposition of multiple drivers (weakened poleward winds, an extremely warm GOA, and El Niño) contributed heavily to the CCLME anomalies, and additional mechanisms are also likely at play [e.g., reemergence, where anomalies are sequestered beneath the mixed layer in spring/summer and reemerge when the mixed layer deepens in winter (Fig. ES6.3)]. Nonetheless, climate model ensembles suggest that anthropogenic warming increased the likelihood of the 2014–16 SST extremes through both a shift to a warmer mean state and an increase in temperature variability (Fig. 6.1; DM2016).

Marine resource management decisions will benefit greatly from mechanistic understanding,

risk assessments, and attribution studies of extreme events (Oliver et al. 2017; Webb and Werner 2018). To that end, we find that the recent extreme ocean temperatures off the U.S. West Coast, which significantly impacted many marine species and fisheries, were caused by the confluence of multiple complementary natural drivers and were likely exacerbated by long-term anthropogenic warming.

ACKNOWLEDGMENTS. We thank Desiree Tommasi, the editors, and two anonymous reviewers for their insightful comments on our manuscript, and Michael O’Farrell for providing a summary of 2016 and 2017 U.S. West Coast salmon fisheries. Contributing support for this work was provided by the National Oceanic and Atmospheric Administration’s Integrated Ecosystem Assessment (NOAA IEA) Program. The scientific results and conclusions, as well as any views or opinions expressed herein, are those of the authors and do not necessarily reflect the views of NOAA or the Department of Commerce.

REFERENCES

- Alexander, M. A., I. Blade, M. Newman, J. R. Lanzante, N.-C. Lau, and J. D. Scott, 2002: The atmospheric bridge: The influence of ENSO teleconnections on air-sea interaction over the global oceans. *J. Climate*, **15**, 2205–2231.
- Banzon, V., T. M. Smith, T. M. Chin, C. Liu, and W. Hankins, 2016: A long-term record of blended satellite and in situ sea-surface temperature for climate monitoring, modeling and environmental studies. *Earth Syst. Sci. Data*, **8**, 165–176, doi:10.5194/essd-8-165-2016
- Bond, N. A., M. F. Cronin, H. Freeland, and N. Mantua, 2015: Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophys. Res. Lett.*, **42**, 3414–3420, doi:10.1002/2015GL063306.
- Brady, R. X., M. A. Alexander, N. S. Lovenduski, and R. R. Rykaczewski, 2017: Emergent anthropogenic trends in California Current upwelling. *Geophys. Res. Lett.*, **44**, 5044–5052, doi:10.1002/2017GL072945.
- Cavole, L. M., and Coauthors, 2016: Biological impacts of the 2013–2015 warm-water anomaly in the Northeast Pacific. *Oceanography*, **29** (2), 273–285, doi:10.5670/oceanog.2016.32.

- Chavez, F. P., and Coauthors, 2002: Biological and chemical consequences of the 1997–1998 El Niño in central California waters. *Prog. Oceanogr.*, **54**, 205–232, doi:10.1016/S0079-6611(02)00050-2.
- Checkley, Jr., D. M., and J. A. Barth: 2009: Patterns and processes in the California Current System. *Prog. Oceanogr.*, **83**, 49–64, doi:10.1016/j.pocean.2009.07.028.
- Daly, E. A., R. D. Brodeur, and T. D. Auth, 2017: Anomalous ocean conditions in 2015: Impacts on spring Chinook salmon and their prey field. *Marine Ecol. Prog. Ser.*, **566**, 169–182, doi:10.3354/meps12021.
- Di Lorenzo, E., and N. J. Mantua, 2016: Multi-year persistence of the 2014/15 North Pacific marine heatwave. *Nat. Climate Change*, **6**, 1042–1047, doi:10.1038/nclimate3082.
- Frischknecht, M., M. Münnich, and N. Gruber, 2017: Local atmospheric forcing driving an unexpected California Current System response during the 2015–2016 El Niño. *Geophys. Res. Lett.*, **44**, 304–311, doi:10.1002/2016GL071316.
- Gentemann, C. L., M. R. Fewings, and M. García-Reyes, 2017: Satellite sea surface temperatures along the West Coast of the United States during the 2014–2016 northeast Pacific marine heat wave. *Geophys. Res. Lett.*, **44**, 312–319, doi:10.1002/2016GL071039.
- Jacox, M. G., J. Fiechter, A. M. Moore, and C. A. Edwards, 2015: ENSO and the California Current coastal upwelling response. *J. Geophys. Res. Oceans*, **120**, 1691–1702, doi:10.1002/2014JC010650.
- , E. L. Hazen, K. D. Zaba, D. L. Rudnick, C. A. Edwards, A. M. Moore, and S. J. Bograd, 2016: Impacts of the 2015–2016 El Niño on the California Current System: Early assessment and comparison to past events. *Geophys. Res. Lett.*, **43**, 7072–7080, doi:10.1002/2016GL069716.
- Johnstone, J. A., and N. J. Mantua, 2014: Atmospheric controls on northeast Pacific temperature trends and variations, 1900–2012. *Proc. Natl. Acad. Sci. USA*, **111**, 14360–14365, doi:10.1073/pnas.1318371111.
- Kay, J. E., and Coauthors, 2015: The Community Earth System Model (CESM) Large Ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bull. Amer. Meteor. Soc.*, **96**, 1333–1349, doi:10.1175/BAMS-D-13-00255.1.
- Kilduff, D. P., E. Di Lorenzo, L. W. Botsford, and S. L. H. Teo, 2015: Changing central Pacific El Niños reduce stability of North American salmon survival rates. *Proc. Natl. Acad. Sci. USA*, **112**, 10962–10966, doi:10.1073/pnas.1503190112.
- Lamarque, J.-F., and Coauthors, 2010: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application. *Atmos. Chem. Phys.*, **10**, 7017–7039, doi:10.5194/acp-10-7017-2010.
- , G. P. Kyle, M. Meinshausen, K. Riahi, S. J. Smith, D. P. van Vuuren, A. J. Conley, and F. Vitt, 2011: Global and regional evolution of short-lived radiatively-active gases and aerosols in the representative concentration pathways. *Climatic Change*, **109**, 191–212, doi:10.1007/s10584-011-0155-0.
- Oliver, E. C. J., J. A. Benthuysen, N. L. Bindoff, A. J. Hobday, N. J. Holbrook, C. N. Mundy, and S. E. Perkins-Kirkpatrick, 2017: The unprecedented 2015/16 Tasman Sea marine heatwave. *Nat. Comms.*, **8**, 16101, doi:10.1038/ncomms16101.
- Pearcy, W.G. 1992: *Ocean Ecology of North Pacific Salmonids*. Washington Sea Grant Program, 179 pp.
- Peterson, W. T., and F. B. Schwing, 2003: A new climate regime in northeast Pacific ecosystems. *Geophys. Res. Lett.*, **30**, 1896, doi:10.1029/2003GL017528
- , N. Bond, and M. Robert, 2016: The Blob (Part three): Going, going, gone? *PICES Press*, **24**, 46–48. [Available online at www.pices.int/publications/pices_press/volume24/PPJan2016.pdf.]
- PFMC, 2017a: Review of 2016 ocean salmon fisheries: Stock assessment and fishery evaluation document for the Pacific coast salmon fishery management plan. Pacific Fishery Management Council, 342 pp. [Available online at www.pcouncil.org/wp-content/uploads/2017/03/Review_of_2016_Ocean_Salmon_Fisheries_03032017.pdf.]
- , 2017b: Preseason report I: Stock abundance analysis and environmental assessment part 1 for 2017 ocean salmon fishery regulations. Pacific Fishery Management Council, 131 pp. [Available online at www.pcouncil.org/salmon/stock-assessment-and-fishery-evaluation-safe-documents/preseason-reports/2017-preseason-report-i/.]
- Rayner, N.A., D. E. Parker, E. B. Horton, C. K. Folland, L. V. Alexander, D. P. Rowell, E. C. Kent, and A. Kaplan, 2003: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *J. Geophys. Res.*, **108**, 4407, doi:10.1029/2002JD002670.
- Reynolds, R. W., T. M. Smith, C. Liu, D. B. Chelton, K. S. Casey, and M. G. Schlax, 2007: Daily high-resolution-blended analyses for sea surface temperature. *J. Climate*, **20**, 5473–5496, doi:10.1175/2007JCLI1824.1.

- Sharma, S., L. A. Vélez-Espino, A. C. Wertheimer, N. Mantua, and R. C. Francis, 2012: Relating spatial and temporal scales of climate and ocean variability to survival of Pacific Northwest Chinook salmon (*Oncorhynchus tshawytscha*). *Fisheries Oceanogr.*, **22**, 14–31, doi:10.1111/fog.12001.
- Stott, P. A., D. A. Stone, and M. R. Allen, 2004: Human contribution to the European heatwave of 2003. *Nature*, **432**, 610–614, doi:10.1038/nature03089.
- Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design. *Bull. Amer. Meteor. Soc.*, **93**, 485–498, doi:10.1175/BAMS-D-1100094.1.
- Webb, R. S., and F. E. Werner, 2018: Explaining extreme ocean conditions impacting living marine resources [in “Explaining Extreme Events of 2016 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **99** (1), S7–S10, doi:10.1175/BAMS-D-17-0265.1.
- Welch, C., 2016: The blob that cooked the Pacific. *Nat. Geogr.*, **230** (3), 54+ (~11 pages).
- Weller, E., S.-K. Min, D. Lee, W. Cai, S.-W. Yeh, and J.-S. Kug, 2015: Human contribution to the 2014 record high sea surface temperatures over the western tropical and northeast Pacific Ocean [in “Explaining Extreme Events of 2014 from a Climate Perspective”]. *Bull. Amer. Meteor. Soc.*, **96** (12), S100–S104, doi:10.1175/BAMS-D-15-00055.1.
- Wells, B. K., and Coauthors, 2017: Environmental conditions and prey-switching by a seabird predator impacts juvenile salmon survival. *J. Mar. Syst.*, **174**, 54–63, doi:10.1016/j.jmarsys.2017.05.008.

Table I.I. SUMMARY of RESULTS

ANTHROPOGENIC INFLUENCE ON EVENT			
	INCREASE	DECREASE	NOT FOUND OR UNCERTAIN
Heat	Ch. 3: Global Ch. 7: Arctic Ch. 15: France Ch. 19: Asia		
Cold		Ch. 23: China Ch. 24: China	
Heat & Dryness	Ch. 25: Thailand		
Marine Heat	Ch. 4: Central Equatorial Pacific Ch. 5: Central Equatorial Pacific Ch. 6: Pacific Northwest Ch. 8: North Pacific Ocean/Alaska Ch. 9: North Pacific Ocean/Alaska Ch. 9: Australia		Ch. 4: Eastern Equatorial Pacific
Heavy Precipitation	Ch. 20: South China Ch. 21: China (Wuhan) Ch. 22: China (Yangtze River)		Ch. 10: California (failed rains) Ch. 26: Australia Ch. 27: Australia
Frost	Ch. 29: Australia		
Winter Storm			Ch. 11: Mid-Atlantic U.S. Storm "Jonas"
Drought	Ch. 17: Southern Africa Ch. 18: Southern Africa		Ch. 13: Brazil
Atmospheric Circulation			Ch. 15: Europe
Stagnant Air			Ch. 14: Western Europe
Wildfires	Ch. 12: Canada & Australia (Vapor Pressure Deficits)		
Coral Bleaching	Ch. 5: Central Equatorial Pacific Ch. 28: Great Barrier Reef		
Ecosystem Function		Ch. 5: Central Equatorial Pacific (Chl- <i>a</i> and primary production, sea bird abundance, reef fish abundance) Ch. 18: Southern Africa (Crop Yields)	
El Niño	Ch. 18: Southern Africa		Ch. 4: Equatorial Pacific (Amplitude)
TOTAL	18	3	9

		METHOD USED	Total Events
Heat		Ch. 3: CMIP5 multimodel coupled model assessment with piCont, historicalNat, and historical forcings Ch. 7: CMIP5 multimodel coupled model assessment with piCont, historicalNat, and historical forcings Ch. 15: Flow analogues conditional on circulation types Ch. 19: MIROC-AGCM atmosphere only model conditioned on SST patterns	
Cold		Ch. 23: HadGEM3-A (GA6) atmosphere only model conditioned on SST and SIC for 2016 and data fitted to GEV distribution Ch. 24: CMIP5 multimodel coupled model assessment	
Heat & Dryness		Ch. 25: HadGEM3-A N216 Atmosphere only model conditioned on SST patterns	
Marine Heat		Ch. 4: SST observations; SGS and GEV distributions; modeling with LIM and CGCMs (NCAR CESM-LE and GFDL FLOR-FA) Ch. 5: Observational extrapolation (OISST, HadISST, ERSST v4) Ch. 6: Observational extrapolation; CMIP5 multimodel coupled model assessment Ch. 8: Observational extrapolation; CMIP5 multimodel coupled model assessment Ch. 9: Observational extrapolation; CMIP5 multimodel coupled model assessment	
Heavy Precipitation		Ch. 10: CAM5 AMIP atmosphere only model conditioned on SST patterns and CESM1 CMIP single coupled model assessment Ch. 20: Observational extrapolation; CMIP5 and CESM multimodel coupled model assessment; auto-regressive models Ch. 21: Observational extrapolation; HadGEM3-A atmosphere only model conditioned on SST patterns; CMIP5 multimodel coupled model assessment with ROF Ch. 22: Observational extrapolation, CMIP5 multimodel coupled model assessment Ch. 26: BoM seasonal forecast attribution system and seasonal forecasts Ch. 27: CMIP5 multimodel coupled model assessment	
Frost		Ch. 29: weather@home multimodel atmosphere only models conditioned on SST patterns; BoM seasonal forecast attribution system	
Winter Storm		Ch. 11: ECHAM5 atmosphere only model conditioned on SST patterns	
Drought		Ch. 13: Observational extrapolation; weather@home multimodel atmosphere only models conditioned on SST patterns; HadGEM3-A and CMIP5 multimodel coupled model assessment; hydrological modeling Ch. 17: Observational extrapolation; CMIP5 multimodel coupled model assessment; VIC land surface hydrological model, optimal fingerprint method Ch. 18: Observational extrapolation; weather@home multimodel atmosphere only models conditioned on SSTs, CMIP5 multimodel coupled model assessment	
Atmospheric Circulation		Ch. 15: Flow analogues distances analysis conditioned on circulation types	
Stagnant Air		Ch. 14: Observational extrapolation; Multimodel atmosphere only models conditioned on SST patterns including: HadGEM3-A model; EURO-CORDEX ensemble; EC-EARTH+RACMO ensemble	
Wildfires		Ch. 12: HadAM3 atmosphere only model conditioned on SSTs and SIC for 2015/16	
Coral Bleaching		Ch. 5: Observations from NOAA Pacific Reef Assessment and Monitoring Program surveys Ch. 28: CMIP5 multimodel coupled model assessment; Observations of climatic and environmental conditions (NASA GES DISC, HadCRUT4, NOAA OISSTV2)	
Ecosystem Function		Ch. 5: Observations of reef fish from NOAA Pacific Reef Assessment and Monitoring Program surveys; visual observations of seabirds from USFWS surveys. Ch. 18: Empirical yield/rainfall model	
El Niño		Ch. 4: SST observations; SGS and GEV distributions; modeling with LIM and CGCMs (NCAR CESM-LE and GFDL FLOR-FA) Ch. 18: Observational extrapolation; weather@home multimodel atmosphere only models conditioned on SSTs, CMIP5 multimodel coupled model assessment	