

King Saud University
College of Science
Department of Mathematics

151 Math Exercises

(3,1)

Methods of Proof

- 1-Direct Proof
- 2- Proof by Contraposition
- 3- Proof by Contradiction
- 4- Proof by Cases

By:

Malek Zein AL-Abidin

$$\frac{1443 \ H}{2022 \ G}$$

Direct Proof: A direct proof shows that a conditional statement $p \rightarrow q$ is true by showing that if p is true, then q must also be true, so that the combination p true and q false never occurs. In a direct proof, we assume that p is true and use axioms, definitions, and previously proven theorems, together with rules of inference, to show that q must also be true

DEFINITION 1 The integer n is *even* if there exists an integer k such that $n = 2k$, and n is *odd* if there exists an integer k such that $n = 2k + 1$. (Note that every integer is either even or odd, and no integer is both even and odd.) Two integers have the *same parity* when both are even or both are odd; they have *opposite parity* when one is even and the other is odd.

EXAMPLE 1 Give a direct proof of the theorem “If n is an odd integer, then n^2 is odd.”

Solution: $n = 2k + 1$, $k \in \mathbb{Z}$

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$$

$$n^2 = 2h + 1 \quad : \quad h = (2k^2 + 2k) \in \mathbb{Z}$$

$\therefore n^2$ is odd

EXAMPLE 2 Give a direct proof that if m and n are both perfect squares, then mn is also a perfect square. (An integer a is a **perfect square** if there is an integer b such that $a = b^2$.)

Solution: We assume that m and n are both perfect squares. By the definition of a perfect square, $\exists s, t \in \mathbb{Z}$ such that $m = s^2$ and $n = t^2$. $mn = s^2t^2 = (st)^2 = h^2$: $h = st \in \mathbb{Z}$
 $\therefore mn = h^2$. $\therefore mn$ is a perfect square .

DEFINITION 3 The real number r is **rational** if there exist integers p and q with $q \neq 0$ such that $r = p/q$.

A real number that is not rational is called **irrational**.

EXAMPLE 3 Prove that the sum of two rational numbers is rational.

Solution:

Let $r = p/q$ where p and q are integers, with $q \neq 0$, and $s = t/u$ where t and u are integers, with $u \neq 0$.

$$r + s = \frac{p}{q} + \frac{t}{u} = \frac{pu + qt}{qu} \text{ where } pu + qt, qu \in \mathbb{Z} : qu \neq 0$$

$\therefore r + s$ is rational number

DEFINITION 4 Let $a, b \in \mathbb{Z} : a \neq 0$. a is a divisor to b , $a|b$ if there exist integer c such that $b = ac$

DEFINITION 5 Let $a, b \in \mathbb{Z}$ and $n \in \mathbb{Z}^+$. a is congruent to b modulo n , $a \equiv b \pmod{n}$
if $n|(a - b)$

EXAMPLE 4 : $25 \equiv 7 \pmod{9}$, $3 \equiv -15 \pmod{9}$

$$SEG = 2K + 1 + 2h + 1 : h, k \in \mathbb{Z}$$

Exercises

1. Use a direct proof to show that the sum of two odd integers is even.

Solution:

$$\begin{aligned} S+G &= 2K + 2h + 2 = \\ S+G &= 2(K+h+1) = 2M \\ 2M \text{ is even } &\quad M \in \mathbb{Z} \\ \text{done} \# & \end{aligned}$$

2. Use a direct proof to show that : if n is an even number then $(3n + 5)$ is odd .

Solution:

$$\begin{aligned} n &= 2K : K \in \mathbb{Z} \\ 3(2K) + 5 &= 6K + 5 \\ = 6K + 4 + 1 &= 2(3K + 2) + 1 \\ = 2M + 1 & \quad M \in \mathbb{Z} \\ \text{is odd} \quad \text{done} \# & \end{aligned}$$

3. Use a direct proof to show that : if n is an odd number then $(5n + 6)$ is odd .

Solution:

3. Use a direct proof to show that : if n is an odd number then $(5n+3)$ is odd.

Solution:

$$\begin{aligned} n &= 2k+1, \quad k \in \mathbb{Z} \\ 5(2k+1) + 6 &= 10k + 5 + 6 \\ &\quad \text{1-4} \\ &= 10k + 11 \\ &= 2(5k+3) + 1 \quad ? \text{ is odd} \\ &\quad \text{done} \end{aligned}$$

4. Suppose a , b and c are integers. Use a direct proof to show that if a divides $(b - 1)$ and a divides $(c - 1)$, then a divides $(bc - 1)$

Solution:

ii. Suppose a , b and c are integers. Use a direct proof to show that if a divides $(b - 1)$ and a divides $(c - 1)$, then a divides $(bc - 1)$

Solution: $a \mid b-1 \Rightarrow \exists k \in \mathbb{Z} \text{ such that } b = ak + 1$
 $a \mid c-1 \Rightarrow \exists m \in \mathbb{Z} \text{ such that } c = am + 1$

$bc - 1 = (ak + 1)(am + 1) - 1$
 $= \underbrace{a^2 km}_{\in \mathbb{Z}} + \underbrace{ak + am + 1}_{\in \mathbb{Z}} - 1$
 $= a(cmk + k + m)$
 $\therefore a \mid bc - 1$

$\boxed{bc-1 = a(mk + k + m)}$

5. Prove that if $m + n$ and $n + p$ are even integers, where m, n , and p are integers, then $m + p$ is even.

Solution:

$$\begin{aligned}
 m+n &= 2k \Rightarrow m = 2k - n \quad \text{for } k \in \mathbb{Z} \\
 n+p &= 2h \Rightarrow p = 2h - n \quad \text{for } h \in \mathbb{Z} \\
 m+p &= 2k + 2h - 2n \\
 m+p &= 2(k+h-1) \\
 m+p \text{ is even} &\quad \text{for } m \in \mathbb{Z} \\
 \text{done} &\quad \text{#}
 \end{aligned}$$

6. Use a direct proof to show that the product of two odd numbers is odd.

Solution:

$$\begin{aligned}
 S \cdot G &= (2k+1) \cdot (2h+1) : b, k \in \mathbb{Z} \\
 S \cdot G &= \underbrace{4kh}_{\text{for } m \in \mathbb{Z}} + \underbrace{2k+2h+1}_{\text{for } m \in \mathbb{Z}} \\
 &= 2(2kh+k+h) + 1 \\
 2m+1 \text{ is odd} &\quad \text{done} \quad \text{#}
 \end{aligned}$$

7. Use a direct proof to show that every odd integer is the difference of two squares.

Solution:

$$\because (n \text{ is odd}) \Rightarrow n = 2k + 1 : k \in \mathbb{Z}$$

$$(k+1)^2 - k^2 = k^2 + 2k + 1 - k^2 = 2k + 1 = n$$

8. Use a direct proof to show that $x^2 + x$ is an even number where x is integer.

Solution:

$$\begin{aligned} & \text{(even)} \quad x = 2k : k \in \mathbb{Z} \\ & x^2 + x = 4k^2 + 2k \\ & = 2(x^2 + k) \\ & \quad \text{MORE} \\ & = 2m \quad (\text{even}) \end{aligned}$$

$$\begin{aligned} & \text{(odd)} \quad x = 2h+1 : h \in \mathbb{Z} \\ & x^2 + x = (2h+1)^2 + 2h+1 \\ & = 4h^2 + 4h + 1 + 2h + 1 \\ & = 2(2h^2 + 2h + 1) \\ & \quad \text{MORE} \\ & = 2t \quad (\text{even}) \quad \# \end{aligned}$$

9. Use a direct proof to show that if x is an odd integer, then $x^2 = 8m + 1$ where m is integer.

Solution:

$$\begin{aligned} & x = 2k+1 : k \in \mathbb{Z} \\ & x^2 = 4k^2 + 4k + 1 = 4k(k+1) + 1 \\ & \quad \text{ONE OF THESE IS EVEN} \\ & \quad \text{THE OTHER IS ODD} \end{aligned}$$

$$4(2m) + 1 = 8m + 1$$

10. Let n be an integer. Use a direct proof to show that if $3 \mid (n - 1)$, then

$$3 \mid (n^2 - n + 6)$$

Solution:

$$3k = n - 1 \Rightarrow n = 3k + 1$$

$$n^2 = 9k^2 + 6k + 1$$

$$3n = n^2 - 6$$

$$n^2 - 0.16 = 9K^2 + 6K + 1 - 1 + 3K$$

$$I^2 - A + b = \overbrace{3C_2 K^2 + 2K + K}^{\text{neglected}} \\ \Rightarrow 3 \sqrt{A^2 - A + b} \text{ done}$$

11. Use a direct proof to prove that if n is an odd number, then $n^2 \equiv 1 \pmod{4}$. (S)

Solution:

11. Use a direct proof to prove that if n is an odd number, then $n^2 \equiv 1 \pmod{4}$.

Solution:

$$\begin{aligned} n &= 2k+1 : k \in \mathbb{Z} \quad n^2 = 4k^2 + 4k + 1 \\ n^2 - 1 &= 4k^2 + 4k + 1 - 1 \quad | 4 \mid n^2 - 2 \\ n^2 - 1 &= 4(k^2 + k) = 4n \quad | n^2 - 1 = 4n \end{aligned}$$

$k \in \mathbb{Z}$

$\Rightarrow 4 \mid \tilde{a}^n - 1 \Rightarrow \text{den eff}$

12. Use a direct proof to prove that if n is an even number, then $n^2 \equiv 0 \pmod{4}$. (*)

Solution:

$$\begin{aligned} n &= 2k \cdot k \in \mathbb{Z} \quad n^2 = 4k^2 \\ n^2 &\equiv 4k^2 \equiv 4(k^2) \Rightarrow 4M \end{aligned}$$

$4 \mid n^2 - 0$

$\Rightarrow 4 \mid n^2 \Rightarrow (*)$ done \dagger

13. Use a direct proof to prove that, if 3 is not a divisor of n , then $n^2 \equiv 1 \pmod{3}$. (x)

Solution:

$$\begin{aligned} n^2 - 1 &= 9k + 6k + 1 - 1 \\ &= 3(3k^2 + 2k) = 3M \end{aligned}$$

$n \in \mathbb{Z}$

$3 \mid n^2 - 1$

$3 \mid n^2 - 1$

$3 \mid n^2 - 1$

$n^2 \equiv 1 + 3k$

$= 9k^2 + 6k + 1$

$k \in \mathbb{Z}$

$\therefore (*)$ done \dagger

14. If $n \in \mathbb{Z}^+$ and $a, b, c \in \mathbb{Z}$ where $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$ prove that :
- (i) $a + c \equiv b + d \pmod{n}$ (ii) $ac \equiv bd \pmod{n}$

Solution:

Proof by Contraposition Proofs by contraposition make use of the fact that the conditional statement $p \rightarrow q$ is equivalent to its contrapositive, $\neg q \rightarrow \neg p$. This means that the conditional statement $p \rightarrow q$ can be proved by showing that its contrapositive, $\neg q \rightarrow \neg p$, is true. In a proof by contraposition of $p \rightarrow q$, we take $\neg q$ as a premise, and using axioms, definitions, and previously proven theorems, together with rules of inference, we show that $\neg p$ must follow. the proof by contraposition can succeed when we cannot easily find a direct proof.

EXAMPLE 1 Prove that if n is an integer and $3n + 2$ is odd, then n is odd.

Solution:

Assume that n is even. Then, $n = 2k : k \in \mathbb{Z}$. Substituting $2k$ for n , we find that $3n + 2 = 3(2k) + 2 = 6k + 2 = 2(3k + 1) = 2h : h = (3k + 1) \in \mathbb{Z}$.

This tells us that $3n + 2$ is even (because it is a multiple of 2), and therefore not odd. This is the negation of the premise of the theorem.

EXAMPLE 2 Prove that if $n = ab$, where a and b are positive integers, then

$$a \leq \sqrt{n} \quad \text{or} \quad b \leq \sqrt{n} .$$

Solution:

Assume that $a > \sqrt{n}$ and $b > \sqrt{n} \Rightarrow ab > \sqrt{n} \cdot \sqrt{n} = n$

$$\therefore ab > n \Rightarrow \therefore ab \neq n$$

EXAMPLE 3 Prove that if n is an integer and n^2 is odd, then n is odd.

Solution:

Assume that n is even integer, $n = 2k : k \in \mathbb{Z}$

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2) = 2h ; h = (2k^2) \in \mathbb{Z}$$

$$\therefore n^2 \text{ is even} .$$

Exercises

1. Use a proof by contraposition to show that if $x + y \geq 2$, where x and y are real numbers,

then $x \geq 1$ or $y \geq 1$.

Solution:

$$\begin{aligned} y < 1 \text{ and } x < 1 \Rightarrow \\ x + y < 2 \rightarrow ?P \end{aligned}$$

2. Prove that if m and n are integers and mn is even, then m is even or n is even.

Solution:

$$\begin{aligned} m = 2k+1 : k \in \mathbb{Z} \text{ and } n = 2h+1 : h \in \mathbb{Z} \\ mn = (2k+1)(2h+1) = \cancel{2k}h + \cancel{2k} + 2h + 1 \\ mn = \cancel{2}(\cancel{2k}h + \cancel{k}h) + 1 \rightarrow mn = 2m' + 1 \\ \text{done} \quad \text{is odd} \end{aligned}$$

3. Show that if n is an integer and $n^3 + 5$ is odd, then n is even using a proof by contraposition.

Solution:

$$\begin{aligned} n^3 &= (2k+1)(2k+1)^2 + 5 \\ n^3 + 5 &= \cancel{2}k^3 + (\cancel{2}k^2 + \cancel{2}k + \cancel{2}) + \cancel{1} + 5 = 2(4k^3 + 6k^2 + 3k + 1) \\ n^3 + 5 &= 2m \quad m \in \mathbb{Z} \\ \therefore n^3 + 5 &\neq 2n \quad \text{done} \end{aligned}$$

4. Prove that if n is an integer and $3n + 2$ is even, then n is even, using a proof by contraposition.

Solution:

$$\begin{aligned} n &= 2k + 1 \quad k \in \mathbb{Z} \\ 3(2k+1) + 2 &= 6k + 3 + 2 \\ 6k + 4 + 1 &= 2(\underbrace{3k+2}_{n \in \mathbb{Z}}) + 1 = 2m + 1 \\ \therefore ? \rightarrow ?P \quad \text{done} \# & \quad \text{(odd)} \end{aligned}$$

5. Prove that if n is a positive integer, then n is even if and only if $7n + 4$ is even.

Solution:

6. Prove that if a is an integer where $5 \nmid a$, then $5 \nmid (a + 20)$ using a proof by contraposition.

Solution:

$$\begin{aligned} a &= 5c - 20 = 5(c - 4) = 5m \\ \Rightarrow 5 \mid a &\Rightarrow ?P \\ \therefore ?Q \rightarrow ?P \quad \text{done} \# & \end{aligned}$$

7. Use a proof by contraposition to show that if $2|mn : m, n \in \mathbb{N}$, then $2|m$ or $2|n$.

Solution: $\neg Q = 2 \nmid mn \Rightarrow m = 2k+1 : k \in \mathbb{Z}$

$\neg Q \rightarrow \neg m = 2k+1 : k \in \mathbb{Z}$

$m \cdot n = (2k+1)(2h+1) = 4kh + 2k + 2h + 1$

$= 2(2kh + kh) + 1 = 2M + 1 \Rightarrow \neg P$

$\Rightarrow 2 \nmid mn \therefore \neg Q \rightarrow \neg P \text{ done } \#$

8. Use a proof by contraposition to show that if xy is even number where $x, y \in \mathbb{Z}$, then x is even or y is even.

Solution:

9. Use a proof by contraposition to show that if $x^2 + y^2 = z^2 : x, y, z \in \mathbb{Z}$, then one of these numbers x, y, z is at least should be even.

Solution: $\neg Q : x, y, z \text{ all odd}$

$x^2 = (2k+1)^2 = 4k^2 + 4k + 1 : k \in \mathbb{Z}$

$y^2 = (2h+1)^2 = 4h^2 + 4h + 1 : h \in \mathbb{Z}$

$z^2 = (2l+1)^2 = 4l^2 + 4l + 1 : l \in \mathbb{Z}$

$x^2 + y^2 = 4k^2 + 4k + 4h^2 + 4h + 2 = 2(2k^2 + 2k + 2h^2 + 2h + 1)$

$x^2 + y^2 = 2H \text{ even} \therefore x^2 + y^2 \neq z^2 \text{ for } H \in \mathbb{Z}$

$\therefore \neg Q \rightarrow \neg P \text{ done } \#$

- 10.** Use a proof by contraposition to show that if $3mn + 2$ is irrational number, then m is irrational or n is irrational.

Solution:

$\exists \Delta \delta \cap \mathbb{Q} : n = \frac{c}{d} \cdot c, d \in \mathbb{Z}$

$$3mn + 2 = 3 \frac{c}{d} \frac{c}{d} + 2 = \frac{3cc}{bd} + 2$$

$\cancel{(3cc+2d)} \in \mathbb{Q} : bd \neq 0, (*) \in \mathbb{Z}$

$3mn + 2$ is rational \Rightarrow SP

$\therefore \exists q \rightarrow \text{SP done} \#$

- 11.** Use a proof by contraposition to show that if $a^2 + b^2$ is odd, then a is even or b is even.

Solution:

$$\exists a = 2k + 1 \in \mathbb{Z}$$

$$b = 2h + 1 : h \in \mathbb{Z}$$

$$a^2 + b^2 = (2k+1)^2 + (2h+1)^2 = 4k^2 + 4k + 1 + 4h^2 + 4h + 1$$

$$= 2(2k^2 + 2k + 2h^2 + 2h + 1) = 2m \quad m \in \mathbb{Z}$$

$\therefore \exists q \rightarrow \text{SP done} \#$

- 12.** Use a proof by contraposition to show that if $x + y < 15$, where x and y are real numbers, then $x < 8$ or $y < 8$.

Solution:

$$\begin{aligned} & \neg(x \geq 8 \wedge y \geq 8) \rightarrow x+y \geq 15 \\ & \neg(x+y \geq 15) \rightarrow x+y < 15 \\ & \therefore x+y < 15 \rightarrow x < 8 \text{ or } y < 8 \end{aligned}$$

- 13.** Prove that if m is an integer where $3 \nmid m$, then $3 \nmid (m+1)^2 + 2m^2 + 5$ using a proof by contraposition.

Solution:

$$\begin{aligned} & \neg(3 \mid (m+1)^2 + 2m^2 + 5) \rightarrow 3 \nmid h \\ & 3 \nmid (m+1)^2 + 2m^2 + 5 : h \in \mathbb{Z} \\ & = m^2 + 2m + 1 + 2m^2 + 5 = 3h \\ & 2m = 3h - 3m^2 - 6 = 3(h - m^2 - 2) \in \mathbb{Z} \\ & \exists k [2m = 3k \rightarrow 3 \mid m] \rightarrow \neg(3 \mid (m+1)^2 + 2m^2 + 5) \end{aligned}$$

- 14.** Use a proof by contraposition to show that if $3n^2 + 4n + 3$ is even, then n is odd.

Solution:

15. Let $6|m : m \in \mathbb{Z}$. Use a proof by contraposition to show that if $3 \nmid (m+n) : n \in \mathbb{Z}$, then $3 \nmid n$.

Solution:

$$m+n = 6h + 3k = 3(2h+k) = 3M$$

$$\Rightarrow \therefore 3|(m+n) \equiv 7P \quad M \in \mathbb{Z}$$

$\therefore 7q \rightarrow 7P$ done $\#$

②

16. Prove that if n is a positive integer, then n is odd if and only if $5n+6$ is odd.

Solution: If n is odd if $5n+6$ is odd then

then $5n+6$ odd

$$n = 2k+1 : k \in \mathbb{Z}$$

$$5(2k+1) + 6 =$$

$$10k + 5 + 6 =$$

$$= 10k + 6 + 4 + 1$$

$$= 2(5k+3+2)+1$$

$$= 2M+1 \text{ is odd} \quad M \in \mathbb{Z}$$

done $\#$

n is odd

$$7q = n = 2h : h \in \mathbb{Z}$$

$$5(2h) + 6 = 10h + 6$$

$$2(5h+3) = 2H \text{ even}$$

$H \in \mathbb{Z}$

$$7P = 2H$$

$$\therefore 7q \rightarrow 7P$$

done

$\#$

If $3n+2$ is odd then $3n+2$ is even
 if and only if $9n+5$ is even

17. Prove that if n is integer, then $3n + 2$ is odd if and only if $9n + 5$ is even .

Solution: (CDP)

$$3n+2 = 2k+1 : k \in \mathbb{Z}$$

$$3n = 2k-1$$

$$\text{by } 9 \Rightarrow 9n = 6k - 3$$

$$\begin{aligned} 9n+5 &= 6k - 3 + 5 \\ &= 6k + 2 \end{aligned}$$

$$\begin{aligned} 9n+5 &= 2(3k+1) \\ &\Rightarrow 2m \quad m \in \mathbb{Z} \end{aligned}$$

even $\#$

(contradiction)

let $q = 3n+2$ is even

$$2h : h \in \mathbb{Z}$$

$$\begin{aligned} 2h+1+2 &\Rightarrow 9n = 6h-6 \\ &\Rightarrow 9n+5 = 6h-6+6+5 \end{aligned}$$

$$\begin{aligned} 9n+5 &= 6h-6+4+1 \\ &= 2(3h-1)+1 \\ &\quad h \in \mathbb{Z} \end{aligned}$$

$$\begin{aligned} 9n+5 &= 2h+1 \\ &\quad \text{odd} \end{aligned}$$

$7q \rightarrow 7p$

18. Use a proof by contraposition to show that if $3x - y = z$ where $x, y, z \in \mathbb{Z}$, then x is even or y is even or z is even .

Solution:

19. Use a proof by contraposition to show that if $4n^2 + n - 3$ is even, then n is odd.

Solution:

20. Use a proof by contraposition to show that if $a + b \leq 4$, where a and b are real numbers, then $a \leq 1$ or $b \leq 3$.

Solution:

- 21.** Use a proof by contraposition to show that if $x + y < 9$, where x and y are real numbers, then $x < 4$ or $y < 5$.

Solution:

- 22.** Use a proof by contraposition to show that if $m \nmid n$, then $m \nmid l$ or $m \neq n + l$ where $l, m, n \in \mathbb{Z}$.

Solution:

$$\begin{aligned} & \text{Let } m \mid l \text{ and } m \mid n+l \\ & \Rightarrow \exists k \in \mathbb{Z} \text{ such that } m = n + mk \\ & \Rightarrow \exists k \in \mathbb{Z} \text{ such that } l = mk \\ & \therefore m \mid l \end{aligned}$$

$\therefore \neg(m \nmid l) \rightarrow m \mid l \quad \text{done} \#$

- 23.** Use a proof by contraposition to show that if $a + b = 4$ where $a, b \in \mathbb{R}$, then $a \neq 2$ or $b \neq 3$.

Solution:

24. Use a proof by contraposition to show that if $mn = l$, then $l \geq 0$ or $m \geq 0$ or $n \geq 0$: $l, m, n \in \mathbb{Z}$

Solution:

$$\begin{aligned} & \text{Assume } l < 0 \quad (+) \\ & \text{Then } mn < 0 \quad (+) \\ & \text{So } m > 0 \quad (+) \\ & \text{And } n < 0 \quad (+) \\ & \text{Therefore } mn < 0 \quad (+) \end{aligned}$$

25. Use a proof by contraposition to show that if $a + b - c$ is even then a is even or b is even or c is even, where $a, b, c \in \mathbb{Z}$.

Solution:

26. Use a proof by contraposition to show that if $n^2 - 1$ is odd, then n is even : $n \in \mathbb{Z}$.

Solution:

27. Let m be a positive integer .Use a proof by contraposition to show that if $m \nmid 12$ or $m \nmid 21$, then $m \neq 3$.

28. Let $m \in \mathbb{Z}$, then $3 \mid m$ if and only if $6 \mid 2m + 24$.

Solution:

(D P)

$$3 \mid m \Rightarrow m = 3h \quad h \in \mathbb{Z}$$

$$\Rightarrow 2m + 24$$

$$= 2(3h) + 24$$

$$6h + 24$$

$$= 6(h + 4) \quad h \in \mathbb{Z}$$

$$= 6m$$

$$2m + 24 = 6m$$

$$6 \mid 2m + 24$$

$$6 \mid 2m + 24 \Rightarrow 6h = 2m + 24$$

$$2m = \underline{6h} - \underline{24} \quad \div 2$$

$$m = 3h - 12 = 3(h - 4) \quad h \in \mathbb{Z}$$

$$m = 3h \Rightarrow 3 \mid m$$

29. Use a proof by contraposition to show that if $3n^2$ is even, then n is even.

Solution:

30. Use a proof by contraposition to show that

if $5x + 3y - 4z \geq 27$ where $x, y, z \in \mathbb{R}$, then $x \geq 2$ or $y \geq 3$ or $z \leq -2$.

Solution:

$$(5x) \wedge x < 2 = 5x < 10$$

$$(3) \wedge y < 3 = 3y < 9$$

$$(-4) \wedge z > -2 \rightarrow -4z < 8$$

$\underbrace{\quad}_{\text{done}} \text{ (181)}$

$$5x + 3y - 4z < 27$$

$$\therefore \neg P \rightarrow P \text{ done } \#$$

31. Let a, b, c be real numbers. Prove by contraposition that :

if $5a - 2b + 3c > 7$ then $a > 2$ or $b < 3$ or $c > 1$.

Solution:

32. Let a, b, c be real numbers . Prove by contraposition that :

$$\text{if } a + 2b - 3c > 7 \text{ then } a > 4 \text{ or } b > 3 \text{ or } c < 1 .$$

Solution:

33. Use a proof by contraposition to show that

$$\text{if } 2x + 3y - 4z \geq 21 \text{ where } x, y, z \in \mathbb{R} , \text{ then } x \geq 2 \text{ or } y \geq 3 \text{ or } z \leq -2 .$$

Solution:

Proofs by Contradiction Suppose we want to prove that a statement p is true. Furthermore, suppose that we can find a contradiction q such that $\neg p \rightarrow q$ is true. Because q is false, but $\neg p \rightarrow q$ is true, we can conclude that $\neg p$ is false, which means that p is true. How can we find a contradiction q that might help us prove that p is true in this way? Because the statement $r \wedge \neg r$ is a contradiction whenever r is a proposition, we can prove that p is true if we can show that $\neg p \rightarrow (r \wedge \neg r)$ is true for some proposition r . Proofs of this type are called **proofs by contradiction**

EXAMPLE 1 Prove that $\sqrt{2}$ is irrational by giving a proof by contradiction.

Proof: Let p be the proposition “ $\sqrt{2}$ is irrational.” We will show that assuming that $\neg p$ is true leads to a contradiction. If $\sqrt{2}$ is rational, there exist integers a and b with $\sqrt{2} = a/b$, where $b \neq 0$ and $\gcd(a, b) = 1$; a and b have no common factors (so that the fraction a/b is in lowest terms.)

$$\sqrt{2} = a/b \stackrel{(\cdot)^2}{\Rightarrow} 2 = a^2/b^2 \Rightarrow a^2 = 2b^2 \quad (1)$$

$$\Rightarrow a^2 \text{ is even} \Rightarrow 2|a^2 \Rightarrow 2|a \Rightarrow a = 2k : k \in \mathbb{Z} \xrightarrow{\text{by subst. into (1)}} (2k)^2 = 2b^2$$

$$\Rightarrow 4k^2 = 2b^2 \stackrel{\div 2}{\Rightarrow} 2k^2 = b^2 \Rightarrow b^2 \text{ is even too} \Rightarrow 2|b^2 \Rightarrow 2|b \Rightarrow 2|a \text{ and } 2|b$$

This is contradiction with the assumption that $\gcd(a, b) = 1$. $\therefore p$ is true.

\therefore “ $\sqrt{2}$ is irrational.”

EXAMPLE 2 Give a proof by contradiction of the theorem “If $3n + 2$ is odd, then n is odd.”

Solution: Let p be “ $3n + 2$ is odd” and q be “ n is odd.” To construct a proof by contradiction, assume that both p and $\neg q$ are true. That is, assume that $3n + 2$ is odd and that n is not odd.

Because n is not odd, we know that it is even. Because n is even, there is an integer k such that $n = 2k$. This implies that $3n + 2 = 3(2k) + 2 = 6k + 2 = 2(3k + 1)$. Because $3n + 2$ is $2t$, where $t = 3k + 1$, $\therefore 3n + 2$ is even. Note that the statement “ $3n + 2$ is even” is equivalent to the statement $\neg p$, because an integer is even if and only if it is not odd. Because both p and $\neg p$ are true, we have a contradiction. This completes the proof by contradiction, proving that if $3n + 2$ is odd, then n is odd.

$$\begin{aligned} 3(2k) + 2 \\ 6k + 2 \\ = 2(3k + 1) \end{aligned}$$

- Exercises
1. Use a proof by contradiction to prove that the sum of an irrational number and a rational number is irrational.

Solution:

$$\text{Assume } i + j \in \mathbb{Q} \Rightarrow i + j = \frac{z}{y} \quad (i \notin \mathbb{Q})$$

$$i + j = \frac{z}{y} \Rightarrow j = \frac{z}{y} - i = \frac{z - y\alpha}{y} \quad (\alpha \neq 0)$$

$$z - y\alpha \in \mathbb{Z}$$

this is a contradiction with

assumption $j \notin \mathbb{Q}$

done #

2. Show that if n is an integer and $n^3 + 5$ is odd, then n is even using a proof by contradiction.

Solution:

$$\text{Assume } n^3 + 5 \text{ is odd} \Rightarrow n = 2k+1$$

KER

if n is true

$$n^3 = 8k^3 + (2k^2 + 6k + 1) =$$

$$8k^3 + 12k^2 + 6k + 1 = 2(4k^3 + 6k^2 + 3k) + 1$$

$= 2m + 1$ is even? then this $m \in \mathbb{Z}$ contradiction

with assumption

$$n^3 + 5 \text{ is odd}$$

done #

3. Prove that if n is an integer and $3n + 2$ is even, then n is even using a proof by contradiction.

Solution:

$$\begin{aligned}
 & 3(2k+1) + 2 = 6k+5 \\
 & = 6k+4 + 1 \quad \text{1 is odd} \\
 & 2m+1 \text{ is odd} \quad \text{m is even} \quad \text{then this is} \\
 & \text{a contradiction with assumption} \\
 & 3n+2 \text{ is even done} \#
 \end{aligned}$$

4. Prove that $\sqrt{3}$ is irrational by giving a proof by contradiction.

Solution:

$$\begin{aligned}
 \sqrt{3} &= \frac{a}{b} \quad a, b \in \mathbb{Z} \text{ and } \gcd(a, b) = 1 \\
 \Rightarrow 3 &= \frac{a^2}{b^2} \Rightarrow a^2 - 3b^2 \Rightarrow 3|a^2 \Rightarrow 3|a \\
 (\exists m: m \in \mathbb{Z}) \Rightarrow a^2 - 3b^2 &= 3m^2 \\
 3|b^2 \Rightarrow 3|b &\Rightarrow 3|a \wedge 3|b \\
 \text{this is a contradiction where} \\
 \gcd(a, b) &= 1 \\
 &\text{done} \#
 \end{aligned}$$

5. Prove that $\sqrt{5}$ is irrational by giving a proof by contradiction.

Solution:

6. Let $\sqrt{5}$ is irrational, prove that $2 - 3\sqrt{5}$ is irrational number. $(\text{CSS HM } 2 - 3\sqrt{5} \in \mathbb{Q})$

Solution: $2 - 3\sqrt{5} = \frac{a}{b}$ $a, b \in \mathbb{Z}$

$$= 3\sqrt{5} = 2 - \frac{a}{b} = \frac{2b - a}{b} \stackrel{?}{=} \sqrt{5} \stackrel{?}{=} \frac{2b - a}{3b}$$

$3b \neq 0$
 $(*) \in \mathbb{Q}$

this is a contradiction where
 $\sqrt{5} \notin \mathbb{Q}$

done ~~#~~

7. Let $\sqrt{2}$ is irrational, prove that $1 + 3\sqrt{2}$ is irrational number.

Solution:

8. Let $\sqrt{6}$ is irrational, prove that $-3 + 2\sqrt{6}$ is irrational number.

Solution:

ex 2

9. Let $\sqrt{5}$ is irrational, prove that $\frac{\sqrt{5}-3}{2}$ is irrational number.

Solution:

$$\frac{\sqrt{5}-3}{2} \geq \frac{c}{b} \Rightarrow \sqrt{5} \geq \frac{2c+d}{b} = \sqrt{5} = \frac{3c+d}{b} \quad \text{b} \neq 0$$

$$\sqrt{5} = \frac{2c+3d}{b} \in \mathbb{Q} \Rightarrow \sqrt{5} \in \mathbb{Q}: \text{this is a contradiction where } \sqrt{5} \notin \mathbb{Q}$$

10. Let $\sqrt{2}$ is irrational, prove that $\frac{3+\sqrt{8}}{3}$ is irrational number.

$$\frac{3+\sqrt{8}}{3} = \frac{c}{b} \Rightarrow 3+\sqrt{8} = \frac{3c}{b} - \sqrt{8} = \frac{3c-3b}{b} \quad b \neq 0$$

$$\frac{3c-3b}{b} = 2\sqrt{2} = \frac{3c-3b}{b} - \sqrt{2} = \frac{3c-3b}{b} \quad \text{b} \neq 0$$

this is a contradiction where $\sqrt{2} \notin \mathbb{Q}$
done #

11. Let m, n are rational numbers : $n \neq 0$ and $\sqrt{5}$ is irrational, prove that $\frac{\sqrt{5}}{2n} - m$ is irrational number.

Solution:

$$\frac{\sqrt{5}}{2n} - m \in \mathbb{Q} \Rightarrow \frac{\sqrt{5}}{2n} - m = \frac{c}{b} \quad b \neq 0$$

$$\frac{\sqrt{5}}{2n} = \frac{c}{b} + m = \frac{c+mb}{b} \quad b \neq 0$$

$$\sqrt{5} = \frac{b(c+mb)}{b} \quad (*) \in \mathbb{Z}$$

$\sqrt{5} \in \mathbb{Q}$: this is a contradiction where $\sqrt{5} \notin \mathbb{Q}$
done #

12. Let $x, y, z \in \mathbb{R}$: $x + y + z = 20$, use a proof by contradiction to show that $x \geq 10$ or $y \geq 6$ or $z \geq 4$.

Solution:

$$\begin{array}{l} \neg x < 10 \\ \neg y < 6 \\ \neg z < 4 \end{array}$$

$$x + y + z < 20$$



this is a contradiction

where $x + y + z \leq 20$ done

~~#~~

13. Let $x, y, z \in \mathbb{R}$: $x + y + z = 21$, use a proof by contradiction to show that $x \geq 8$ or $y \geq 7$ or $z \geq 6$.

Solution: Suppose $x < 8$,

$$\begin{array}{l} \neg y < 7 \\ \neg z < 6 \end{array}$$

$$x + y + z < 21$$



this is a contradiction
where $x + y + z = 21$

- 14.** Use a proof by contradiction to prove that the product of an irrational number and a non-zero rational number is irrational.

Solution:

$$x = \frac{c \neq 0}{d \neq 0} : c, d \in \mathbb{Z} \text{ (sum)} \quad xy \in \mathbb{Q} \Rightarrow xy = \frac{c}{d}$$

$$xy = \frac{c}{d} \Rightarrow y = \frac{c}{d} \cdot \frac{d}{b} = \frac{cd}{db} \quad \begin{cases} c \in \mathbb{Z} \\ cd \in \mathbb{Z} \end{cases}$$

$y = \frac{cd}{db}$ this is a contradiction

where $y \notin \mathbb{Q}$ done #

- 15.** Use a proof by contradiction to prove that $\nexists m, n \in \mathbb{Z}$, such that $4n + 6m = 11$

Solution: (Assume) $m, n \in \mathbb{Z}$

$$4n + 6m = 11 \Rightarrow 2(2n + 3m) = 11$$

$11 = 2t$ even $\Rightarrow 2 \mid 11$ $t \in \mathbb{Z}$

this is cont.

16. Let $x, y \in \mathbb{R}$: $xy \leq 0$, use a proof by contradiction to show that
 $x \geq 0$ or $y \geq 0$.

Solution:

17. Let a and b be positive real numbers such that $a^3b^2 > 72$. Give a proof by contradiction to show that $a > 2$ or $b > 3$.

18. Prove or disprove that the product of two irrational numbers is irrational.

Solution:

19. Prove or disprove that the sum of two irrational numbers is irrational.

Solution:

20. Use a proof by contradiction to prove that $x + \frac{1}{x} \geq 2$ for $\forall x \in \mathbb{R}^*$.

Solution:

21. Use a proof by contradiction to prove that $\nexists a, b \in \mathbb{Z}^+$, such that

$$a + b = 500 \text{ and } \gcd(a, b) = 7 \quad (\text{SS un } 3 \text{ cpl})$$

Solution:

$$a + b = 500 \quad (\text{and } 9 \mid \gcd(a, b))$$

$$\Rightarrow \exists k \in \mathbb{Z}^+ : a = 7k \quad \text{and} \quad b = 7h \quad (k, h \in \mathbb{Z}^+)$$

$$a + b = 7k + 7h = 500 \Rightarrow 7(k+h) = 500$$

$$= 7 \cdot 71 = 500 \Rightarrow 7 \mid 500 \text{ cont}$$

where $7 \nmid 500$ done $\#$

22. Prove or disprove that if $a^2|b^3$ then $a|b : a, b \in \mathbb{N}$.

Solution:

23. Let m be an odd integer. Give a proof by contradiction to show that $m^2 - 3$ is not a multiple of 4. *let m is odd and suppose 3|m²-3*

Solution:

$$\Rightarrow m^2 - 3 = 4h : h \in \mathbb{Z}$$

$$m = 2k + 1 : k \in \mathbb{Z} \Rightarrow m^2 = 4k^2 + 4k + 1$$

$$4k^2 + 4k + 1 - 3 = 4h = 4k^2 + 4k - 4h = 2$$

$$2 = 4(k^2 + k + h) = 4m \Rightarrow 4|2$$

this is a contradiction

where $2 \not| 4$ but $4|2$

done ~~#~~

Let a be a real number such that a^2 is irrational. Use a proof by contradiction to show that $a + 1$ is irrational.

Solution:

$$a + 1 = \frac{d}{b} \quad ; \quad d, b \in \mathbb{Z}$$

$$d \neq 0$$

$$a + 1 = \frac{d}{b} \Rightarrow a = \frac{d}{b} - 1 = \frac{d-b}{b}$$

$$a^2 = \left(\frac{d-b}{b} \right)^2 = \frac{(d-b)^2}{b^2} =$$

$$\frac{d^2 - 2db + b^2}{b^2}$$



this is a contradiction

where $a^2 \notin \mathbb{Q}$

done



PROOF BY CASES A proof by cases must cover all possible cases that arise in a theorem. We illustrate proof by cases with a couple of examples. In each example, you should check that all possible cases are covered.

EXAMPLE 1 Prove that if n is an integer, then $n^2 \geq n$.

Solution: We can prove that $n^2 \geq n$ for every integer by considering three cases, when $n = 0$, when $n \geq 1$, and when $n \leq -1$. We split the proof into three cases because it is straight forward to prove the result by considering zero, positive integers, and negative integers separately.

Case (i): When $n = 0$, because $0^2 = 0$, we see that $0^2 \geq 0$. It follows that $n^2 \geq n$ is true in this case.

Case (ii): When $n \geq 1$, when we multiply both sides of the inequality $n \geq 1$ by the positive Integer n , we obtain $n \cdot n \geq n \cdot 1$. This implies that $n^2 \geq n$ for $n \geq 1$.

Case (iii): In this case $n \leq -1$. However, $n^2 \geq 0$. It follows that $n^2 \geq n$.

Because the inequality $n^2 \geq n$ holds in all three cases, we can conclude that if n is an integer, then $n^2 \geq n$.

EXAMPLE 4 Use a proof by cases to show that $|xy| = |x||y|$, where x and y are real

numbers. (Recall that $|a| = \begin{cases} a & : a \geq 0 \\ -a & : a < 0 \end{cases}$)

Solution: In our proof of this theorem, we remove absolute values using the fact that $|a| = a$ when $a \geq 0$ and $|a| = -a$ when $a < 0$. Because both $|x|$ and $|y|$ occur in our formula, we will need four cases: (i) x and y both nonnegative,

(ii) x nonnegative and y is negative,

(iii) x negative and y nonnegative

(iv) x negative and y negative.

We denote by $p1, p2, p3$, and $p4$, the proposition stating the assumption for each of these four cases, respectively.

(Note that we can remove the absolute value signs by making the appropriate choice of signs within each case.)

Case (i): We see that $p1 \rightarrow q$ because $xy \geq 0$ when $x \geq 0$ and $y \geq 0$,

$$\text{so that } |xy| = xy = |x||y|.$$

Case (ii): To see that $p2 \rightarrow q$, note that if $x \geq 0$ and $y < 0$, then $xy \leq 0$,

$$\text{so that } |xy| = -xy = x(-y) = |x||y|. \text{ (Here, because } y < 0, \text{ we have } |y| = -y\text{.)}$$

Case (iii): To see that $p3 \rightarrow q$, we follow the same reasoning as the previous case with the roles of x and y reversed.

Case (iv): To see that $p4 \rightarrow q$, note that when $x < 0$ and $y < 0$, it follows that $xy > 0$.

$$\text{Hence, } |xy| = xy = (-x)(-y) = |x||y|.$$

Because $|xy| = |x||y|$ holds in each of the four cases and these cases exhaust all possibilities, we can conclude that $|xy| = |x||y|$, whenever x and y are real numbers.

EXAMPLE 5 Show that if x and y are integers and both xy and $x + y$ are even, then both x and y are even.

Solution: We will use proof by contraposition, the notion of without loss of generality, and proof by cases. First, suppose that x and y are not both even. That is, assume that x is odd or that y is odd (or both). Without loss of generality, we assume that x is odd, so that $x = 2m + 1$ for some integer k . To complete the proof, we need to show that

xy is odd or $x + y$ is odd. Consider two cases:

- (i) y even
- (ii) y odd.

In (i), $y = 2n$ for some integer n , so that $x + y = (2m + 1) + 2n = 2(m + n) + 1$ is odd.

In (ii), $y = 2n + 1$ for some integer n , so that $xy = (2m + 1)(2n + 1) = 4mn + 2m + 2n + 1 = 2(2mn + m + n) + 1$ is odd. This completes the proof by contraposition.

(Note that our use of without loss of generality within the proof is justified because the proof when y is odd can be obtained by simply interchanging the roles of x and y in the proof we have given.)

Exercises

1. Prove that if x and y are real numbers, then $\max(x, y) + \min(x, y) = x + y$.

[*Hint:* Use a proof by cases, with the two cases corresponding to $x \geq y$ and $x < y$, respectively.]

Solution: If $x \leq y$, then $\max(x, y) + \min(x, y) = y + x = x + y$.

If $x \geq y$, then $\max(x, y) + \min(x, y) = x + y$.

Because these are the only two cases, the equality always holds .

2. Use a proof by cases, to prove that $n^2 + 1 \geq 2n$ when n is a positive integer with $1 \leq n \leq 4$.

Solution: $1^2 + 1 = 2 \geq 2 = 2(1)$;

$$2^2 + 1 = 5 \geq 4 = 2(2);$$

$$3^2 + 1 = 10 \geq 8 = 2(3);$$

$$4^2 + 1 = 17 \geq 16 = 2(4)$$