# Package 'MultinomFourier'

October 13, 2026

**Type** Package

**Title** p-Values of Multinomial Goodness-of-Fit Tests by Fourier Series Inversion

**Version** 1.0

**Author** Charles Thraves [aut], Pablo Rivas [aut, cre], Felipe Subiabre [aut]

**Maintainer** Pablo Rivas <pablo.rivas.g@ug.uchile.cl>

**Description** Computes p-values of multinomial goodness-of-fit tests based on an additive statistic: the log-likelihood ratio, Pearson's chi-squared, the Cressie-Read power divergence, and the exact multinomial test. The main function, pval_fourier(), writes the p-value as a Fourier series in the moment-generating function of the statistic and sums it until an adaptive stopping rule is met, which makes small p-values of large instances computable. The function pval_exact() computes the exact p-value by branch pruning with bounds from a dynamic program, and pval_flexible() chooses between the two according to the size of the instance.

**License** GPL (>= 2)

**Encoding** UTF-8

**Imports** parallel

**Suggests** knitr,
rmarkdown,
ggplot2

**VignetteBuilder** knitr

**Config/roxygen2/version** 8.1.0

## Contents


MultinomFourier-package . . . . . 2
print.multinomfourier . . . . . 3
pval_exact . . . . . 3
pval_flexible . . . . . 6
pval_fourier . . . . . 8

**Index** **13**

`MultinomFourier-package` *MultinomFourier: p-values of multinomial goodness-of-fit tests*

## Description

Computes p-values of multinomial goodness-of-fit tests based on an additive statistic: the log-likelihood ratio, Pearson's chi-squared, the Cressie-Read power divergence, and the exact multinomial test. The main function, pval_fourier(), writes the p-value as a Fourier series in the moment-generating function of the statistic and sums it until an adaptive stopping rule is met, which makes small p-values of large instances computable. The function pval_exact() computes the exact p-value by branch pruning with bounds from a dynamic program, and pval_flexible() chooses between the two according to the size of the instance.

## Details

`MultinomFourier` computes the p-value $P\{T(Y) \geq T(x)\}$ of a multinomial goodness-of-fit test, where $x$ is the observation (the vector of observed counts in the $K$ categories, with $N = \sum_k x_k$ trials), $Y \sim \text{Multinomial}(N, p)$ under the null hypothesis with category probabilities $p$, and $T$ is an additive statistic: the log-likelihood ratio $G^2$, Pearson's $\mathcal{X}^2$, the Cressie–Read power divergence, or the probability mass function of the multinomial (exact multinomial test). It is aimed at small p-values, far in the tail, and at instances where the support of the multinomial is too large to sum over.

The package provides three functions:

- `pval_fourier`: writes the p-value as a Fourier series whose terms are values of the moment-generating function of the statistic, and sums it until an adaptive stopping rule is met. Its cost grows slowly with $N$ and $K$; its accuracy depends on the statistic (fast convergence for $G^2$ and the probability mass function, possibly none for $\mathcal{X}^2$).
- `pval_exact`: exact p-value by branch pruning with bounds from a dynamic program, without enumerating the support. Feasible for a moderate number of categories and trials.
- `pval_flexible`: chooses between the two according to the size of the support.

All three return an object of class `"multinomfourier"`; check its `status` component before using the p-value of `pval_fourier`.

## Author(s)

**Maintainer**: Pablo Rivas <pablo.rivas.g@ug.uchile.cl>

Authors:

- Pablo Rivas <pablo.rivas.g@ug.uchile.cl>
- Charles Thraves
- Felipe Subiabre

## References

Subiabre, F. and Thraves, C. Efficient computation of p-values for multinomial tests. Manuscript.

## See Also

Report bugs at https://github.com/PabRvss/MultinomFourier/issues.

## print.multinomfourier *Print method for multinomfourier objects*

### Description

Prints the method, statistic, p-value, status and total runtime and, for the Fourier method, also the tilt gamma, the time spent choosing it, the period half-width W and the number of terms.

### Usage

```
## S3 method for class 'multinomfourier'
print(x, ...)
```

### Arguments

| | |
|---|---|
| x | Object of class "multinomfourier" returned by pval_fourier, pval_exact or pval_flexible. |
| ... | Additional arguments (currently unused). |

### Value

Invisibly returns x.

### Examples

```
# Note: Reported p-values in comments are illustrative and may vary slightly
# across versions or due to changes in numerical criteria.
p <- c(0.1, 0.2, 0.3, 0.4)
x <- c(35, 30, 60, 75)

# Fourier method: also shows gamma, W and the number of terms
fit_fourier <- pval_fourier(x, p, stat = "llr")
fit_fourier              # same as print(fit_fourier)

# Exact method
fit_exact <- pval_exact(x, p, stat = "llr")
fit_exact
```

## pval_exact *Exact multinomial test p-value*

### Description

Computes the exact p-value $P\{T(Y) \geq T(x)\}$, $Y \sim \text{Multinomial}(N, p)$, of a multinomial goodness-of-fit test, up to floating-point rounding. Instead of enumerating the support, it prunes whole groups of outcomes using bounds on the statistic (see Details). pval_exhaustive() is a deprecated alias.

## Usage

```
pval_exact(
  x,
  p = rep(1/length(x), length(x)),
  stat = c("llr", "pd", "chi2", "pmf"),
  lambda = NULL,
  rescale_p = FALSE,
  n_threads = min(2L, parallel::detectCores()),
  max_time = NULL,
  verbose = FALSE,
  ...
)

pval_exhaustive(...)
```

## Arguments

| | |
|---|---|
| x | Vector of observed counts, one per category (non-negative integers, without missing values). Their sum is the number of trials $N$. |
| p | Vector of category probabilities under the null hypothesis, all strictly positive. They must sum to 1 (up to a tolerance of `sqrt(.Machine$double.eps)`), unless `rescale_p = TRUE`. Default: equiprobable categories. |
| stat | Test statistic: `"llr"` (log-likelihood ratio $G^2$, the default), `"chi2"` (Pearson's $\mathcal{X}^2$), `"pd"` (Cressie–Read power divergence with parameter `lambda`) or `"pmf"` (probability mass function of the multinomial, i.e. the exact multinomial test). See Details. |
| lambda | Parameter $\lambda$ of the power-divergence statistic; used only when `stat = "pd"`, and then required (there is no default). Must be a single finite real number strictly greater than -1 ($\lambda > -1$); values $\lambda \leq -1$ produce an error. A warning is issued if it is given for another statistic. |
| rescale_p | Logical; if `TRUE`, `p` is rescaled to `p / sum(p)`, so it can be given as any vector of positive weights. If `FALSE` (default) and `p` does not sum to 1, an error is given. Same argument and default as `rescale.p` in `chisq.test`. |
| n_threads | Number of threads (positive integer, capped at the number of available cores). Default `min(2L, parallel::detectCores())`. |
| max_time | Optional time limit in seconds. `NULL` (default) means no limit. The limit is only checked between blocks of values of the first category, so a run can exceed it by a wide margin; if it is reached, `pval` is `NA` and `status` is 1. |
| verbose | Logical; if `TRUE`, prints progress information. Default `FALSE`. |
| ... | Advanced options passed to the C engine. Unknown names produce a warning. |

## Details

The statistics are those of `pval_fourier`. Because the statistic is additive, a dynamic program over the $(N+1)K$ states (category, trials left) first computes the exact minimum and maximum that the remaining categories can add to it, at cost $O(KN^2)$. Counts are then assigned one category at a time, in decreasing order of probability; whenever the bounds show that every completion of a partial assignment is (or none is) at least as extreme as the observed data, its whole probability is added (or it is discarded) without visiting it. Only partial assignments near the boundary $T(y) = T(x)$ are expanded.

The cost still grows quickly with the number of categories $K$. As a rough guide, in our tests with the log-likelihood ratio on 8 threads, instances with $K \leq 6$ and $N \leq 1000$ took under five minutes, while $K \geq 8$ with $N \geq 500$ did not finish within an hour. For large instances use `pval_fourier`.

## Value

An object of class `"multinomfourier"`: a list with components

**x, p** The input counts and null probabilities (`p` after rescaling, if `rescale_p = TRUE`).

**pval** The exact p-value, up to floating-point rounding; `NA` if `max_time` was reached.

**stat** Name of the test statistic, e.g. `"Log-Likelihood Ratio"` or `"power-divergence (lambda = 0.6667)"`.

**method** `"exact"`.

**status** `0`: the computation finished; `1`: `max_time` was reached before it finished.

**message** Human-readable version of `status`. For `status = 0` it reads `"Converged"`, a label shared with `pval_fourier`.

**time** Total wall-clock time in seconds.

The components that describe the Fourier series are also present, so that results of both methods have the same structure, but are always `NA`: `gamma`, `time_gamma`, `W`, `n_terms`, `Q`, `D_Q`, `err_rel_ext`, `err_rel_est`, `n_eff` and `nu_max_ratio`.

## See Also

`pval_fourier`, `pval_flexible`.

## Examples

```
# Note: Reported p-values in comments are illustrative and may vary slightly
# across versions or due to changes in numerical criteria.
p <- c(0.1, 0.2, 0.3, 0.4)
x <- c(35, 30, 60, 75)   # N = 200

# Log-likelihood ratio
fit <- pval_exact(x, p, stat = "llr")
fit
fit$pval                 # 0.00684
fit$status               # 0: computation finished

# Probability mass function (exact multinomial test)
fit_pmf <- pval_exact(x, p, stat = "pmf")
fit_pmf$pval             # 0.00578

# Pearson's chi-square
fit_chi2 <- pval_exact(x, p, stat = "chi2")
fit_chi2$pval            # 0.00294
```

pval_flexible *Multinomial test p-value, choosing the method automatically*

## Description

Computes the p-value with `pval_exact` when $\log_{10}$ of the support size is strictly less than `enum_cutoff`, and with `pval_fourier` when it is greater than or equal to `enum_cutoff`.

## Usage

```
pval_flexible(
  x,
  p = rep(1/length(x), length(x)),
  stat = c("llr", "pd", "chi2", "pmf"),
  lambda = NULL,
  rescale_p = FALSE,
  enum_cutoff = 10,
  undersampling = 1,
  rel_eps = 0.001,
  B = 50,
  max_terms = 10000,
  max_time = NULL,
  exact_terms = FALSE,
  n_threads = min(2L, parallel::detectCores()),
  precompute = TRUE,
  verbose = FALSE,
  ...
)
```

## Arguments

| | |
|---|---|
| `x` | Vector of observed counts, one per category (non-negative integers, without missing values). Their sum is the number of trials $N$. |
| `p` | Vector of category probabilities under the null hypothesis, all strictly positive. They must sum to 1 (up to a tolerance of `sqrt(.Machine$double.eps)`), unless `rescale_p = TRUE`. Default: equiprobable categories. |
| `stat` | Test statistic: `"llr"` (log-likelihood ratio $G^2$, the default), `"chi2"` (Pearson's $\mathcal{X}^2$), `"pd"` (Cressie–Read power divergence with parameter `lambda`) or `"pmf"` (probability mass function of the multinomial, i.e. the exact multinomial test). See Details. |
| `lambda` | Parameter $\lambda$ of the power-divergence statistic; used only when `stat = "pd"`, and then required (there is no default). Must be a single finite real number strictly greater than -1 ($\lambda > -1$); values $\lambda \leq -1$ produce an error. A warning is issued if it is given for another statistic. |
| `rescale_p` | Logical; if `TRUE`, `p` is rescaled to `p / sum(p)`, so it can be given as any vector of positive weights. If `FALSE` (default) and `p` does not sum to 1, an error is given. Same argument and default as `rescale.p` in `chisq.test`. |
| `enum_cutoff` | Threshold on $\log_{10}$ of the support size $\binom{N+K-1}{K-1}$ (the number of possible count vectors). Below it ($<$ enum_cutoff) the exact method is used; at or above it ($\geq$ enum_cutoff) the Fourier method is used. Default 10. See Details. |

`undersampling` Positive factor dividing the half-width $W$ of the Fourier period. The default 1 uses the smallest $W$ for which the series is exact. Values below 1 enlarge $W$ (still exact, but more terms are needed); values above 1 shrink $W$ below that minimum, so the series converges to a wrong value (aliasing). Intended for experimentation only.

`rel_eps` Relative tolerance of the stopping rule ($\geq 0$). Default 0.001. It controls how stable the partial sums must be before the series is truncated; it is *not* a bound on the error of the returned p-value.

`B` Non-negative integer used by the term-size stopping rule: the series stops after `B` consecutive terms each smaller, relative to the partial sum, than `rel_eps / B`. Default 50; `B = 0` uses an internal value.

`max_terms` Maximum number of series terms (positive integer). Default 10000. If it is reached before the stopping rule is met, `status` is 2.

`max_time` Optional time limit in seconds, passed to the chosen method.

`exact_terms` Logical; if `FALSE` (default), evaluates the Fourier terms using the fast Poissonized Cauchy-integral evaluator. If `TRUE`, evaluates each term exactly via polynomial convolution.

`n_threads` Number of threads, passed to whichever method is chosen. Default `min(2L, parallel::detectCores())`.

`precompute` Logical; if `TRUE` (default), precomputes the FFT plan when evaluating terms via polynomial convolution (`exact_terms = TRUE`).

`verbose` Logical; if `TRUE`, prints the partial sums and diagnostic information while running. Default `FALSE`.

`...` Advanced options passed to the C engine, for example `print_freq` (how often partial sums are printed when `verbose = TRUE`; default 100) or `extrapolate` (0: no extrapolation; 1: report the extrapolated value; 2: also stop on it, the default). Unknown names produce a warning.

## Details

The cost of `pval_exact` depends strongly on the observed counts, not only on the support size, and it is highest far in the tail, where small p-values are computed. With counts drawn from the null hypothesis it takes about one second on 2 threads at a support of $10^{12}$ to $10^{13}$ outcomes (log-likelihood ratio, $K = 5, 6, 7$); with counts far in the tail it can take tens of seconds at a support of $10^{11}$ (for example $K = 6$, $N = 400$, $p \approx 10^{-33}$). The default `enum_cutoff = 10` is a simple, conservative value that keeps the exact method for instances where it is fast in both situations.

When the Fourier method is chosen and does not meet its stopping rule (`status != 0`), the result is returned as is; there is no automatic fallback to the exact method.

## Value

An object of class `"multinomfourier"`, as described in `pval_fourier`; its `method` component (`"exact"` or `"fourier"`) shows which method was used.

## See Also

`pval_fourier`, `pval_exact`.

## Examples

```
# Note: Reported p-values in comments are illustrative and may vary slightly
# across versions or due to changes in numerical criteria.
# Small instance (K = 4, N = 200, about 10^6 outcomes): exact method
p <- c(0.1, 0.2, 0.3, 0.4)
x <- c(35, 30, 60, 75)
fit <- pval_flexible(x, p, stat = "llr")
fit
fit$method              # "exact"
fit$pval                # 0.00684
fit$status              # 0: computation finished

# Large instance (K = 8, N = 2000, about 10^19 outcomes): Fourier method
x_large <- c(240, 260, 250, 270, 230, 250, 245, 255)
fit_large <- pval_flexible(x_large, stat = "llr")
fit_large$method        # "fourier"
fit_large$pval          # 0.761
fit_large$status        # 0: stopping rule met
```

---

`pval_fourier` *Multinomial test p-value by Fourier series inversion*

---

## Description

Computes the p-value of a multinomial goodness-of-fit test,

$$p = P\{T(Y) \geq T(x)\}, \qquad Y \sim \text{Multinomial}(N, p), \quad N = \sum_k x_k,$$

for an additive test statistic $T$, by summing a truncated Fourier series whose terms are values of the moment-generating function of the statistic. The series is truncated by an adaptive stopping rule, so the result is an approximation whose accuracy depends on the statistic and the instance; always check the returned `status`.

## Usage

```
pval_fourier(
  x,
  p = rep(1/length(x), length(x)),
  stat = c("llr", "pd", "chi2", "pmf"),
  lambda = NULL,
  rescale_p = FALSE,
  undersampling = 1,
  rel_eps = 0.001,
  B = 50,
  max_terms = 10000,
  max_time = NULL,
  n_threads = min(2L, parallel::detectCores()),
  exact_terms = FALSE,
  precompute = TRUE,
  verbose = FALSE,
  ...
)
```

## Arguments

| | |
|---|---|
| x | Vector of observed counts, one per category (non-negative integers, without missing values). Their sum is the number of trials $N$. |
| p | Vector of category probabilities under the null hypothesis, all strictly positive. They must sum to 1 (up to a tolerance of `sqrt(.Machine$double.eps)`), unless `rescale_p = TRUE`. Default: equiprobable categories. |
| stat | Test statistic: `"llr"` (log-likelihood ratio $G^2$, the default), `"chi2"` (Pearson's $\mathcal{X}^2$), `"pd"` (Cressie–Read power divergence with parameter `lambda`) or `"pmf"` (probability mass function of the multinomial, i.e. the exact multinomial test). See Details. |
| lambda | Parameter $\lambda$ of the power-divergence statistic; used only when `stat = "pd"`, and then required (there is no default). Must be a single finite real number strictly greater than -1 ($\lambda > -1$); values $\lambda \leq -1$ produce an error. A warning is issued if it is given for another statistic. |
| rescale_p | Logical; if `TRUE`, `p` is rescaled to `p / sum(p)`, so it can be given as any vector of positive weights. If `FALSE` (default) and `p` does not sum to 1, an error is given. Same argument and default as `rescale.p` in `chisq.test`. |
| undersampling | Positive factor dividing the half-width $W$ of the Fourier period. The default 1 uses the smallest $W$ for which the series is exact. Values below 1 enlarge $W$ (still exact, but more terms are needed); values above 1 shrink $W$ below that minimum, so the series converges to a wrong value (aliasing). Intended for experimentation only. |
| rel_eps | Relative tolerance of the stopping rule ($\geq 0$). Default 0.001. It controls how stable the partial sums must be before the series is truncated; it is *not* a bound on the error of the returned p-value. |
| B | Non-negative integer used by the term-size stopping rule: the series stops after `B` consecutive terms each smaller, relative to the partial sum, than `rel_eps / B`. Default 50; `B = 0` uses an internal value. |
| max_terms | Maximum number of series terms (positive integer). Default 10000. If it is reached before the stopping rule is met, `status` is 2. |
| max_time | Optional time limit in seconds. `NULL` (default) means no limit. When it is reached, `status` is 1. |
| n_threads | Number of threads used to evaluate the series terms (positive integer, capped at the number of available cores). Default `min(2L, parallel::detectCores())`. |
| exact_terms | Logical; if `FALSE` (default), evaluates the Fourier terms using the fast Poissonized Cauchy-integral evaluator. If `TRUE`, evaluates each term exactly via polynomial convolution. |
| precompute | Logical; if `TRUE` (default), precomputes the FFT plan when evaluating terms via polynomial convolution (`exact_terms = TRUE`). |
| verbose | Logical; if `TRUE`, prints the partial sums and diagnostic information while running. Default `FALSE`. |
| ... | Advanced options passed to the C engine, for example `print_freq` (how often partial sums are printed when `verbose = TRUE`; default 100) or `extrapolate` (0: no extrapolation; 1: report the extrapolated value; 2: also stop on it, the default). Unknown names produce a warning. |

## Details

**Statistics.** All supported statistics are additive, $T(y) = \sum_k f_k(y_k)$, which is what the method requires. With $E_k = Np_k$:

- `"llr"`: $G^2 = 2\sum_k y_k \log(y_k/E_k)$.
- `"chi2"`: $\mathcal{X}^2 = \sum_k (y_k - E_k)^2/E_k$.
- `"pd"`: $2I^\lambda = \frac{2}{\lambda(\lambda+1)} \sum_k y_k[(y_k/E_k)^\lambda - 1]$ (Cressie and Read, 1984); $\lambda \to 0$ gives $G^2$ and $\lambda = 1$ gives $\mathcal{X}^2$.
- `"pmf"`: the probability mass function $P(y)$ of the multinomial under the null hypothesis (exact multinomial test). Small values are the extreme ones, so the p-value is $P\{P(Y) \le P(x)\}$, the total probability of the outcomes that are at most as likely as the observed one.

**Which statistics converge.** The p-value is written exactly as a Fourier series for every additive statistic, but the number of terms needed for a given accuracy depends on the statistic. For `"llr"` and `"pmf"` the series typically converges in a few hundred to a few thousand terms. For `"chi2"`, and often for `"pd"` unless $\lambda$ is close to 0 ($\lambda = 2/3$ has failed on real data), the values of the statistic lie on or near a lattice, the terms do not decay, and the series can fail to converge within `max_terms` (`status = 2`) or even return a negative value. In that case use `pval_exact` when it is feasible.

**Stopping rule and reported value.** The partial sums are accelerated with Wynn's epsilon algorithm (in the safeguarded form of QUADPACK's DQELG) and smoothed with a running median of five. The series stops as soon as either (i) `B` consecutive terms are each smaller than `rel_eps / B` relative to the partial sum, or (ii) the smoothed extrapolated values vary by at most a relative `rel_eps` over a trailing window covering the last half of the terms. The reported p-value is the last smoothed extrapolated value.

**Term evaluation.** With `exact_terms = FALSE` (default), terms are evaluated via a Poissonized Cauchy-integral evaluator, which is significantly faster on large instances. With `exact_terms = TRUE`, terms are evaluated via exact polynomial convolution.

## Value

An object of class `"multinomfourier"`: a list with components

**x, p** The input counts and null probabilities (`p` after rescaling, if `rescale_p = TRUE`).

**pval** The computed p-value.

**stat** Name of the test statistic, e.g. `"Log-Likelihood Ratio"` or `"power-divergence (lambda = 0.6667)"`.

**method** `"fourier"`.

**time** Total wall-clock time in seconds.

**gamma** The exponential tilt $\gamma$ applied to the statistic so that the series terms stay within floating-point range (a saddlepoint choice).

**time_gamma** Time in seconds spent choosing `gamma`.

**W** Half-width of the Fourier period, $\max(s_{\max}, -s_{\min})$ over the range $[s_{\min}, s_{\max}]$ of the statistic minus its observed value, divided by `undersampling`.

**n_terms** Number of series terms summed.

**status** `0`: the stopping rule was met (the message reads `"Converged"`); `1`: `max_time` was reached; `2`: `max_terms` was reached without meeting the stopping rule, so `pval` is unreliable; `3`: the search for `gamma` failed; `4`: numerical overflow in the evaluation of the terms. With a status other than 0, `pval` is the last value reached or `NA`.

**message** Human-readable version of `status`.

**err_rel_ext** Relative spread of the smoothed extrapolated values over the final window, i.e. the quantity that rule (ii) compares with `rel_eps`. When the series stopped by rule (i) it can be larger than `rel_eps`.

**err_rel_est** Estimate of the relative truncation error of the raw partial sum (not of the extrapolated value that is reported), based on a random-phase argument; in our tests it tends to underestimate the error.

**n_eff** Effective number of support points implied by the size of the late terms; used by `err_rel_est`.

**nu_max_ratio** Largest size of the late transform values relative to the first one; values near 1 mean the terms are not decaying (a lattice-like statistic).

**Q** Number of equispaced points on the saddlepoint circle used to extract the multinomial coefficient in each term of the series (a $Q$-point trapezoid rule for Cauchy's integral). It is chosen automatically, by doubling from a power of two until the aliasing diagnostic `D_Q` falls below $10^{-10}$. `NA` with `exact_terms = TRUE`.

**D_Q** Aliasing diagnostic $D(Q)$. $D(Q)$ is a computable lower bound on $\alpha(Q)$, which provides a uniform upper bound on the aliasing error of each Fourier-series term. For the probability-mass statistic (`"pmf"`), $D(Q)$ approximates $\alpha(Q)$ to first order. For the log-likelihood-ratio statistic (`"llr"`), $D(Q)$ remains a lower bound on $\alpha(Q)$, but no guarantee of closeness is available. Stopping the doubling of $Q$ when $D(Q)$ falls below a threshold does not by itself guarantee that the aliasing error is strictly below that threshold. `D_Q` measures an error separate from the truncation of the series (reported by `err_rel_ext` and `err_rel_est`). It is `0` when `Q` is large enough for the Cauchy formula to be exact (no aliasing possible), and `NA` with `exact_terms = TRUE`.

The last six components are diagnostics of the series. The four about truncation (`err_rel_ext`, `err_rel_est`, `n_eff` and `nu_max_ratio`) are heuristics, not error bounds. The names `Gamma` and `Time_gamma` used by earlier versions still work, with a deprecation warning.

## References

Subiabre, F. and Thraves, C. Efficient computation of p-values for multinomial tests. Manuscript.

Cressie, N. and Read, T. R. C. (1984). Multinomial goodness-of-fit tests. *Journal of the Royal Statistical Society, Series B*, 46(3), 440–464.

## See Also

`pval_exact` for the exact p-value, `pval_flexible` to choose between the two automatically.

## Examples

```
# Note: Reported p-values and term counts in comments are illustrative and
# may vary slightly across versions or due to changes in numerical criteria.
p <- c(0.1, 0.2, 0.3, 0.4)
x <- c(35, 30, 60, 75)   # N = 200

# Log-likelihood ratio
fit <- pval_fourier(x, p, stat = "llr")
fit
fit$pval                 # 0.00688
fit$status               # 0: stopping rule met
fit$n_terms              # 1208
```

```
# Probability mass function (exact multinomial test)
fit_pmf <- pval_fourier(x, p, stat = "pmf")
fit_pmf$pval             # 0.00577
fit_pmf$status           # 0: stopping rule met

# Power divergence: lambda is required
fit_pd <- pval_fourier(x, p, stat = "pd", lambda = 2/3)
fit_pd$pval              # 0.00384
fit_pd$status            # 0: stopping rule met
```

# Index

∗ **package**
MultinomFourier-package, 2

chisq.test, *4*, *6*, *9*

MultinomFourier
(MultinomFourier-package), 2
MultinomFourier-package, 2

print.multinomfourier, 3
pval_exact, *2*, 3, *3*, *6*, *7*, *10*, *11*
pval_exhaustive (pval_exact), 3
pval_flexible, *2*, *3*, *5*, 6, *11*
pval_fourier, *2–7*, 8