The cost of precision-fermented whey and egg proteins

A Humbird-style techno-economic analysis of β-lactoglobulin and ovalbumin, from stoichiometry to the physical limit, with a lactoferrin contrast case

Pablo Antonio Moreno Casares

Version 2.1 (audited; AI pre-reviewed) for human review · numbers generated 2026-10-01 22:02 · repository `pf-tea`

Status. Not yet peer-reviewed. The model code, the evidence extraction and this text were produced with an AI assistant (Claude, Anthropic) under the author's direction. Version 1 was audited by eight independent AI reviewers, each owning one area and required to support every finding with a computation, quote or code line; version 2 was then read by four simulated expert reviewers (a fermentation scientist, a bioprocess engineer, a sceptic and the method's author's perspective), also AI. Every finding, decision and effect is logged in `docs/AUDIT.md` and `docs/REVIEW.md`, and this version implements them. No human expert has reviewed the analysis yet. Every number is computed by the model and read from `results/analyses.json` at compile time; every input traces to a source and an evidence grade in `data/` (Appendix A). Inputs graded E are guesses; they are listed in Section 6.

Summary

Precision fermentation (PF) uses an engineered microbe to secrete a single animal protein. We estimate what food-grade β-lactoglobulin (BLG, the main whey protein) and ovalbumin (OVA, the main egg-white protein) cost to make, and what performance would bring them to the price of the proteins they replace. The method is Humbird's (2021): element- and electron-balanced black-box stoichiometry, an oxygen-transfer-limited fed-batch fermenter, and his capital factors and fixed costs, with which his baker's-yeast workbook lands on that product's market-implied cost. We add a secreted product, a downstream train and a cation-exchange step, and check the model against three published designs.

The base case is a secreting **Komagataella phaffii** strain at 30 g/L after a 72 h production phase, the design point of Intelligen's (2025) OVA process, in 200 m³ industrial fermenters making 10 kt/yr of protein in the US Midwest (2025 dollars). With a capital charge of 14.6% of investment per year its minimum selling price is **$21.2/kg BLG** and **$21.3/kg OVA**. With the lighter accounting of NREL-style design reports it is $16.4 and $16.5; with the aseptic, cell-culture-grade fermenters that version 2 priced, $30.9 and $31.1. At the average titer that private companies' cost models assume (42 g/L, GFI 2025) BLG costs $15.8/kg. At the best public titers it costs $43/kg BLG (13.9 g/L, with an ethanol-fed production phase that the model prices as glucose) and $246/kg OVA (2 g/L, partly clipped). Whey protein traded at $9–$18 per kg protein in 2015–2025 (export unit values in 2025 dollars) and dried egg white at $10–$21.

Five results stand out:

1. **Cheap protein needs fast secretion more than high titer.** At the base cell density, $20/kg BLG needs 29–32 g/L in 48–72 h, a specific productivity of 5–7 mg protein per g cells per hour against 4.7 in the base design. With today's biology and capital, the cheapest design the model finds costs $7.1/kg if cells could secrete 40 mg/g/h (the ceiling we allow), and $12.0 at 10 mg/g/h. No published **K. phaffii** process is near either rate. At the best rate demonstrated in a public glucose fed-batch, 2.1 mg/g/h (Prielhofer et al. 2018), the cheapest design costs $36.2/kg, and even the best case of point 3 $24.5/kg, above the whey price band.
2. **The titer ceiling is a design ceiling.** With Intelligen's 500 g/L glucose feed and a batch that must start at least 30% full, a 200 m³ fermenter at 107 g/L cells reaches about 70 g/L in a 72 h production phase; with a 600 g/L feed, about 105 g/L.
3. **A best case, not a floor.** With every assumed limit at its best at once, 400 m³ vessels and 100 kt/yr, BLG costs $4.1/kg and OVA $4.2/kg ($3.3 and $3.4 with design-report accounting). That assumes secretion at 40 mg/g/h; at 10 mg/g/h the BLG case costs $8.1, at 5 mg/g/h $12.1.
4. **Capital: the vessel price and the fixed-cost rule.** Aseptic cell-culture vessels instead of industrial ones would add $9.8/kg to the base; the lighter design-report accounting would remove $4.7, mostly through fixed costs of under 2% of investment a year instead of Humbird's 9%. Published NREL-style designs come

out 1.8× cheaper than our model of the same process, for those reasons; Intelligen's food-PF design comes out within −4% of ours.

5. **A high-value contrast.** At a design titer of 10 g/L, PF lactoferrin costs $136–$178/kg, below the reported range for dairy lactoferrin ($250–$1,585/kg in 2025 dollars). Intelligen's own model of the same route costs 2.8× ours to produce, mostly through its unit prices (Table 12). For high-value proteins PF can compete on cost if that titer is reached; the commodity proteins are the hard case.

The least-tested assumption is secretion without growth. The production phase follows Intelligen's design, whose cells secrete 4.7 mg/g/h while barely growing. In the one near-zero-growth study we found (a **K. phaffii** strain secreting an antibody fragment), secretion fell as growth slowed and settled about five times below the trend at faster growth (Rebnegger et al. 2024, who link part of the fall to the promoter used). Across the literature **K. phaffii** secretes fastest while growing: a review compiles maximal rates of 0.004–0.69 mg/g/h, those on glucose peaking at growth rates of 0.15–0.20/h (Looser et al. 2015), and the best glucose fed-batch we found reached 2.1 mg/g/h near 0.1/h (Prielhofer et al. 2018). Growing while producing does not rescue the design: at that strain's ratio, 21 mg of protein per g of new cells, 30 g/L would need 1.4 kg of new cells per litre. No public study we found shows secretion at the design rate without growth.

Scenario	Titer (g/L)	BLG, price-anchored	BLG, design-report	OVA, price-anchored	OVA, design-report
Best public academic strain	13.9 / 2	$43	$32	$246	$176
Base case (72 h production)	30	$21.2	$16.4	$21.3	$16.5
Industry benchmark (titer in private cost models)	42	$15.8	$12.3	$15.9	$12.4
Physical-limit best case (100 kt/yr, secretion 40 mg/g/h)	140 / 140	$4.1	$3.3	$4.2	$3.4
Base case with aseptic cell-culture vessels (v2′s basis)	30	$30.9	—	$31.1	—
Incumbent, per kg protein (2015–2025, 2025$)			whey $9–$18		egg white $10–$21

Table 1: Minimum selling price ($/kg target protein, 2025) with a capital charge of 14.6% of total capital investment per year. The capital conventions and the vessel price basis are defined in Section 2.5. 10 kt/yr and US Midwest unless noted. Academic BLG: Yao et al. (2026), whose production phase runs on ethanol (priced as glucose); on glucose, Li et al. (2026) reach 5.14 g/L ($107/kg). Academic OVA sums full-length and clipped protein; at 1 g/L of intact OVA it costs $489/kg.




Figure 1: Every case against its market, on one axis: filled dots use price-anchored capital (the base), hollow dots design-report capital; the shaded band is the incumbent's price range in 2025 dollars. Model figures are production costs with a return on capital, not selling prices.

Contents

Summary 1
1 Question and approach 5
2 Methods 6
2.1 Plant, products and functional unit 6
2.2 Stoichiometry and energetics 6
2.3 The fed-batch fermenter and its feasibility ceiling 8
2.4 Downstream processing 9
2.5 Capital: vessel prices and two accounting conventions 9
2.6 Operating costs, labour and money 10
2.7 Evidence grades, scenarios and uncertainty 11
3 Calibration 11
4 Results 13
4.1 Base case 13
4.2 Q1: cost against titer, with every public and claimed data point 14
4.3 Q2: what drives the cost 15
4.4 Q3: what must be true for $10 and $20 per kg 17
4.5 Q4: the physical-limit best case 19
4.6 Q5: claimed titers against the fermenter's limits 20
4.7 Q6: scale and siting 21
4.8 Q7: milk and egg equivalents 21
4.9 Q8: a high-value contrast, lactoferrin 22
5 Discussion 24

5.1 Answers, in brief ... 24
5.2 Why published PF costs are lower ... 24
5.3 For funders ... 24
5.4 For EU policy ... 25
5.5 What would change these conclusions ... 25
6 Limitations ... 25
7 Scoring the pre-registered priors ... 26
8 Reproducibility ... 26
References ... 26
A Parameters (generated from `data/pf/base.yaml`) ... 29
B Evidence base ... 41

1 Question and approach

PF companies now sell BLG (Perfect Day, Remilk, Vivici) and OVA (EVERY), and others are entering (Onego Bio). Whether these proteins can displace animal proteins at scale or stay a premium niche depends on what they cost to make. Published estimates disagree by a factor of several, and company claims are rarely checked against the physics of aerobic fermentation. This study asks:

> What does it cost to make food-grade BLG and OVA by precision fermentation with today's public strain performance, and at the physical limits of aerobic fermentation? What combination of titer, rate, yield, downstream design and scale would bring them to price parity with whey and egg-white proteins?

#	Sub-question	Answered in
Q1	What do BLG and OVA cost at public (academic) and claimed (industry) performance?	Section 4.2, Figure 3
Q2	What drives the cost, and how good is the evidence for each driver?	Section 4.3, Figure 5, Figure 6, Figure 7
Q3	Which combinations of performance reach $10 and $20 per kg protein?	Section 4.4, Figure 8
Q4	What is the cost at the physical limits?	Section 4.5, Figure 9
Q5	Are published company claims consistent with the fermenter's limits?	Section 4.6, Table 10
Q6	How much do siting (US vs EU) and scale (1–100 kt/yr) matter?	Section 4.7, Figure 10
Q7	*Added after v1:* what do these costs mean per litre of milk and per egg?	Section 4.8, Figure 11
Q8	*Added after v1:* how does a high-value protein (lactoferrin) compare?	Section 4.9, Figure 12

Table 2: Q1–Q6 were fixed before any model output existed (`docs/SCOPE.md`); Q7–Q8 were added at the author's request after the first draft.

Why whey and egg white are the benchmarks. BLG is the main protein of whey, and the PF BLG on the market is sold as a whey-protein ingredient for bars, drinks and dairy analogues (FDA GRN 1056 specifies a concentrate of at least 70% protein). A buyer's alternative is whey protein concentrate or isolate, so their price is the one PF BLG must meet; the same logic sets dried egg white against OVA. It is a hard comparison. Whey is a by-product of cheese, sold for whatever clears the market once cheese has paid for the milk, and it is among the cheapest animal proteins; if PF displaced much of it, its price would fall further. Two consequences follow. Price parity is not the only commercial route: pure, animal-free or functional proteins can sell at a premium, as the lactoferrin case shows (Section 4.9). And displacing whey changes milk output far less than one-for-one, because whey carries only a small part of the milk's value (Section 4.8).

Why Humbird's method. Humbird's (2021) cultured-meat analysis builds costs from conservation laws: element and electron balances, oxygen transfer, heat removal. His full cost model reproduces the market-implied cost of one fermented commodity, baker's yeast at about $1.90/kg; capital-related charges are less than half of that cost, so the check constrains his capital factors only loosely (Section 3). We port his baker's-yeast workbook exactly and extend it with a secreted protein, a downstream train, and the specifics of **K. phaffii** and **Trichoderma reesei**.

Rules we followed (written protocol, `PLAN.md`):

- Four scenarios are never merged: academic, industry, required and physical limit.
- Every input carries an evidence grade, A to E.
- The model structure is frozen before the calibration anchors are run; misses are recorded, not tuned away.
- Priors about the results were written down before the model existed; they are scored in Section 7.
- Model-to-model agreement is a consistency check, not validation.
- After the first draft, eight independent reviewers audited it; their findings and every resulting change are logged in `docs/AUDIT.md` (R1–R8, P-1 to P-13). Version 2 was then read by four simulated expert reviewers (AI); their reviews are kept verbatim in `docs/ai-review/` and answered comment by comment in `docs/REVIEW.md`.

2 Methods

2.1 Plant, products and functional unit

The functional unit is 1 kg of target protein in a spray-dried powder that meets the product specification. The base specification is a **concentrate**, following Remilk's GRAS notice for BLG (FDA GRN 1056): at least 70 wt% protein, at least 80% of it BLG, at most 7% moisture. OVA uses the same specification by analogy. The **isolate** variant (99% target protein, dry basis) adds an ammonium-sulfate precipitation, following Intelligen's (2025) OVA design. The lactoferrin contrast case follows the recombinant bovine lactoferrin notices (FDA GRN 1219 and 1284): at least 90% protein, at least 95% of it lactoferrin.

The plant is greenfield, single-product and nth-of-a-kind, built to food (not pharmaceutical) good manufacturing practice. The base is 10 kt/yr of protein in the US Midwest, swept from 1 to 100 kt/yr, with a north-west Europe (Netherlands) variant. Money is 2025 US dollars. Excluded costs: R&D, strain licences, regulatory approval, sales and distribution, land, first-of-a-kind premiums, carbon pricing, start-up and commissioning, working capital (except in the discounted-cash-flow price), failed batches, per-lot release testing and packaging. A reviewer (AI) put the last four at roughly $1–2/kg together. Figure 2 shows the base-case plant.




Figure 2: Base-case process: **K. phaffii** secreting BLG at 30 g/L; flows per kg of BLG.

2.2 Stoichiometry and energetics

Each metabolic process is one element-balanced "black-box" reaction on C, H, O and N: glucose + NH_3 + O_2 → biomass (or protein) + CO_2 + H_2O. Electrons are tracked through the degree of reduction γ per C-mol (glucose 4, **K. phaffii** biomass 4.06, BLG 4.24, OVA 4.18). This closes the O_2 and CO_2 balances once the carbon split is fixed.

Growth. Biomass yield comes from the Gibbs-energy dissipation correlation of Heijnen and van Dijken (1992), as in Humbird's workbook. The energy dissipated per C-mol of biomass formed is

$$D = 200 + 18(6 - c)^{1.8} + \exp\left\{\left[(3.8 - \gamma_s)^2\right]^{0.16}(3.6 + 0.4c)\right\} \quad \text{kJ/C-mol},$$

where c and γ_s are the carbon number and degree of reduction per C-mol of the substrate (valid for $1 \le c \le 6$). The catabolic reaction (glucose to CO_2) is run just enough to supply D. For glucose and our **K. phaffii** biomass formula (Carnicer et al. 2009) this gives a maximum biomass yield of 0.57 g/g; the measured value is 0.50–0.56 g/g (`data/evidence/hosts.yaml`, grade A). Biomass is counted as ash-free organic matter; ignoring the 7% ash overstates the yield per g of dry cells, but the same convention understates maintenance per g, and the two nearly cancel in glucose per g of cells (audit R1-F3).

Maintenance. In the growth phase maintenance follows Tijhuis et al. (1993), $m_G = 4.5\exp[-69000/R(1/T - 1/298)]$ kJ per C-mol biomass per hour: 0.0174 g glucose per g cells per hour at 30 °C, close to the 0.0147–0.0161 measured for protein-producing **K. phaffii** in glucose chemostats (Maurer et al. 2006; Rebnegger et al. 2016). In the production phase, where cells barely grow, maintenance falls: Rebnegger

et al. (2016) found it threefold lower at near-zero growth. We use the value measured for a protein-secreting strain at low growth, 0.0044 g/g/h (Rebnegger et al. 2024; range 0.0023–0.0161).

Product. The protein's elemental formula comes from its mature UniProt sequence:
- BLG (P02754, 162 residues): $C_{817}H_{1312}N_{206}O_{248}S_9$, 18.3 kDa;
- OVA (P01012, 385 residues): $C_{1901}H_{2997}N_{499}O_{575}S_{22}$, 42.75 kDa;
- bovine lactoferrin (P24627, 689 residues): $C_{3358}H_{5251}N_{945}O_{1002}S_{38}$, 76.1 kDa.

Glycans and phosphates are left out of the formulas: 3–6% of the mass of glycosylated OVA and 5–10% of lactoferrin, which changes the mass balance by less than 1%.

The product reaction makes protein from glucose and ammonia and respires enough extra glucose to pay for synthesis, so the product-reaction yield Y_{ps} fixes its O_2 demand. **No measured BLG or OVA yield on sugar exists**; the design values are lumped yields that already include maintenance and some growth (Intelligen 0.28 g/g; Behm et al. 2022 0.223 g/g). We therefore derive Y_{ps} so that the model reproduces Intelligen's expression phase (their Eq. 2: 100 g glucose ⟶ 28 g OVA + 8 g cells + 4 g host proteins):

$$\frac{1}{Y_{ps}} = \frac{100}{28} - \frac{8/28}{Y_x} - \frac{m_{s,\mathrm{prod}}}{q_p} = 3.571 - 0.501 - 0.919$$

with our growth yield $Y_x = 0.5705$ g/g and Intelligen's specific productivity $q_p = 0.00475$ g/g/h, giving $Y_{ps} = 0.465$ g/g. The growth rate in production (0.0014/h) and the cell density at its start (107 g/L) follow from the same design and its volume balance. With chemostat-level maintenance Intelligen's Eq. 2 would be impossible: maintenance alone would use all of its glucose. Three cautions:
- **The inversion is sensitive to maintenance.** Maintenance and product enter as a ratio against a total of only 3.571: at the low end of the measured maintenance (0.0023 g/g/h) it gives $Y_{ps} = 0.39$, and above about 0.0146 it has no solution. With the product made at the energetic ceiling (below), Intelligen's expression phase leaves room for at most 0.0062 g/g/h of production-phase maintenance: above the one measurement at near-zero growth (0.0044, for a strain secreting far less) and below every maintenance coefficient compiled for growing cultures on methanol or glycerol, 0.009–0.032 (Looser et al. 2015, which gives none for glucose).
- **It charges Intelligen's host-cell protein to the target.** Read with the 4 g of host protein as product, the same equation gives 0.53 g/g.
- **It assumes secretion without growth.** The base production phase takes up glucose at only 0.016 g per g cells per hour, a few times maintenance, while secreting 4.7 mg/g/h. In the **K. phaffii** chemostat studies on file, secretion rises with growth rate, linearly or up to an optimum near 0.08–0.1/h (Rebnegger et al. 2014; Garrigós-Martínez et al. 2019; Nieto-Taype et al. 2020). The best glucose fed-batch we found, run with a glucose-regulated promoter, peaked at 2.1 mg/g/h near 0.1/h (Prielhofer et al. 2018: 6.4 g/L in 60 h). A review of **K. phaffii** cultivation compiles maximal secretion rates of 0.004–0.69 mg/g/h; its four entries on glucose (0.05–0.49) all peak at growth rates of 0.15–0.20/h, and it calls such secretion growth-associated (Looser et al. 2015). In the one near-zero-growth study we found secretion settled about fivefold below the trend at faster growth (Rebnegger et al. 2024, 0.02 mg/g/h at doubling times of two months, partly attributed to the promoter). Intelligen's design point is not a measurement.

The model's production phase then runs at an effective yield of 0.277 g/g at the base design point, close to Intelligen's 0.28. The effective yield is an output: it rises with productivity, to 0.32 at 42 g/L and 0.52 in the physical-limit case, and neither is anchored to data. The v1 reading (0.28 as the reaction yield, plus maintenance and growth on top) is the low end of the yield range.

Energetic ceiling. Counting ATP gives an upper bound on Y_{ps}. Each residue costs the peptide bond (4 ATP; Martin et al. 2025), the net cost of making the amino acid from glucose and ammonium (0.29 ATP on average: Stouthamer 1973 as recomputed by Martin et al. 2025, whose note is that the common amino acids yield ATP) and, for the protein's sulfur, sulfate assimilation (3 ATP plus 8 electrons per S). That is 4.79 ATP per residue for BLG, paid by respiring glucose at 4 + 12 × P/O ATP per glucose, with P/O = 1.48 for **K. phaffii** (a genome-scale-model value, Chung et al. 2010), i.e. 21.8 ATP per glucose. The ceiling is 0.563 g/g for BLG and 0.570 for OVA (0.582 and 0.590 counting peptide bonds alone, as in version 2). Membrane transport, secretion, folding and turnover are not counted, so it stays an upper bound. The base yield is 83% of it. A different estimate, which

treats protein synthesis as if it dissipated energy like growth (Heijnen's correlation applied to the protein), gives 0.50 g/g for BLG; it is not a bound, but it suggests the real ceiling sits below the ATP count.

Heat. Heat release is computed from formation enthalpies of the dissolved species: about 480–510 kJ per mol O_2 for growth and maintenance and 520–570 for the product reaction. The fermenter uses its own computed heat when it slows growth to the heat-removal limit and when a design's peak heat is checked; the 460 kJ per mol O_2 of Roels (1983, via Almeida Benalcázar et al. 2025; calorimetry gives 455 ± 15, Maskow et al. 2010) only screens the O_2 limit.

> **Check (reader):** The dissipation formula and its units; the conversion of m_G to 0.0174 g/g/h; the Y_{ps} derivation (`tests/test_model.py::test_y_ps_reproduces_intelligen_eq2`); the ATP ceiling (`src/pftea/plant.py`, `atp_per_residue`; `tests/test_model.py::test_energetic_ceiling_components`). The cost structure rests on these.

2.3 The fed-batch fermenter and its feasibility ceiling

Each production batch has two phases:

1. **Growth** from an inoculum of 0.88 g/L (5% of seed broth at 17.6 g/L, the density in Intelligen's lactoferrin flowsheet) at the maximum specific rate, slowed so that the oxygen uptake rate (OUR) never exceeds what the vessel can transfer and the heat never exceeds what it can remove, up to the production cell density X_{prod}.
2. **Production** for t_{prod} hours at a small growth rate and a constant specific productivity q_p. No product is made during growth, a simplification for constitutive promoters.

The initial liquid volume is solved so that the broth ends at 80% of the vessel, and q_p so that the harvest titer equals the target. A volume balance tracks the feeds (glucose at 500 g/L, about 42 wt%, Intelligen's feed; ammonia at 30 wt%, i.e. 268 g/L), O_2 uptake, CO_2 release and evaporation.

Oxygen transfer. The transfer coefficient follows van 't Riet's correlation for non-coalescing broths, $k_L a = 0.002(P/V)^{0.7} U_s^{0.2} 1.025^{T-20}$ (s^{-1}, SI units), as in Humbird's microbial fermenters, with a log-mean driving force between the bottom (hydrostatic pressure, fresh air) and the top (back-pressure, depleted gas). For each demanded OUR the model picks the air rate and agitation power that minimise compressor plus refrigeration-weighted agitator power, within 5,000 W/m³ and a superficial gas velocity of 0.1 m/s (grade D). In the base vessel the correlation alone would allow about 376 mol O_2/m³/h, but that extrapolates it well beyond the k_La of about 800 h^{-1} that Humbird gives as the practical upper end; the 25 kW/m³ heat-removal ceiling allows about 187. The design cap is Humbird's value for an air-sparged stirred tank, "an OTR approaching 200 mol O_2/m³-h is common", on his own basis, per m³ of cell-free liquid (his Eq. 4.2): 129 mol/m³/h per m³ of broth at the base cell density, 147 at 80 g/L and 100 at 150 g/L. Growth runs at it. (Version 2 used 150 per m³ of broth; 150 is Humbird's limit for bubble columns.) An industrial designer may plan on 100–120 per m³ of broth with air at these cell densities (review BE-7), and our aeration power is lower than NREL's and Intelligen's (Section 3).

Compressor, and a check on aeration power. The compressor raises the air adiabatically to the bottom pressure at 80% efficiency. Humbird's workbook formula puts $k = 1.4$ where $(k-1)/k = 0.286$ belongs and gives about 1.6× that power; we keep it only in the port of his workbook, and in our model it would add $0.02/kg (review H1). With his formula, our aeration reproduces his own 2021 aeration sheet: 0.50 kWh of compressor and agitator power per kg O_2 at 150 mol/m³/h in a 200 m³ vessel, against his 0.50; with the corrected formula, 0.39. Humbird, Davis and McMillan's (2017) process simulations imply about twice his 2021 figure (their Fig. 3 total less the chilling of reaction heat, our reading of a chart); their model could not hold dissolved O_2 down, as production plants do. Aeration electricity costs $0.15 per kg of protein in the base, so doubling it would add about as much; holding dissolved O_2 at 50% of air saturation adds $0.26/kg, and leaving out the temperature term of the k_La correlation, as the 2017 paper does, $0.07/kg.

Other limits. A design is **infeasible**, and never costed as if it worked, when it needs more O_2 or cooling than above, starts below 30% of the vessel volume (the lowest start in a published design, Intelligen's 30.5%), secretes more than 40 mg of protein per g of cells per hour (a judgment, grade D; the base runs at 4.7, and no published **K. phaffii** process is near 40; compiled maxima for secreted proteins are below 1, Looser et al. 2015), exceeds the product's solubility, or has a yield above the energetic ceiling.

Why a ceiling exists. Each kg of BLG made costs about 133 mol O_2 and 9.2 kg of glucose across the batch, including growth and maintenance. The broth must end at 80% of the vessel, so every extra kg of product means more feed and a batch that starts emptier. In the base design (107 g/L cells, a 500 g/L feed) that, not O_2, is what caps the titer: at 70 g/L (72 h) the batch would have to start below 30% of the vessel. Longer production phases lower the ceiling (76 g/L at 36 h, 60 g/L at 120 h), because maintenance keeps consuming feed. The ceiling depends on the design choices behind it (Section 4.4).

What the fermenter model leaves out. Broth viscosity and foam do not lower k_La: the correlation is for water-like broths at every cell density and titer, while Humbird's own method stops a fed-batch at a viscosity (2 cP) his broth model reaches near 80 g/L cells. At the high cell densities and titers of the best cases the O_2 limit is therefore generous (review FS-6, HM-4). Specific productivity is constant through the production phase, whereas real secreting cultures often lose productivity after two or three days; and no product is made during growth, which is conservative for constitutive promoters.

2.4 Downstream processing

The train (Figure 2) follows Intelligen's OVA design, with evidence-graded performance (`data/evidence/dsp.yaml`):

- **Cell removal.** Dilute to 45 g/L cells, then two disk-stack centrifuge stages with re-dilution of the paste. Each paste holds 50% extracellular liquid by volume (Petrides: 40–60%), so a fraction $\varphi = eV_p/(V_1 - V_{wc})$ of the product is trapped per stage and the recovery is $1 - \varphi^2$ (96.9% in the base). The wash water is counted.
- **Microfiltration polish** at 20 L/m²/h (grade D, range 10–50) and 97.5% recovery (Behm et al. 2022).
- **Ultrafiltration and diafiltration.** Concentrate at 30 L/m²/h to 50 g/L, diafilter 3 volumes there at 5 L/m²/h (the only diafiltration flux on file, EVERY patent US 2025/0134130, measured on a redissolved precipitate at 10 °C, so probably conservative for clarified broth), then concentrate to the spray-dryer feed at the same low flux. Rejection 0.995. Skids cost \$263 per m² of membrane installed (a dairy plant, Mercier-Bouchard et al. 2017), converted to a purchase cost before the installation factors are applied.
- **Dryer feed.** Dairy plants feed high-protein whey concentrates to the dryer at 27–37% solids by membranes, not evaporators (Tetra Pak); egg white is ultrafiltered to about 23% (Tsai et al. 1977). We use 30% for BLG and 23% for OVA.
- **Spray dryer** at 4.87 MJ per kg water evaporated (industry survey, grade A); purchase cost from a supplier-based SuperPro model (Yver et al. 2012).
- **Isolate only:** ammonium-sulfate precipitation (350 g/L; 89% recovery), washed basket centrifuges; the dryer is fed the 20%-solids cake.
- **Lactoferrin only:** cation-exchange capture between microfiltration and ultrafiltration, as in the dairy process and every commercial PF lactoferrin (Section 4.9).
- **Cleaning in place** (7.3 kg solution per kg product, Intelligen; its chemicals are inside Humbird's maintenance charge) and **spent-cell disposal** (\$458/t dry cells in the US, a landfill tipping fee; \$211/t in the EU, an anaerobic-digestion gate fee; both per wet tonne converted at the 15% dry solids of the model's centrifuge paste, with no dewatering or inactivation costed). Sale as feed or fertiliser would make it free or a credit.

The target protein share of the concentrate powder is 85% on a dry basis (grade D; the GRN 1056 minimum is 56%). The base overall recovery is 90%.

OVA solubility. Intelligen stops its ultrafiltration at 50 g/L, "its practical solubility limit", without a source; vendor data sheets give 40–50 mg/mL as quality-control concentrations. OVA dissolves at 200 g/L in water (Wang et al. 2025) and is processed at about 110 g/L in ultrafiltered egg white (Tsai et al. 1977). We therefore do not treat 50 g/L as a limit, and show it as a sensitivity. No data exist at pH 5, the fermentation pH, which is close to OVA's isoelectric point.

2.5 Capital: vessel prices and two accounting conventions

Fermenter prices. Fermenters are the largest equipment item. The base prices them from a vendor quote: NREL 2011's 300 m³ stainless fermenter with internal cooling coils and an agitator, \$0.99M in 2007 dollars, scaled with exponent 0.7, i.e. \$1M per 200 m³ in 2025 dollars. It is an industrial enzyme-plant vessel, the closest public analogue of a food-grade PF fermenter. It is also at the vessel price level of Humbird's own baker's-yeast workbook, whose 200 m³ bubble column costs about \$0.8M (2018), and consistent with Intelligen's food-PF

design (Section 3). Version 2 used Humbird's quote curve for the aseptic stirred tank he designed for cultured meat: 316L, full vacuum, built to the ASME bioprocessing standard. That vessel costs $3M per 200 m³ (2025 dollars), 2.9× the vendor quote. Humbird himself separates aseptic industrial fermentation, which tolerates a few foreign organisms, from aseptic animal-cell culture, which does not. We keep his aseptic vessels as a named high case ("aseptic vessels", +$9.8/kg on the base BLG). Seed fermenters (4.7 and 0.47 m³, 11 of each so that a batch can start every few hours) are priced on the same curve, scaled far below its quote. They are 2.6% of fermenter cost; Humbird's aseptic quotes, which do cover that size, are 2.2× the curve there and would add $0.18/kg.

Price-anchored capital (Humbird method; base).

- Other equipment comes from Humbird's cost curves (Aspen ACCE quotes, 2018 dollars). Costs are escalated to 2025 with a factor of 1.33 (CEPCI and the BLS machinery price index; grade B, range 1.23–1.52). The BLS index for heavy-gauge metal tanks rose 1.92× over the same years; escalating the fermenters with it would add $2.3/kg.
- The equipment list is abbreviated (it omits CIP and feed tanks per fermenter, sterile filter housings, sugar storage, biomass inactivation, powder packing and effluent pretreatment), so we add 20% for missing items, the allowance of Humbird's yeast workbook for its own abbreviated list (his fuller cell-culture sheets use 10%).
- Installation, buildings, engineering, fees and contingency follow the "commodity aerobic" facility class that Humbird tabulates (a food-grade plant with the process in a sanitary building). None of his own sheets uses that class: his cultured-meat sheets use the cGMP biopharmaceutical class and his yeast workbook a flat Lang factor of 4.5. The effective Lang factor (investment over equipment cost) here is 6.5, high because buildings are costed by floor area while the fermenters are now cheaper.
- Maintenance is 4% of total capital investment (TCI) per year, including CIP chemicals, and insurance 5%, Humbird's whole fixed-cost rule. There is no separate plant-overhead, quality-control or local-tax line, so the 9% stands for all of them; insurance alone is usually well under 1%.
- What the anchor supports: with these factors and fixed costs Humbird's yeast workbook lands in the cost band implied by the market price (Section 3), and a lighter, NREL-like set does not. Capital-related charges are less than half of the yeast cost, so the check is loose: it favours Humbird's set but cannot pin it.

Design-report capital (NREL-style; sensitivity). The convention of NREL's 2011 design report (whose lead author is also Humbird): the same fermenter quote; NREL's installation factors (1.7 for fermenters, 1.8 otherwise) plus indirect costs of 60% of direct cost; no separate warehouse; maintenance of 3% of installed process-area cost and insurance of 0.7% of fixed capital.

On the base BLG plant the two conventions differ by $4.7/kg: $3.3 of it comes from the fixed-cost rule alone, and $1.4 from the installation and indirect factors.

A Netherlands plant costs 1.09× the same plant in the US Midwest (IEAGHG 2018, same plants costed in both places), applied to capital in both conventions.

2.6 Operating costs, labour and money

Operating costs. Raw materials, utilities and consumables come from the mass and energy balance and the prices in Appendix A: EIA, Eurostat and BLS statistics, trade unit values and published designs (grades A–B).

Labour. Humbird's operator-attention model: operators per fermenter by the fraction of time a batch needs attention, fixed staffing for continuous areas, 4.5 shifts per position, one supervisor per ten staff at 1.4× salary, and a 100% labour burden (Humbird). The base plant employs 124 people, operators and supervisors only; salaried staff are inside the fixed-cost rule (Section 2.5).

Headline metric. $\mathrm{MSP}_{\mathrm{CCF}}$: annual operating cost plus a capital charge, divided by annual output. The charge is the capital recovery factor $i/(1-(1+i)^{-n}) = 0.146$ per year at $i = 7.5\%$ and $n = 10$ years (Humbird). It is a cost with a return on capital, not a market price. We also compute the cost of production (COP: operating cost plus depreciation, no return) and an NREL-style discounted-cash-flow price $\mathrm{MSP}_{\mathrm{DCF}}$ (10% IRR, 30 years, 21% tax, 7-year MACRS, construction over three years, working capital). For base BLG these are $18.9 and $20.7.

2.7 Evidence grades, scenarios and uncertainty

Every parameter and evidence item is graded (Table 3). The four scenarios combine grades differently and are never averaged:

- **academic:** the best public strain performance, read in full text. BLG: 13.88 g/L in **K. phaffii** (Yao et al. 2026), whose production phase runs on ethanol with glutamate feeding; the model has one carbon source and prices it as glucose (per electron the two cost about the same), so this case is flagged: ethanol is more reduced than glucose, so the O_2 demand and heat per kg of product are higher than modelled, and the feasibility check is not conservative for this case. The modelled batch ends at 88 g/L cells against about 120 in Yao's run, because the model's volume balance dilutes; it is a cost estimate for 13.88 g/L in 108 h, not a simulation of that run. On glucose, Li et al. (2026) reach 5.14 g/L in 132 h. Their feed table (supplementary Table S4) gives the only public check on a BLG yield on glucose: 1–2% overall, depending on whether its feed rates refer to the starting or the current broth volume (our arithmetic; the volume profile is not reported). Our model of that run gives 2.5%, so it is, if anything, optimistic on sugar there. OVA: 2 g/L in **T. reesei** (Aro et al. 2023), with no growth in production. That is the sum of full-length and clipped OVA, and the purified product was mostly the clipped 36 kDa form, so intact OVA was probably well under 2 g/L; a 1 g/L variant is shown. The **T. reesei** production phase uses the yeast maintenance correlation, which is a lower bound on its sugar use. Yield and maintenance keep the base values: no measured product yield exists.
- **industry:** 42 g/L, the average that Hawkwood computed over private companies' models of unspecified secreted proteins for GFI (2025) (grade C), with everything else at the base case.
- **required:** the inverse problem: the titer needed for $10, $20 and $50/kg at each production time.
- **physical limit:** a best case at assumed limits, not a floor set by physics. Every lever moved from the base: product yield 0.535 g/g (95% of BLG's energetic ceiling); the lowest measured maintenance in both phases; no growth in production; a 600 g/L glucose feed (Shemesh et al. 2024; 730 g/L has been fed in a 2.6 L laboratory vessel, Heyland et al. 2010, which would raise the ceiling somewhat further); 400 m^3 vessels; O_2 transfer capped at 250 mol/m^3/h per m^3 of cell-free liquid (head pressure, not costed; grade D) and heat removal at 40 kW/m^3 (external exchangers, not costed; grade D); an 8 h turnaround; a 90% concentrate; contract sugar; and 100 kt/yr. A search over cell density and production time puts the titer at its feasibility ceiling, where cost is lowest, and reports which limits bind. Secretion stays capped at 40 mg/g/h, and the result is shown against that cap from 5 to 120 mg/g/h.

Grade	Meaning
A	Measured, primary source or official statistic (peer-reviewed data, EIA, Eurostat, BLS, USDA)
B	Design value or patent example from a named source (NREL, Intelligen, Humbird, patents)
C	Company claim, press release, industry survey, secondary transcription
D	Analogy, judgment or calculation from graded inputs
E	Guess; listed in Section 6

Table 3: Evidence grades (`data/evidence/README.md`).

Uncertainty. Two tools:

- **Tornado.** Each parameter with a graded range moves to its low and high value, one at a time. An end that breaks a physical limit is moved to the last feasible value, so no bar quotes the cost of an impossible plant. The accounting convention and the fermenter price basis are shown as their own (structural) bars.
- **Monte Carlo.** Triangular draws over each scenario's uncertain parameters, with host-specific ranges. Draws that break a physical limit, or for which no plant can be built, count as infeasible; we report the feasible fraction, the limit each infeasible draw breaks, and percentiles over feasible plants and over all draws with a cost. Draws are independent: no data exist to set correlations, and the feasibility filter is the only coupling. Structural choices (conventions, vessel price basis, minimum fill, secretion ceiling) are not sampled; they are shown one at a time.

3 Calibration

The structure was frozen first. Each anchor then changes only its own process inputs, prices, cost year and financing, and misses are recorded in `docs/VALIDATION.md`. `tests/test_calibration.py` pins each recorded value to within five points, so a change that moves an anchor cannot pass silently.

Anchor (type)	Quantity	Ours	Theirs	Diff.
1. Humbird baker's yeast (software + reality)	cost, $/kg dry cells (2018)	1.890	1.890	+0.00%
2. NREL 2011 on-site cellulase (consistency)	glucose, kg/kg protein	3.87	4.17	−7.1%
	O_2, mol/kg protein	83.1	84	−1.0%
	electricity, kWh/kg	4.9	9	−45%
	cost, $/kg protein (2007), price-anchored	7.81	4.24	+84%
	cost, design-report (NREL's own) capital	6.08	4.24	+43%
3. Intelligen 2025 OVA (consistency)	cost, $/kg powder, price-anchored	42.00	43.52	−4%
	total capital investment	$114M	$92M	+24%
	glucose, kg/kg powder	10.6	13.0	−19%
	electricity, kWh/kg powder	9.6	27.1	−65%
4. Voutilainen 2021 recombinant case (consistency)	sugar, kg/kg protein (by construction)	6.28	6.15	+2.2%
	cost, €/t protein, their DCF, price-anchored	16,345	9,007	+81%
	their fixed costs, price-anchored capital	12,329	9,007	+37%
	their fixed costs, design-report capital	11,423	9,007	+27%

Table 4: Calibration anchors. Anchor 1 is Humbird's own workbook, ported line by line; anchors 2–4 are other models run on their own inputs. Diagnoses: `docs/VALIDATION.md`.

What the anchors say.

1. **The port is exact.** It reproduces Humbird's baker's-yeast cost to 0.004%, and that cost sits on his market-implied $1.90/kg. The market band itself depends on how his 33% margin is read: $1.47–$1.88/kg if it is a share of the price, $1.65–$2.11 if it is a mark-up on cost. With a lighter, NREL-like capital set (investment 3.2× listed equipment, fixed costs 1.9% a year) the workbook gives $1.37/kg, below both bands; with Humbird's factors but NREL-like fixed costs, $1.62; with our PF capital charge (7.5% over 10 years instead of his 10% over 20), $2.00. The anchor favours Humbird's factors and fixed costs, loosely. It says nothing about fermenter prices beyond its own bubble column.
2. **Sugar and O_2 agree with the design that states them independently** (NREL, −7% and −1%). Voutilainen's sugar is matched by construction, and Intelligen's expression phase is, too, because our product yield is derived from it; its growth phase uses 0.40 g/g, below our 0.53 effective and the measured 0.50–0.56. **Electricity does not agree**: ours is 45–65% lower, because our aeration is power-optimised while NREL sizes agitators for peak load and Intelligen aerates at 1.2–1.6 vvm and runs a fertiliser section; a real plant may need more aeration power than ours. These are consistency checks, not calibration fits.
3. **Capital now agrees; fixed costs do not.** With vendor-quote fermenters our investment for Voutilainen's plant is within +2% of theirs, and for Intelligen's +24% above (their Lang factor is 4, ours higher). Their total costs are still 1.8× lower than ours (NREL, Voutilainen), mainly because their fixed costs are 1.5–2% of capital per year instead of 9%, and their capital charges lighter. With their conventions our model still lands above theirs, by +27% (Voutilainen) and +43% (NREL); the rest is mostly labour, buildings and cleaning.
4. **Intelligen's total is within −4%** (their cost recast into our capital charge and fixed costs), with compensating differences: our capital is higher and our plant needs 6 whole fermenters where theirs has five; our glucose and electricity are lower.

We keep Humbird's factors and fixed costs as the base because his workbook with them lands in the yeast market band and a lighter set does not; that is a loose test, and the design-report convention is shown everywhere as a named structural sensitivity, not as an error band. NREL itself notes that enzyme producers quoted costs about 47% above its estimate.

4 Results

4.1 Base case

Cost item, $/kg protein	BLG	OVA	Physical intensity, per kg BLG	
capital charge (14.6%/yr)	$7.14	$7.20	glucose	10.3 kg
maintenance	$1.96	$1.98	oxygen	4.8 kg
insurance (Humbird's 5%, all overheads)	$2.45	$2.47	electricity	9.4 kWh
glucose	$4.11	$4.10	natural gas (drying, steam)	14 MJ
labour	$1.55	$1.55	spent cells (dry)	3.5 kg
spent-cell disposal	$1.60	$1.60	overall yield on glucose	0.097 g/g
electricity	$0.81	$0.81	fermenter volume	720 m³ per kt/yr
media, ammonia and DSP chemicals	$0.64	$0.64	staff	12.4 per kt/yr
gas, steam, water, membranes	$0.91	$0.93	total capital investment	$490M
MSP_{CCF}	**$21.17**	**$21.28**	production fermenters	36 × 200 m³
per kg powder	$17.10	$17.18	harvest	94 g/L cells

Table 5: Base case: **K. phaffii**, 30 g/L, 72 h production, 10 kt/yr, US Midwest, vendor-quote fermenters, price-anchored capital.

Capital-related charges (capital charge, maintenance and insurance) are 55% of the BLG price; glucose is 19% and spent-cell disposal 8%.

- **Why so much capital.** At 30 g/L the plant needs 36 production fermenters of 200 m³. Each batch occupies a vessel for about 121 h including turnaround and yields 4,299 kg of protein.
- **Why glucose is modest.** The overall yield of protein on glucose is only 0.097 g/g, because the cells, their maintenance and the product reaction all consume sugar; at $0.40/kg glucose it is still only 19% of the price.

The product made per m³ of fermenter per hour is the lever the rest of the paper explores.

4.2 Q1: cost against titer, with every public and claimed data point




Figure 3: Minimum selling price against harvest titer for three production times. Vertical lines mark the best public titers, the average titers assumed in published and private cost models (GFI 2025), and design or claimed titers. Lines end at the feasibility ceiling of the fermenter.

Cost falls roughly as 1/titer until the ceiling (Figure 3):

- **Best public academic strains:** BLG at 13.88 g/L costs $43/kg (ethanol-fed production priced as glucose), and $107/kg at 5.14 g/L on glucose. OVA at 2 g/L in **T. reesei** costs $246/kg. BLG is then 2.4–5.0× the whey band and OVA 12–25× the egg-white band.
- **The titer private cost models assume (42 g/L):** $15.8 (BLG) and $15.9 (OVA), or $12.3 (BLG) and $12.4 (OVA) with design-report capital.

The incumbents. Official trade unit values form a consistent series, deflated to 2025 dollars with the US consumer price index:

- **Whey protein >80%** (WPC80 and WPI mixed; US and EU exports): $9–$18 per kg protein in 2015–2025, taking the traded product as 80% protein, the minimum for its trade code. If the mix averaged 88% (more isolate), the band would be $8–$16.
- **Dried egg albumin** (EU exports; 81.1% protein, USDA): $10–$21. Egg white is not always dearer than whey: per kg of protein it was cheaper than the average whey export price in 5 of 11 years (2017, 2019, 2021, 2024, 2025).
- **A 2026 whey spike.** WPC80 and WPI rose to about $34 per kg protein at the 2026 peak (USDA Dairy Market News narrative quotes, not a formal series). At that price the base case ($21.2) would undercut whey, as would the industry benchmark ($15.8). Against 2015–2025 prices, the industry benchmark is inside the band with price-anchored capital and inside it with design-report capital.

Other protein ingredients (Figure 4). Per kg of protein, plant protein concentrates and isolates trade far below whey and egg white ($3.8–$10.6 for pea and soy, GFI 2025), and casein, gelatin and collagen sit around $10–$12. PF BLG and OVA do not compete with plant proteins on price per gram of protein; they compete with the dairy and egg ingredients whose functions (gelling, foaming, whipping) they reproduce.




Figure 4: Protein ingredients per kg of protein against this model's costs. Bars: market prices (export unit values, 2015–2025 in 2025 dollars; GFI 2025 ranges, nominal). Lines: modelled production costs.

Parity in production cost is necessary but not sufficient for commercial parity: PF proteins also compete on functionality and price per unit of function.

4.3 Q2: what drives the cost




Figure 5: From NREL's on-site cellulase to our base case, one change at a time. The start is NREL's 2011 design re-priced to 2025 with design-report capital.

Why a food protein costs more than an industrial enzyme. Figure 5 walks from an NREL-style crude enzyme ($7.6/kg) to the base BLG concentrate ($21.2/kg). The steps:

- capital: price-anchored convention (Humbird factors and fixed costs): +$2.7
- food-grade indoor sanitary plant: +$1.4
- host: K. phaffii stoichiometry and broth (no maintenance): −$0.1
- high-density yeast process: 107 g/L cells, yeast media (no maintenance): +$0.6
- maintenance energy (Tijhuis in growth, measured m_s in production): +$0.5

- schedule and vessel: 72 h production, 12 h turnaround, 200 m3: −$1.9
- titer 50 -> 30 g/L: +$5.1
- scale 4.5 -> 10 kt/yr: −$1.7
- product: BLG, downstream to a dried concentrate: +$7.0

The OVA isolate variant adds precipitation and a second recovery loss: +$40.2 over the BLG concentrate. The order of the steps changes their sizes, not the total. The first 4 designs of the walk (marked † in Figure 5) break our secretion ceiling: NREL's design makes 50 g/L of cellulase with only 6.6 g/L of cells, which implies about 117 mg/g/h. Their costs are shown for the walk only.




Figure 6: One-at-a-time sensitivity around the industry benchmark (BLG, 42 g/L), coloured by the evidence grade of each parameter's range. † marks an end moved to the feasibility boundary.

Tornado (Figure 6). Around the industry benchmark the largest swings are:

1. titer (g/L): $12.9–$25.9 (grade C)
2. fermenter price (vendor quote vs aseptic cell-culture spec): $15.8–$22.9 (a convention)
3. cell density in production (g/L): $12.0–$17.3 (grade B), one end truncated at the feasibility boundary
4. production phase (h): $14.5–$18.5 (grade B)
5. capital convention (design-report vs price-anchored): $12.3–$15.8 (a convention)
6. turnaround between batches (h): $15.6–$17.8 (grade B)
7. glucose ($/kg): $15.3–$17.5 (grade B)

What sets fermenter productivity leads: titer, production time and cell density decide how much protein a vessel makes per hour and so how much capital each kg carries. Three points deserve emphasis:

- **Yield matters through the ceiling, not the sugar bill.** At a fixed titer the product yield moves the price by $0.7 (rank 20), because most sugar goes to making and maintaining cells. But it sets the sugar feed (and O_2) each gram of protein needs, and so the ceiling: at 72 h the ceiling is 48 g/L at 0.28, 57 g/L at 0.35, 63 g/L at 0.40, 70 g/L at 0.47, 73 g/L at 0.50, 76 g/L at 0.54 g/g.
- **Vessel size.** 100–500 m^3 moves the benchmark by $1.8 (rank 10).
- **The structural choices.** The fermenter price basis (vendor quote against aseptic cell-culture vessels) moves it by $7.1 (rank 2), and the accounting convention by $3.5 (rank 5).
- **The minimum starting volume** does not appear: raising it above 0.35 makes the benchmark design infeasible, so its bar is truncated to almost nothing; Section 4.4 shows what it does to the ceiling.

Few strain-performance ranges are grade A; they rest on design values and company-reported averages, because no public BLG or OVA fed-batch dataset reports titer, rate and yield together.




Figure 7: Monte Carlo over each scenario's evidence ranges (triangular), feasible plants only; the black tick is the scenario's own value.

Monte Carlo (Figure 7). Over the evidence ranges the industry benchmark's median is $18 for BLG (p10–p90 $14–$23) over feasible plants, and $19 ($15–$24) over all draws with a cost. 49% of draws are feasible. The rest fail as follows: 164 on initial fill, 22 on no plant (model error), 18 on OUR. The failing draws are those with many cells at the start of production (111 against 83 g/L on average) and fast growth in production (0.0090 against 0.0049/h): they need so much feed that the batch would have to start below 30% full. The scenario values sit at these percentiles of their own feasible distributions: industry BLG 29, industry OVA 29, academic BLG 13, academic OVA 22. For the academic strains the p10 is $43 (BLG) and $236 (OVA).

4.4 Q3: what must be true for $10 and $20 per kg




Figure 8: Titer needed for a target price, against production time, at the base cell density. Grey: designs the fermenter cannot run with a 500 g/L glucose feed. Dotted: the ceiling with a 600 g/L feed.

Production phase (h)	36	48	60	72	96	120
Ceiling, 500 g/L feed (base)	76	74	72	70	65	60
Ceiling, 600 g/L feed	97	109	107	105	102	98
$10/kg, price-anchored	64	68	—	—	—	—
$20/kg, price-anchored	27	29	30	32	35	39
$50/kg, price-anchored	10	11	11	12	13	14
$20/kg: implied q_p (mg/g/h)	8	7	6	5	4	4
$10/kg, price-anchored, 600 g/L feed	67	70	73	78	86	92
implied q_p (mg/g/h)	23	19	16	15	13	11
$10/kg, design-report	47	50	52	54	59	—
$20/kg, design-report	21	22	23	24	26	28
$50/kg, design-report	8	8	9	9	10	11

Table 6: BLG titer (g/L) needed for each target price at the base cell density (107 g/L), searched only below the feasibility ceiling; grey rows give the specific productivity that titer implies (the base design runs at 4.7). — means the target is not reachable at that production time (`results/analyses.json`, key `required`).

Figure 8 and Table 6 answer the inverse question at the base cell density:

- **$50/kg** needs 10–14 g/L depending on production time: 1.9–2.8× the best public BLG titer on glucose.
- **$20/kg** needs about 29–32 g/L in 48–72 h with price-anchored capital (design-report: 22–24 g/L), a specific productivity of 5–7 mg/g/h.
- **$10/kg** needs 64–68 g/L with price-anchored capital. With a 600 g/L feed it needs 67–92 g/L in 36–120 h, at 11–23 mg/g/h. With design-report capital it needs 47–59 g/L in 36–96 h.

The gate is secretion. Cell density is a free design choice, and at a fixed titer fewer cells cost less; what fewer cells cannot escape is the specific productivity. Table 7 lets the model choose cell density, production time and titer freely, with everything else at the base (yield, maintenance, 200 m³ vessels, 10 kt/yr, price-anchored capital), for four secretion ceilings. With secretion at 40 mg/g/h the cheapest BLG design costs $7.1/kg; at 10 mg/g/h it costs $12.0. The base design runs at 4.7 mg/g/h, Intelligen's design value; the best public BLG fed-batch implies about 1 (Yao et al. 2026: 13.88 g/L in 108 h at roughly 120 g/L cells). Fungal cellulase processes make 0.2–0.4 g protein per litre per hour (Ellilä et al. 2017), which at the 20–50 g/L of cells such processes run is roughly 5–20 mg/g/h (our arithmetic), for native enzymes, not animal proteins.

Secretion ceiling (mg/g/h)	$/kg BLG	Cells (g/L)	Production (h)	Titer (g/L)	Limits that bind
40	$7.1	70	63	98	O2 transfer, specific productivity
20	$8.7	90	81	87	O2 transfer, specific productivity, minimum initial fill
10	$12.0	100	90	62	O2 transfer, specific productivity
5	$18.6	100	90	36	O2 transfer, specific productivity
2.1 (best demonstrated)	$36.2	114	90	19	O2 transfer, specific productivity

Table 7: What process design alone can do: the cheapest BLG design with base biology, vessel, scale and price-anchored capital, when cell density, production time and titer are free, for five secretion ceilings (a coarse search; review SC-1). 2.1 mg/g/h is the best secretion rate we found demonstrated in a **K. phaffii** glucose fed-batch (Prielhofer et al. 2018).

How firm is the ceiling? It is a design ceiling, not a law of physics (Table 8). In the base it comes from the vessel's volume: the minimum starting volume (30%, the lowest start in a published design), the glucose feed (500 g/L, Intelligen's) and the cell density at the start of production (107 g/L) are all design choices. A 600 g/L feed lifts the 72 h ceiling to 105 g/L (minimum starting volume); 80 g/L of cells lifts it to 97 g/L; a 20% start to 121 g/L, still set by the minimum starting volume. Only with no volume limit at all does another limit take over: secretion ceiling at 127 g/L. The other way, a 35% start brings the 72 h ceiling down to 44 g/L, and at 40% 19 g/L (NREL's design starts at 50%). The ceiling also rises with the product yield (Section 4.3) and with lower production-phase maintenance: at 72 h it is 73 g/L at 0.0023, 70 g/L at 0.00437, 64 g/L at 0.008, 51 g/L at 0.0161 g/g/h.

Design choice	36 h	72 h	120 h
base	76 g/L (minimum starting volume)	70 g/L (minimum starting volume)	60 g/L (minimum starting volume)
glucose feed 600 g/L	97 g/L (secretion ceiling)	105 g/L (minimum starting volume)	98 g/L (minimum starting volume)
80 g/L cells at the start of production	75 g/L (secretion ceiling)	97 g/L (minimum starting volume)	91 g/L (minimum starting volume)
minimum fill 0.20	90 g/L (secretion ceiling)	121 g/L (minimum starting volume)	114 g/L (minimum starting volume)
minimum fill 0.35	52 g/L (minimum starting volume)	44 g/L (minimum starting volume)	33 g/L (minimum starting volume)
minimum fill 0.40	27 g/L (minimum starting volume)	19 g/L (minimum starting volume)	6 g/L (minimum starting volume)
minimum fill 0.50 (NREL)	—	—	—
no minimum fill	90 g/L (secretion ceiling)	127 g/L (secretion ceiling)	152 g/L (O2 transfer)
no minimum fill, O2 cap 150 (bubble column)	75 g/L (O2 transfer)	112 g/L (O2 transfer)	138 g/L (O2 transfer)

Table 8: BLG titer ceiling (g/L) by production time, and the limit that fails just above it. Base: 107 g/L cells, a 30% minimum start, a 500 g/L glucose feed, and an O_2 cap of 200 mol/m³/h per m³ of cell-free liquid.

4.5 Q4: the physical-limit best case




Figure 9: Cost structure from today's benchmarks to the physical-limit best case. The best academic strains are off this scale.

The search (Figure 9) finds the cheapest feasible design once every lever listed in Section 2.7 is at its assumed best. For BLG it has 83 g/L cells, a 84 h production phase and 140 g/L, at a specific productivity of 40 mg/g/h. The limits active there are O2 transfer, specific productivity, minimum initial fill.

It costs $4.1/kg BLG with price-anchored capital and $3.3 with design-report capital; for OVA $4.2 and $3.4.

- **It is set by an assumed secretion rate.** The binding biological limit is the 40 mg/g/h ceiling, a judgment (grade D) about 9× the base design and above any published ***K. phaffii*** value. Table 9 shows the case at other ceilings.
- **Capital still dominates:** 56% of the BLG case. Glucose ($0.84) is the largest non-capital item.
- **Engineering levers are part of it.** 400 m³ vessels, 100 kt/yr, contract sugar and the 600 g/L feed are not biology. With the base 500 g/L feed the BLG case costs $4.5; with aseptic cell-culture vessels, $5.7.
- **OVA and the 50 g/L question.** If Intelligen's 50 g/L were a real solubility limit at fermentation pH, the OVA case would cost $8.3 instead of $4.2.

Secretion ceiling (mg/g/h)	$/kg BLG	Cells (g/L)	Production (h)	Titer (g/L)
2.1 (best demonstrated)	$24.5	150	90	25
5	$12.1	150	90	53
10	$8.1	150	81	81
20	$5.4	114	90	118
40 (assumed)	$4.1	83	84	140
120 (NREL's design, implied)	$3.4	45	72	162

Table 9: The BLG physical-limit case at other secretion ceilings (price-anchored capital; the sensitivities use a coarser search). The base design runs at 4.7 mg/g/h.

The case is not a forecast and not a lower bound: design-report accounting, continuous operation or further relaxed design choices would all go lower. It is what the model gives once every assumed limit is at its best at the same time. Its yield sits at 95% of an energetic ceiling that leaves out transport, secretion and turnover costs. Its O_2 limit assumes water-like broth at 83 g/L cells and 140 g/L of dissolved protein, where viscosity and foam would lower k_La. And its heat removal (up to 25–40 kW/m³, i.e. about 16 MW per 400 m³ vessel) needs external heat-exchange loops that are not costed. The BLG case is below the 2015–2025 whey band; the OVA case is below the egg-white band.

4.6 Q5: claimed titers against the fermenter's limits

Claim or benchmark	Titer	Modelled as	At its own time: $/kg	Design-report	Shortest feasible time	q_p there	$/kg there
average titer in private companies' cost models (GFI 2025)	42 g/L	*K. phaffii*, 107 g/L cells	$15.8 (72 h)	$12.3	13 h	38	$12.3
average titer in published cost models (GFI 2025)	24 g/L	*K. phaffii*, 107 g/L cells	$25.9 (72 h)	$20.1	12 h	21	$19.9
Behm 2022 BLG design titer (T. reesei, 120 h)	50 g/L	*T. reesei*, 35 g/L cells	$13.5 (120 h)	$9.9	48 h	40	$10.2
T. reesei homologous secreted protein (Behm 2022, grade C)	100 g/L	*T. reesei*, 35 g/L cells	infeasible at 72 h	—	143 h	40	$8.7
Onego platform claim (protein not specified; tested as OVA; press, grade C)	120 g/L	*T. reesei*, 35 g/L cells	infeasible at 72 h	—	not reachable	—	n/a

Table 10: Claimed and benchmark titers in a 200 m³ vessel: the cost at the claim's own production time (Behm's design: 120 h; otherwise 72 h), and the shortest production phase in which the titer is feasible (O_2 transfer, heat removal, minimum starting volume, 40 mg/g/h secretion ceiling), with the specific productivity and price-anchored cost there. The shortest-time columns are a best case: secretion runs at or near its ceiling. The Onego figure is a platform claim for an unnamed protein, tested as OVA.

Of the 5 claims and benchmarks, 4 are feasible if the production phase is right (Table 10). The price of a high titer is time or secretion rate: at its own 72 h the 42 g/L average costs $15.8, the industry benchmark of Section 4.2; squeezed into the shortest feasible production phase it costs $12.3, but only by secreting 38 mg/g/h. A claimed titer alone therefore does not settle the cost. The check is physical; whether a strain can sustain the implied productivity is a separate, biological question. Onego platform claim (protein not specified; tested as OVA; press, grade C) cannot be reached at any production phase up to 20 days in our model: either the O_2 demand exceeds what the broth can take up, or the batch would have to start below 30% of the vessel because the feed fills it first. Such claims need better O_2 transfer than our assumptions (for **T. reesei**, a k_La half that of yeast broth, grade D), a denser feed, or a different process (continuous, repeated fed-batch or cell recycle), none of which is modelled here.

4.7 Q6: scale and siting




Figure 10: Industry-benchmark BLG against plant output, US Midwest and north-west Europe, both capital conventions.

Scale (Figure 10). Going from 1 to 10 kt/yr cuts the price by $15.2, and going from 10 to 100 kt/yr saves a further $1.3. Beyond about 10 kt/yr the plant adds 200 m³ fermenters rather than larger ones, so only the downstream train and utilities keep scaling; larger vessels are a separate lever (Section 4.3).

Siting. A Netherlands plant costs $3.1/kg more than one in the US Midwest (base BLG). One at a time, US-level inputs would remove:

- glucose ($0.61 against $0.40/kg): $2.2;
- electricity and gas ($0.146 against $0.086/kWh; $19.6 against $4.8/GJ): $0.8;
- construction cost (1.09×): $1.0;
- wages: nothing (Dutch and Iowa operator pay are within a few per cent)
- spent-cell disposal (anaerobic digestion in the EU against US landfill): −$0.9, i.e. the EU is cheaper here;
- water and wastewater: +$0.0.

The one-at-a-time effects need not add up to the gap exactly, because the inputs interact.

4.8 Q7: milk and egg equivalents



Figure 11: The cost of replacing the protein of a litre of milk (left) and the egg-white protein of an egg (right) with PF protein, against titer (72 h production, price-anchored capital), beside milk and egg prices in 2025 dollars.

A litre of cow's milk holds 33.7 g of protein, 3.3 g of it BLG; a large egg holds 3.64 g of egg-white protein, 1.96 g of it OVA (`data/evidence/milk_egg.yaml`). At the base cost, the PF protein needed to replace all the protein of a litre of milk costs $0.71, against $0.48 for the litre at the US farm gate in 2025 and $1.07 at retail.

Replacing one egg's white costs $0.077, against $0.311 per egg wholesale in 2025, a year of avian-influenza prices ($0.096–$0.311 over 2015–2025).

Dividing a source's price by its protein gives the price per kg of protein at which PF would match it: US farm milk sold its protein at $14–$19/kg in 2015–2025, and wholesale eggs their egg-white protein at $26–$86/kg. With a 72 h production phase (ceiling 70 g/L), BLG matches the 2025 farm-milk protein price at 48 g/L, and OVA matches the lowest egg-white protein price of the decade at 24 g/L.

These are upper bounds on what the protein is worth: charging the whole price of milk or eggs to their protein makes them look dearer per gram than they are, since fat and lactose carry much of the milk price and the yolk part of the egg's. Bought as the incumbent ingredient, the same protein is much cheaper: the white of one egg, as dried egg white, cost $0.035–$0.077 in 2015–2025. Whey is a by-product of cheese: the whey protein of a litre of milk (6.6 g) at WPC80/WPI prices was worth 11%–21% of the US farm milk price in the same year, so replacing whey protein cuts milk output far less than the "litre of milk" framing suggests. For animal welfare, egg white is the stronger lever: per kg of target protein it involves roughly 50–60 times more animal-years than whey BLG (about 0.6 kg OVA per hen-year against about 35 kg BLG per cow-year). These last two figures are our order-of-magnitude arithmetic, not model outputs.

4.9 Q8: a high-value contrast, lactoferrin

Lactoferrin, an iron-binding milk protein, sells as a supplement and infant-formula ingredient at tens of times the price of whey (14–183× across the two price ranges). Two recombinant bovine lactoferrins made in **K. phaffii** without methanol have FDA "no questions" letters (GRN 1219 TurtleTree, 2025; GRN 1284 All G, 2026). We model it at 250 t/yr (Intelligen's design makes 251 t/yr of powder; the world market is roughly 0.3–1.6 kt/yr, and sources disagree by five times), with the dairy downstream route: microfiltration, cation exchange, UF/DF and spray drying to 90% lactoferrin. The chromatography step uses a binding capacity of 75 g/L of resin and 16 column volumes of buffer per cycle (EVERY patent), elution at 1 M NaCl, 83.2% step recovery, and resin at $1,250/L (Petrides) lasting 1,000 cycles (industrial lactoferrin columns, Pires et al. 2025). The column is loaded with dilute microfiltered broth, 26 column volumes per cycle, so a cycle takes 2.1 h at 20 column volumes per hour, the flow of Intelligen's design (500 cm/h through a 25 cm bed of a fast-flow agarose resin; version 2.1 guessed 5). One guess (grade E) remains: the column hardware per litre of resin.




Figure 12: Lactoferrin cost against titer, the reported dairy lactoferrin price range (2025 dollars), and Intelligen's own unit production costs for its 10 g/L design, with and without chromatography (costs of production, not selling prices; Table 12).

Case (250 t/yr)	Titer (g/L)	Price-anchored	Design-report
Design titer, cation exchange (Intelligen's design)	10	$178	$136
Best public methanol-free titer (porcine LF, Yen 2024)	2.8	$394	$294
Design titer, UF/DF only (optimistic)	10	$131	$92
Intelligen's model, cost of production, UF/DF only (case a)	10		$299 per kg
Intelligen's model, cost of production, cation exchange (case b)	10		$441 per kg
Dairy lactoferrin, reported 2013–2025 (grade C; 2025 $)			$250–$1,585 per kg

Table 11: Lactoferrin, $/kg lactoferrin (2025). Resin life is the largest downstream lever: at 125 cycles (biopharma practice) the design case costs $326. Scale: $291 at 100 t/yr, $178 at 250 t/yr, $145 at 500 t/yr, $127 at 1000 t/yr, $123 at 1588 t/yr, $119 at 2500 t/yr, $117 at 5000 t/yr, $115 at 10000 t/yr.

$/kg lactoferrin (2025)	Intelligen a	Ours, their design a	Intelligen b	Ours, their design b
Raw materials	$101.2	$14.8	$123.5	$22.4
Labour	$41.6	$19.7	$47.2	$24.8
Facility (maintenance, insurance, depreciation)	$91.6	$61.3	$122.5	$73.9
Laboratory, QC and QA	$11.6	$0.0	$13.2	$0.0
Consumables (resin, membranes)	$29.8	$0.6	$104.8	$81.6
Waste treatment	$6.5	$5.9	$11.1	$5.9
Utilities	$17.1	$3.1	$18.7	$3.1
Cost of production	**$299.3**	**$105.5**	**$441.0**	**$211.6**
Our base design (Table 11)		$113.3		$158.6
Total capital investment	$126M	$73M	$150M	$79M

Table 12: Intelligen's lactoferrin model against ours, in SuperPro's operating-cost categories. Intelligen's figures are its 2024 costs (Tables 2–5 of the report) per kg of lactoferrin, deflated to 2025; "ours, their design" runs our model with the design inputs their report states (scale, six 100 m³ fermenters, diavolumes, column flow, loading and buffers, step recovery, resin at $1,000/L lasting 150 cycles, electricity at $0.10/kWh) and our own unit prices. Our facility line is maintenance, insurance and ten-year depreciation. `data/pf/scenarios/lactoferrin.yaml`, variants `lf_intelligen_a` and `lf_intelligen_b`; review L.

At the design titer PF lactoferrin costs $136–$178 per kg (design-report to price-anchored capital), below the reported range for dairy lactoferrin in 2025 dollars. That range comes from reports and trade press, partly from a PF-lactoferrin seller (TurtleTree, 2023), and the price is volatile (it fell to about $200/kg nominal in the 2016 glut).

Against Intelligen's own model. Intelligen published the design our case takes its titer from (Da Gama Ferreira et al. 2026): six 100 m³ fermenters, glycerol growth and methanol induction to 10 g/L, and two downstream routes. Its unit production costs, per kg of lactoferrin in 2025 dollars, are $299 without chromatography and $441 with cation exchange. They include depreciation and no return on capital, so they compare with our cost of production, $113 and $159: theirs are 2.6× and 2.8× ours. Their design inputs explain a small part of the gap. With their output, vessels, diavolumes, column flow and loading, buffers, step recovery, resin price and life, and electricity price, our cost of production becomes $105 and $212 (Table 12); the largest single input is their short resin life (150 cycles against our 1,000). The rest is their unit costs, category by category: raw materials (glycerol at about $1.8/kg, and a part of the line that the report does not itemise), capital (their investment is 1.9× ours for the same design; their equipment list is not published), labour, membranes, utilities, and a quality-control line that we do not have. We keep our sourced prices; theirs are SuperPro inputs without a stated basis. Our cost sits below the reported price range and Intelligen's inside it, but only at the design titer: at the best public methanol-free titer the cost is $394/kg. No PF lactoferrin price has been disclosed, so this is a cost comparison only. The contrast with BLG is the point: the same fermentation and accounting that leave the base BLG above the whey band put lactoferrin below its incumbent's range at a design titer, because that incumbent is scarce rather than a by-product.

5 Discussion

5.1 Answers, in brief

- **Q1.** Today's public strains make BLG and OVA at $43 and $246 per kg: not competitive. The titer private cost models assume gives $15.8 (BLG) and $15.9 (OVA), or $12.3 and $12.4 with design-report capital. Against 2015–2025 prices, BLG is then inside the whey band (inside it with design-report capital) and OVA inside the egg-white band.
- **Q2.** Fermenter productivity (protein per hour of vessel time) drives the cost, then the capital basis (vessel prices and fixed-cost rule); sugar and energy prices come second. The product yield acts mainly through the feasibility ceiling. The strain parameters are the weakest evidence: design values (B) and survey averages (C).
- **Q3.** With price-anchored capital, $20/kg needs 29 g/L at 48 h or 32 g/L at 72 h at the base cell density. What $10/kg needs above all is secretion: with today's biology and capital, process design alone reaches $7.1 at 40 mg/g/h and $12.0 at 10.
- **Q4.** The physical-limit best case is $4.1 (BLG) and $4.2 (OVA) per kg, bound by O2 transfer, specific productivity, minimum initial fill at a secretion ceiling of 10 mg/g/h the BLG case is $8.1.
- **Q5.** 4 of 5 claimed titers are feasible in our fed-batch model, some only with long production phases or secretion near the ceiling.
- **Q6.** Scale matters up to about 10 kt/yr. EU siting adds about $3.1/kg.
- **Q7.** The base BLG replaces the protein of a litre of milk for $0.71, while the litre costs $0.48 at the farm gate; bought as WPC80/WPI the same whey protein is a fraction of that.
- **Q8.** At a design titer, PF lactoferrin costs $178/kg, below the reported range for dairy lactoferrin; Intelligen's own model costs 2.8× ours to produce.

5.2 Why published PF costs are lower

Published PF cost estimates for food proteins span roughly $12–39/kg: GFI Europe and Arthur D. Little (2025) give $24–33/kg for their "P1" molecules today and $12/kg for α-lactalbumin after improvements; Intelligen (2025) $39/kg for an OVA isolate at 1 kt/yr. Of the designs we can rebuild from their own inputs, Intelligen's OVA comes within −4% of our model once recast into our capital charge; NREL's cellulase and Voutilainen's recombinant case come out 1.8× lower. With vendor-quote fermenters our capital matches theirs; the gap is mostly accounting: fixed costs of 1.5–2% of capital a year instead of Humbird's 9%, and lighter capital charges. Intelligen's lactoferrin model runs the other way: its cost of production is 2.8× ours, through its unit prices for raw materials, labour and consumables and a capital estimate 1.9× ours for the same design (Table 12). Published costs differ from ours in both directions, for reasons that can be traced line by line.

Neither convention is "correct". Humbird's reproduces the cost of one fermented commodity in his own workbook; the design-report one matches how process-design studies cost greenfield enzyme plants. We read our base as a central estimate on Humbird's method and design-report capital as an optimistic one, not as a confidence interval.

5.3 For funders

In order of expected cost reduction per unit of evidence gained (our judgment):

1. **Secretion rate without growth.** How much BLG or OVA a strain secretes per gram of cells per hour when it barely grows, over a production phase of two to five days at 1–10 m³. It sets both today's cost and the best case; the one near-zero-growth study we found saw secretion fall with growth, and every compiled glucose process secretes fastest at 0.1–0.2/h (Looser et al. 2015; Prielhofer et al. 2018).
2. **Public fed-batch data.** Titer, productivity and yield measured **together** for BLG and OVA; no public dataset reports all three, and the product yield is currently derived, not measured.
3. **The design ceiling.** The workable starting volume, the most concentrated feed and O_2 transfer at production scale and cell density move the titer ceiling itself.
4. **Real capital and fixed costs.** Published capital, staffing and running costs of recent PF plants (for example Liberation Labs' 600 m³ Richmond plant) would settle the fermenter price basis and the fixed-cost rule.
5. **Downstream data.** Diafiltration flux at scale (one patent value, 5 L/m²/h, sizes most of the membrane area) and dryer-feed solids for PF proteins.

On the evidence here, PF BLG at the titer private cost models assume would undercut whey in a spike year such as 2026. With design-report capital ($12.3) it would have undercut whey in 8 of the 21 annual export prices of 2015–2025; with price-anchored capital ($15.8) in 3. High-value proteins such as lactoferrin cost less than most reported incumbent prices at a design titer. OVA's position against egg white depends on the year, because egg-white prices swing more than whey's.

> **Check (reader):** A judgment, not a model output, written by the AI assistant and not yet endorsed by the author (please revise): the probability that a PF plant built with 2026 technology makes BLG below $10/kg protein is about 0.15, with low confidence (unchanged from version 2: the cheaper vessels raise it, the secretion requirement lowers it). The model reaches $10 only if cells secrete several times faster than any published **K. phaffii** process, sustained over days, or with the lighter design-report accounting and high titers together.

5.4 For EU policy

At Dutch sugar, energy, wage and construction prices the base BLG plant costs $3.1/kg more than in the US Midwest. Cheaper capital is a large lever: a capital charge at 4% over 20 years instead of 7.5% over 10 cuts $3.9 from the EU cost; of that, a 20-year recovery alone cuts $2.5 and a 4% rate alone $1.2. US-level glucose would remove $2.2 and US-level energy $0.8. These are cost effects on one plant; the model says nothing about which instruments would deliver them.

5.5 What would change these conclusions

- A BLG or OVA strain that secretes ≥10 mg/g/h over days at near-zero growth, or evidence that it cannot, would move both today's cost and the best case more than any other single result.
- A peer-reviewed BLG or OVA fed-batch at ≥40 g/L in ≤48 h with a measured yield would test the derived product yield and the ceiling.
- The ceiling rests on design choices: a 30% minimum starting volume, a 500 g/L glucose feed and 107 g/L of cells. Industrial practice with lower starts, denser feeds or fewer cells lifts it; then secretion and O_2 transfer set it.
- Published capital and running costs of recent PF plants would settle the fermenter price basis and the fixed-cost rule.
- Measured OVA solubility at pH 5, and protease control for OVA in **T. reesei**.

6 Limitations

- **Strain parameters.** Few are grade A. The product yield is derived from a design's lumped stoichiometry, not measured; specific productivities are design values or survey averages, and the base production phase assumes secretion without growth (Section 2.3). 12 titer rows in `data/titers.csv` are not yet verified in full text.
- **One carbon source.** Methanol-induced and ethanol-fed production phases are priced as glucose; for ethanol this understates O_2 and heat.
- **The feasibility limits are design choices**: the minimum starting volume (30%), the feed concentration (500 g/L glucose) and the cell density, which together set the base ceiling; the O_2 cap and its basis (200 mol/m³/h per m³ of cell-free liquid); the heat-removal ceiling (25 kW/m³, grade D); and the secretion ceiling (40 mg/g/h), a judgment.
- **Directions of the simplifications.** k_La ignores viscosity and foam (optimistic at high cell density and titer); specific productivity is constant for up to 120 h (optimistic for long production phases); no product is made during growth (conservative for constitutive promoters); the energetic ceiling leaves out transport, secretion and turnover (optimistic).
- **OVA** at fermentation pH: solubility is unmeasured (50 g/L is shown as a sensitivity), and **T. reesei** clips much of it.
- **Continuous processes** and cell recycle are not modelled.
- **Chromatography** is modelled only for lactoferrin, with guessed column hardware (the flow rate is Intelligen's design value) and no itemised buffer tanks (inside the missing-equipment allowance).

- **Not costed:** start-up and commissioning, working capital (except in the DCF price), failed batches, QC release testing, packaging, dewatering and inactivation of spent cells, and external heat-exchange loops for the largest vessels.
- **EU wastewater** is priced as in the US; the Dutch levy is per kg of oxygen demand, which is not modelled.
- **Guesses (grade E)** remain for: `hosts.treesei.mu_max_per_h`, `dsp.cex_hardware_usd_per_L_resin`, `utilities.sterilizer_peak_factor`, `utilities.boiler_peak_factor`, `labour.fermenters_per_continuous_crew`, `capital.dsp_area_factor`, `capital.compressor_building_m2_each`.

7 Scoring the pre-registered priors

The priors were written in `docs/SCOPE.md` by the AI assistant (Claude) on 2026-09-26, before any model output existed, and before the author had seen them. They were committed to git together with the first outputs, so the repository cannot prove the order; the session record shows it but is not public.

Prior (2026-09-26)	Stated	Outcome (v2)	Score
1a. Intelligen configuration, $/kg OVA (80% interval)	$38–60	$44.7/kg OVA	inside
1b. Base case, $/kg protein (80% interval)	$28–50	$21.2 (BLG), $21.3 (OVA)	outside
1 (superseded). Purified BLG/OVA at ~30 g/L and 1–10 kt/yr	$20–40	10 kt/yr: $21.2 concentrate, $26.0 OVA isolate; 1 kt/yr: $44.7 (Intelligen configuration)	inside at 10 kt/yr, outside at 1 kt/yr
2. P(physical floor < $10/kg)	0.65	BLG $4.1 at the assumed 40 mg/g/h (yes); at 10 mg/g/h $8.1 (yes)	conditional: Brier 0.12 if secretion reaches 40 mg/g/h, 0.12 if only 10
3. P(a public BLG/OVA yield on sugar turns up)	0.20	no measured yield; design values exist (Behm 0.223, Intelligen 0.28 lumped)	Brier 0.04 (measured); 0.64 if design values count
4. P(first-try calibration within tolerance): yeast / NREL / Intelligen	0.6 / 0.6 / 0.5	yes / no (+136%) / yes (−3.6%)	Brier 0.16 / 0.36 / 0.25
5. P(public v1 by 2027-04-30 / 2027-07-31)	0.5 / 0.8	open	—

Table 13: Pre-registered priors and outcomes. Prior 4 is scored on the first run (2026-09-26), before the changes recorded in `docs/VALIDATION.md` and `docs/AUDIT.md`. Prior 3 meant a measured strain yield; both readings are scored. Prior 2 is scored once (BLG and OVA are one event), and its outcome depends on the secretion ceiling chosen during the audit, so it is scored conditionally.

The base-case interval prior missed low: after the review moved the fermenters from aseptic cell-culture prices to an industrial vendor quote, the base case ($21.2) fell below the $28–50 interval; with aseptic vessels it would be $30.9, inside. The superseded prior was too optimistic at 1 kt/yr, as its replacement said. The calibration prior was about right on average; the NREL miss came from capital conventions, which it did not anticipate.

8 Reproducibility

```
.venv/bin/python -m pytest                      # model, calibration and invariant tests
.venv/bin/python scripts/run_analyses.py        # all results -> results/analyses.json
.venv/bin/python scripts/make_figures.py        # docs/figures/*.png
typst compile --root . report/pf_tea_report.typ # this document
```

Inputs are in `data/pf/base.yaml` (every value with a range, grade and source), with scenario overrides in `data/pf/scenarios/`. The evidence files are `data/evidence/*.yaml`, with verbatim quotes and locators. Calibration cases are in `data/calibration/`. The audit is `docs/AUDIT.md`.

References

- Almeida Benalcázar E, et al. Single cell protein production from ethanol: model-based bioreactor operation at industrial scale. Biotechnol Bioeng 122:1441 (2025). doi:10.1002/bit.28969 (reports Roels 1983: 460 kJ per mol O_2).

- Aro N, et al. Production of bovine β-lactoglobulin and hen egg ovalbumin by *Trichoderma reesei* using precision fermentation technology and testing of their techno-functional properties. Food Res Int 163:112131 (2023). doi:10.1016/j.foodres.2022.112131
- Behm K, et al. Comparison of carbon footprint and water scarcity footprint of milk protein produced by cellular agriculture and the dairy industry. Int J LCA 27:1017 (2022). doi:10.1007/s11367-022-02087-0
- Carnicer M, et al. Macromolecular and elemental composition analysis and extracellular metabolite balances of *Pichia pastoris* growing at different oxygen levels. Microb Cell Fact 8:65 (2009). doi:10.1186/1475-2859-8-65
- Chung BKS, et al. Genome-scale metabolic reconstruction and in silico analysis of methylotrophic yeast *Pichia pastoris* for strain improvement. Microb Cell Fact 9:50 (2010). doi:10.1186/1475-2859-9-50
- Da Gama Ferreira R, Misailidis N, Parisis V, Petrides D. Egg white protein (ovalbumin) production via precision fermentation: process modeling and cost analysis using SuperPro Designer. Intelligen (2025). doi:10.13140/RG.2.2.29080.87048
- Da Gama Ferreira R, Misailidis N, Petrides D. Lactoferrin production via precision fermentation: process modeling and evaluation using SuperPro Designer. Intelligen, February 2026 version.
- The EVERY Company (Kale A, Patnaik R). Purified protein compositions and methods of production, US 2024/0083940 A1; Recombinant protein recovery methods, US 2025/0134130 A1; Ovalbumin protein compositions, US 12,096,784 B2.
- Ellilä S, Fonseca L, Uchima C, et al. Development of a low-cost cellulase production process using *Trichoderma reesei* for Brazilian biorefineries. Biotechnol Biofuels 10:30 (2017). doi:10.1186/s13068-017-0717-0
- FDA GRAS Notices 1056 (β-lactoglobulin, Remilk), 1219 (bovine lactoferrin, TurtleTree) and 1284 (bovine lactoferrin, All G).
- GFI & Hawkwood Biotech. Driving down costs of fermentation-derived ingredients (2025). doi:10.62468/trxj5734
- GFI Europe & Arthur D. Little. Pathfinding towards precision fermentation viability (2025).
- Garrigós-Martínez J, Nieto-Taype MA, Gasset-Franch A, et al. Specific growth rate governs *AOX1* gene expression, affecting the production kinetics of *Pichia pastoris* (*Komagataella phaffii*) P_{AOX1}-driven recombinant producer strains with different target gene dosage. Microb Cell Fact 18:187 (2019). doi:10.1186/s12934-019-1240-8
- Heijnen JJ, van Dijken JP. In search of a thermodynamic description of biomass yields for the chemotrophic growth of microorganisms. Biotechnol Bioeng 39:833 (1992).
- Heyland J, Fu J, Blank LM, Schmid A. Quantitative physiology of *Pichia pastoris* during glucose-limited high-cell density fed-batch cultivation for recombinant protein production. Biotechnol Bioeng 107:357 (2010). doi:10.1002/bit.22836
- Humbird D. Scale-up economics for cultured meat. Biotechnol Bioeng 118:3239 (2021). doi:10.1002/bit.27848 (and SI workbook)
- Humbird D, Davis R, McMillan JD. Aeration costs in stirred-tank and bubble column bioreactors. Biochem Eng J 127:161 (2017). doi:10.1016/j.bej.2017.08.006
- Humbird D, Davis R, Tao L, et al. Process design and economics for biochemical conversion of lignocellulosic biomass to ethanol. NREL/TP-5100-47764 (2011).
- IEAGHG (Amec Foster Wheeler). Effects of plant location on the costs of CO_2 capture. Technical Report 2018/04 (2018).
- Li Y, et al. Transcriptional reprogramming with endogenous cis- and trans-elements for efficient food protein expression from safe substrates in *Komagataella phaffii*. Metab Eng (2026). doi:10.1016/j.ymben.2026.102555
- Liu WC, et al. Scaling-up fermentation of *Pichia pastoris* to demonstration-scale using new methanol-feeding strategy and increased air pressure instead of pure oxygen supplement. Sci Rep 6:18439 (2016).
- Looser V, Bruhlmann B, Bumbak F, et al. Cultivation strategies to enhance productivity of *Pichia pastoris*: a review. Biotechnol Adv 33:1177 (2015). doi:10.1016/j.biotechadv.2015.05.008
- Martin WF, et al. ATP requirements for growth reveal the bioenergetic impact of mitochondrial symbiosis. Biochim Biophys Acta Bioenerg (2025). doi:10.1016/j.bbabio.2025.149564
- Maskow T, Kemp R, Buchholz F, et al. What heat is telling us about microbial conversions in nature and technology. Microb Biotechnol 3:269 (2010).
- Maurer M, et al. Versatile modeling and optimization of fed batch processes for the production of secreted heterologous proteins with *Pichia pastoris*. Microb Cell Fact 5:37 (2006).
- Nieto-Taype MA, Garcia-Ortega X, Garrigós-Martínez J, et al. Rationale-based selection of optimal operating strategies and gene dosage impact on recombinant protein production in *Komagataella phaffii* (*Pichia pastoris*). Microb Biotechnol 13:315 (2020). doi:10.1111/1751-7915.13498
- Mercier-Bouchard D, et al. Process efficiency of casein separation from milk using polymeric spiral-wound microfiltration membranes. J Dairy Sci 100:8838 (2017). doi:10.3168/jds.2017-13015
- Petrides D. Bioprocess design and economics. In Harrison RG, et al., Bioseparations Science and Engineering, 2nd ed. Oxford University Press (2015), ch. 11 (Intelligen web version).
- Pires MC, et al. Column ageing of agarose-based chromatography in industrial lactoferrin purification. J Chromatogr A 1760:466274 (2025).
- Prielhofer R, Reichinger M, Wagner N, et al. Superior protein titers in half the fermentation time: promoter and process engineering for the glucose-regulated *GTH1* promoter of *Pichia pastoris*. Biotechnol Bioeng 115:2479 (2018). doi:10.1002/bit.26800
- Rebnegger C, Graf AB, Valli M, et al. In *Pichia pastoris*, growth rate regulates protein synthesis and secretion, mating and stress response. Biotechnol J 9:511 (2014). doi:10.1002/biot.201300334

- Rebnegger C, et al. *Pichia pastoris* exhibits high viability and a low maintenance energy requirement at near-zero specific growth rates. Appl Environ Microbiol 82:4570 (2016).
- Rebnegger C, et al. Protein production dynamics and physiological adaptation of recombinant *Komagataella phaffii* at near-zero growth rates. Microb Cell Fact 23:44 (2024).
- Tetra Pak. Dairy Processing Handbook (online), chapters 7.6, 18 and 19.
- Tijhuis L, van Loosdrecht MCM, Heijnen JJ. A thermodynamically based correlation for maintenance Gibbs energy requirements in aerobic and anaerobic chemotrophic growth. Biotechnol Bioeng 42:509 (1993).
- Tsai LS, Ijichi K, Harris MW. Concentration of egg white by ultrafiltration. J Food Prot 40:449 (1977).
- van ‘t Riet K. Review of measuring methods and results in nonviscous gas-liquid mass transfer in stirred vessels. Ind Eng Chem Process Des Dev 18:357 (1979).
- Voutilainen E, Pihlajaniemi V, Parviainen T. Economic comparison of food protein production with single-cell organisms from lignocellulose side-streams. Bioresour Technol Rep 14:100683 (2021).
- Wang L, et al. The thermodynamic and gelation properties of ovalbumin and lysozyme. Gels 11:470 (2025).
- Yao C, Zhu H, An Q, Zhang M, Zheng Y, Cai M. Omics-guided global rewiring of yeast metabolism enables high-level production of β-lactoglobulin. Food Biosci 82:109383 (2026).
- Yen CC, Wu PY, Ou-Yang H, et al. Production of bioactive porcine lactoferrin through a novel glucose-inducible expression system in *Pichia pastoris*. Int J Mol Sci 25:1818 (2024).
- Yver AL, et al. Fractionation of whey protein isolate with supercritical carbon dioxide: process modeling and cost estimation. Int J Mol Sci 13:240 (2012).
- Price, wage and trade statistics: US EIA, Eurostat (nrg_pc_203/205, lc_lci_lev, earn_ses22, Comext DS-045409), US BLS (OEWS, PPI, CPI-U), USDA (NASS, ERS, AMS Dairy Market News, FoodData Central), UN Comtrade, ECB, EC agri-food portal, WRAP and EREF. Full locators in `data/evidence/`.

A Parameters (generated from `data/pf/base.yaml`)

Parameter	Value	Range	Gr.	Source
plant.product	blg			blg | ova
plant.spec	concentrate			concentrate (GRN 1056-type, UF/DF + dry) | isolate (adds precipitation) | cex (adds cation-exchange capture; lactoferrin) | crude (whole broth)
plant.annual_target_protein_t	10000	1000–100000	C	gfi2025costs Table 2: private-sector volumes 2,500-25,000+ t/yr; SCOPE base
plant.operating_days	330	320–350	B	NREL 2011 uses 8,410 h/yr (96%); Humbird A.1 350 d; Intelligen ~342 batches x 24 h; 330 d allows for campaigns and cleaning
plant.location	us_midwest			us_midwest | eu_nw
strain.host	kphaffii			
strain.titer_g_L	30	5–60	B	intelligen_ovalbumin2025 design basis 30 g/L; Every2024-HY ovomucoid 30.19 g/L (patent Ex. 9); gfi2025costs public avg 24, private avg 42. For scale: the highest secreted titer in the Looser 2015 review's compilation is 19.55 g/L (glucose oxidase, methanol/ mannitol, 168 h, 100 -> 245 g/L cells; Table 4). Tornado ends beyond the feasibility boundary are truncated to it
strain.t_prod_h	72	48–120	B	intelligen_ovalbumin2025 p.6: expression phase 72 h; NREL 2011: 96 h production
strain.X_prod_g_L	107	40–150	B	derived from intelligen_ovalbumin2025: cells at the start of expression = 92 g/L x 107 m3 at harvest (p.7) - 8/28 x 3,210 kg rOVA made (Eq. 2) = 8,927 kg, in 107 m3 less the expression-phase additions (11,464 kg glucose as 500 g/L solution 22.9 m3 + 30% ammonia 3.5 m3 - net O2/CO2 exchange 2.2 m3) = 82.8 m3 -> 108 g/L; feeding then dilutes it to 92 g/L at harvest. Range: K. phaffii glucose fed-batch 93-169 g/L at harvest (Shemesh 2024, A); patents 100-200 (Every2024-HY, B); hosts.yaml
strain.X_max_g_L	150	100–200	A	K. phaffii glucose fed-batch 93-169 g/L (Shemesh 2024); typical 150; patents up to 200 (hosts.yaml). Growth stops here. Heyland 2010 grew at constant rate only up to ~155 g/L, slowing above it (218 g/L after 65 h in 2.6 L), and states such densities are reachable only in small fermenters because O2 transfer and heat removal limit larger ones (hosts.yaml kphaffii_glucose_hcd_fedbatch_heyland2010)
strain.mu_prod_per_h	0.0014	0–0.02	B	derived from intelligen_ovalbumin2025 Eq. 2: 917 kg cells made on ~8,930 kg in 72 h (8 g per 28 g rOVA at 30 g/L x 107 m3; 92 g/L at harvest) -> ln(9,844/8,927)/72 = 0.00136/h. Range: 0 (non-growing production) to 0.02 (K. phaffii production-phase growth rates, hosts.yaml; D). Growth stops at X_max: from 107 g/L the high end reaches 150 g/L after ~17 h and the phase continues without growth
strain.Y_ps_g_g	0.465	0.28–0.535	B	derived so that the model reproduces intelligen_ovalbumin2025 Eq. 2 (expression phase: 100 g glucose -> 28 g rOVA + 8 g cells + 4 g HCP, maintenance included). Glucose per g product in expression = 1/Y_ps + (8/28)/Y_x + m_s,prod/q_p = 100/28, with Y_x = 0.5705 g/g (the model's Heijnen growth yield for K. phaffii), m_s,prod = 0.00437 g/g/h (hosts.kphaffii.m_s_prod_g_g_h) and q_p = 3,210 kg / 675,213 kg.h = 0.00475 g/g/h (Intelligen) -> 1/Y_ps = 3.5714 - 0.5008 - 0.9192 = 2.151, Y_ps = 0.465 (audit R2-Q2/R6-C2: 0.28 is a lumped yield, not a reaction yield). The glucose of Intelligen's 4 g of host-cell protein is charged to the target (read with the HCP as product,

Parameter	Value	Range	Gr.	Source
				the yield would be 0.53; review HM-6). The inversion is sensitive to m_s,prod: at 0.0023 it gives 0.39 (review FS-4). low = 0.28 (the lumped value read as a reaction yield, as in v1); high = 0.535 (0.95 x the energetic ceiling 0.564 of BLG with amino-acid synthesis and sulfate assimilation counted, product.energetic_ceiling; review FS-2/SC-2; D). Checks: Behm 2022 design 0.223 g/g overall (T. reesei BLG); NREL cellulase 0.24 g/g overall
strain.q_p_max_mg_g_h	40	5–40	D	judgment: a physiological ceiling on specific secretion, applied as a feasibility limit. No K. phaffii measurement is near it: Intelligen's design runs at 4.75 mg/g/h (low, B); the best public BLG fed-batch implies ~1 (Yao 2026, 13.88 g/L in 108 h at ~120 g/L cells); a VHH-secreting strain peaked at 0.64 and fell to 0.019 at near-zero growth (Rebnegger 2024, hosts.yaml); the Looser 2015 review compiles q_P,max of secreted proteins from 0.004 to 0.69, the glucose (GAP) ones 0.05-0.49 at mu 0.15-0.20/h, growth-coupled; the best glucose fed-batch on file secreted 2.1 near mu 0.1/h (Prielhofer 2018, hosts.yaml; the report's 'best demonstrated' case); fungal enzyme processes reach ~5-20 (Ellila 2017 productivities at 20-50 g/L cells; review FS-3). Our model of NREL's T. reesei cellulase design implies ~118 (a model output at 6.6 g/L cells, not physiological evidence; shown only as a sensitivity). Binds only in the physical-limit and required-titer analyses, where the report shows the floor at 5-120
strain.X0_g_L	0.88	0.88–15	B	derived: inoculum 5% v/v (Intelligen OVA p.6) x the seed density fermenter.seed_final_X_g_L = 17.6 g/L (Intelligen's lactoferrin flowsheet; one number for both, review FS-11). v2.1 used 3 (from a guessed 60 g/L seed). Humbird A.1 starts at 15 g/L (high)
hosts.kphaffii.biomass_formula	(table in file)		A	Carnicer et al. 2009 Microb Cell Fact 8:65, Table 4: Fab-producing K. phaffii CH1.665N0.134O0.602S0.0039 (evidence hosts.yaml; re-check table cell)
hosts.kphaffii.mu_max_per_h	0.2	0.15–0.28	A	hosts.yaml: mu_max ~0.2 1/h on sugar vs 0.11-0.12 on methanol. Glucose batch (Heyland 2010 Table I and p.360): 0.28 for the non-producing parent (high: a producer cannot outgrow it; also Looser 2015 Table 2, glucose) and 0.15-0.16 for the producing strain (low); both papers note the better producers grow more slowly
hosts.kphaffii.temperature_C	30		B	intelligen_ovalbumin2025 p.6: 30 C
hosts.kphaffii.maint_factor	1	0.57–1.67	A	growth phase: multiplier on Tijhuis 1993 maintenance (0.0174 g glucose/g DCW/h at 30 C, Humbird A.1 method). Producing K. phaffii in glucose chemostats: 0.0147-0.0161 (Rebnegger 2016 citing Rebnegger 2014; Maurer 2006) = 0.85-0.93x. low = 0.0100 (non-producing strain, Rebnegger 2016) / 0.0174; high = 0.029 (5-copy lipase clone, Nieto-Taype 2020) / 0.0174
hosts.kphaffii.m_s_prod_g_g_h	0.00437	0.0023–0.0161	A	production phase (mu ~ 0): Rebnegger et al. 2024 Microb Cell Fact 23:44, VHH-secreting K. phaffii, glucose, 30 C, fitted over mu <= 0.05/h: 'm S (4.37 +/- 0.27 mg g-1 h-1)'. Maintenance falls at near-zero growth (Rebnegger 2016: retentostat 0.0031 vs chemostat 0.0100 for the same strain). low = 0.0023 (1-copy lipase clone, Nieto-Taype 2020); high = 0.0161 (Maurer 2006 chemostat, producing Fab strain). Not raised to Looser 2015 Table 2's 0.009-0.032 (growing cultures on methanol and glycerol; no glucose value)

Parameter	Value	Range	Gr.	Source
				or to the 5-copy lipase clone's 0.029 (Nieto-Taype 2020, chemostat; used for the growth phase, maint_factor high): maintenance falls at near-zero growth (Rebnegger 2016). The base expression phase can carry at most ~0.006 with the product made at the energetic ceiling (results constants ms_prod_max_at_ceiling; review H2). Evidence: placeholders.yaml nietotaype2020 and ms_heyland2010_looser2015 entries
hosts.kphaffii.po_ratio	1.48		B	Chung et al. 2010 Microb Cell Fact 9:50: 'P/O ratio (1.48 mol ATP/mol O)', a genome-scale-model constraint (hosts.yaml kphaffii_po_ratio_chung2010); no measured P/O found
hosts.kphaffii.kla_factor	1		B	yeast broth treated as non-coalescing water-like (Humbird Fig1.3)
hosts.kphaffii.cell_removal	centrifuge			
hosts.treesei.biomass_formula	(table in file)		B	humbird2011 p.38: T. reesei cell mass CH1.645O0.445N0.205S0.005
hosts.treesei.mu_max_per_h	0.1	0.05–0.2	E	no sourced T. reesei mu_max found (hosts.yaml gap entry); fungi grow slower than yeast
hosts.treesei.temperature_C	28		B	humbird2011 Table 19: 28 C
hosts.treesei.maint_factor	1	0.5–2	D	as for yeast (Tijhuis in both phases; no T. reesei production-phase m_s found)
hosts.treesei.m_s_prod_g_g_h	—			no evidence: the production phase uses Tijhuis x maint_factor
hosts.treesei.po_ratio	1		B	Castillo et al. 2016 Biotechnol Biofuels 9:252: 'the P/O ratio was set to 1', a metabolic-model assumption (hosts.yaml treesei_po_ratio_castillo2016); no measured value
hosts.treesei.kla_factor	0.5	0.3–0.8	D	viscous mycelial broth lowers kLa vs water-like correlations (humbird2011 used a fungal-specific correlation); evidence file
hosts.treesei.cell_removal	filter_press			
energetics.atp_per_peptide_bond	4		A	Martin et al. 2025 BBA Bioenerg 10.1016/j.bbabio.2025.149564: 'The cost assumed for protein synthesis at the ribosome is uncontested, 4 ATP per peptide bond' (hosts.yaml atp_per_peptide_bond)
energetics.atp_aa_synthesis_per_residue	0.29	0–1.2	A	Stouthamer 1973 as recomputed by Martin et al. 2025 (PMC7617979, section 1.2): synthesis of amino acids from glucose and NH4+ adds '1.4 mmol ATP for amino acid synthesis from glucose' per 4.78 mmol of amino acids, 'or average 0.28 ATP per amino acid' weighted by E. coli composition (0.29 = 1.4/4.78). high = the unweighted average over amino acids, 1.2 (same source, Table 2); low = 0 (the common amino acids yield net ATP). Membrane transport (~5 mmol ATP/g cells, Stouthamer) and secretion costs are not counted, so the ceiling stays an upper bound
energetics.sulfur_atp_per_S	3		B	textbook sulfate assimilation: ATP sulfurylase (ATP -> AMP + PPi, 2 ATP-equivalents) + APS kinase (1 ATP). Not re-read for this project. The 8 electrons of sulfate -> sulfide are charged separately at the host's P/O (4 NADH-equivalents), sulfur_electrons_per_S
energetics.sulfur_electrons_per_S	8		A	oxidation state +6 (sulfate) to -2 (cysteine/methionine sulfur)
energetics.substrate_atp_per_glucose	4		A	textbook stoichiometry: 2 ATP in glycolysis + 2 GTP/ATP from succinyl-CoA in the TCA cycle per glucose. Oxidative phosphorylation adds P/O x 12 (24 electrons = 12 O atoms = 6 O2 per glucose), so 4 + 12 x 1.48 = 21.8 ATP per glucose for K. phaffii. Not re-read for this project
fermenter.volume_m3	200	100–500	B	engineering.yaml: industrial aerobic vessels 200-1,000 m3 (Humbird, B), lysine 500 m3 (B), baker's yeast 200 m3 (B); PF designs 136 m3 (Intelligen, B), 150 m3

Parameter	Value	Range	Gr.	Source
				(Liberation Labs, C), 80-120 m3 culture volume (GFI Europe, C); Humbird STRFERM quotes to 196 m3 (larger sizes extrapolate with capital.fermenter_exponent_above_table)
fermenter.h_to_d	2		B	Humbird Equip Table10 STRFERM 200,000: 196.5 m3, D 5 m
fermenter.fill_max	0.8		B	Humbird Fig1.3!C8; NREL 2011 80%
fermenter.min_initial_fill	0.3	0.3–0.5	B	lowest start fill in a published design: Intelligen OVA starts the growth phase at 41,523 L in a 136 m3 vessel = 30.5% (p.6); NREL 2011 starts at 50% (p.37, high); Impossible US 9,938,327: 40% of the maximum working volume (= 32% of vessel). No design text states a physical minimum (impeller submergence). A batch that must start below this fill is infeasible
fermenter.pv_max_W_m3	5000	2500–5000	B	Humbird 2021 p.5 'an agitation rate up to ~5,000 W/m3' design limit in high-density fermentation (B); low = NREL 300 m3 T. reesei maximum 600 kW on 240 m3 of liquid = 2,500 W/m3 (B, derived)
fermenter.us_max_m_s	0.1	0.05–0.23	D	judgment: no design text states a superficial-velocity limit. Humbird's own 200 m3 aeration model at OUR 150 runs 0.050 m/s at the top (Fig1.3 row 197, B); NREL averages 0.033 m/s (B). high = Humbird 2021 p.5 'sparge rate up to ~2 vvm' in a 200 m3 vessel (8 m liquid height, top at 1.33 bar, 30 C) = 0.23 m/s (derived); Intelligen runs 1.6 vvm (B)
fermenter.otr_cap_mol_m3_h	200	150–250	B	Humbird 2021: 'an OTR approaching 200 mol O2/m3-h is common in an air-sparged STR fermentor' (text p.28), on his basis OUR = qO2 / cell-free liquid volume (Eq. 4.2, p.49); low = 'Bubble columns have a practical OTR limit of 150 mol O2/m3-h' (A.1, p.91); high = physical-limit scenario (head pressure, D). Per m3 of broth this is 129 at 107 g/L cells, 147 at 80 and 100 at 150 (review HM-3/BE-7: an industrial designer would use 100-120 per m3 of broth with air at ~100 g/L cells)
fermenter.otr_cap_basis	cell_free_liquid			cell_free_liquid (Humbird's basis, Eq. 4.2; base since the Phase F review) | broth (the cap applied per m3 of broth, the v2 base with 150)
fermenter.heat_per_o2_kJ	460	430–500	A	Roels 1983 via Almeida Benalcazar et al. 2025 (engineering.yaml heat_per_o2_roels_design_value); calorimetry 455 +/- 15, range 430-480 (Maskow et al. 2010, engineering.yaml oxycaloric_equivalent); Cooney/Shuler-Kargi 500 (high). Used only in the OTR_max heat check; the batch heat comes from formation enthalpies (~480-510)
fermenter.heat_removal_max_kW_m3	25	15–40	D	no production-scale ceiling is published. Evidence: 'several million kcal' per hour from 200-300 m3 SCP yeast fermenters = ~8-30 kW/m3 (US 3,986,934, B, derived); internal coils alone ~17 kW/m3 in a 600 m3 column (Almeida Benalcazar 2025, derived with its U 1.4 kW/m2/K), so external exchangers are needed; 42 kW/m3 claimed at pilot scale with tube baffles (US 4,670,397, B). 25 = 1.2 x the OTR-150 load (19.2 metabolic + ~1.2 agitation)
fermenter.turnaround_h	12	8–48	B	intelligen_ovalbumin2025 (5 fermenters, 24 h cycle, 108 h culture -> ~12 h); Humbird A.1 8 h; NREL 48 h
fermenter.feed_glucose_g_L	500	500–600	B	intelligen_ovalbumin2025 p.6: 500 g/L glucose solution (about 42 wt%; not '50 wt%', review FS-1/BE-2/HM-1). high = Shemesh 2024 K. phaffii fed-batches: 'Dextrose or glycerol as a carbon source was fed from a 60% (w/v) sterile solution' (600 g/L, A). Concentrated feeds are common in industry (Aro 2023: '55 % glucose'; Nielsen 2025 DoE up to 52 w/w%), and the titer ceiling depends on this choice (report

Parameter	Value	Range	Gr.	Source
				Table 7). Heyland 2010 fed 730 g/L (73% w/v) to a 2.6 L lab fermenter; not used as the high, which already comes from lab fed-batches (Shemesh 2024), while no production-scale design on file feeds more than 500 g/L. A 730 g/L feed would raise the physical-limit ceiling somewhat further (not computed)
fermenter.nh3_feed_g_L	268		B	intelligen_ovalbumin2025 p.6: 'aqueous ammonia (30% w/w)' at 0.895 kg/L (interpolated between 0.91 at 25% and 0.88 at 35% w/w: GESTIS via Wikipedia 'Ammonia solution') = 268 g NH3/L (review BE-2: v2 counted it at 1 kg/L)
fermenter.seed_ratio	0.1		B	NREL 2011 p.39: seed vessels 10% of the next; Intelligen 5% inoculum
fermenter.seed_time_h	24	15–36	B	intelligen 15 h per seed step; NREL 2-day seed cycle incl. cleaning
fermenter.seed_final_X_g_L	17.6		B	Intelligen lactoferrin design (Feb 2026 report, flowsheet of case a): seed broth leaves the first two seed fermenters at 17.36 and 17.77 g/L biomass (15-16 h fed-batch steps, 5% transfers); the ovalbumin report (the base process) does not state it. v2.1 used 60 g/L (D, a guess)
fermenter.DO_top_frac	0.1258		B	Humbird 2021 A.1!C92 (the dissolved-O2 setpoint of his aeration method). Humbird, Davis & McMillan 2017 p.10: in practice fed-batch aeration and agitation hold DO near zero, where transfer is most efficient; their own simulations could not set DO, and imply about three times this model's aeration energy per kg O2. Review H1 reproduced their 200 m3 stirred-tank curve with this model's equations only by holding DO near 0.5 together with their geometry, 1 atm headspace, no kLa temperature term and a 72%-efficient compressor with line losses (a reconstruction, not a published value); the base sensitivity at 0.5 alone raises aeration energy by about half
fermenter.kla_temperature_factor	1.025	1–1.025	B	Humbird 2021 Fig1.3!C11:F11: van 't Riet kLa x 1.025^(T-20). low = Humbird, Davis & McMillan 2017 Eq. 2, the same correlation (0.002, 0.7, 0.2) with no temperature term (review H1)
fermenter.compressor_formula	adiabatic			adiabatic: P = (D/3600) R T_in (r^a - 1) / (a eta), a = (k-1)/k = 0.286, r = p_btm,max / 1 atm (the textbook form; review H1) | humbird_a1: A.1!BD as written, which puts k = 1.4 where a belongs (1.6x the power at r = 2.1)
fermenter.compressor_efficiency	0.8		B	Humbird 2021 A.1!BD: the compressor power is divided by 0.8
media.salts_kg_per_m3	22.3	14.7–38	B	derived from intelligen_ovalbumin2025 Table 3: 'Basal Salts' 816,341 kg/yr over 342 batches x 107 m3 final broth = 22.3 kg/m3 (audit R6-m1). low = Table 1 basal medium 38 g/L in the 41.5 m3 initial volume per 107 m3 final (14.7); high = Table 1 applied to the final broth (38, the v1 value)
media.salts_price_usd_kg	0.2	0.15–0.4	B	intelligen_ovalbumin2025 Table 3 'Basal Salts' $0.200/kg
media.h3po4_kg_per_kg_cells	0.047		B	intelligen Eq.(1): 1.87 H3PO4 per 40 cells
media.h3po4_price_usd_kg	0.85		B	intelligen Table 3: H3PO4 (85%) $0.850/kg
media.antifoam_kg_per_m3	0.2	0.02–1	B	Behm 2022 Table 2: antifoam 0.2 g/L (T. reesei BLG); US 5,260,202 (Delta Biotechnology): 'Typical total usage rates ... 0.02-0.2 g per liter' in sensor-controlled yeast fed-batch (low); high = NREL 2011 corn oil 0.1% v/v, ~1.0 kg/m3 (P-420 13.4 kg/h over ~12.9 m3/h broth, derived)
media.antifoam_price_usd_kg	1.12	0.5–2	C	prices.yaml: Voutilainen 2021 design value EUR 1,000/t x 1.12 (no market price found); range is a guess

Parameter	Value	Range	Gr.	Source
media.methanol_kg_per_kg_product	0			methanol-free base; Intelligen uses 0.28 kg/kg (Table 3: 280 t per 1,000 t). Cost only: its O2 and heat are not added to the stoichiometry
media.methanol_price_usd_kg	0.91		B	intelligen_ovalbumin2025 Table 3: $0.910/kg
dsp.dilution_to_g_L_cells	45		B	intelligen p.7: broth diluted to 45 g/L biomass before centrifugation; the first underflow is re-diluted to 45 g/L
dsp.centrifuge_stages	2		B	intelligen p.7
dsp.paste_extracellular_frac	0.5	0.4–0.6	B	Petrides 2015 s.11.3.1 p.11-11: 'the centrifuge paste contains 40 to 60% v/v extracellular liquid' (dsp.yaml centrifuge_paste_liquid_fraction). Wet cells taken as 30% dry matter (Humbird A.1!C67)
dsp.recovery_cell_removal_override	—			calibration anchors only: fixes the cell-removal recovery instead of the paste model
dsp.centrifuge_kWh_m3	1.35		B	Humbird SI Equip!AB40 'Power demand 1.35 kWh/m3' (dsp.yaml); Westfalia HSB400 ~1.4 (Milledge & Heaven 2011, C)
dsp.mf_flux_LMH	20	10–50	D	no primary-clarification MF flux on file for yeast broth: EVERY US2025/0134130's 10 +/- 2 LMH is a final bioburden filtration (audit R6-M1, low); microalgae MF 20-100 (Slegers, D)
dsp.mf_recovery	0.975		B	Behm 2022 (T. reesei BLG design): 'microfiltration assuming 100% fungus rejection and 97.5% permeate recovery' (dsp.yaml mf_permeate_recovery_fungal_behm)
dsp.uf_flux_LMH	30	20–50	B	intelligen p.8: 30 L/m2/h concentration flux; Petrides 20-50
dsp.uf_rejection	0.995		B	intelligen p.8: rejection 0.995 for rOvalbumin
dsp.diafiltration_conc_g_L	50		B	EVERY US2025/0134130: diafiltration of rOVA at 50 g/L (dsp.yaml); the protein is diafiltered at this concentration, then concentrated further
dsp.df_flux_LMH	5	3–7	B	EVERY US2025/0134130: DF at 50 g/L '5 +/- 2 LMH' (dsp.yaml); also used for the final concentration to the dryer feed (audit R3-F2)
dsp.diafiltration_volumes	3	2–6	D	salt and sugar washout to meet the protein spec (evidence file); EVERY uses 6-8 DV after resolubilising a precipitate
dsp.dryer_feed_solids_blg	0.3	0.27–0.37	C	whey protein practice (Tetra Pak Dairy Processing Handbook): WPC80 UF retentates reach 27-32% DM and are dried at ~36% ('~36% for WPC80', ch.19); WPI and >60%-protein WPC are brought to 35-37% DM by NF/high-concentration membranes, not evaporators (ch.18). value = UF-only midpoint; high = NF. The final concentration step is costed at the DF flux
dsp.dryer_feed_solids_ova	0.23	0.2–0.3	C	egg-white practice applied to rOVA (close analogy): egg white is ultrafiltered industrially from 11.5% to 23% solids (Tsai et al. 1977 J Food Prot 40:449, 24 plant batches, A as data); UF to 20-30% DM before spray drying (EP 2 926 674 B1 citing DD 290128, B). 23% solids is ~200 g/L protein; OVA dissolves at 200 g/L in water (Wang 2025, A), so the Intelligen/vendor '50 g/L practical solubility' is not a limit (audit P-8)
dsp.dryer_feed_solids_lf	0.27	0.1–0.37	D	analogy: no lactoferrin-specific value; whey-protein UF retentates reach 27-32% DM (Tetra Pak, C), so the UF-only low end is used; high as for BLG and egg white. low = Intelligen's lactoferrin design, which dries a ~10% feed and calls 100 g/L the protein's solubility limit (pp.7, 10; no citation given)
dsp.cex_dbc_g_L	75	50–100	B	EVERY US 2024/0083940 A1 [0700]: 'The ligand density would accommodate protein binding capacity between 50-100 g protein/L resin' (dsp.yaml iex_binding_capacity_cex_every; a design statement, not a measured dynamic capacity for lactoferrin)

Parameter	Value	Range	Gr.	Source
dsp.cex_cv_per_cycle	16	16–25	B	EVERY US 2024/0083940 A1 Table 4: equilibration 4 + wash 2-3 + elution 3-4 + CIP 3 + regeneration 3 column volumes (midpoints; dsp.yaml iex_column_volumes_per_cycle_every). high = Intelligen's lactoferrin design, 10 + 5 + 5 + 5 bed volumes (p.9)
dsp.cex_elution_cv	3.5		B	same Table 4: elution 3-4 CV of '25 mM Citrate buffer with 1M NaCl'
dsp.cex_elution_nacl_kg_m3	58.4		B	1 M NaCl (EVERY Table 4) = 58.44 kg/m3; the dairy process elutes with 10-15% NaCl (Mao 2023), higher
dsp.cex_cip_cv	3		B	EVERY Table 4: CIP 3 CV of 1 M NaOH
dsp.cex_cip_naoh_kg_m3	40		B	1 M NaOH (EVERY Table 4) = 40.0 kg/m3
dsp.cex_regen_cv	3		B	EVERY Table 4: regeneration 3 CV of '20% EtOH and 150 mM NaCl', listed per cycle in 'a typical process' (often only for storage; per cycle is conservative)
dsp.cex_regen_etoh_kg_m3	157.8		B	20% v/v ethanol x 789 kg/m3 (EVERY Table 4)
dsp.cex_regen_nacl_kg_m3	8.77		B	150 mM NaCl (EVERY Table 4) = 8.77 kg/m3
dsp.cex_flow_cv_per_h	20	5–20	B	Intelligen lactoferrin design (Feb 2026 report p.10): SP Sepharose Big Beads at 500 cm/h through a 25 cm bed = 20 column volumes per hour, loading 33 min, cycle 108 min (lactoferrin.yaml lf_cost_intelligen2026_case_b_cex; review L). low = 5, the v2.1 guess for a packed bed (E). The cycle time follows: (buffer CVs + load CVs) / flow, with load CVs = capacity / feed concentration (review BE-12). Sets only the resin volume and column hardware (capital); buffer tanks are not itemised (in-line dilution from concentrates in Intelligen's design) and fall under the missing-equipment allowance
dsp.cex_recovery	0.832	0.6–0.97	B	EVERY US 2025/0134130 Ex. 5: 'Chromatography purification recovered about 83.2% POI' (dsp.yaml iex_cex_step_recovery_every). low: Helaina's whole LF train 'recovery of the purification is >60%' (Lu 2024, C); high: lactoferrin retention 97% in Intelligen's design (p.9, B)
dsp.cex_resin_usd_L	1250	100–2000	B	Petrides 2015 p.11-33: protein IEX/HIC resins '$500 to $2,000 per liter' (midpoint); low = polymeric resins 'under $100 per liter' (both dsp.yaml). Industrial lactoferrin uses agarose SP Sepharose Big Beads (Pires 2025), a protein resin
dsp.cex_resin_life_cycles	1000	125–1000	A	Pires et al. 2025 J Chromatogr A 1760:466274 (industrial lactoferrin CEX): 'After over a thousand cycles, the column must be replaced' (a lower bound; dsp.yaml iex_resin_lifetime_industrial_lactoferrin). low = Petrides' biopharma 50-200 cycles (midpoint, B)
dsp.cex_hardware_usd_per_L_resin	1000	300–3000	E	guess: columns, skid and automation per litre of packed resin; no public cost found (lactoferrin research pass). Petrides' membrane-system curve (biopharma) is the nearest analogue. Resin first fill is added to TCI unfactored
dsp.cex_diafiltration_volumes	4		D	derived: 1 M NaCl in the eluate is washed out by DF to ~2% of protein at 50 g/L after 4 volumes (58.4 x e^-4 = 1.1 g/L), meeting ash <3.5% (GRN 1284); 3 volumes would leave ~6%; Intelligen's lactoferrin design uses 5 (p.10)
dsp.cex_target_frac_dry	0.9	0.77–0.94	A	lactoferrin share of the powder: GRN 1284 (All G) batches 93.6-95.2% protein x 95.7-99.4% LF of protein (~0.90-0.94); low = GRN 1219 (TurtleTree) batches 80.5-83.5% protein x >=95% (~0.77-0.79)
dsp.nacl_usd_kg	0.184		A	prices.yaml salt_food_grade_eu_import_unit_value: EUR 0.1629/kg (2025) x ECB 1.1300
dsp.naoh_usd_kg	0.585		A	prices.yaml sodium_hydroxide_eu_import_unit_value: EUR 0.5177/kg NaOH (100% basis, 2025) x ECB 1.1300

Parameter	Value	Range	Gr.	Source
dsp.ethanol_usd_kg	0.915		A	prices.yaml ethanol_undenatured_eu_import_unit_value: EUR 0.8099/kg (2025) x ECB 1.1300
dsp.isolate_uf_end_g_L	50		B	intelligen p.8: UF to 50 g/L before precipitation (their design; not a solubility limit, see dryer_feed_solids_ova)
dsp.precip_recovery	0.891		B	intelligen p.8: 90% precipitated x 99% retained by basket centrifuges
dsp.ammonium_sulfate_kg_per_m3	350		B	intelligen p.8: 350 g/L
dsp.ammonium_sulfate_usd_kg	0.18	0.18–0.33	B	intelligen Table 3: $0.180/kg; EU import UV 2025 EUR 0.19 (A); US import UV $0.33 (A, high)
dsp.ammonium_sulfate_recycled_frac	0			Intelligen sells it on as fertiliser; base assumes no recovery credit
dsp.isolate_cake_solids	0.2		B	intelligen p.8-9: 'a soft cake with a moisture content of approximately 80%', fed to the spray dryer (dsp.yaml precipitate_cake_retention_intelligen; audit R3-F4)
dsp.isolate_wash_volumes	6		B	intelligen p.9: 'The protein cake is then washed with 6 volumes of purified water' (dsp.yaml precipitate_wash_volumes_intelligen); applied to the retentate volume
dsp.dryer_MJ_per_kg_water	4.87	3.4–6	A	dsp.yaml: Baker & McKenzie survey average 4.87 GJ/t water evaporated; milk powder 3.4 (3-stage) to 4.9 (1-stage), Ramirez et al.
dsp.dryer_fan_kWh_kg_water	0.05		A	derived from Baker & McKenzie 2005 (dsp.yaml spray_dryer_sec_survey): average 4.87 GJ/t with a fuel-to-electricity ratio of about 27 -> 0.18 GJ/t = 0.05 kWh/kg
dsp.dryer_recovery	0.98		D	cyclone + bag filter losses (evidence file)
dsp.spray_dryer_cost_2018usd	923650	361740–1377300	B	Yver et al. 2012 Int J Mol Sci 13:240 Table 3: spray dryer $897k (2011$, supplier-based SuperPro model) drying ~790 kg/h water, x CEPCI 603.1/585.7 to 2018$. low = BioSTEAM SprayDryer correlation (CEPCI 567, unsourced in its code) at 790 kg/h, x 603.1/567.3. high = Watson Dairy Consulting (C): mid-size milk dryer GBP 5-10M at 1,000 kg/h powder (~1,000 kg/h water) scaled to 790 kg/h at 0.68, x ECB 2025 1.3189 $/GBP, read as an installed package and divided by the model's TCI/PEC factor 4.6 and the 2018->2025 index 1.33 (audit R3-F5)
dsp.spray_dryer_ref_kg_h	790		B	Yver et al. 2012: 10% WPI feed, ~816 kg/h water less ~30 kg/h to the precipitate (derived)
dsp.spray_dryer_exponent	0.62	0.6–0.68	B	Yver et al. 2012 s.3.2.4: halving the feed concentration (~2.1x the water) needed a 'spray dryer (59% cost increase)' -> ln1.59/ln2.1 = 0.62 (derived). low = six-tenths rule; high = Watson Dairy Consulting bands GBP 5-10M (1,000 kg/h) vs 15-30M (5,000 kg/h) -> ln3/ln5 = 0.68 (C)
dsp.spray_dryer_max_kg_h	10000		C	largest single dryers offered: SiccaDania 200-10,000 kg/h powder, Tetra Pak 15,000, GEA 25,000 (Watson Dairy Consulting supplier table); 10,000 kg/h water is conservative
dsp.concentrate_target_frac_dry	0.85	0.7–0.92	D	target protein share of dry powder after UF/DF of a secreted protein; GRN 1056 minimum is 0.56 (0.70 x 0.80)
dsp.isolate_target_frac_dry	0.99		B	intelligen p.9: 'a purity of 99% on a dry basis' (audit R6-m2)
dsp.powder_moisture	0.05		B	intelligen p.9: 5% moisture; GRN 1056 max 7%
dsp.membrane_price_usd_m2	45	45–272	B	dairy spiral-wound elements: CAD 300 per 6.69 m2 module (Mercier-Bouchard 2017 J Dairy Sci Table 4 fn 3) = CAD 44.8/m2 / 1.2966 CAD per USD (ECB 2017) x CPI-U 2017->2025 1.3134 = $45/m2. high = Petrides 2015 'under $200/m2' for industrial membranes x CPI-U 2015->2025 1.3583. Flat-sheet PES quote $171/m2 (2023$; Chen 2023) lies between

Parameter	Value	Range	Gr.	Source
dsp.membrane_life_y	0.5	0.25–2	B	dsp.yaml: Petrides 2015, industrial membranes last >2000 operating hours (lower bound)
dsp.membrane_skid_usd_m2	263	85–526	B	an INSTALLED cost: the model divides it by the convention's installation factor to get a purchased cost (review BE-8; v2 applied installation on top). Mercier-Bouchard 2017 J Dairy Sci Table 4 fn 2: 'capital investment costs (excluding membranes) of Can$321.02/m2 for SW membranes' in a 5,185 m2 dairy MF plant (the row reads 'Installation capital investment') / 1.2966 x CEPCI 603.1/567.5. low = SuperPro help: food and water installations '$70-100/m2' of membrane (midpoint; year unknown). high = 2x, the low end of SuperPro's 2-5x multiplier for units built to biopharma specifications. Ceramic MF would be ~$3,900/m2 (same paper), not used by EVERY's process
dsp.membrane_kWh_m3_permeate	5	1–13	D	dsp.yaml: UF ~13 kWh/m3 permeate (A, upper end); typical crossflow UF lower
dsp.cip_solution_kg_per_kg	7.33		B	intelligen_ovalbumin2025 Table 3: CIP-Acid 3,430,444 + CIP-Caustic 3,901,202 kg/yr for 1,000 t/yr powder (dsp.yaml cip_chemicals_price_intelligen); counted as process water and wastewater
dsp.cip_price_usd_kg	0.023		B	intelligen_ovalbumin2025 Table 3: CIP-Acid and CIP-Caustic $0.023/kg
utilities.electricity_usd_kWh	US 0.0862; EU 0.1456	US 0.072 / EU 0.1297–US 0.1014 / EU 0.1631	A	prices.yaml: EIA industrial average 2025 US 8.62 c/kWh (Iowa 7.20 = low, Illinois 10.14 = high). EU = Eurostat nrg_pc_205 NL band IF (70-150 GWh/yr, the size of a 10 kt plant) 2025 mean EUR 0.1289/kWh excl. VAT x ECB 1.1300; EU range = band IF in BE (low, $0.1297) and DE (high, $0.1631), 2025 means
utilities.natural_gas_usd_GJ	US 4.78; EU 19.61	US 3.34 / EU 12.97–US 6.38 / EU 19.61	A	prices.yaml: EIA US industrial 2025 $5.23/Mcf / 1.036 MMBtu/Mcf (low = Henry Hub $3.52/MMBtu, high = Illinois $7.05/Mcf). EU = Eurostat nrg_pc_203 NL band I4 (0.1-1 PJ/yr) 2025 mean (S1 18.89, S2 15.81) EUR 17.35/GJ GCV excl. VAT x ECB 1.1300 (audit R5-7: NL for gas as for electricity); EU range = band I4 in BE (low, $12.97) and NL (high); DE $17.77, DK $16.81
utilities.glucose_usd_kg	US 0.4; EU 0.61	US 0.33 / EU 0.49–US 0.61 / EU 0.674	B	prices.yaml: GFI/Hawkwood 2025 Midwest DE95 $0.40/kg (B); low = Humbird contract $0.26 (2019) x CPI-U 1.2593 = $0.33; high = IMARC US dextrose Jun 2025 $611/t (C). EU (grade C, proxy; audit R5-2): EU white-sugar ex-factory average 2025 EUR 537/t x 1.13 = $0.61/kg (EC agri-food portal, A as data; sugar used as the EU sweetener price reference, not as feedstock: K. phaffii does not use sucrose). EU low = Comext glucose-syrup export UV 2025 EUR 0.4305/kg as traded x 1.13 (a lower bound on the dry basis); EU high = IMARC Germany dextrose Jun 2025 $674/t (C)
utilities.ammonia_usd_kg	0.61	0.38–0.86	B	prices.yaml: US anhydrous NH3 import unit value 2025 $0.611/kg (UN Comtrade, A proxy; EU import UV EUR 0.53 = $0.60); low = Humbird SI $0.30 (2019) x CPI-U 1.2593; high = USDA Iowa retail 2025 $784/short ton. Priced per kg NH3 contained (fed as 30% w/w)
utilities.water_usd_m3	US 1.5; EU 1.62	US 0.33 / EU 1.62–US 3 / EU 1.62	B	prices.yaml: GFI/Hawkwood 2025 Midwest $1.50/m3 (B); low = Humbird well water $0.26 (2019) x CPI-U 1.2593; high = Intelligen $3/t; EU = VEMW 2025 NL large-user average EUR 1.43/m3 (C; no EU range on file)
utilities.wastewater_usd_m3	1	0.4–4	B	dsp.yaml: $0.2-0.5/m3 below 1 g/L BOD (Petrides); Chicago $0.42/kg BOD implies ~$4/m3 at 10 g/L (A). EU priced as US: the NL levy is per kg COD and effluent COD is not modelled (audit R5-8, a limitation)

Parameter	Value	Range	Gr.	Source
utilities.steam_gas_GJ_per_t	3.29		D	physics: LP steam ~2,740 kJ/kg from feed water at 25 C (105 kJ/kg), steam tables, at an 80% boiler efficiency (judgment) = 3.29 GJ of gas per t steam, no condensate return (audit R5-6). Steam price = gas price x this
utilities.steam_usd_t	—			override only (calibration anchors); otherwise derived from the gas price
utilities.biomass_disposal_usd_t_dry	US 458; EU 211	US 0 / EU 0–US 458 / EU 686	C	spent GM biomass is today mostly sterilised and landfilled (Garnica-Garcia 2026 New Biotechnol, abstract). Fees per wet tonne are converted at 15% dry solids, the model's own centrifuge paste (30% dry wet cells x 50% extracellular liquid; review BE-9: v2 used Humbird's 20% cream yeast, i.e. $343 and $158, while no dewatering is costed). US = EREF 2024 national average MSW landfill tipping fee $62.28/short ton (Waste Advantage, C) = $458/t DM, excl. transport. EU = WRAP 2024/25 UK anaerobic-digestion gate fee median GBP 24/t wet (C) x ECB 2025 1.3189 = $211/t DM (the Kalundborg route: Novonesis/Novo Nordisk biomass to biogas). low = 0: a free outlet as feed (21 CFR 573.587 allows dried K. pastoris in broiler feed) or fertiliser; credits are possible but unquantified. high: EU WRAP range top GBP 78/t wet = $686; US: no higher source
utilities.rt_refrig	0.2629248030187212		B	Humbird RawUtil!BY4 (1.24 hp/ton, Couper)
utilities.cooling_water_dT_C	8		B	Humbird A.1!C158
utilities.building_W_m2	50		B	Humbird Equip Table9: Sanitary 50 W/m2
utilities.sterilizer_peak_factor	2		E	guess: HTST sterilizer sized for twice the average media + feed flow (sizing only; v1 hard-coded value, audit R3-F13)
utilities.boiler_peak_factor	3		E	guess: boiler sized for three times the average steam demand, SIP being intermittent (sizing only; v1 hard-coded value, audit R3-F13)
labour.operator_salary_usd	US 60430; EU 59150	US 58320 / EU 55250–US 79970 / EU 80870	A	prices.yaml: BLS OEWS May 2025 Iowa 51-9011 chemical equipment operators $60,430 (US mean $60,710; Humbird's basis was $50k in 2018); low = Iowa 51-8091 $58,320, high = US 51-8091 chemical plant operators $79,970. EU (grade B, derived): Eurostat SES 2022 NL ISCO OC8 gross earnings EUR 43,822 x NL manufacturing (C) hourly labour cost 2022->2024 (47.3/42.0) x NL industry (B-E) 2024->2025 (50.7/47.8; C not yet published) x ECB 2025 1.1300 = $59,150 (audit R5-9). EU range = the same chain on DE (EUR 40,934) and DK (EUR 59,910) OC8 earnings
labour.burden	1		B	Humbird A.1!C246: 100% labour burden
labour.shifts_per_position	4.5		B	Humbird A.1!C225
labour.employees_per_supervisor	10		B	Humbird A.1!E242
labour.supervisor_factor	1.4		B	Humbird RawUtil!BG24: supervisors paid 1.4x the operator salary
labour.attention_production	0.1		B	Humbird A.1!C231: 10% of online+offline time
labour.attention_seed	0.1		B	Humbird A.1!C230 (large seed)
labour.worker_efficiency	0.8		B	Humbird A.1!C224
labour.continuous_fte_per_shift	(table in file)		D	Humbird A.1 continuous-area FTE rules (0.25-1 per area) extended to the DSP areas; isolate_extra only for the isolate, chromatography only for the cex route
labour.fermenters_per_continuous_crew	12		E	guess: one crew for the continuous areas per ~12 production fermenters (v1 hard-coded value, audit R3-F13); Humbird's rules are per plant of 6-8 fermenters
capital.cost_index_2018_to_2025	1.33	1.229–1.524	B	prices.yaml: CEPCI 2018->2023 x1.323 (B) and 2025 = 1.016 x 2024 (publisher) -> ~x1.33-1.35; BLS PPI machinery & equipment WPU11 x1.327 (A). low = BLS PPI chemical manufacturing PCU325325 x1.229 (A), the index Humbird's Equip sheet uses (290.2 in 2018); high = PPI pumps & compressors WPU1141

Parameter	Value	Range	Gr.	Source
				x1.524 (A). Equipment-specific PPIs run higher still: heavy-gauge tanks x1.92, heat exchangers x1.64
capital.location_factor	US 1; EU 1.09	US 1 / EU 1.075–US 1 / EU 1.09	B	IEAGHG 2018/04 (Amec Foster Wheeler), Chapter E Tables 5-6: the same plants costed in the Netherlands and the US Midwest at 3Q2016 prices and 1.13 $/EUR: TPC ratio NL/US 1.087-1.090 (coal cases; value, high) and 1.075-1.082 (gas-turbine cases, whose turbine is world-priced; low). Applied to TCI. A PF plant (imported stainless equipment) may sit at the low end
capital.basis	humbird			humbird ('price-anchored capital (Humbird method)', base: Humbird's facility-class factors, 9%/yr fixed costs and capital charge; with his factors and fixed costs his baker's-yeast workbook lands inside the market-implied cost band, which the lighter NREL stack falls below) | nrel ('design-report capital (NREL-style)': NREL 2011 installation + Table 27 indirects, 3% ISBL maintenance + 0.7% FCI insurance): structural sensitivity
capital.fermenter_quote	vendor			vendor (base since the Phase F review: capital.vendor_fermenter, an industrial enzyme-plant fermenter, at the vessel-price level of the yeast anchor, whose A.1 bubble column is $0.78M at 199 m3 in 2018$, and of Intelligen's food-PF design) | acce_aseptic (Humbird 2021 STRFERM: ACCE bare vessel + agitator to the ASME BPE-2016 aseptic cell-culture specification, 316L and full vacuum, text p.37; 2.9x the vendor quote at 300 m3; structural sensitivity). Review BE-1/HM-2/SC-3
capital.seed_fermenter_quote	—			null = seed vessels priced like the production fermenters (base) | acce_aseptic: Humbird's STRFERM quotes, which cover 1-20 m3, instead of the 300 m3 vendor quote scaled down (sensitivity)
capital.vendor_fermenter	(table in file)		B	humbird2011 App. A: F-400 $3,628,673 + A-400 $5,255,007 for 9 units (2007$) = $987,076 per 300 m3 fermenter with agitator (F-400: '80,000 gal, 1 atm, 28 C, Internal coil, SS316'), x CEPCI 603.1/525.4 to 2018$ (audit R4-17/R5-5); exponent 0.7 (Area 300 fermentors). Used for production and seed fermenters in both conventions
capital.nrel_basis	(table in file)		B	humbird2011: install F-400 2.0 and A-400 1.5, cost-weighted 1.70; other equipment 1.8 (Table 25 skidded equipment). Table 27: site development 9% + piping 4.5% + other 4% of ISBL (the 4% warehouse is inside this, so no warehouse building on this basis); prorateable 10% + field 10% + home office 20% + contingency 10% + other 10% of TDC. Table 31: maintenance 3% of ISBL installed, insurance 0.7% of FCI; maintenance_frac_fci is for other design reports (Voutilainen 1.5% of FCI)
capital.fermenter_exponent_above_table	0.69	0.6–0.998	B	used only with fermenter_quote acce_aseptic. Humbird SI Equip!BM74:BO74: power fit of the STRFERM bare-vessel quotes, exponent 0.69, used beyond the largest quote (196 m3). low = six-tenths rule (D); high = the last quoted segment 100 -> 196 m3 (0.998), the v1 extrapolation (audit R3-F3)
capital.facility_class	commodity_aerobic			commodity_aerobic (food-grade, indoor sanitary) | commodity_anaerobic (outdoor, lysine-type)
capital.facility_classes	(table in file)		B	Humbird Equip Table14 (Petrides classes, as tabulated by Humbird): ISBL non-bioreactor DC x PEC, OSBL DC x PEC, E&C x TDC, fees & contingency x TPC, fermentation-area building class. Humbird tabulates the commodity-aerobic column but none of his sheets uses it (his 4FB/4P sheets use cGMP biopharm; A.1

Parameter	Value	Range	Gr.	Source
				uses a flat Lang factor of 4.5). The model applies the ISBL factor to the fermenters too, where Humbird enters bioreactors at their installed ACCE cost (Table14 bioreactor install factor 1): with the aseptic quote and the 20% allowance this gives $7.5M per 200 m3 installed (2018$), against his own installed aseptic STR at $6.8M ($2.4M bare; text Table 3.3). Review HM-5
capital.missing_frac	0.2	0.1–0.2	B	Humbird A.1 (baker's yeast): 20% missing capital on an abbreviated equipment list (TCI = Lang 4.5 x PEC / 0.8), the case closest to this model's list, which omits CIP and feed tanks per fermenter, sterile filter housings, glucose storage, biomass inactivation, powder packing and effluent pretreatment (review BE-4). low = Humbird 4FB sheets: 10% on a fuller list (the v2 value). As a fraction of the total: allowance = m/(1-m) x listed PEC
capital.packing_factor	0.15		B	Humbird 4FB-020!D232: fraction of area occupied by fermenters
capital.building_usd_m2_level_2018	(table in file)		B	Humbird Equip Table9 (ACCE 2016$ escalated to 2018)
capital.shop_area_frac	0.25		B	Humbird SI 4FB-020 row 237: 'Shop | % of ferm area | 0.25'
capital.warehouse_area_frac	0.25		B	Humbird SI 4FB-020 row 238: 'Warehouse | % of ferm area | 0.25'
capital.dsp_area_factor	0.5	0.3–1	E	guess: downstream + drying building area as a fraction of fermentation area; no source found (placeholders item 2). Check: Petrides Table 11.3 puts buildings at 0.45 x equipment purchase cost for enzyme plants (range 0.1-3.0); design.building_to_pec reports the model's ratio
capital.office_m2_per_employee	15		B	Humbird 4FB-020!D235
capital.lab_m2	400	150–600	D	analogy: Humbird 2021 SI 4FB-020 rows 233-234, a 6.8 kt/yr cultured-meat plant has 441 m2 of labs (cell lab 294 + QC lab 147). low = the QC lab alone; high = judgment
capital.compressor_building_m2_each	20		E	guess: floor area per air compressor (v1 hard-coded value, audit R3-F13)
finance.ccf_rate	0.075		B	humbird2021 p.39: i = 7.5%, n = 10 y -> CCF ~15% TCI/y
finance.ccf_life_y	10		B	humbird2021 p.39
finance.maintenance_frac_tci	0.04		B	humbird2021 p.46: 'Annual maintenance is here taken to be 4% TCI/y, which is assumed to include CIP chemicals in addition to the repair and preventative maintenance of rotating equipment, etc.'
finance.insurance_tax_frac_tci	0.05		B	humbird2021 p.46: 'Annual insurance is taken to be 5% TCI/y.' Humbird's whole fixed-cost ('overhead') rule is these two; the model has no separate plant-overhead, QC or local-tax line, so 9% stands for all of them (typical insurance alone is ~0.4-1%; review BE-3)
finance.fixed_cost_rule	—			null = follow capital.basis (humbird: 4% + 5% of TCI; nrel: 3% of installed ISBL + 0.7% of FCI); set to show the two rules on one plant (report, capital section)
finance.maintenance_includes_cip_chemicals	yes		B	humbird2021 p.46 (quote under maintenance_frac_tci): with the price-anchored convention the CIP chemicals are not charged again (review HM-8)
finance.dcf_irr	0.1		B	humbird2011 Table 33
finance.dcf_life_y	30		B	humbird2011 Table 33 (construction 8/60/32% and 7-y MACRS are fixed in finance.py with the same source)
finance.tax_rate	0.21		A	US federal corporate rate since 2018 (NREL 2011 used 35%)
finance.working_capital_frac_fci	0.05		B	humbird2011 Table 33

B Evidence base

Evidence file	A	B	C	D	E	Total
data/evidence/hosts.yaml	66	19	3	4	8	100
data/evidence/dsp.yaml	7	72	5	9	0	93
data/evidence/engineering.yaml	38	38	8	1	0	85
data/evidence/prices.yaml	66	23	15	0	0	104
data/evidence/placeholders.yaml	13	30	13	0	1	57
data/evidence/milk_egg.yaml	46	0	0	0	0	46
data/evidence/lactoferrin.yaml	22	7	6	0	0	35

Table 14: Evidence entries by grade. Each entry carries a source, a locator and, where a document was read, a verbatim quote.