



Design for CKMR



Mark Bravington, CSIRO: Aug 2022

O&A
www.csiro.au




Design in CKMR

If You...

- do not collect *enough* samples, or
- do not collect the right *types* of sample, or
- do not measure their stuff adequately


... **then CKMR will fail**



How to reap these benefits?

A page of (mostly) nice things

- Absolute **N** ; **m**; **stock structure** insights
 - **Samplees already caught for you**
 - pretty cheap (\$■ per sample... *small* ppn of catch)
 - Auto-back-dated; quick/good mixing
 - “Let the fish come to you”:
 - don’t have to sample *every* fishery, *every* location
 - Can’t cheat
 - Conditioned on *what* samples you’ve got...
 - not *why* you’ve got them, nor how many
 - Model *fish biology* not *human psychology*
 - Minimally subjective
 - “automatic” assessment; easy MSE-testing
 - Vague good news / precise bad news
 - Precision not good enough? Just \$pend a bit more
- Direct info only about adults




Design in CKMR

If You...

- do not collect *enough* samples, or
- do not collect the right *types* of sample, or
- do not measure their stuff adequately


... **then** CKMR will **fail**



“*enough samples*” really means “enough kin-pairs” ...

Can't control latter, but can plan former and types, and stuff via... **DESIGN**


At the end of the project...




At the end of the project...




At the end of the project...




At the end of the project...



At the end of the project...



At the end of the project...



DESIGN FLOWCHART



Levels of design

0. Qualitative considerations..?
1. Aim for at least XXX kin-pairs
 - by calculating $E[\text{total POPs}]$ & $E[\text{total HSPs}]$
(the laugh test)
2. Variance straight from model code
3. Inverse design
4. Optimal design
5. “Long CKMR”
6. Role of simulation

Design: qualitative sampling needs

What I reckon... for *Fish*

1. POPs & HSPs

ie ads & “juves”

how old can a “juve” be?

2. Accurate-enuf juve age

3. Ads: age or length both is best

and sex, of course

4. Full size range of ads

5. Reasonable # juve cohorts

6. Adequate spatial coverage

to allow for checks; a priori important

Design: qualitative sampling needs

What I reckon... for *Fish*

1. POPs & HSPs

ie ads & “juves”

how old can a “juve” be?

2. Accurate-enuf juve age

3. Ads: age or length both is best and sex, of course

4. Full size range of ads

5. Reasonable # juve cohorts

6. Adequate spatial coverage to allow for checks; a priori important

What I reckon... for *Mammals*

1. Are you sure?

2. Fish version is best

3. At a pinch, “juve” HSPs only or even POPs only-- less likely ?

4. Accurate-enuf juve age

5. Plenty of juve cohorts

NB eg skip-spawning

6. Adequate spatial coverage

Design: beyond the qualitative

IJAD

KISS

FFS

Design: beyond the qualitative

IJAD

It's Just A Design...

KISS

FFS

Design: beyond the qualitative

IJAD

It's Just A Design...

KISS

Keep It Simple, Stupid!

FFS

Design: beyond the qualitative

IJAD

It's Just A Design...

KISS

Keep It Simple, Stupid!

FFS

Fast and Fairly Simple

Design: beyond the qualitative

1. I would *not* approach *practical* design for “Highly Mobile Pelagic Thing” using the fully-spatial ideas of yesterday!
 - instead, much cruder box-based model (details=?)
 - make sure enuf samples to crudely estimate “rates”
2. BTW there is a bit more to the eqns/concepts for HMPT than we discussed yesterday

IJAD	It's Just A Design...
KISS	Keep It Simple, Stupid!
FFS	Fast and Fairly Simple

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
 - $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
 - $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
 - $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
 - $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$
- $\#(\text{POPs and/or HSPs}) \sim \text{Poisson}$; so need *at least* 50 for good CV

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
 - $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
 - $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$
- $\#(\text{POPs and/or HSPs}) \sim \text{Poisson}$; so need *at least* 50 for good CV
- Sample sizes: m_z with covariate z
- Group "alike" comparisons (same z , z')

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
 - $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
 - $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$
- $\#(\text{POPs and/or HSPs}) \sim \text{Poisson}$; so need *at least* 50 for good CV

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
- $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
- $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$
- $\#(\text{POPs and/or HSPs}) \sim \text{Poisson}$; so need *at least* 50 for good CV

• Sample sizes: m_z with covariate z

Wanna get 50–100

• Group "alike" comparisons (same z, z')

Each comp has prob $\sim O(1/N)$

m samples, so $\sim m^2$ comps

... so $m^2 * \text{"const"} / N = 50\text{--}100$

... so $m = \text{"const (a different one)"} * \text{sqrt}(N)$

usually 1

guess at demographic params

this kinship prob

comps of this "type"

CKMR design: the rock-bottom line

- For a precise estimate*, must find a fair number of kin-pairs
- $\mathbb{P}[K_{ij} \neq \text{UP}] \sim \bar{N}_{\text{adult}}^{-1}$
- $\hat{N}_{\text{adult}} \propto (\#\text{POPs and/or HSPs})^{-1}$
- $\#(\text{POPs and/or HSPs}) \sim \text{Poisson}$; so need *at least* 50 for good CV
- Sample sizes: m_z with covariate z
- Group "alike" comparisons (same z, z')

$$\mathbb{E}[\#\text{POPs}|\theta_0] = \sum_{z,z'} \overset{\text{usually 1}}{c_{zz'}} \times \underbrace{(m_z m_{z'})}_{\text{\# comps of this "type"}} \times \underbrace{p_{\text{POP}zz'}(\theta_0)}_{\text{this kinship prob}} \overset{\text{guess at demographic params}}{\quad}$$

CKMR design: the rock-bottom line

“If we sample X of these and Y of those--- will we get enuf kin?”

Software `microscoping::ckmr_laugh_test` for stock-assessed

It's rough--- but it's ready.

CKMR design: the rock-bottom line

“If we sample X of these and Y of those--- will we get enuf kin?”

Software `microscoping::ckmr_laugh_test` for stock-assessed

It's rough--- but it's ready.

“PASSING THE LAUGH TEST” IS NOT “DOING A DESIGN”

CKMR design: total $\text{Exp}[\text{Kin}]$ is not enough

Even if you get lots of kin-pairs, does not guarantee good CV

- depends on covars
- and measuring stuff adequately!

CKMR design: total $\text{Exp}[\text{Kin}]$ is not enough


Even if you get lots of kin-pairs, does not guarantee good CV

- depends on covars
- and measuring stuff adequately!

- To *properly* evaluate a design: choose a *goal* $g(\theta)$ e.g. trend in biomass
 - check that sample sizes $\{m_{z \in \mathcal{Z}}\}$ will give low enough $\mathbb{V} \left[g(\hat{\theta}) \right]$

"Generic" process for evaluation:

That's a **lot** of work !!



CKMR design: an easier way

- Big populations, sparse sampling: pairwise comps independent
- Each pair comp is Poisson: expected Fisher Info is

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^\top$$

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

$$\mathbb{V} \left[g(\hat{\theta}) \right] \approx \left[\frac{dg}{d\theta} \right]^\top H^{-1} \left[\frac{dg}{d\theta} \right]$$

Quadratic
dependence
on sample size

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

Big populations, sparse sampling: pairwise comps independent

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^\top$$

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

$$\mathbb{V} \left[g(\hat{\theta}) \right] \approx \left[\frac{dg}{d\theta} \right]^\top H^{-1} \left[\frac{dg}{d\theta} \right]$$

Quadratic
dependence
on sample size

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

Big populations, sparse sampling: pairwise comps independent

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^\top$$

- to code the lglik you expect to use for real but you **can** take shortcuts !

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

Quadratic
dependence
on sample size

$$\mathbb{V} \left[g(\hat{\theta}) \right] \approx \left[\frac{dg}{d\theta} \right]^\top H^{-1} \left[\frac{dg}{d\theta} \right]$$

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

Big populations, sparse sampling: pairwise comps independent

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^\top$$

- to code the lglik you expect to use for real but you **can** take shortcuts !
- prior guesstimates of pop dyn stuff eg from stock assessment

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

Quadratic
variance
on sample size

$$\mathbb{V} \left[g(\hat{\theta}) \right] \approx \left[\frac{dg}{d\theta} \right]^\top H^{-1} \left[\frac{dg}{d\theta} \right]$$

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

Big populations, sparse sampling: pairwise comps independent

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^T$$

- to code the lglik you expect to use for real but you **can** take shortcuts !
- prior guesstimates of pop dyn stuff eg from stock assessment
- some idea of what you *want* to find out!

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

Quadratic
variance
on sample size

$$\nabla [g(\hat{\theta})] \approx \left[\frac{dg}{d\theta} \right] H^{-1} \left[\frac{dg}{d\theta} \right]^T$$

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

Big populations, sparse sampling: pairwise comps independent

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^T$$

- to code the lglik you expect to use for real but you **can** take shortcuts !
- prior guesstimates of pop dyn stuff eg from stock assessment
- some idea of what you *want* to find out!
- sample sizes / sampling schemes to try out

Quadratic
variance
on sample size

Numerical derivatives are OK for p and g

NO simulation;

NO estimation;

NO repetition !

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: an easier way

What on does all *that* mean?!

- Each pair comp is Poisson: expected Fisher Info is

Well, you just need:

- to code the lglik you expect to use for real but you **can** take shortcuts !
- prior guesstimates of pop dyn stuff eg from stock assessment
- some idea of what you *want* to find out!
- sample sizes / sampling schemes to try out

Quadratic
variance
on sample size

You **don't** need to *simulate*, nor to actually *fit* your model;
there's a generic "glue framework" that does the maths

Other data: priors into H_0 ; pure multinomials (age samples)

CKMR design: NAtl makos

CKMR design: NAtl makos etc

Q: *But that's just under one scenario about "truth". Shouldn't I explore others?*

A: Arguably, that's not really necessary...

CKMR design: Inverse design

What if *no* prior stock assessment ?

eg SAtl makos; Blue-Eye Trevalla

Optimal design

- Everybody knows: "*don't trust optimal designs*"
- Not always appropriate to try

NEVERTHELESS

- it lets you know when to stop trying
- "CKMR design space" is a big place. Optimal designs are *beacons*
 - ie several optimal designs for different goals
- You learn a lot about where the "information" comes from

Optimal design for CKMR

$$H(z, z') = 4 \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right] \left[\frac{d\sqrt{p_{zz'}}}{d\theta} \right]^\top$$

$$H = H_0 + \sum_{z, z'} m_z m_{z'} H(z, z')$$

$$\mathbb{V}[g(\hat{\theta})] \approx \left[\frac{dg}{d\theta} \right]^\top H^{-1} \left[\frac{dg}{d\theta} \right]$$


$$V(m) \triangleq \mathbb{V}[\hat{g}(\theta) | m]$$

- H is sum of outer products: closed form inverse Sherman-Woodbury
- dV/dm , d^2V/dm^2 easily found by Automatic Differentiation
- Cost constraint (or target variance for minimum cost)
- Replace $V(m)$ by 2nd order approx; solve Quadratic Program; repeat
 - reliable and quick
 - NB *non-convex* QP

Optimal design: example

- "Fish-like": age structure, fecundity depends on age, survival depends on year
- Cross-cohort HSPs with 0-year-olds; POPs (all age classes)
- 10-year trend in abundance, 5 ages (5 fixed effects, 10 random effects)

Opt. for
BIOMASS




At these two optima :

3X variance (biomass)

2X variance (survival)

Opt. for
SURVIVAL



- Optimal design does depend on goal
- Runtime: ~ 5 seconds

Summary

- CKMR can handle big problems (big population sizes)
 - but you'll need a lot of samples
- *Bare minimum* due diligence: will your design give enough kin-pairs ?
 - need to compute kinship probs based on a guess at parameters
 - maybe `ckmr_laugh_test` will do...
 - or maybe you need to code up your planned `lgk`. If so, then...
- ... not much extra work to properly evaluate a design
 - just supply a goal function (in R is OK)
- & almost no extra work to consider *optimal* designs

Design in CKMR: niggling concerns laid to rest

Q: I have 1000 samples but $1000 \times 1000 / 2$ comparisons. Are they *really* all independent? Really?

A: Yeah pretty much, prolly.

At least, if yr popn is Big "sparse sampling"

Design in CKMR: niggling concerns laid to rest

Q: I have 1000 samples but $1000 \times 1000 / 2$ comparisons. Are they *really* all independent? Really?

A: Yeah pretty much, proolly.

At least, if yr popn is Big "sparse sampling"

How to approach the thought-experiment:

- think about Maternal descent Qs ie separately from P descent
- implications & chance of knowing result of "earlier" comp

Design in CKMR: niggling concerns laid to rest

Q: I have 1000 samples but $1000 \times 1000 / 2$ comparisons. Are they *really* all independent? Really?

A: Yeah pretty much, prolly.

At least, if yr popn is Big "sparse sampling"

How to approach the thought-experiment:

- think about Maternal descent Qs ie separately from P descent
- implications & chance of knowing result of "earlier" comp

Post hoc diagnostic: # triads etc.

Larval sampling? different, tricky; *not* the first resort!

“Long CKMR”

To date, fisheries interest in CKMR has come from The Desperate

- *“how on earth can we ground-truth the assessment?”*

Then... lots of work to get to that point.

But what happens next?!

“Long CKMR”

To date, fisheries interest in CKMR has come from The Desperate

- *“how on earth can we ground-truth the assessment?”*

Then... lots of work to get to that point.

But what happens next?!

CKMR is easy once you've gotten your first “N”

?Use in ongoing MPs?

Needs simulation--- can't do via nifty variance calcs

Role of simulation in CKMR

MUCH LESS THAN YOU THINK!

- You *don't* need simulation for most Designs
 - and you *shouldn't* do it because it will wa\$te time


Role of simulation in CKMR

MUCH LESS THAN YOU THINK!

- You *don't* need simulation for most Designs
 - and you *shouldn't* do it because it will waste time

If you are estimating Something, and you just want to check bias:

- ... even if you *do* have to **Simulate & Repeat**
 - ... you *don't* have to **Estimate** each time



... just **calculate** instead...

$$\left. \frac{d\Lambda(y^*)}{d\theta} \right|_{\hat{\theta}^*} = 0$$

\Rightarrow

$$\hat{\theta}^* - \theta_0 \approx - \left[\left. \frac{d^2\Lambda(y^*)}{d\theta^2} \right|_{\theta_0} \right]^{-1} \left[\left. \frac{d\Lambda(y^*)}{d\theta} \right|_{\theta_0} \right]$$

Role of simulation in CKMR

MUCH LESS THAN YOU THINK!

- You *don't* need simulation for most Designs
 - and you *shouldn't* do it because it will wa\$te time

Role of simulation in CKMR

MUCH LESS THAN YOU THINK!

- You *don't* need simulation for most Designs
 - and you *shouldn't* do it because it will wa\$te time
- For MSE, can just simulate “from” the CKMR model:
`dpois() ---> qpois() ed`
Bozo et al. *could* have done this (no IBM) to check Pyro Enfum

Role of simulation in CKMR

MUCH LESS THAN YOU THINK!

- You *don't* need simulation for most Designs
 - and you *shouldn't* do it because it will wa\$te time
- For MSE, can just simulate “from” the CKMR model:
`dpois() ---> qpois() ed`
Bozo et al. *could* have done this (no IBM) to check Pyro Enfum
- IBM sim *is* required if you plan to fit a very dicey model...
- CKMR IBMs are somewhat painful
 - tho packages can help: `fishSim`, `kinsimmer`, `CKMRsim`