

Quasi-Polynomial Growth of Betti Numbers Over Local Rings

Nicholas Packauskas

Department of Mathematics
SUNY Cortland

DG Methods in Commutative Algebra and Representation Theory
May 2nd, 2020

Set up

Throughout, (R, \mathfrak{m}, k) will denote a commutative local ring with maximal ideal \mathfrak{m} and residue field k and M will denote a finitely generated R -module.

Set up

Throughout, (R, \mathfrak{m}, k) will denote a commutative local ring with maximal ideal \mathfrak{m} and residue field k and M will denote a finitely generated R -module.

Definition

R is a **complete intersection** if its completion $\hat{R} \cong Q/I$ where (Q, \mathfrak{n}, k) is a regular local ring and $I \subseteq \mathfrak{n}^2$ is generated by a regular sequence f_1, \dots, f_c .

Set up

Throughout, (R, \mathfrak{m}, k) will denote a commutative local ring with maximal ideal \mathfrak{m} and residue field k and M will denote a finitely generated R -module.

Definition

R is a **complete intersection** if its completion $\hat{R} \cong Q/I$ where (Q, \mathfrak{n}, k) is a regular local ring and $I \subseteq \mathfrak{n}^2$ is generated by a regular sequence f_1, \dots, f_c .

- The codimension of R is $c := \dim Q - \dim R$.

Set up

Throughout, (R, \mathfrak{m}, k) will denote a commutative local ring with maximal ideal \mathfrak{m} and residue field k and M will denote a finitely generated R -module.

Definition

R is a **complete intersection** if its completion $\hat{R} \cong Q/I$ where (Q, \mathfrak{n}, k) is a regular local ring and $I \subseteq \mathfrak{n}^2$ is generated by a regular sequence f_1, \dots, f_c .

- The codimension of R is $c := \dim Q - \dim R$.
- The embedding dimension of R is $e := \dim_k \mathfrak{m}/\mathfrak{m}^2 = \dim Q$.

Set up

Throughout, (R, \mathfrak{m}, k) will denote a commutative local ring with maximal ideal \mathfrak{m} and residue field k and M will denote a finitely generated R -module.

Definition

R is a **complete intersection** if its completion $\hat{R} \cong Q/I$ where (Q, \mathfrak{n}, k) is a regular local ring and $I \subseteq \mathfrak{n}^2$ is generated by a regular sequence f_1, \dots, f_c .

- The codimension of R is $c := \dim Q - \dim R$.
- The embedding dimension of R is $e := \dim_k \mathfrak{m}/\mathfrak{m}^2 = \dim Q$.

Let M be a finitely generated R -module. We will examine the growth of the Betti numbers $\beta_i^R(M) := \dim_k \operatorname{Ext}_R^i(M, k)$ of finitely generated modules over complete intersection rings.

Background

Theorem (Tate '57)

Let R be a complete intersection ring of codimension c and embedding dimension e . Then the Poincaré series of the residue field is given by

$$P_k^R(t) := \sum_{i \geq 0} \beta_i^R(k) t^i = \frac{(1+t)^e}{(1-t^2)^c} = \frac{(1+t)^{e-c}}{(1-t)^c}$$

Background

Theorem (Tate '57)

Let R be a complete intersection ring of codimension c and embedding dimension e . Then the Poincaré series of the residue field is given by

$$P_k^R(t) := \sum_{i \geq 0} \beta_i^R(k) t^i = \frac{(1+t)^e}{(1-t^2)^c} = \frac{(1+t)^{e-c}}{(1-t)^c}$$

In particular, there exists a polynomial $b_k(n) \in \mathbb{Q}[n]$ of degree $c - 1$ such that $\beta_n^R(k) = b_k(n)$ for large values of n .

Theorem (Gulliksen '74, Avramov '89)

Let R be a complete intersection of codimension c and M be a finitely generated R -module. Then we can write

$$P_M^R(t) = \frac{p(t)}{(1 - t^2)^c}$$

Theorem (Gulliksen '74, Avramov '89)

Let R be a complete intersection of codimension c and M be a finitely generated R -module. Then we can write

$$P_M^R(t) = \frac{p(t)}{(1-t^2)^c} = \frac{h(t)}{(1-t)^d(1+t)^b}$$

for some nonnegative integer $d \leq c$. polynomial $h(t)$ with $h(1) \neq 0$ and $b < d$.

Theorem (Gulliksen '74, Avramov '89)

Let R be a complete intersection of codimension c and M be a finitely generated R -module. Then we can write

$$P_M^R(t) = \frac{p(t)}{(1-t^2)^c} = \frac{h(t)}{(1-t)^d(1+t)^b}$$

for some nonnegative integer $d \leq c$. polynomial $h(t)$ with $h(1) \neq 0$ and $b < d$.

Equivalently, there exists two integer valued polynomials b_+^M, b_-^M with $\deg b_+^M = \deg b_-^M < d$ such that for $i \gg 0$

$$\beta_i^M(t) = \begin{cases} b_+^M(i) & i \text{ is even} \\ b_-^M(i) & i \text{ is odd.} \end{cases}$$

Further, b_-^M and b_+^M have the same leading term. The order of the pole at $t = -1$ determines how many of the terms (from leading term down) b_-^M and b_+^M have in common.

For a complete intersection R , when does $b_-^M = b_+^M$?

For a complete intersection R , when does $b_-^M = b_+^M$?

- It is always the case that $b_-^k = b_+^k$.

For a complete intersection R , when does $b_-^M = b_+^M$?

- It is always the case that $b_-^k = b_+^k$.
- (Avramov '94) If R is a complete intersection of minimal multiplicity ($= 2^c$) then $b_-^M = b_+^M$ for all finitely generated modules M .

For a complete intersection R , when does $b_-^M = b_+^M$?

- It is always the case that $b_-^k = b_+^k$.
- (Avramov '94) If R is a complete intersection of minimal multiplicity ($= 2^c$) then $b_-^M = b_+^M$ for all finitely generated modules M .
- (Avramov, Seceleanu, Yang '16) Let R be a graded complete intersection of codimension c . Then $b_-^M = b_+^M$ for all finitely generated M if and only if at least $c - 1$ of the minimal generators of the defining ideal are quadrics.

Quadratic Codimension

Let (R, \mathfrak{m}, k) be a local ring, and $\hat{R} \cong Q/I$ a minimal Cohen presentation of R , so that (Q, \mathfrak{n}, k) is a regular local ring. Set I_2^\square to be the image of $I \subseteq \mathfrak{n}^2/\mathfrak{n}^3$, and let $I^\square \subseteq \text{gr}_{\mathfrak{n}} Q$ be the ideal generated by I_2^\square .

Quadratic Codimension

Let (R, \mathfrak{m}, k) be a local ring, and $\hat{R} \cong Q/I$ a minimal Cohen presentation of R , so that (Q, \mathfrak{n}, k) is a regular local ring. Set I_2^\square to be the image of $I \subseteq \mathfrak{n}^2/\mathfrak{n}^3$, and let $I^\square \subseteq \text{gr}_{\mathfrak{n}} Q$ be the ideal generated by I_2^\square .

Definition

The **quadratic codimension** q of a local ring R is the height of the ideal I^\square defined above.

Note that $q \leq c$.

Quadratic Codimension

Let (R, \mathfrak{m}, k) be a local ring, and $\hat{R} \cong Q/I$ a minimal Cohen presentation of R , so that (Q, \mathfrak{n}, k) is a regular local ring. Set I_2^\square to be the image of $I \subseteq \mathfrak{n}^2/\mathfrak{n}^3$, and let $I^\square \subseteq \text{gr}_{\mathfrak{n}} Q$ be the ideal generated by I_2^\square .

Definition

The **quadratic codimension** q of a local ring R is the height of the ideal I^\square defined above.

Note that $q \leq c$.

Example

Let $R = k[x, y]/(xy, x^2 + y^3)$. Then $I^\square = (\overline{xy}, \overline{x}^2)$ which has height one in $\text{gr}_{\mathfrak{n}} Q = k[\overline{x}, \overline{y}]$. Thus, for R , $c = 2$ and $q = 1$.

Quadratic Codimension

Let (R, \mathfrak{m}, k) be a local ring, and $\hat{R} \cong Q/I$ a minimal Cohen presentation of R , so that (Q, \mathfrak{n}, k) is a regular local ring. Set I_2^\square to be the image of $I \subseteq \mathfrak{n}^2/\mathfrak{n}^3$, and let $I^\square \subseteq \text{gr}_{\mathfrak{n}} Q$ be the ideal generated by I_2^\square .

Definition

The **quadratic codimension** q of a local ring R is the height of the ideal I^\square defined above.

Note that $q \leq c$.

Example

Let $R = k[x, y]/(xy, x^2 + y^3)$. Then $I^\square = (\overline{xy}, \overline{x^2})$ which has height one in $\text{gr}_{\mathfrak{n}} Q = k[\overline{x}, \overline{y}]$. Thus, for R , $c = 2$ and $q = 1$.

In the graded setting, the quadratic codimension of a complete intersection ring measures the number of generators in a minimal generating set of I which are quadric.

The Main Result

Theorem (Avramov, P. , Walker)

Let (R, \mathfrak{m}, k) be a complete intersection of codimension c and quadratic codimension q . Then for each finitely generated R -module M , we can write

$$P_M^R(t) = \frac{g_M(t)}{(1-t)^c(1+t)^b}$$

where $g_M(t) \in \mathbb{Z}[t]$, and $b = \max\{0, c - q - 1\}$.

This puts an upper bound on the degree of the discrepancy between b_-^M and b_+^M , that is, $\deg b_-^M - b_+^M \leq c - q - 1$.

Homotopy Lie Algebras of Local Rings

Recall that $\mathrm{Ext}_R^*(M, k)$ is a (left) module over the associative algebra $\mathrm{Ext}_R^*(k, k)$ via the composition product. We can associate a graded Lie algebra over k to R , called its homotopy Lie algebra.

Homotopy Lie Algebras of Local Rings

Recall that $\text{Ext}_R^*(M, k)$ is a (left) module over the associative algebra $\text{Ext}_R^*(k, k)$ via the composition product. We can associate a graded Lie algebra over k to R , called its homotopy Lie algebra.

Theorem (Milnor-Moore '65, André '71, Sjödin '80)

For a local ring R , there exists a graded Lie algebra $\pi^(R)$ with the following properties:*

- $\dim_k \pi^i(R) = \varepsilon_i(R)$ for each $i \in \mathbb{Z}$.
- *There is an isomorphism of associative algebras*

$$U(\pi^*(R)) \cong \text{Ext}_R^*(k, k)$$

where $U(\pi^(R))$ is the universal enveloping algebra of $\pi^*(R)$.*

Homotopy Lie Algebras over Complete Intersections

For complete intersections, $\mathrm{Ext}_R^*(M, k)$ is finitely generated over $\mathrm{Ext}_R^*(k, k)$ and the homotopy Lie algebra is completely understood:

Homotopy Lie Algebras over Complete Intersections

For complete intersections, $\text{Ext}_R^*(M, k)$ is finitely generated over $\text{Ext}_R^*(k, k)$ and the homotopy Lie algebra is completely understood:

Theorem (Sjödin '76)

Let (R, \mathfrak{m}, k) be a complete intersection of codimension c and embedding dimension e . Let $\hat{R} \cong Q/I$, with $\mathfrak{n} = (x_1, \dots, x_e)$, $I = (f_1, \dots, f_c)$ and write $f_h = \sum_{i,j} a_{ij}^h x_i x_j$ with $a_{ij}^h \in Q$.

- $\dim_k \pi^i(R) = \begin{cases} e & i = 1 \\ c & i = 2 \\ 0 & i \neq 1, 2 \end{cases}$.
- There are k bases ξ_1, \dots, ξ_e and χ_1, \dots, χ_c such that

$$[\xi_i, \xi_j] = - \sum_{h=1}^c \bar{a}_{ij}^h \chi_h, \quad \xi^{[2]} = - \sum_{h=1}^c \bar{a}_{ii}^h \chi_h$$

Where \bar{a}_{ij}^h is the image of a_{ij}^h in k .

The Special Subalgebra

The heavy lifting of the proof is done by constructing a Lie sub-algebra of the homotopy Lie algebra with nice properties.

The Special Subalgebra

The heavy lifting of the proof is done by constructing a Lie sub-algebra of the homotopy Lie algebra with nice properties.

Theorem (Avramov, P., Walker)

Let R be a complete intersection of codimension c and quadratic codimension q . Then there is a Lie sub-algebra of $\lambda^ \subseteq \pi^*(R)$ with the following properties:*

- $\dim_k \lambda^1 = q$
- $\lambda^i = \pi^i(R)$ for $i \neq 1$.
- $\text{Ext}_R^*(k, k)$ is free of finite rank as a left $U(\lambda^*)$ -module.
- $U(\lambda^*)$ has finite global dimension.

Idea of the Proof

Theorem (Avramov, P. , Walker)

Let (R, \mathfrak{m}, k) be a complete intersection of codimension c and quadratic codimension q . Then for each finitely generated R -module, we can write

$$P_M^R(t) = \frac{g_M(t)}{(1-t)^c(1+t)^b}$$

where $g_M(t) \in \mathbb{Z}[t]$, and $b = \max\{0, c - q - 1\}$

Sketch of the Proof:

Set $\mathcal{L} := U(\lambda^*)$, Then $\mathcal{M} = \text{Ext}_R^*(M, k)$ is a finite \mathcal{L} module, and there is a finite free resolution

$$0 \leftarrow \text{Ext}_R^*(M, k) \leftarrow \mathcal{F}_0 \leftarrow \cdots \leftarrow \mathcal{F}_r \leftarrow 0$$

Therefore

$$\begin{aligned}
 P_M^R(t) &= H_{\mathcal{M}}(t) = \sum_{i=0}^r (-1)^i H_{\mathcal{F}_i}(t) \\
 &= \sum_{i=0}^r (-1)^i (\text{rank } \mathcal{F}_i) t^i H_{\mathcal{L}}(t) \\
 &= h_{\mathcal{M}}(t) H_{\mathcal{L}}(t)
 \end{aligned}$$

For some polynomial $h_{\mathcal{M}}(t) \in \mathbb{Z}[t]$.

Therefore

$$\begin{aligned}
 P_M^R(t) &= H_{\mathcal{M}}(t) = \sum_{i=0}^r (-1)^i H_{\mathcal{F}_i}(t) \\
 &= \sum_{i=0}^r (-1)^i (\text{rank } \mathcal{F}_i) t^i H_{\mathcal{L}}(t) \\
 &= h_{\mathcal{M}}(t) H_{\mathcal{L}}(t)
 \end{aligned}$$

For some polynomial $h_{\mathcal{M}}(t) \in \mathbb{Z}[t]$. Now, by the graded version of the Poincaré Birkhoff-Witt Theorem,

$$H_{\mathcal{L}}(t) = \frac{(1+t)^q}{(1-t^2)^c} = \frac{1}{(1-t)^c (1+t)^{c-q}}$$

Therefore

$$\begin{aligned}
 P_M^R(t) &= H_{\mathcal{M}}(t) = \sum_{i=0}^r (-1)^i H_{\mathcal{F}_i}(t) \\
 &= \sum_{i=0}^r (-1)^i (\text{rank } \mathcal{F}_i) t^i H_{\mathcal{L}}(t) \\
 &= h_{\mathcal{M}}(t) H_{\mathcal{L}}(t)
 \end{aligned}$$

For some polynomial $h_{\mathcal{M}}(t) \in \mathbb{Z}[t]$. Now, by the graded version of the Poincaré Birkhoff-Witt Theorem,

$$H_{\mathcal{L}}(t) = \frac{(1+t)^q}{(1-t^2)^c} = \frac{1}{(1-t)^c(1+t)^{c-q}}$$

Thus

$$P_M^R(t) = \frac{h_{\mathcal{M}}(t)}{(1-t)^c(1+t)^{c-q}}$$

We have

$$P_M^R(t) = \frac{h_{\mathcal{M}}(t)}{(1-t)^c(1+t)^{c-q}}.$$

If $c = q$ was previously known (Avramov '94). If $c > q$, then one can reduce to case where R is Artinian, induce on the length of M , and use Tate's theorem for $P_k^R(t)$ to show that $h_{\mathcal{M}}(-1) = 0$. Hence, one may write

$$P_M^R(t) = \frac{g_M(t)}{(1-t)^c(1+t)^{c-q-1}}$$

with $g_M(t) \in \mathbb{Z}[t]$.

□

An Example

Example

Let $R = k[x, y, z]/(x^2, y^3, z^3)$.

An Example

Example

Let $R = k[x, y, z]/(x^2, y^3, z^3)$. Then R is a graded complete intersection of codimension 3 and quadratic codimension 1. The theorem implies that if $b_-^M \neq b_-^M$ then they can only differ in their constant term.

An Example

Example

Let $R = k[x, y, z]/(x^2, y^3, z^3)$. Then R is a graded complete intersection of codimension 3 and quadratic codimension 1. The theorem implies that if $b_-^M \neq b_-^M$ then they can only differ in their constant term. Taking $M = (x, y, z)^2$ and $k = \mathbb{Z}/101\mathbb{Z}$, one can compute

$$P_M^R(t) = \frac{1 + 3t}{(1 - t)^3(1 + t)}.$$

An Example

Example









Let $R = k[x, y, z]/(x^2, y^3, z^3)$. Then R is a graded complete intersection of codimension 3 and quadratic codimension 1. The theorem implies that if $b_-^M \neq b_-^M$ then they can only differ in their constant term. Taking $M = (x, y, z)^2$ and $k = \mathbb{Z}/101\mathbb{Z}$, one can compute

$$P_M^R(t) = \frac{1 + 3t}{(1 - t)^3(1 + t)}.$$

For $n > 3$, one has

$$\beta_n(M) = \begin{cases} n^2 + \frac{5}{2}n + \frac{3}{2} & n \text{ is even} \\ n^2 + \frac{5}{2}n + 1 & n \text{ is odd} \end{cases}.$$

Bibliography

-  L. L. Avramov, *Modules of finite virtual projective dimension*, Invent. Math. **96** (1989), 71-101
-  L. L. Avramov, *Local rings over which all modules have rational Poincaré series*, J. Pure Appl. Algebra **91** (1994). 29-48
-  L. L. Avramov, N. Packauskas, M. Walker, *Quasipolynomial growth of Betti sequences over complete intersections*, (In preparation)
-  L. L. Avramov, A. Seceleanu, Z. Yang, *Polynomial growth of Betti sequences over local rings* (In preparation)
-  T.H. Gulliksen, *A homological characterization of local complete intersections*, Compositio Math. **23** (1971), 251-255
-  G. Sjödin, *A set of generators for $\text{Ext}_R(k, k)$* , Math. Scand. **38** (1976), 1-12
-  G. Sjödin *Hopf Algebras and derivations*, J. Algebra **64** (1980), 218-229
-  J. Tate, *Homology of noetherian and local rings*, Illinois J. Math. **1** (1957), 14-25