Ex.No-1

AIM:

To calculate the values for the mathematical formulas using NumPy library

## INTEGRATED DEVELOPMENT ENVIRONMENT (IDE) REQUIRED:

JUPYTER NOTEBOOK

### **REQUIRED LIBRARIES FOR PYTHON:**

Numpy

#### **PROCEDURE:**

### A) Euclidean distance

The mathematical formula for calculating the Euclidean distance between 2 points in 2D space:

$$d(p,q) = \sqrt[2]{(q_1-p_1)^2+(q_2-p_2)^2}$$

### **B)**Dot Product

$$u = \begin{bmatrix} 5 \\ 12 \end{bmatrix}, \quad v = \begin{bmatrix} 8 \\ 6 \end{bmatrix}$$

Dot product is 
$$u \cdot v = u_1 imes v_1 + u_2 imes v_2$$
 
$$= 5 imes 8 + 12 imes 6$$
 
$$= 112$$

### C) Solving a System of Linear Equations

A system of linear equations can be represented in matrix form as AX=B, where A is the matrix of coefficients, X is the column vector of variables, and B is the column vector of solutions. To solve for **X**, we can use:  $X=A^{-1}B$  assuming A is invertible.

#### **PROGRAM:**

# A) Calculating the Euclidean Distance Between Two Points

importnumpy as np

defeuclidean distance(p, q):

```
returnp.sqrt(np.sum((q - p) ** 2))
# Example usage
p = np.array([1, 2])
q = np.array([4, 6])
distance = euclidean distance(p, q)
print("Output for Calculating the Euclidean Distance Between Two Points is: ",distance)
B)Calculating the Dot Product of Two Vectors
importnumpy as np
A = np.array([1, 3, -5])
B = np.array([4, -2, -1])
dot product = np.dot(A, B)
print("Output for dot product of two vectors A and B is ",dot product)
C) Solving a System of Linear Equations
importnumpy as np
# Coefficients matrix A and result vector b
A = np.array([[3, 1], [1, 2]])
b = np.array([9,
8]) \# Solve for x
x = np.linalg.solve(A, b)
print("Output solution of System of Linear Equations is ",x)
Output:
A)Output for Calculating the Euclidean Distance between Two Points is: 5.0. Exercise 2 –B)Output for
dot product of two vectors A and B is 3
C)Output solution of System of Linear Equations is [2. 3.]
```

#### **Result:**

The programs were run successfully