## Linear equations in two variables

Paidisetty Rithik(paidisettyrithik@sriprakashschools.com)

August 2, 2023

## $10^{th}$ Maths - Chapter 3

This is Problem-3.5 from Exercise 3.2

1. On comparing the ratios  $\frac{a1}{a2}$ ,  $\frac{b1}{b2}$  and  $\frac{c1}{c2}$ , find out whether the following pairs of linear equations are consistent, or inconsistent

$$\frac{4}{3}x + 2y = 8\tag{1}$$

$$2x + 3y = 12 \tag{2}$$

## **Solution:**

This can also be written as:

$$\begin{pmatrix} \frac{4}{3} & 2\\ 2 & 3 \end{pmatrix} \begin{pmatrix} x\\ y \end{pmatrix} = \begin{pmatrix} 8\\ 12 \end{pmatrix} \tag{3}$$

$$x = \frac{\begin{vmatrix} \mathbf{b} & \mathbf{a}_2 \end{vmatrix}}{\begin{vmatrix} \mathbf{a}_1 & \mathbf{a}_2 \end{vmatrix}} = \frac{\begin{vmatrix} 8 & 12 \\ 2 & 3 \end{vmatrix}}{\begin{vmatrix} \frac{4}{3} & 2 \\ 2 & 3 \end{vmatrix}} = \frac{(8)(3) - (2)(12)}{(\frac{4}{3})(3) - (2)(2)} = \frac{24 - 24}{4 - 4} = 0 \quad (4)$$

$$y = \frac{\begin{vmatrix} \mathbf{a}_1 & \mathbf{b} \end{vmatrix}}{\begin{vmatrix} \mathbf{a}_1 & \mathbf{a}_2 \end{vmatrix}} = \frac{\begin{vmatrix} \frac{4}{3} & 2 \\ 8 & 12 \end{vmatrix}}{\begin{vmatrix} \frac{4}{3} & 2 \\ 2 & 3 \end{vmatrix}} = \frac{\left(\frac{4}{3}\right)(12) - (8)(2)}{\left(\frac{4}{3}\right)(3) - (2)(2)} = \frac{16 - 16}{4 - 4} = 0 \quad (5)$$

(6)

Therefore the equation is consistent as  $\frac{a1}{a2} {=} \frac{b1}{b2} {=} \frac{c1}{c2}$