PakEmis: a national greenhouse gas emissions inventory for Pakistan, July 2015–June 2024

Rehan Ahmad[1]

[1]Pakistan Air Quality Initiative (PAQI), Karachi, Pakistan

Correspondence: Rehan Ahmad (rehan@pakairquality.com)

Abstract. Pakistan's greenhouse gas emissions are estimated from activity records kept by different institutions at different levels of detail, and these records cannot always be combined directly. We present PakEmis, a national inventory of CO_2, CH_4 and N_2O outside land use, land-use change and forestry, covering nine fiscal years from July 2015 to June 2024. The inventory compiles Pakistani regulatory, statistical, survey and operator records, reconciles the fuel and biomass quantities that sectors share, and applies IPCC methods with Pakistani activity statistics. Annual estimates are linked to monthly profiles and a 0.05° grid, and to provincial and district tables, with the evidence behind each allocation recorded. Under the reference assumptions, the estimated categories total $444.96\,\mathrm{Mt}\,CO_2e$ in 2023–24 using IPCC Fifth Assessment Report 100-year global warming potentials, with a conditional $95\,\%$ Monte Carlo interval of $400.30\,\mathrm{Mt}$ to $504.97\,\mathrm{Mt}\,CO_2e$, and the estimate rises by $51.12\,\mathrm{Mt}$ over the period. The reference counts the dung that household surveys record as burned; assigning to households all the dung that the agricultural account diverts from manure management gives $450.20\,\mathrm{Mt}\,CO_2e$. For public power, a different calorific basis for the same stations' coal gives $443.37\,\mathrm{Mt}\,CO_2e$, adding the K-Electric system, whose coverage by the national fuel statistic is not established, gives $448.97\,\mathrm{Mt}\,CO_2e$, and the aggregate coal account used in earlier releases gives $462.74\,\mathrm{Mt}\,CO_2e$; these alternatives differ in kind, are not combined into a range, and are reported beside the parameter interval rather than inside it. Monthly allocation is limited by the available records: $54.65\,\%$ of reference-year emissions use flat or weakly constrained profiles. Fluorinated gases, several industrial sources and industrial wastewater are not estimated. The dataset identifier and citation are given in the data availability section.

1 Introduction

Pakistan's greenhouse gas emissions are estimated from activities recorded by different institutions and at different levels of detail. Fuel accounts describe energy supply and consumption, agricultural statistics describe livestock and crop production, and household surveys provide information on domestic fuel use. These records offer a basis for a national inventory, but they cannot always be combined directly.

Four difficulties recur. A fuel account describes fuel supplied to a customer class, while an inventory needs the fuel burned by a source: the regulator's aggregate coal energy for public power, for example, implies more coal than the country imported and produced, whereas the stations' own metered tonnage does not (Sect. 4). Quantities overlap: gas that leaks from the network is also counted in delivered gas, petrol sold to the road customer class also fuels household generators, and dung removed from

manure management is burned as household fuel. Reporting periods differ: energy statistics follow the July–June fiscal year, livestock statistics and international comparators follow the calendar year, and household surveys describe a survey round. Geographical information is uneven: a fuel account constrains national combustion without saying where the fuel was burned, and monthly sales can precede consumption.

Pakistan's official reporting provides the national context. The First Biennial Update Report includes an inventory for FY2017–18, and the Biennial Transparency Report provides a category-level inventory for FY2020–21, and the associated Common Reporting Tables report calendar year 2022 (Government of Pakistan, 2022, 2024, 2026). These submissions also identify coverage gaps that affect comparisons: residential biomass, for example, is reported as not estimated in the Common Reporting Tables, so two residential totals can differ because of the fuels included even when the estimates for the fuels common to both agree. International inventories add further estimates: EDGAR covers most sectors, FAOSTAT provides agricultural emissions, and Climate TRACE combines sectoral and asset-level information (Crippa et al., 2025; FAO, 2025; Tubiello et al., 2013; Climate TRACE, 2026). These products differ in method and source boundary, and they can share activity statistics or emission factors with a national estimate, so agreement between two totals may reflect common inputs rather than independent confirmation.

PakEmis addresses these difficulties through the work it performs on the records. We compile the regulatory, statistical, survey and operator records into activity tables that retain their original period, units and source; we reconcile the quantities that sectors share so that each is used once; we estimate CO_2, CH_4 and N_2O with IPCC methods (IPCC, 2006, 2019); and we allocate the annual estimates to months, administrative areas and a 0.05° grid while recording the evidence each allocation rests on. Where the records support more than one reading of a quantity, as they do for public-power coal energy and for household dung, we report the alternatives beside the reference calculation rather than selecting one silently. Categories that cannot be estimated and emissions that cannot be placed remain identified as such.

The paper describes the source records and methods (Sect. 2), reports the inventory and its sensitivity to the readings of the evidence (Sect. 3), evaluates it against internal checks and external inventories (Sect. 4), and describes the products and their recommended use (Sect. 5).

2 Data and methods

2.1 Scope and source records

The inventory covers nine fiscal years from July 2015 to June 2024. Each fiscal year runs from July to June; 2023–24 is the reference year, for which the monthly grid and the uncertainty analysis are built. The inventory estimates CO_2, CH_4 and N_2O for agriculture, electricity generation, transport, residential fuel, industrial combustion, industrial processes and product use (IPPU), oil and gas fugitives, and waste. Land use, land-use change and forestry (LULUCF) are outside its scope. Fluorinated gases, coal-mining fugitives, industrial wastewater and several other industrial sources are not estimated; a category–gas matrix records these gaps separately from zeros and from emissions included elsewhere.

Activity comes from regulatory reports, agricultural and population statistics, household surveys, and operator records (Table 1). Energy inputs come principally from the National Electric Power Regulatory Authority (NEPRA), the Hydrocarbon Development Institute of Pakistan (HDIP), the Oil and Gas Regulatory Authority (OGRA) and the Oil Companies Advisory Council (OCAC). The Pakistan Bureau of Statistics (PBS) supplies census, household and industrial data, and the National Fertilizer Development Centre (NFDC) supplies fertiliser records. Each compiled input retains its original period, units, source edition and conversion; the full source register and the sector-specific calculations are in the technical documentation.

A fiscal-year label does not make every input a fiscal-year observation. Reference livestock numbers are the 2024 agricultural census enumeration; household fuel use is the adjacent HIES 2024–25 round; and the historical livestock series keeps its calendar-year basis (PBS, 2026a, b). Crop production and fertiliser offtake are fiscal records. Approximately $77\,\%$ of the reference agricultural estimate comes from enteric fermentation and manure management, so the historical livestock reconstruction affects the national series materially, and its annual values are not successive fiscal-year census observations.

2.2 Emission estimates and shared material flows

For a source s, gas g and year y, the basic estimate is

$$E_{s,g,y} = A_{s,y}\,\mathrm{EF}_{s,g}, \tag{1}$$

where A is activity and EF is the corresponding emission factor. Most sources use Tier 1 methods from the IPCC Guidelines with Pakistani activity statistics and relevant updates from the 2019 Refinement (IPCC, 2006, 2019). Fuel mass is converted to energy on the calorific basis its factor requires; livestock, soil and waste calculations use the animal, nitrogen or organic-carbon basis of their methods. Quantities are calculated by source and then aggregated to reporting categories.

We retain native-gas mass and apply global warming potentials (GWPs) at reporting. The headline AR5 100-year values are 1 for CO_2, 28 for biogenic CH_4, 30 for fossil CH_4 and 265 for N_2O (IPCC, 2013); AR6 equivalents are provided separately (IPCC, 2021). Biomass-combustion CO_2 is a memo item outside the total, while its CH_4 and N_2O are included, following inventory reporting conventions.

2.2.1 Public-power coal.

The reference public-power coal energy is calculated from each generating station's metered coal tonnage as reported in the regulator's station tables, converted at the IPCC 2006 net calorific value for the rank the station burns (lignite at the Thar stations, imported coal elsewhere) and summed over the enumerated stations. For the years before the per-station tables begin, the same account is reconstructed from the editions that print each station's mass or its generation and fuel intensity, with the two commissioning years carried as a bounded lane. Two other readings of the same quantity exist and are reported in Sect. 3.3: the aggregate coal energy the regulator publishes as "fuel consumed", which on the fleet's rank composition requires more coal than Pakistan's recorded imports and production supplied in 2023–24 unless stocks were drawn down (Sect. 4), and the energy implied by each station's own heat rate and generation, which is derived from generation and therefore cannot test it. Captive industrial generation is estimated separately and contributes to industrial combustion.

Table 1. Principal source records. Editions are those held and transcribed; the full per-file register with page and table citations is `outputs/pakemis_source_register.csv`. Fuel energy is obtained from mass or volume at the IPCC 2006 net calorific value for the fuel or coal rank unless the row states another basis.

Quantity	Publication and edition	Reporting period	Units as published	Conversion or use
Station fuel and generation	NEPRA State of Industry Report 2018, 2023 and 2024 (per-station tables)	2015–16 to 2023–24	tonnes, MMCFT, MMBTU; GWh	coal to TJ at the rank default; gas at the declared heating value
Coal by consumer; supply	Finance Division Economic Survey Statistical Supplement 2024–25 (Tables 14.1, 14.3); HDIP Energy Yearbook (Table 4.4)	fiscal years	kt; TOE	supply bound and consumer allocation (Sect. 4)
Gas sales by sector	OGRA State of the Regulated Petroleum Industry, 2015–16 to 2023–24 editions	fiscal years	MMCFD, MMCF	delivered gas; leakage deducted before combustion
Petroleum sales by customer class	OCAC sectoral sales, annual (2016–17 onward) and monthly (July 2021 onward)	fiscal years; months	tonnes by product	road, off-road, generator and aviation partitions; monthly profiles
Livestock	PBS 7th Agricultural Census 2024; FAOSTAT historical stocks	2024 enumeration; calendar years	head	Tier 1 factors; historical series on the calendar basis
Household fuel	PBS HIES 2018–19 and 2024–25	survey rounds	kg per household per month, design-weighted	national firewood and dung-cake consumption
Fertiliser	NFDC Fertilizer Review 2023–24 (monthly offtake)	fiscal years; months	nutrient tonnes	managed soils and urea; monthly profile
Crops and residues	Pakistan Economic Survey 2023–24 agriculture appendix; PBS crop dashboard	fiscal years	tonnes; hectares	rice area-days, residue ledger
Industrial production	PBS Monthly Minerals Production and Quantum Index of Manufacturing	fiscal years; months	tonnes; index	cement, lime, steel; monthly profiles
Railways; aviation	Pakistan Railways operator records; PCAA Major Traffic Flows	fiscal years	thousand litres; movements	diesel combustion; domestic flights
Waste	PBS population census; World Bank What a Waste 2.0; JMP sanitation shares	census years; 2018; 2023	persons; kg per person per day; shares	first-order decay; wastewater pathways

Combustion estimates combine fuel records with the factors for each use. Industrial fuels include metered gas, petroleum products and recorded coal use in brick kilns and cement production. Road and off-road estimates partition the petroleum customer classes; railway fuel comes from operator records; aviation covers domestic activity. Household electricity is excluded at the point of consumption because generation emissions are already included. IPPU covers cement, ammonia, iron and steel, and lubricant use, with process carbon separated from the fuel burned to operate the plants.

Agricultural estimates cover enteric fermentation, manure management, managed soils, rice, residue burning and urea application. Livestock populations and per-head factors determine enteric methane. Excreted nitrogen is partitioned among grazing, manure management and fuel diversion, with management losses deducted before soil application; direct and indirect soil N_2O are then calculated from the nitrogen reaching each pathway. The reference direct factor for non-rice cropland is the dry-climate value of 0.005 kg N_2O–N per kilogram of nitrogen; flooded rice and grazing use separate factors. Rice methane depends on cultivated area, cultivation duration and water management. Crop residues are divided among soil return, field burning, fuel and other destinations before emissions are assigned.

Residential fuels combine domestic sales with weighted HIES quantities. Dung connects the household account to the agricultural manure calculation, and the two records disagree: the agricultural diversion implies $58.74\,\mathrm{Mt}$ of air-dry dung cake, whereas the survey estimate of household demand is $10.93\,\mathrm{Mt}$. The reference counts household demand as household combustion in every year, with the quantity per household interpolated between the 2018–19 and 2024–25 survey rounds. This leaves approximately $47.8\,\mathrm{Mt}$ with other or unresolved destinations, to which no emission is applied. Configuration L instead assigns the whole diverted quantity to households. Balancing the agricultural diversion establishes how much dung left manure management; it does not establish how much households burned, which is why the residential estimate follows the survey and the destinations of the remainder are evaluated as scenarios with no probabilities assigned (Sect. 3.3).

Waste estimates use urban and rural population, generation rates, collection shares and treatment pathways. Solid-waste methane follows first-order decay of historical deposits with a deposit history extending to 1960, so each year's estimate depends on the deposits accumulated to that year. Open burning is applied to a share of uncollected waste, and the unburned remainder retains an unresolved fate. Domestic wastewater methane follows organic load and sanitation pathways. N_2O follows the 2019 Refinement (IPCC, 2019): nitrogen is allocated to the same sanitation pathways, reduced by the fraction each treatment type removes, and emitted at treatment plants and from effluent discharged by every pathway. Historical generation and collection parameters are held fixed and sanitation pathway shares follow the annual JMP estimates, so the waste series principally reflects demographic change, the change in sanitation and accumulated deposits.

Oil and gas fugitives are estimated by production, processing, transmission and distribution segment from throughput and infrastructure records; operating practice and coverage remain uncertain and are carried as separate configurations. Gas assigned to leakage is deducted from the delivered gas used in combustion. Other shared quantities are handled in the same way: petrol assigned to residential generators is deducted from the customer-class quantity otherwise used for road transport, dung removed from manure management enters the fuel balance, and retail diesel is partitioned between road and off-road use. These deductions prevent a quantity from being counted twice; the activity and destination assumptions behind them are evaluated in Sect. 4.

2.3 Monthly and spatial products

Monthly and spatial estimates are derived from the annual source estimates. Monthly profiles distribute emissions through the year, while spatial weights assign them to supported locations. Emissions without sufficient location information remain




Figure 1. Construction of the data products. Source records give four harmonised inputs: reconciled activity, emission factors and methods, monthly profiles by source and spatial weights by source. Annual estimates by source and gas are made from the activity and the factors, and feed both allocations. Temporal allocation gives the monthly tables. Spatial allocation gives the annual grids, the district tables and the province tables from one allocation. Mass without a supported location below the province is held in the unallocated account: it completes the grids and the district tables, and the province tables already contain it. The monthly grid for 2023–24 applies the same spatial weights to the monthly tables. The identities in the footer hold for each source, gas and period (Eq. 2); the closure arithmetic is tabulated in the technical documentation.

unallocated. For a monthly profile p and spatial weight w,

$$E_{s,g,y,m,i} = E_{s,g,y}\, p_{s,y,m}\, w_{s,y,i},$$
$$\sum_{m} p_{s,y,m} = 1, \qquad \sum_{i} w_{s,y,i} = 1, \tag{2}$$

for the streams that can be placed (Fig. 1). The spatial weight carries no month index: within a year the spatial pattern of a stream is fixed and its months differ in magnitude only, which is how the products are built. A national total reconstructed from the products therefore includes both the placed and the unallocated mass. The reference monthly grid combines timing and location; the historical grids are annual.

Temporal profiles use plant dispatch, petroleum sales, fertiliser offtake, industrial production, fire detections or process assumptions, depending on the source. Each stream-year is assigned one of four evidence classes: observed source-month activity (class TE1), reconstructed monthly administrative activity (TE2), a process or activity-shape profile (TE3), or a flat or weakly constrained profile (TE4). A flat profile assumes a constant daily rate and weights months by their length, including leap days; it expresses unresolved seasonality rather than an observation. Sales and production profiles keep their own meaning: fertiliser offtake, for example, precedes application and the soil emissions that follow.

Spatial products use a fixed 0.05° grid. Facilities are placed at supported locations; distributed sources use administrative activity or proxies such as crop, livestock, settlement and infrastructure patterns. Administrative summaries use a 152-unit district frame based on geoBoundaries 2019. Historical weights are taken from the same year where a record supports them; otherwise the weight is a timeless structural surface, the reference year's pattern rescaled, or a declared substitute, and the class is recorded for every stream-year. The maps therefore combine annual changes in the source estimates with spatial evidence of varying age.

The units are chosen for reuse. Gridded values are native-gas mass per cell and period, not flux density; conversion to kg m^{-2} s^{-1} requires the cell area and the period duration. Country fractions must be handled consistently with the allocation, and the unallocated mass must be retained when reproducing a national total. Administrative and gridded views describe the same inventory and must not be added together.

2.4 Uncertainty and evaluation

Three kinds of uncertainty are kept apart, because they answer different questions:

1. parameter uncertainty within the selected configuration — activity, emission factors and conversion factors, sampled with their stated distributions;

2. alternative interpretations of the source evidence — the public-power coal account, household dung use, sewer conditions and fugitive practice, each reported as a named configuration with no probability assigned;

3. unestimated categories and allocation limitations — recorded in the category–gas matrix and in the evidence class of every placement.

The national parameter interval describes the first of these only.

A 100 000-realization Monte Carlo calculation samples the parameters and combines them with the waste and fugitive sectors' own distributions. Shared inputs receive the same draw wherever they enter the account. Two settings test fuel-carbon factors shared only within a calculation module or across all combustion modules. National percentiles are taken after summing each realization. A common subset of realizations supports the monthly and provincial products, which remain conditional on their selected allocation. Distributions, seeds, dependence groups and the registry-based sensitivity analysis are described in the technical documentation.

Changes over time are calculated from paired realisations because errors shared between years can cancel in their difference. The reference dependence setting holds persistent parameters, such as emission factors, fixed across years while annual activity varies; a fully time-correlated setting and a time-independent setting bound the sensitivity to that choice.

Evaluation examines magnitude, timing and location. Internal tests check the shared-quantity deductions, gas aggregation and conservation across the products. External comparisons state the benchmark version, period, source boundary, gas and reporting metric; we compare native-gas mass where it is available, align GWPs for CO_2 equivalents, and keep fiscal–calendar offsets visible, since an average of adjacent calendar years only approximates a fiscal year. Shared statistics and factors are identified for each comparison.

3 Inventory results

3.1 Composition and historical change

Estimated emissions total $444.96\,\mathrm{Mt}$ CO_2e in 2023–24 under the reference configuration and AR5 GWPs. By calculation module, agriculture contributes $223.45\,\mathrm{Mt}$, the power module $45.59\,\mathrm{Mt}$, transport $41.93\,\mathrm{Mt}$ and waste $37.45\,\mathrm{Mt}$ (Fig. 2a). These modules are groups of calculations, not IPCC reporting sectors: the power module includes captive industrial generation, which IPCC reporting assigns to industrial combustion, and transport includes off-road machinery. The category–gas account under IPCC codes is given in the technical documentation.

Methane contributes $242.93\,\mathrm{Mt}$ CO_2e, $54.6\,\%$ of the total; CO_2 and N_2O contribute 160.94 and $41.08\,\mathrm{Mt}$ CO_2e. Enteric fermentation is the largest category at $151.66\,\mathrm{Mt}$, $34.1\,\%$ of the inventory, which is why livestock populations and emission factors govern both the total and its uncertainty. Reporting the same native-gas emissions with AR6 GWPs gives $438.19\,\mathrm{Mt}$ CO_2e; the $6.76\,\mathrm{Mt}$ difference is a change of metric, not of emissions.

The historical estimate rises from $393.84\,\mathrm{Mt}$ in 2015–16 to $444.96\,\mathrm{Mt}$ in 2023–24 (Fig. 2b). The trajectory is uneven: it reaches $483.45\,\mathrm{Mt}$ in 2020–21 and falls to $438.34\,\mathrm{Mt}$ in 2022–23, and the position of the peak is not robust, since 2021–22 is $483.45\,\mathrm{Mt}$, the same to two decimal places and marginally above it. Estimated emissions increase by $51.12\,\mathrm{Mt}$ CO_2e ($13.0\,\%$) over the study period. The conditional interval excludes zero under the reference and fully correlated parameter settings, but includes zero under time independence: the reference setting gives $15.94\,\mathrm{Mt}$ to $89.57\,\mathrm{Mt}$, the time-correlated setting $40.08\,\mathrm{Mt}$ to $64.01\,\mathrm{Mt}$, and the time-independent setting $-10.01\,\mathrm{Mt}$ to $114.07\,\mathrm{Mt}$ (Fig. 2c). Individual year-to-year fluctuations also depend on reconstructed activity, principally the livestock series and the industrial coal allocation.

3.2 Uncertainty and allocation evidence

The deterministic reference estimate is $444.96\,\mathrm{Mt}$ CO_2e; the Monte Carlo median is $446.52\,\mathrm{Mt}$, with a conditional $95\,\%$ interval of $400.30\,\mathrm{Mt}$ to $504.97\,\mathrm{Mt}$ (Fig. 3b), approximately $-10.4\,\%$ and $13.1\,\%$ around the median. Sharing fuel-carbon factors across modules changes the interval only slightly, to $400.27\,\mathrm{Mt}$ to $505.22\,\mathrm{Mt}$; this applies to the tested factor uncertainties and dependence structure and does not imply that dependence is unimportant for other inputs or for changes over time. In the




Figure 2. Estimated emissions under the reference configuration, in CO_2 equivalents on AR5 100-year GWPs. (a) Totals for 2023–24 by calculation module; the power module includes captive generation and transport includes off-road machinery, so the modules are not IPCC reporting sectors. (b) The nine annual estimates over the common ten-module scope, each for a fiscal year running from July to June; the vertical axis does not start at zero. A parameter interval exists for the reference year only (Fig. 3b), so none is drawn for the individual years. (c) The change from 2015–16 to 2023–24: the difference between the two deterministic estimates, with the 95 % interval of that difference (2.5th to 97.5th percentile) under the three temporal-dependence settings, from paired realizations (`pakemis_trend_uncertainty.csv`). The three rows are intervals for one change, not three trajectories.

Approach 1 category screen, which assumes independence between categories, enteric fermentation accounts for 63.0 % of variance and domestic wastewater for 13.0 %. In the released ensemble, which carries the dependence between categories, the share of the variance of the national total is 59.8 % for enteric fermentation, 14.0 % for managed soils and 9.0 % for domestic wastewater, and the upper 2.5 % of draws has nearly the same composition. These are shares for categories; the importance of individual parameters is not estimated.

The monthly product is less strongly constrained than the annual total. Observed source-month activity supports 0.94 % of reference-year emissions, reconstructed administrative activity 25.29 % and process profiles 19.12 %, while 54.65 % is flat or weakly constrained (Fig. 3a). Across the nine-year panel, observed and reconstructed monthly activity together support 18.79 % and the flat share is 60.84 %. Twelve monthly values therefore do not mean twelve observations, and monthly analysis should be confined to the streams whose class supports it.




Figure 3. Evidence behind the monthly allocation, and the national parameter interval, for 2023–24. (a) Share of reference-year emissions (CO_2e, AR5 GWPs) by temporal evidence class; the class states what supports the monthly shape of a source and is not a measure of error. (b) The deterministic reference estimate beside the conditional national $95\,\%$ Monte Carlo interval (2.5th to 97.5th percentile) under within-module parameter sharing, with its median. Sharing fuel-carbon factors across modules moves the interval from $400.30\,\mathrm{Mt}$ to $504.97\,\mathrm{Mt}$ to $400.27\,\mathrm{Mt}$ to $505.22\,\mathrm{Mt}$: $-0.03\,\mathrm{Mt}$ at the lower endpoint and $0.25\,\mathrm{Mt}$ at the upper, on a width of $105\,\mathrm{Mt}$. The interval describes parameter uncertainty in annual magnitude only.

The reference spatial calculation places 199 of 202 streams; three aviation cruise streams, approximately $0.06\,\%$ of the total, remain unallocated below their supported administrative level. This measures allocation completeness ($99.94\,\%$ of the mass is placed), not positional accuracy: most of the placed mass rides administrative shares or proxy surfaces, and by mass $62.2\,\%$ of the reference year sits on a timeless structural surface, $30.2\,\%$ on that year's observed activity and $7.5\,\%$ on a declared substitute (`pakemis_spatial_basis_shares.csv`). The reference-year grid itself is shown in Fig. 4, the evidence behind its placement in Fig. 5, and the surfaces that placed it with the concentration of the placed mass in Fig. 6.

3.3 Sensitivity to source interpretation

3.3.1 Household dung.

The reference counts the household consumption that the survey records. Assigning to households all the dung in the agricultural diversion (configuration L) changes residential emissions from $24.22\,\mathrm{Mt}$ to $29.47\,\mathrm{Mt}$ and the national total from 444.96 to $450.20\,\mathrm{Mt}$. The diverted dung that the reference leaves without a destination could also have remained in manure management, in which case the diversion and its nitrogen are revised and the national total is $449.44\,\mathrm{Mt}$, or been applied to fields, giving $448.44\,\mathrm{Mt}$; burning by users other than households gives the configuration L total. The survey quantity is itself uncertain: per household the 2024–25 round reports 1.5 times the 2018–19 rate, and holding the earlier rate or taking the later round as measured gives residential emissions of 23.85 or $24.33\,\mathrm{Mt}$.




Figure 4. Placed mass of (a) CO_2, (b) CH_4 and (c) N_2O on the 0.05° grid for 2023–24 (July 2023 to June 2024) under the reference configuration. Each cell shows the mass of the native gas emitted in it over the fiscal year, in Gg; the value is a cell total and is not divided by cell area, so it is not a flux density. The colour scales are logarithmic and separate for each gas, so equal colours on two maps are not equal masses; each scale spans the cells that together hold all but a thousandth of the gas's placed mass, and the cells below it share one pale colour. Cells outside every district carry no value and are blank, cells with exactly zero mass are grey, and a cell cut by the coast or the border holds only the mass of its in-country part. The headings give the placed mass; the national estimate also includes the mass retained at province level, which is not on the maps (Fig. 5). Boundaries are from geoBoundaries, represented year 2019 (Runfola et al., 2020). Gilgit-Baltistan and Azad Jammu and Kashmir, outlined with a dashed line, are gridded and lie outside the national statistical tables; the technical documentation defines these terms. The companion CSV files carry the plotted values.

3.3.2 Public-power coal energy and coverage.

Three kinds of alternative are carried for public power, and they are not combined into a range. The reference — station tonnage at the rank-default net calorific value — gives $40.04\,\mathrm{Mt}$ for public power. The fuel-property alternative keeps the same stations: station tonnage at the calorific value implied by each station's own heat rate and generation gives $38.45\,\mathrm{Mt}$, but that calorific value is derived from the generation it is then compared with, so it cannot test the reference. The coverage alternative adds the fuel of the K-Electric system: the reference with coverage extended to K-Electric gives $44.05\,\mathrm{Mt}$. Whether the national fuel statistic used for gas and oil already includes K-Electric is not established; the comparison of station sums with that statistic suggests it does not for gas and that it does for furnace oil in four of five years. The K-Electric fuel is therefore reported beside the reference and not added to it. The regulator's aggregate "fuel consumed" energy, which earlier releases used, gives $57.82\,\mathrm{Mt}$; it is retained as the previously published account, and it exceeds the supply bound in Sect. 4.3 unless coal stocks fell.




Figure 5. Evidence behind the spatial weights, by gas and fiscal year. Bars show the share of the national mass of (a) CO_2, (b) CH_4 and (c) N_2O in each year that was placed by each class of evidence; the denominator includes the mass retained at province level. Road transport and industrial liquid fuels are classed as a year's own observed activity in 2021–22 and 2022–23 only; in the reference year they rest on a structural surface, and in the years before 2021–22 on that same pattern rescaled to the year's national mass, so the hatched and light-blue segments of panel (a) are one road and built-up pattern under two classes. Percentages are printed where a segment is large enough to carry one; the companion table carries every value.

The corresponding national totals are 443.37 Mt for the fuel-property alternative, 448.97 Mt with K-Electric and 462.74 Mt for the previously published account, against 444.96 Mt for the reference. Agreement with an external total cannot select among them, because the readings differ in fuel properties and system coverage rather than in emissions physics.

3.3.3 Sewer conditions and fugitive practice.

Changing the sewer methane correction factor from 0.5 to 0.1 lowers domestic wastewater from 24.46 to 15.39 Mt. Oil and gas fugitive practice and coverage configurations span 9.601 Mt to 24.694 Mt. The reference, 23.499 Mt, keeps the higher published practice factors and excludes from the post-meter leakage base the gas that fertiliser plants use as chemical feedstock, because the leakage factor is defined on gas consumed in combustion; the gas the same plants burn as process fuel stays in the base. The top of the range is the configuration that includes the feedstock, as earlier releases did. None of these alternatives carries a probability, and they are not combined into one interval.




Figure 6. The surfaces that placed the reference year's mass, and the concentration of the placed mass. (a) Share of each gas's national mass in 2023–24 placed by each kind of surface, with the source and vintage of each raster surface; a register of facilities or sites that falls back to a raster where it holds no location is divided between its located share and that raster. A vintage is not a measure of spatial error. (b) Cumulative share of placed mass against the valid cells ranked from the highest mass, all valid cells included, beside the line of equal mass per cell. (c) The same curves against the number of cells on a logarithmic axis, with a marker where half of each gas's placed mass is reached. The cell counts belong to this grid, mask and allocation; a finer display grid would add no evidence about source locations.

3.3.4 Wastewater nitrous oxide method.

The reference follows the 2019 Refinement; the 2006 equation, which earlier releases used, treats the nitrogen of the whole population as effluent and removes none of it. For 2023–24 the Refinement gives 9.71 and the 2006 equation 11.45 Gg N_2O, a difference of 0.46 Mt CO_2e. On the same draws the conditional 95 % interval of the national total is 399.81 Mt to 505.41 Mt under the Refinement and 401.02 Mt to 524.35 Mt under the 2006 equation, mainly because the published range of the effluent factor ends at 0.075 in the Refinement and at 0.25 in the 2006 Guidelines. The Refinement is used because it allocates nitrogen

to the sanitation pathways already used for methane and accounts for nitrogen removed before discharge. Three of its inputs are assumed and not measured: that $1\,\%$ of sewered flow is treated, that nitrogen leaving septic tanks and latrines reaches water, and that receiving waters are not nutrient-impacted. If nitrogen from those decentralised systems goes to soil the waste estimate falls by $1.59\,\mathrm{Mt}$; if every discharge is to nutrient-impacted water it rises by $7.12\,\mathrm{Mt}$.

3.3.5 A parameter endpoint, kept separate.

Raising the wastewater effluent N_2O factor from 0.005 to 0.075 kg N_2O–N per kilogram of nitrogen, the upper end of its published range, increases the national total by approximately $35.6\,\mathrm{Mt}$. That range is already sampled in the waste uncertainty specification, so the endpoint is a stress test of a parameter the ensemble represents; it is not a structural alternative and not an additional national upper bound.

4 Technical validation

4.1 Transcription and date alignment of monthly records

Monthly reconstruction checks establish that the records were transcribed and dated correctly; they do not independently establish when fuel was burned. Across four OCAC customer classes and three fiscal years, 132 of 144 reconstructed monthly values agree with separately printed values to the tonne, the maximum difference being $0.14\,\%$, and an injected one-month displacement is detected in 141 cases. Cement dispatches and the PBS production profile differ by a mean of 0.22 percentage points of annual activity across 60 months, which supports the seasonal shape while keeping sales and production distinct.

4.2 Comparison with Pakistan's own submission and with EDGAR

The most direct comparison is with Pakistan's First Biennial Transparency Report, because its inventory year 2020–21 is a Pakistani fiscal year by its own statement and it reports on AR5 GWPs, so no calendar blend and no GWP-family conversion is needed (Fig. 7). On the 16 categories both inventories estimate on a like-for-like boundary, the totals agree to 1.04; with the other-sectors aggregate, which the two inventories bound differently, they agree to 1.00, and the ratio of the two totals excluding LULUCF is 0.993. 10 categories lie within a factor of two of the submission and six do not: open burning of waste (17.74) and oil and gas fugitives (6.91) above the band; managed soils (0.44), solid-waste disposal (0.49), lubricant use (0.43) and indirect manure N_2O (0.37) below it. Enteric fermentation is 1.19 times the submitted value while manure management from the same herd is 0.99; by species, dairy cattle, goats, sheep and camels agree to the submission's printed precision and the difference sits in buffalo and other cattle (`pakemis_enteric_btr_bridge.csv`). One convention remains unmatched: this inventory weights fossil CH_4 at AR5's 30 and biogenic CH_4 at 28, whereas the submission prints CO_2e without a fossil split, so each ratio is also published at a uniform 28, under which the fugitives ratio becomes 6.47 and the national ratio 0.990. 14 further subjects cannot be expressed as ratios because one side is not estimated, is zero, is booked elsewhere or is a wider aggregate; they are listed in Table 2.

Table 2. Subjects of the comparison with Pakistan's First Biennial Transparency Report (inventory year 2020–21) that cannot be expressed as a ratio, with the mass each side reports in Mt CO_2e (AR5). A blank cell means the account does not estimate the subject.

Subject	BTR	PakEmis	Why not a ratio
Residue burning, against the submission's cropland non-CO_2	3.16	0.58	submission carries it only inside a wider aggregate
Solid-waste fate (disposal + burning)	17.83	12.08	submission carries it only inside a wider aggregate
Other sectors	36.80	24.70	scope differs between the two accounts
Glass production	0.03	–	estimated only by the submission
Industrial wastewater	2.22	–	estimated only by the submission
Lime production	1.33	–	estimated only by the submission
Other industrial processes	1.35	–	estimated only by the submission
Other process uses of carbonates	0.27	–	estimated only by the submission
Solid-fuel fugitives	0.73	–	estimated only by the submission
Waste incineration	0.11	–	estimated only by the submission
Metal industry	0.00	0.38	reported as zero by the submission
ODS substitutes (F-gases)	0.00	–	reported as zero by the submission
Field burning of residues	–	0.58	reported under another category by the submission
Land (LULUCF)	34.82	–	outside this inventory's scope

EDGAR provides a calendar-year comparator across more sectors (Crippa et al., 2025). Electricity and residential emissions are 0.70 and 0.77 of EDGAR's calendar-2024 estimates, with the fiscal–calendar offset retained. Industrial combustion is $28.13\,\mathrm{Mt}$ CO_2e against $43.98\,\mathrm{Mt}$ in EDGAR; both accounts represent the activity, their difference changes sign in earlier years, and the two can share upstream energy statistics, so the $15.85\,\mathrm{Mt}$ difference cannot be assigned to missing emissions. The $11.54\,\mathrm{Mt}$ EDGAR estimate for fluorinated gases, by contrast, concerns a category PakEmis does not estimate. A transport ratio is withheld because road transport plus mixed off-road machinery does not match EDGAR's multimodal boundary.




Figure 7. Ratio of PakEmis estimates to Pakistan's First Biennial Transparency Report for inventory year 2020–21, a fiscal year on both sides, in CO_2 equivalents on AR5 100-year GWPs. Each point is the PakEmis estimate divided by the submitted value for a category both inventories estimate on a like-for-like boundary. Three totals are set apart below: the sum of the plotted categories, the same sum with the other-sectors aggregate that the two inventories bound differently, and the national totals excluding land use. The band from half to double is for orientation and is not an acceptance threshold. The inventories are not matched on the weighting of fossil CH_4, which PakEmis weights at 30: an open marker and a value in brackets give the ratio recomputed at a uniform 28 wherever the printed ratio changes. Because the two inventories can draw on the same activity statistics and default factors, agreement is not independent validation. Subjects that cannot be expressed as a ratio are listed in Table 2.

Railways provide a matched-year comparison with the earlier submission. The First Biennial Update Report gives $4.06\,\mathrm{Mt}\ CO_2e$ for FY2017–18 (Government of Pakistan, 2022); the PakEmis method applied to the operator's $156\,396$ thousand litres of diesel for that year gives $0.4664\,\mathrm{Mt}\ CO_2e$, a factor of 8.7 lower, and 2023–24 consumption of $132\,588$ thousand litres gives $0.394\,\mathrm{Mt}$. The $18\,\%$ change in reported consumption between the two years cannot explain the discrepancy, and the Common Reporting Tables' $56\,609\,\mathrm{TJ}$ of railway activity for calendar 2022 points to the original activity and factor cells as the place to look (Government of Pakistan, 2026).

4.3 The coal supply bound

Under the stated coal-rank and calorific assumptions, the aggregate energy estimate requires more coal than reported annual imports and production. Expressed as coal at the generating fleet's own rank composition, the regulator's aggregate energy for 2023–24 requires $29.53\,\mathrm{Mt}$ of coal against the $24.72\,\mathrm{Mt}$ the national accounts report as imports plus production, $119.5\,\%$ of the supply before any coal reaches a cement plant, a brick kiln or a household. Five tests ask whether the records could reverse this: a supply figure that understates the same twelve months, an inventory drawdown, the assumed lignite share, the coverage of the consumer allocation, and the calorific basis. Each is reported with the change in the records it would require, in the technical supplement. The supply figure has no stock term and no record of coal stocks is held, so a drawdown cannot be excluded; it would have to be $19\,\%$ of the year's supply for public power alone and $42\,\%$ with the other consumers. The aggregate account is therefore implausible rather than impossible, and under the stated assumptions none of the five tests reverses the bound at a size the records support. The station account used for the reference requires $15.88\,\mathrm{Mt}$ in the same year, $64\,\%$ of the supply and below the $19.20\,\mathrm{Mt}$ the energy yearbook assigns to electricity generation. The aggregate account also implies fleet efficiencies of $57.5\,\%$ in 2020–21 and $18.9\,\%$ in 2023–24, against $32.4\,\%$ to $45.7\,\%$ under the generation-implied (heat-rate) calorific basis. The supply bound and the efficiency band are the grounds for preferring the station account; the calorific basis on which the two station readings still differ is treated in the technical documentation. Close offsets between the power and industrial coal allocations indicate dependence within the national coal balance, not independent confirmation of changes in industrial fuel use.

4.4 Attribution of the managed-soils difference

Managed soils contribute $28.53\,\mathrm{Mt}$ CO_2e in PakEmis and $58.50\,\mathrm{Mt}$ in FAOSTAT, which reports calendar 2023 against our 2023–24 (FAO, 2025). Synthetic-fertiliser nitrogen inputs agree within $0.9\,\%$, as do the combined counts of six common livestock species, so the input scales are similar and the difference lies in the nitrogen pathways and their factors. Pasture deposition contributes $7.71\,\mathrm{Mt}$ here against $27.29\,\mathrm{Mt}$ in FAOSTAT, $65.3\,\%$ of the overall difference. Replacing the reference dry-climate EF_1 of 0.005 with the 2006 aggregate value of 0.010 raises managed soils to $40.43\,\mathrm{Mt}$, and applying the earlier pasture-deposition factors as well raises it to $54.72\,\mathrm{Mt}$ (Fig. 8); together the two substitutions account for $26.19\,\mathrm{Mt}$, approximately $87\,\%$ of the $29.97\,\mathrm{Mt}$ difference, and the remaining $3.78\,\mathrm{Mt}$ lies in the indirect factors and nitrogen partitioning. The substitutions apply to distinct nitrogen pathways and are alternative factor choices, not calibration steps: agreement with FAOSTAT does not identify the factors appropriate to Pakistan, and evidence for a dry-climate factor on irrigated cropland does not validate the corresponding treatment of grazing land (IPCC, 2019).

4.5 Facility locations and historical allocation

An audit of 77 located power stations identifies seven provincial assignments inconsistent with curated or source records, all within $3.4\,\mathrm{km}$ of a district boundary and representing $1.5\,\%$ to $10.3\,\%$ of plant-ledger generation across the audited years; four also disagree with simplified district polygons, so a finer raster alone would not resolve them. The district allocation takes




Figure 8. Managed-soils N_2O in CO_2 equivalents (AR5 100-year GWP) under the reference factors and three alternative factor choices, beside FAOSTAT. The reference uses the 2019 direct soil factor for dry climates and the 2019 pasture-deposition factors. The alternatives apply the 2006 direct factor to synthetic fertiliser, organic manure and crop residues; the 2006 pasture-deposition factors; and both. The two substitutions act on distinct nitrogen pathways, and indirect pathways are unchanged. PakEmis describes fiscal 2023–24 and FAOSTAT calendar 2023. The cases are alternative factor choices and are not calibration steps.

the curated record where one exists, placing a station whose grid cell lies outside its province in the cells of the district its record names, intersected with that province. Across the nine periods this moves between 36 and 109 districts' allocations in a period and $2.4\,\mathrm{Mt}$ to $20.2\,\mathrm{Mt}$ of gross mass in CO_2e (AR5), with a net change of order 1×10^{-8} Mt: it is a reallocation within provinces, and no provincial or national total changes.

Each grid fuel is split between provinces by its own stations in each period. The shares come from the station fuel record joined to the curated locations: coal by station tonnage at the calorific value the national coal account books it at; gas, furnace oil and diesel by the fuel quantity the regulator prints for each station; and bagasse by generation. From 2019–20 the record is the plant fuel ledger. For the four earlier periods it is the same columns of the regulator's 2020 report, and where that report prints no quantity for a station its generation and its own specific consumption stand in, for at most $20\,\%$ of a period's gas energy. The split supplies proportions only, so the national total of every fuel and period is unchanged. Releases before this one applied the reference year's vector to all nine periods. The largest correction is $11.6\,\mathrm{Mt}\,CO_2e$ of coal emissions moved out of one province in a single period (Sindh, 2020–21), and the reference year moved by at most $0.17\,\mathrm{Mt}\,CO_2e$ when coal took the booking's calorific basis; `pakemis_power_provincial_basis_by_year.csv` gives every fuel, period and province before and after. Two fuel-periods keep the reference vector, diesel in 2018–19 and 2019–20, because the station record covers less than half of the national diesel control there; the split is wholly Punjab on either reading. K-Electric's stations are outside

Table 3. Core product coverage. Annual and monthly availability refer to the reference configuration; alternative configurations carry their own product identifiers.

Product	Period and resolution	Main qualification
National tables	Nine fiscal years, annual and monthly	Unestimated categories and the timing evidence class are explicit.
Administrative tables	Annual province and district estimates	Fixed boundary frame; historical allocations partly reconstructed.
Annual grid	Nine fiscal years, 0.05°	Native-gas mass with allocation provenance and an unallocated companion.
Monthly grid	2023–24, 0.05°	Combines source timing and location assumptions.
Uncertainty and alternatives	Reference ensemble; historical changes; named configurations	Parameter intervals and structural alternatives answer different questions.

the split, as they are outside the national fuel control it allocates. No station carries coal outside its commissioning window (`pakemis_power_station_status.csv`). The provincial comparison of SO_2 against EDGAR is given in Appendix A.

5 Data products and recommended use

Users receive five families of products (Table 3): national annual and monthly tables by IPCC category and gas, provincial and district tables, an annual 0.05° grid for each of the nine years, a monthly grid for the reference year, and the uncertainty ensemble with the named alternative configurations. Every table carries the category, gas, period, configuration and notation status, and every placement carries its evidence class, so a user can select the detail a question supports.

How the products should be read follows from how they were built. National annual estimates are the best supported. Monthly values in a category governed by a flat profile express the annual estimate spread by days, not observed seasonality. Historical grids using a fixed or borrowed spatial pattern represent changes in the annual source estimate under that pattern. They should not be interpreted as observations of changing local activity. Near administrative boundaries, a station's provincial assignment can shift district totals (Sect. 4.5). A grid-based analysis should retain the unallocated term and use the stated cell-mass convention. A comparison with another inventory should first match the emitting sources, the period and the gas metric.

Four source questions remain open, each with the size of what it would move. Managed soils is the largest: the reference estimates $28.53\,\mathrm{Mt\,CO_2e}$ for 2023–24, two factor substitutions inside the same method give 40.43 and $54.72\,\mathrm{Mt}$, and FAOSTAT reports $58.50\,\mathrm{Mt}$. The livestock bridge is next: enteric fermentation is $63.0\,\%$ of the Approach 1 variance screen, and against the submission for FY2020–21 it is 1.19 times the submitted value while manure management from the same herd is 0.99 times, with the difference confined to buffalo and other cattle. Crop residues are 0.18 of the submission's cropland non-CO_2 line, which is a wider aggregate than the inventory's own row. Solid waste is a specification question: disposal is

0.49 of the submitted figure and the broader fate account 0.68. None of the four is settled by evidence this release holds, and none receives a provisional correction; each needs activity or factor evidence that we have not acquired or reconciled (`data/shared/requests/data_gaps.csv`).

Reproducibility has two levels. The calculation can be rebuilt from the compiled activity tables with the released software, and the test suite checks the identities between the products. Re-deriving a compiled table from its primary document requires the document. Every document a compiled table cites is listed with the file that holds it and its checksum: 153 of the 217 registered data files cite only documents that are held, 52 are derived inside the repository, and 12 cite at least one document that is not held as cited. Those are a priced energy yearbook, one year-book edition its publisher no longer lists, one superseded edition of a facility tracker, the district tables of the 2017 census, the satellite fire responses behind the burning proxy, and web pages that could not be captured with their content; a held reprint of the same figures is recorded beside each where one exists. A document fetched after its table was compiled is marked as a later copy and compared with the table. Of the eight validation products that are run by hand, 6 can be re-derived and compared without being overwritten, and the other two are listed with the reason. File checksums establish which release a number came from; they do not show that a source transcription was correct.

6 Conclusions

PakEmis combines publicly available activity statistics with IPCC methods and reconciles fuel and biomass quantities shared between sectors. It links annual estimates to monthly and spatial products while recording the evidence supporting their allocation. The reference-year total is 444.96 Mt CO_2e on the AR5 basis, with a conditional parameter interval of 400.30 Mt to 504.97 Mt; the estimated increase of 51.12 Mt over the study period excludes zero under the reference and time-correlated dependence settings and includes it under time independence (-10.01 Mt to 114.07 Mt). Methane is 54.6 % of the total and enteric fermentation a third of it, so the livestock series and its factors govern both the level and its uncertainty.

Three limitations matter most for interpretation. The public-power coal account rests on the stations' metered tonnage because the regulator's aggregate energy exceeds recorded coal supply unless stocks fell; the calorific basis of that tonnage is still an open choice worth 1.6 Mt. The household dung quantity follows the household survey, which leaves most of the dung diverted from manure management without an observed destination; assigning all of it to households raises the national total by 5.2 Mt, and the two survey rounds differ from each other. The monthly and spatial products carry their annual estimates on profiles and patterns that are, for most of the mass, not observations of that year, and the evidence class attached to each placement is the guide to what they support.

7 Data availability

The dataset is PakEmis version 1.0.0rc67 (the release candidate this manuscript describes). The released products, their checksums (`outputs/SHA256SUMS`) and the release manifest naming the repository commit are archived with the software; the dataset DOI, the review link and the version-specific citation are pending registration of the deposit and will be inserted here.

8 Code availability

The source code is at https://github.com/RehanAhmad13/pakistan-emissions-sectors; the commit the release describes is recorded in `outputs/release_manifest.json`. Reproduction is documented in `docs/reproducibility.md`: the inventory rebuilds with `python3 -m pkinv.run_all` and the test suite runs with `python3 -m unittest discover -s tests`. The software archive identifier is pending with the dataset DOI.

Author contributions. RA designed the inventory, compiled the source data, developed the calculation software and tests, produced the data records and wrote the paper.

Competing interests. The author declares no competing interests.

Release provenance. This report documents one release of PakEmis and is generated against it. Where a sector report and the release's generated product catalogue (`outputs/pakemis_product_catalog.csv`) disagree, the catalogue is authoritative: sector reports are revised on their own cadence and may lag the products.

Release version	`1.0.0rc67`
Commit	`1876f398f85c`
Reporting period	FY2023-24 (2023-07-01 to 2024-06-30)
Baseline configuration	S (survey basis)
National total	444.96 Mt CO_2e (AR5) under S, the reference; 450.20 Mt under L
Release manifest	`13462fb4c2d1ea62` (SHA-256 of `outputs/release_manifest.json`, first 16)
Source manifest	`c1be6d314cfce1ea` (SHA-256 of `data/manifests/source_manifest.csv`, first 16)
Data-description article	`paper_national/` (ESSD, in preparation), same release

Appendix A: Provincial sulphur dioxide against EDGAR

Provincial SO_2 reveals differences that national aggregation conceals. EDGAR v8.1 emissions are integrated into common administrative boundaries using fractional grid-cell areas, with adjacent calendar years blended for the fiscal comparison (Crippa et al., 2026). For FY2021–22 the national PakEmis-to-EDGAR ratio is 0.92, but the provincial ratios are 1.25 in Punjab, 1.67 in Khyber Pakhtunkhwa, 2.05 in Balochistan and 0.18 in Sindh (`pakemis_air_provincial_external_comparison.csv`); opposing provincial differences nearly cancel nationally. Neither inventory supplies observed provincial emissions against which these ratios could be read as accuracy scores.

Electricity generation is the principal source of the disagreement: the PakEmis power layer gives $167.0\,\mathrm{Gg}$ against EDGAR's $409.3\,\mathrm{Gg}$, including 4.5 against $196.9\,\mathrm{Gg}$ in Sindh. The PakEmis layer includes oil and diesel; coal SO_2 is a separate screening calculation because desulphurisation performance is unresolved. Adding the coal-screening estimates gives $202.0\,\mathrm{Gg}$ under the controlled case and $362.2\,\mathrm{Gg}$ under the uncontrolled case, so EDGAR lies above the screening range, exceeding its uncon-

trolled endpoint by $47.1\ \mathrm{Gg}$ ($13\,\%$). This identifies a consequential coverage and control assumption without establishing that plants operate without effective desulphurisation.

References

Climate TRACE: Climate TRACE Emissions Inventory and Data Downloads, country- and source-level records accessed through API v7 on 6 August 2026, https://climatetrace.org/data, 2026.

Crippa, M., Guizzardi, D., Pagani, F., Banja, M., Muntean, M., Schaaf, E., Quadrelli, R., Risquez Martin, A., Taghavi-Moharamli, P., Köykkä, J., Grassi, G., Rossi, S., Melo, J., Jacome Felix Oom, D., Branco, A., Suárez-Moreno, M., Sedano, F., San-Miguel, J., Manca, G., Pisoni, E., and Pekar, F.: GHG emissions of all world countries—2025 report, JRC143227, Publications Office of the European Union, Luxembourg, https://doi.org/10.2760/9816914, 2025.

Crippa, M., Guizzardi, D., Pagani, F., Banja, M., Muntean, M., Schaaf, E., Becker, W., Monforti Ferrario, F., Quadrelli, R., Risquez Martin, A., Grassi, G., Rossi, S., Brandao De Melo, J., Jacome Felix Oom, D., Branco, A., San-Miguel, J., and Vignati, E.: EDGAR v8.1 Global Air Pollutant Emissions, European Commission, Joint Research Centre [data set], https://doi.org/10.2905/JRC.E1ZGQER, 2026.

FAO: FAOSTAT: emissions totals, crops and livestock, fertilisers by nutrient, and forestry production and trade, Food and Agriculture Organization of the United Nations, Rome [data sets], https://www.fao.org/faostat/, 2025 snapshot used in the source inventory; catalogue checked 30 September 2026, 2025.

Government of Pakistan: Pakistan's First Biennial Update Report to the United Nations Framework Convention on Climate Change, Ministry of Climate Change, Islamabad, https://unfccc.int/documents/470405, 2022.

Government of Pakistan: Pakistan's First Biennial Transparency Report to the UNFCCC, Ministry of Climate Change and Environmental Coordination, https://unfccc.int/documents/645241, report designated 2024; UNFCCC landing page published 16 June 2025, 2024.

Government of Pakistan: Pakistan: 2024 Common Reporting Table (CRT), UNFCCC submission, https://unfccc.int/documents/655653, submitted 6 February 2026, 2026.

IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, edited by: Eggleston, H. S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K., Institute for Global Environmental Strategies, Hayama, Japan, https://www.ipcc-nggip.iges.or.jp/public/2006gl/, 2006.

IPCC: Climate Change 2013: The Physical Science Basis, Cambridge University Press, Cambridge, https://doi.org/10.1017/CBO9781107415324, 2013.

IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, edited by: Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P., and Federici, S., IPCC, Switzerland, https://www.ipcc-nggip.iges.or.jp/public/2019rf/, 2019.

IPCC: Climate Change 2021: The Physical Science Basis, Cambridge University Press, Cambridge, https://doi.org/10.1017/9781009157896, 2021.

PBS: Agricultural Census 2024, Pakistan Bureau of Statistics, Islamabad, report collection, https://www.pbs.gov.pk/agriculture-census/, accessed 30 September 2026, 2026a.

PBS: Household Integrated Economic Survey (HIES) 2024–25, Pakistan Bureau of Statistics, Islamabad, https://www.pbs.gov.pk/household-integrated-economic-survey-hies-2024-25/, accessed 30 September 2026, 2026b.

Runfola, D., Anderson, A., Baier, H., Crittenden, M., Dowker, E., Fuhrig, S., Goodman, S., Grimsley, G., Layko, R., Melville, G., Mulder, M., Oberman, R., Panganiban, J., Peck, A., Seitz, L., Shea, S., Slevin, H., Youngerman, R., and Hobbs, L.: geoBoundaries: a global database of political administrative boundaries, PLOS ONE, 15, e0231866, https://doi.org/10.1371/journal.pone.0231866, 2020.

Tubiello, F. N., Salvatore, M., Rossi, S., Ferrara, A., Fitton, N., and Smith, P.: The FAOSTAT database of greenhouse gas emissions from agriculture, Environmental Research Letters, 8, 015009, https://doi.org/10.1088/1748-9326/8/1/015009, 2013.