


Generalised Additive Models

Francisco Rodríguez-Sánchez


<https://frodriguezsanchez.net>

How do we model these time series?



Simpson 2018

How do we model these time series?



Simpson 2018

Generalised Linear Model (GLM):

$$y = a + bx$$

Generalised Additive Model (GAM):

$$y = a + s(x)$$

Modelling non-linear time series with GAM

```
isotopes <- readRDS('data/isotope.rds')
```

	Depth	d13C	TotalC	d15N	TotalN	DryWeight	Year
1	0.2	-27.57	806.49	3.05	64.21	8.2	2007.254
2	0.4	-27.67	949.33	3.01	73.26	7.6	2006.510
3	0.8	-27.63	1305.52	2.93	93.25	11.6	2004.941
4	1.2	-27.62	1136.04	2.33	86.09	9.6	2003.269
5	1.6	-27.48	1028.27	2.09	93.80	10.9	2001.496
6	2.0	-27.39	809.91	2.66	79.98	9.9	1999.626

Modelling non-linear time series with GAM

```
library('mgcv')  
m <- gam(d15N ~ s(Year, k = 15), data = isotopes, method = 'REML')
```

Family: gaussian

Link function: identity

Formula:

d15N ~ s(Year, k = 15)

Parametric coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.30958	0.02622	126.2	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value
s(Year)	9.282	11.07	61.33	<2e-16 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

R-sq.(adj) = 0.935 Deviance explained = 94.8%

-REML = 3.9734 Scale est. = 0.03299 n = 48

Visualising fitted GAM



`visreg(m)`



Checking fitted GAM





```
library('DHARMa')  
simulateResiduals(m, plot = TRUE)
```

DHARMa residual



Checking fitted GAM

```
gam.check(m)
```



Including temporal autocorrelation

```
mod <- gamm(d15N ~ s(Year, k = 15), data = isotopes,  
            correlation = corCAR1(form = ~ Year), method = 'REML')
```

Family: gaussian

Link function: identity

Formula:

d15N ~ s(Year, k = 15)

Parametric coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.30909	0.03489	94.84	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value
s(Year)	7.954	7.954	47.44	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Modelling infant mortality

Modelling infant mortality

```
mort <- read.csv('data/UN_GDP_infantmortality.csv')
```

	country	infant.mortality	gdp
1	Afghanistan	154	2848
2	Albania	32	863
3	Algeria	44	1531
4	American.Samoa	11	NA
5	Andorra	NA	NA
6	Angola	124	355

Modelling infant mortality with a GLM

```
library('MASS')  
mort.glm <- glm.nb(infant.mortality ~ gdp, data = mort)
```

Call:

```
glm.nb(formula = infant.mortality ~ gdp, data = mort, init.theta = 2.460991808,  
       link = log)
```

Coefficients:


	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	4.072e+00	5.727e-02	71.11	<2e-16 ***
gdp	-8.675e-05	6.221e-06	-13.95	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Negative Binomial(2.461) family taken to be 1)

Null deviance: 385.83 on 192 degrees of freedom
Residual deviance: 202.51 on 191 degrees of freedom
(14 observations deleted due to missingness)
AIC: 1715

Modelling infant mortality with a GLM



Modelling infant mortality with a GLM (log.gdp)

```
mort$log.gdp <- log(mort$gdp)
mort.glm.log <- glm.nb(infant.mortality ~ log.gdp, data = mort)
```

Call:

```
glm.nb(formula = infant.mortality ~ log.gdp, data = mort, init.theta = 3.119314453,
       link = log)
```

Coefficients:


	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	7.07818	0.20045	35.31	<2e-16 ***
log.gdp	-0.47238	0.02647	-17.85	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Negative Binomial(3.1193) family taken to be 1)

Null deviance: 478.54 on 192 degrees of freedom
Residual deviance: 198.03 on 191 degrees of freedom
(14 observations deleted due to missingness)
AIC: 1667.7

Modelling infant mortality with a GLM (log.gdp)



Modelling infant mortality with a GAM

```
library('mgcv')  
mort.gam <- gam(infant.mortality ~ s(log.gdp), family = nb, data = mort)
```

Family: Negative Binomial(3.251)

Link function: log

Formula:

infant.mortality ~ s(log.gdp)

Parametric coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	3.51137	0.04257	82.49	<2e-16 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Approximate significance of smooth terms:

	edf	Ref.df	Chi.sq	p-value
s(log.gdp)	3.134	3.937	329.9	<2e-16 ***





Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Modelling infant mortality with a GAM



Checking GAM

```
gam.check(mort.gam)
```



Comparing models

```
library('easystats')  
compare_performance(mort.glm, mort.glm.log, mort.gam)
```

Comparison of Model Performance Indices


Name	Model	AIC (weights)	AICc (weights)	BIC (weights)
mort.glm	negbin	1715.0 (<.001)	1715.1 (<.001)	1724.7 (<.001)
mort.glm.log	negbin	1667.7 (0.035)	1667.9 (0.041)	1677.5 (0.816)
mort.gam	gam	1661.1 (0.965)	1661.6 (0.959)	1680.5 (0.184)

Name	RMSE	Sigma	Score_log	Score_spherical	Nagelkerke's R2	R2
mort.glm	31.089	1.000	-4.437	0.048	0.709	
mort.glm.log	30.034	1.000	-4.356	0.053	0.836	
mort.gam	26.249	1.000	-4.296	0.049		0.526

Generalised Additive Mixed Models (GAMM)

Reaction time with sleep deprivation

```
library('lme4')  
data('sleepstudy')
```



Modelling reaction time with sleep deprivation (GAMM)

```
sgamm <- gam(Reaction ~ s(Days, Subject, k = 3, bs = 'fs'),  
             data = sleepstudy, method = 'REML')
```

Family: gaussian

Link function: identity

Formula:

Reaction ~ s(Days, Subject, k = 3, bs = "fs")

Parametric coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	298.51	9.05	32.98	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value
s(Days,Subject)	45.67	53	17.11	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

R-sq.(adj) = 0.835 Deviance explained = 87.7%

-REML = 883.64 Scale est. = 523.2 n = 180

Modelling reaction time with sleep deprivation (GAMM)

```
visreg(sgamma, xvar = 'Days', by = 'Subject')
```

