

## Hypothesis testing

---

## NHST concepts

---

# Null and alternative hypotheses

- Tell me...

# Null and alternative hypotheses

- Tell me...
- **Null hypothesis:** there is no difference between groups.

# Null and alternative hypotheses

- Tell me...
- **Null hypothesis:** there is no difference between groups.
- **Alternative hypothesis:** groups are different.

## Are there any differences? A non-sensical question in ecology

Alejandro Martínez-Abraín

IMEDEA (CSIC-UIB), C/Miquel Marqués 21, 07190 Esporles, Majorca, Spain

---

### ARTICLE INFO

#### Article history:

Received 19 December 2006

Accepted 27 April 2007

Published online 13 June 2007

---

#### Keywords:

### ABSTRACT

One of the main questions that ecologists pose in their investigations includes the analysis of differences in some trait between two or more populations. I argue here that asking whether there are differences or not between populations is biologically irrelevant, since no two living things are ever equal. On the contrary the appropriate question to pose is how large differences are between populations. That is, we urge a shift in interest from statistical significance to biological relevance for proper knowledge accumulation. I empha-

# What is the p-value?

- The probability that the data were produced by random chance alone

<https://pollev.com/franciscorod726>

# What is the p-value?

- The probability that the data were produced by random chance alone
- The probability of getting results at least as extreme as the ones you observed if  $H_0$  was true

<https://pollev.com/franciscorod726>



# What is the p-value?

- The probability that the data were produced by random chance alone
- The probability of getting results at least as extreme as the ones you observed if  $H_0$  was true
- The probability of null hypothesis being true

<https://pollev.com/franciscorod726>

- Very complicated concept: even statisticians fail to describe it well.

- Very complicated concept: even statisticians fail to describe it well.
- Probability of observing data as or more extreme than these *if  $H_0$  was true*.

- Very complicated concept: even statisticians fail to describe it well.
- Probability of observing data as or more extreme than these *if  $H_0$  was true*.
- Low P-value: data unlikely if  $H_0$  was true.

- Very complicated concept: even statisticians fail to describe it well.
- Probability of observing data as or more extreme than these *if  $H_0$  was true*.
- Low P-value: data unlikely if  $H_0$  was true.
- Large P-value: data not unusual if  $H_0$  was true.

If  $p\text{-value} > 0.05$

- the null hypothesis is false, i.e. the alternative hypothesis must be true

<https://pollev.com/franciscorod726>

If  $p\text{-value} > 0.05$

- the null hypothesis is false, i.e. the alternative hypothesis must be true
- the alternative hypothesis is false, i.e. the null hypothesis must be true

<https://pollev.com/franciscorod726>

If  $p\text{-value} > 0.05$

- the null hypothesis is false, i.e. the alternative hypothesis must be true
- the alternative hypothesis is false, i.e. the null hypothesis must be true
- it's unclear if there are differences between groups

<https://pollev.com/franciscorod726>



- the null hypothesis is false, i.e. the alternative hypothesis must be true
- the alternative hypothesis is false, i.e. the null hypothesis must be true
- it's unclear if there are differences between groups
- there is no difference between groups

<https://pollev.com/franciscorod726>

## Are differences *significant*?

- If  $p < 0.05$ , we reject  $H_0$ .

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$
- (which is **NOT** the same as ' $H_0$  is true')

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$
- (which is **NOT** the same as ' $H_0$  is true')
- **CAUTION:**

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$
- (which is **NOT** the same as 'H0 is true')
- **CAUTION:**
- P-value is continuous. We must **avoid binary decisions** based on **arbitrary thresholds**.

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$
- (which is **NOT** the same as 'H0 is true')
- **CAUTION:**
- P-value is continuous. We must **avoid binary decisions** based on **arbitrary thresholds**.

## Are differences *significant*?

- If  $p < 0.05$ , we **reject**  $H_0$ .
- If  $p > 0.05$ , we **fail to reject**  $H_0$
- (which is **NOT** the same as 'H0 is true')
- **CAUTION:**
- P-value is continuous. We must **avoid binary decisions** based on **arbitrary thresholds**.



<https://doi.org/10.1038/d41586-019-00857-9>



## Are the heights of local and non-local students different?

```
t.test(h.sevi, h.out)
```

Welch Two Sample t-test

data: h.sevi and h.out

t = -3.159, df = 11.768, p-value = 0.00842

alternative hypothesis: true difference in means is not equal to 0

95 percent confidence interval:

-21.986075 -4.013925

sample estimates:


mean of x mean of y

165.8 178.8


<https://pollev.com/franciscorod726>

## Rejecting hypotheses: two types of error

**Type I error**  
(false positive)



**Type II error**  
(false negative)



## Rejecting hypotheses: two types of error

| <b>Statistics:<br/>Hypothesis<br/>Test</b> | Null Hypothesis<br>is True | Null Hypothesis<br>is False |
|--|----------------------------|-----------------------------|
|  | <b>Type I Error</b>        | <b>Correct</b>              |
| Reject<br>Null Hypothesis                  |                            |                             |
| Fail to Reject<br>Null Hypothesis          | <b>Correct</b>             | <b>Type II Error</b>        |

**Power:** Probability of detecting true difference (rejecting  $H_0$  when it's false).

<http://rpsychologist.com/d3/NHST/>

## Example: biased coin

```
[1] 1 0 1 0 1 0 0 1 1 0
```

1-sample proportions test without continuity correction

data: sum(coin) out of ntrials, null probability 0.5

X-squared = 0, df = 1, p-value = 1

alternative hypothesis: true p is not equal to 0.5

95 percent confidence interval:

0.2365931 0.7634069

sample estimates:

p

0.5

<https://pollev.com/franciscorod726>

<http://rpsychologist.com/d3/correlation/>

## Common pitfalls and good practice

---

Eur J Epidemiol (2016) 31:337–350  
DOI 10.1007/s10654-016-0149-3



ESSAY

## **Statistical tests, *P* values, confidence intervals, and power: a guide to misinterpretations**

Sander Greenland<sup>1</sup> · Stephen J. Senn<sup>2</sup> · Kenneth J. Rothman<sup>3</sup> · John B. Carlin<sup>4</sup> · Charles Poole<sup>5</sup> · Steven N. Goodman<sup>6</sup> · Douglas G. Altman<sup>7</sup>

<https://doi.org/10.1007/s10654-016-0149-3>





ECOSPHERE

Applied statistics in ecology:  
common pitfalls and simple solutions

E. ASHLEY STEEL,<sup>1,†</sup> MAUREEN C. KENNEDY,<sup>2</sup> PATRICK G. CUNNINGHAM,<sup>3</sup> AND JOHN S. STANOVICK<sup>4</sup>

<https://doi.org/10.1890/ES13-00160.1>

Also <http://www.statisticsonewrong.com/>



Twenty tips for  
interpreting  
scientific claims

Visualisation of data and models  
is key

---





- Always

- Always
- Always

# First things first

- Always
- Always
- Always


## Plot data and models




# Don't use statistics blindly: *Visualise*

All correlations:  $r(50) = 0.5$


(1) Normal x, normal residuals




(2) Uniform x, normal residuals




(3) +skewed x, normal residuals




(4) -skewed x, normal residuals




(5) Normal x, +skewed residuals




(6) Normal x, -skewed residuals




(7) Increasing spread




(8) Decreasing spread




(9) Quadratic trend




(10) Sinusoid relationship




(11) A single positive outlier




(12) A single negative outlier




(13) Bimodal residuals




(14) Two groups



(15) Sampling at the extremes



(16) Coarse data







# Don't use statistics blindly: *Visualise*

All correlations:  $r(50) = 0$


(1) Normal x, normal residuals




(2) Uniform x, normal residuals




(3) +skewed x, normal residuals




(4) -skewed x, normal residuals




(5) Normal x, +skewed residuals




(6) Normal x, -skewed residuals




(7) Increasing spread




(8) Decreasing spread




(9) Quadratic trend




(10) Sinusoid relationship




(11) A single positive outlier




(12) A single negative outlier




(13) Bimodal residuals




(14) Two groups



(15) Sampling at the extremes




(16) Coarse data




# Don't use statistics blindly: *Visualise*

All correlations:  $r(50) = 0.1$


(1) Normal x, normal residuals




(2) Uniform x, normal residuals




(3) +skewed x, normal residuals




(4) -skewed x, normal residuals




(5) Normal x, +skewed residuals




(6) Normal x, -skewed residuals




(7) Increasing spread




(8) Decreasing spread




(9) Quadratic trend




(10) Sinusoid relationship




(11) A single positive outlier




(12) A single negative outlier




(13) Bimodal residuals




(14) Two groups



(15) Sampling at the extremes



(16) Coarse data



*Plot. Check models. Plot. Check assumptions. Plot.*

Lavine 2014 Ecology

## Inference from observational studies

---

- In a sample of 10,000 people, it was found that people eating >2 hamburgers a week had 20% higher probability of heart attack.

- In a sample of 10,000 people, it was found that people eating >2 hamburgers a week had 20% higher probability of heart attack.
- Do hamburgers increase heart attacks?

## News: Hamburgers increase risk of heart attack

- In a sample of 10,000 people, it was found that people eating >2 hamburgers a week had 20% higher probability of heart attack.
- Do hamburgers increase heart attacks?
- <https://pollev.com/franciscorod726>

## Bigger flowers increase reproductive success

- We found that plants with big flowers produced 30% more seeds...




## Bigger flowers increase reproductive success

- We found that plants with big flowers produced 30% more seeds...
- Do big flowers increase reproductive success?


## Bigger flowers increase reproductive success

- We found that plants with big flowers produced 30% more seeds...
- Do big flowers increase reproductive success?
- <https://pollev.com/franciscorod726>

# Correlation vs Causation



<http://tylervigen.com/spurious-correlations>



## NHST and p-values

---

## Are there any differences? A non-sensical question in ecology

Alejandro Martínez-Abraín

IMEDEA (CSIC-UIB), C/Miquel Marqués 21, 07190 Esporles, Majorca, Spain

---

### ARTICLE INFO

#### Article history:

Received 19 December 2006

Accepted 27 April 2007

Published online 13 June 2007


---

#### Keywords:

### ABSTRACT


One of the main questions that ecologists pose in their investigations includes the analysis of differences in some trait between two or more populations. I argue here that asking whether there are differences or not between populations is biologically irrelevant, since no two living things are ever equal. On the contrary the appropriate question to pose is how large differences are between populations. That is, we urge a shift in interest from statistical significance to biological relevance for proper knowledge accumulation. I empha-

## Instead of falsifying null model, compare meaningful models



<https://doi.org/10.1242/jeb.104976>

## P-value depends on sample size







# P-value depends on sample size


Same real difference is detected as significant or not depending on sample size:

Real difference = 40 g



## Statistically significant != biologically important

With big sample size, we can find highly significant but biologically unimportant differences.



## Statistically significant != biologically important

- Statistically significant = unlikely to be zero

## Statistically significant != biologically important

- Statistically significant = unlikely to be zero
- Good read: *significantly misleading*





# Statistically significant != biologically important

- Statistically significant = unlikely to be zero
- Good read: *significantly misleading*
- My suggestion: avoid significant/not significant (and maybe p-values too)

# Statistically significant != biologically important


- Statistically significant = unlikely to be zero
- Good read: *significantly misleading*
- My suggestion: avoid significant/not significant (and maybe p-values too)
- Beyond significance, look at *effect sizes*.

## 'Not significant' does NOT mean 'there is no effect'




- Absence of evidence != Evidence of absence

Failure to reject  $H_0 \neq H_0$  is true



$P \gg 0.05$



$P \ll 0.05$



- “We were **unable to find evidence** against the hypothesis that  $A = B$  **with the current sample size**” ([Harrell](#))

- “We were **unable to find evidence** against the hypothesis that  $A = B$  **with the current sample size**” ([Harrell](#))
- “Differences between groups were **not statistically clear**” ([Dushoff et al](#))

# Is it safe to allow right turn with red lights?

- Right turn not allowed: 308 accidents

<https://www.statisticsonewrong.com/power.html#the-wrong-turn-on-red>

## Is it safe to allow right turn with red lights?

- Right turn not allowed: 308 accidents
- Right turn allowed: 337 accidents

<https://www.statisticsonewrong.com/power.html#the-wrong-turn-on-red>

## Is it safe to allow right turn with red lights?

- Right turn not allowed: 308 accidents
- Right turn allowed: 337 accidents
- No *significant* difference, hence safe

<https://www.statisticsonewrong.com/power.html#the-wrong-turn-on-red>

## Is it safe to allow right turn with red lights?

- Right turn not allowed: 308 accidents
- Right turn allowed: 337 accidents
- No *significant* difference, hence safe
- Misinterpretation of underpowered study cost lives


<https://www.statisticsonewrong.com/power.html#the-wrong-turn-on-red>

**The Difference Between “Significant” and “Not Significant” is not  
Itself Statistically Significant**

Andrew GELMAN and Hal STERN


<http://dx.doi.org/10.1198/000313006X152649>

## Multiple hypothesis testing





# How to make your results significant: *p*-hacking



## How to make your results significant: *p-hacking*

1. Test multiple variables, then report the ones that are significant.

## How to make your results significant: *p-hacking*

1. Test multiple variables, then report the ones that are significant.
2. Artificially choose when to end your experiment.

## How to make your results significant: *p-hacking*

1. Test multiple variables, then report the ones that are significant.
2. Artificially choose when to end your experiment.
3. Add covariates until effects are significant.

## How to make your results significant: *p-hacking*

1. Test multiple variables, then report the ones that are significant.
2. Artificially choose when to end your experiment.
3. Add covariates until effects are significant.
4. Test different conditions (e.g. different levels of a factor) and report the ones you like.

## How to make your results significant: *p-hacking*

1. Test multiple variables, then report the ones that are significant.
  2. Artificially choose when to end your experiment.
  3. Add covariates until effects are significant.
  4. Test different conditions (e.g. different levels of a factor) and report the ones you like.
- To read more: [Simmons et al 2011](#)

## How to make your results significant: *p-hacking*

<https://www.youtube.com/watch?v=ZaNtz76dNSI>

- P-values do not measure the **probability of hypothesis** being true, or the probability that the data were produced by **random chance** alone.

<https://doi.org/10.1080/00031305.2016.1154108>



- P-values do not measure the **probability of hypothesis** being true, or the probability that the data were produced by **random chance** alone.
- Scientific conclusions or policy decisions should NOT be based only on **whether a p-value passes a specific threshold**.

<https://doi.org/10.1080/00031305.2016.1154108>

- P-values do not measure the **probability of hypothesis** being true, or the probability that the data were produced by **random chance** alone.
- Scientific conclusions or policy decisions should NOT be based only on **whether a p-value passes a specific threshold**.
- P-value, or statistical significance, does not measure the **size of an effect** or the **importance** of a result.

<https://doi.org/10.1080/00031305.2016.1154108>

- P-values do not measure the **probability of hypothesis** being true, or the probability that the data were produced by **random chance** alone.
- Scientific conclusions or policy decisions should NOT be based only on **whether a p-value passes a specific threshold**.
- P-value, or statistical significance, does not measure the **size of an effect** or the **importance** of a result.
- By itself, a p-value does NOT provide a good **measure of evidence** regarding a model or hypothesis.

<https://doi.org/10.1080/00031305.2016.1154108>

Aim for estimation of effects and their uncertainty (SE, CI...)

*General Article*

---

## **The New Statistics: Why and How**

**Geoff Cumming**  
La Trobe University



Psychological Science  
2014, Vol. 25(1) 7–29  
© The Author(s) 2013  
Reprints and permissions:  
sagepub.com/journalsPermissions.nav  
DOI: 10.1177/0956797613504966  
pss.sagepub.com



<http://dx.doi.org/10.1177/0956797613504966>

## How many types of errors?

- **Type I:** False positive (incorrect rejection of null hypothesis).

## How many types of errors?

- **Type I:** False positive (incorrect rejection of null hypothesis).
- **Type II:** False negative (failure to reject false null hypothesis).

## How many types of errors?

- **Type I:** False positive (incorrect rejection of null hypothesis).
- **Type II:** False negative (failure to reject false null hypothesis).
- **Type S (Sign):** estimating effect in opposite direction.

## How many types of errors?

- **Type I:** False positive (incorrect rejection of null hypothesis).
- **Type II:** False negative (failure to reject false null hypothesis).
- **Type S (Sign):** estimating effect in opposite direction.
- **Type M (Magnitude):** Misestimating magnitude of the effect (under or overestimating).



# How many types of errors?

- **Type I:** False positive (incorrect rejection of null hypothesis).
- **Type II:** False negative (failure to reject false null hypothesis).
- **Type S (Sign):** estimating effect in opposite direction.
- **Type M (Magnitude):** Misestimating magnitude of the effect (under or overestimating).
- **Beyond Power Calculations: Assessing Type S (Sign) and Type M (Magnitude) Errors**