The importance of sample size & good study

design

The most important aspect of a statistical analysis is not what you do with the data, it's what data you use

H. Stern / A. Gelman

The importance of sample size

• Many studies have too low sample sizes.


The importance of sample size

- · Many studies have too low sample sizes.
- Low sample sizes miss subtle effects, but also prone to bias.

The importance of sample size


- · Many studies have too low sample sizes.
- · Low sample sizes miss subtle effects, but also prone to bias.
- The fallacy of assuming that which does not kill statistical significance makes it stronger (Loken & Gelman 2017).


Low sample sizes are very sensitive to random noise



https://twitter.com/ajstewart_lang/status/1020038488278945797

Low sample sizes may bias inferences about population





Source: statisticalgate.com

Low sample sizes may bias inferences

See The evolution of correlations

Stopping rules




Fig. 2. Illustrative simulation of p values obtained by a researcher who continuously adds an observation to each of two conditions, conducting a t test after each addition. The dotted line highlights the conventional significance criterion of $p \le .05$.

• Plan model/statistical analysis before data collection.

- Plan model/statistical analysis before data collection.
- Do simulations. Power/Sample size/Precision analyses (e.g. see papers like this & this, or software like this & this).

- Plan model/statistical analysis before data collection.
- Do simulations. Power/Sample size/Precision analyses (e.g. see papers like this & this, or software like this & this).
- Justify your sample size (precision, minimal effect size...). See Lakens' book.

- Plan model/statistical analysis before data collection.
- Do simulations. Power/Sample size/Precision analyses (e.g. see papers like this & this, or software like this & this).
- Justify your sample size (precision, minimal effect size...). See Lakens' book.
- Plan to have at least 10-30 observations per predictor.

- Plan model/statistical analysis before data collection.
- Do simulations. Power/Sample size/Precision analyses (e.g. see papers like this & this, or software like this & this).
- Justify your sample size (precision, minimal effect size...). See Lakens' book.
- · Plan to have at least 10-30 observations per predictor.
- Complex models (w/ many predictors, interactions etc) require high sample sizes.

Calculating sample size for Gaussian (Normal) response model:

- expected mean: 30
- · expected sd: 10
- 10 parameters (predictors)
- · expected R-squared: 0.2

```
library(pmsampsize)
pmsampsize(type = "c", parameters = 10, intercept = 30, sd = 10, rsquared = 0.2)
```

NB: Assuming 0.05 acceptable difference in apparent & adjusted R-squared
NB: Assuming MMOE <= 1.1 in estimation of intercept & residual standard deviation
SPP - Subjects per Predictor Parameter

```
Samp size Shrinkage Parameter Rsq SPP
Criteria 1
             313
                   0.900
                            10 0.2 31.3
Criteria 2 161
                0.827
                            10 0.2 16.1
Criteria 3 244 0.876 10 0.2 24.4
Criteria 4*
         313 0.900
                            10 0.2 31.3
Final
                   0.900
                            10 0.2 31.3
            313
```

Minimum sample size required for new model development based on user inputs = 313

```
* 95% CI for intercept = (29.01, 30.99), for sample size n = 313
```

Calculating sample size for binary response model:

- expected prevalence: 0.1
- · 20 parameters (predictors)
- · expected R-squared: 0.2

```
library(pmsampsize)
pmsampsize(type = "b", parameters = 20, prevalence = 0.1, nagrsquared = 0.2)
```

NB: Assuming 0.05 acceptable difference in apparent & adjusted R-squared NB: Assuming 0.05 margin of error in estimation of intercept

NB: Events per Predictor Parameter (EPP) assumes prevalence = 0.1

		Samp_size	Shrinkage	Parameter	CS_Rsq	Max_Rsq	Nag_Rsq	EPP	
Criteria :	1	1774	0.900	20	0.096	0.478	0.201	8.87	
Criteria 2	2	786	0.801	20	0.096	0.478	0.201	3.93	
Criteria 3	3	139	0.900	20	0.096	0.478	0.201	0.70	
Final		1774	0.900	20	0.096	0.478	0.201	8.87	

Minimum sample size required for new model development based on user inputs = 1774, with 178 events (assuming an outcome prevalence = 0.1) and an EPP = 8.87