

Experimental design

How would you evaluate fertilizer effect?

Discuss with partner (5')



Experimental design principles

Replication

Replication!




- Replication is key: we need several samples.

- Replication is key: we need several samples.
- How many? As much as you can! See [Gelman & Carlin 2014](#).

- Replication is key: we need several samples.
- How many? As much as you can! See [Gelman & Carlin 2014](#).
- Traditionally, ecology studies have had **too low sample sizes**.


- Replication is key: we need several samples.
- How many? As much as you can! See [Gelman & Carlin 2014](#).
- Traditionally, ecology studies have had **too low sample sizes**.
- Low sample sizes miss subtle effects, but also [prone to bias](#).

Low sample sizes very sensitive to random noise



https://twitter.com/ajstewart_lang/status/1020038488278945797

Low sample sizes may bias inferences about population



<http://statisticalgate.com/regression-simulation/>

Low sample sizes may bias inferences

See [The evolution of correlations](#)

Stopping rules




Fig. 2. Illustrative simulation of p values obtained by a researcher who continuously adds an observation to each of two conditions, conducting a t test after each addition. The dotted line highlights the conventional significance criterion of $p \leq .05$.

- Plan model/statistical analysis **before** data collection.

- Plan model/statistical analysis **before** data collection.
- **Do simulations.** Power/Sample size/Precision analyses (e.g.[this](#), [this](#) or [this](#)).

- Plan model/statistical analysis **before** data collection.
- **Do simulations.** Power/Sample size/Precision analyses (e.g.[this](#), [this](#) or [this](#)).
- Plan to have at least **10-30 observations per predictor**.

- Plan model/statistical analysis **before** data collection.
- **Do simulations.** Power/Sample size/Precision analyses (e.g.[this](#), [this](#) or [this](#)).
- Plan to have at least **10-30 observations per predictor**.
- Complex models (w/ many predictors, interactions etc) require **high** sample sizes.

Randomization

Randomization



- Haphazard \neq Random

- Haphazard \neq Random
- Stratify: randomize within groups (e.g. species, soil types)

Controls

- Untreated individuals, plots... (assigned randomly, of course).

- Untreated individuals, plots... (assigned randomly, of course).
- **Must differ only in treatment** (i.e. homogeneous environment).

- Untreated individuals, plots... (assigned randomly, of course).
- **Must differ only in treatment** (i.e. homogeneous environment).
- Measure **before & after** treatment.

- Untreated individuals, plots... (assigned randomly, of course).
- **Must differ only in treatment** (i.e. homogeneous environment).
- Measure **before & after** treatment.
- Consider **blind designs** to avoid observer bias.

1. Replication

1. Replication
2. Randomization

Experimental design principles

1. Replication
2. Randomization
3. Controls

- Ruxton & Colegrave. Experimental Design for the Life Sciences. OUP