

Sampling, confidence intervals, likelihood and Bayesian inference

Inference: from samples to population

We rarely measure the whole **population**, but take **samples**.

Then we make inferences from sample to population.



If we sample 30 trees in our neighbourhood...

Can we extrapolate results to

- whole neighbourhood?
- whole city?
- whole country?
- the world?

What's the **suitable population** to make inferences given this sample?

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))
- A 95% CI is **NOT** 95% likely to contain the true parameter value!

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))
- A 95% CI is **NOT** 95% likely to contain the true parameter value!
- Instead, 95% of the CIs obtained with this sampling will contain the true value.

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))
- A 95% CI is **NOT** 95% likely to contain the true parameter value!
- Instead, 95% of the CIs obtained with this sampling will contain the true value.
- Like person who tells truth 95% of the time, but we can't tell if a particular statement is true.

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))
- A 95% CI is **NOT** 95% likely to contain the true parameter value!
- Instead, 95% of the CIs obtained with this sampling will contain the true value.
- Like person who tells truth 95% of the time, but we can't tell if a particular statement is true.
- It's a frequentist, long-run property.

Understanding confidence intervals

- Summarise **uncertainty** in parameter estimates.
- <https://rpsychologist.com/d3/CI/> (or [here](#))
- A 95% CI is **NOT** 95% likely to contain the true parameter value!
- Instead, 95% of the CIs obtained with this sampling will contain the true value.
- Like person who tells truth 95% of the time, but we can't tell if a particular statement is true.
- It's a frequentist, long-run property.
- To read more: [Morey et al \(2015\)](#)

What happens if we increase sample size?

<https://rpsychologist.com/d3/CI/>

- CI width *decreases*...

What happens if we increase sample size?

<https://rpsychologist.com/d3/CI/>

- CI width *decreases*...
- but still 5% of CIs will NOT contain true mean!

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150

<https://pollev.com/franciscorod726>

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150
- We can be 95% confident that X lies between 120 and 150

<https://pollev.com/franciscorod726>

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150
- We can be 95% confident that X lies between 120 and 150
- If we repeated the experiment, 95% of the time X would fall between 120 and 150

<https://pollev.com/franciscorod726>

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150
- We can be 95% confident that X lies between 120 and 150
- If we repeated the experiment, 95% of the time X would fall between 120 and 150
- If we repeated the experiment, 95% of the CIs would contain the true value of X

<https://pollev.com/franciscorod726>

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150
- We can be 95% confident that X lies between 120 and 150
- If we repeated the experiment, 95% of the time X would fall between 120 and 150
- If we repeated the experiment, 95% of the CIs would contain the true value of X
- The probability that X is greater than 0 is at least 95%

<https://pollev.com/franciscorod726>

If the 95% CI of X is (120, 150)...

- There is a 95% probability that X lies between 120 and 150
- We can be 95% confident that X lies between 120 and 150
- If we repeated the experiment, 95% of the time X would fall between 120 and 150
- If we repeated the experiment, 95% of the CIs would contain the true value of X
- The probability that X is greater than 0 is at least 95%
- The probability that X equals 0 is smaller than 5%

<https://pollev.com/franciscorod726>


- Bayesian **credible** intervals do give the probability that true parameter value is contained within them.

- Bayesian **credible** intervals do give the probability that true parameter value is contained within them.
- Frequentist CIs and Bayesian credible intervals can be similar, but not always.

Bayesian inference: prior, posterior, and likelihood


$$P(\text{Unknown}|\text{Data}) \propto P(\text{Data}|\text{Unknown}) \times P(\text{Unknown})$$

$$\text{Posterior} \propto \text{Likelihood} \times \text{Prior}$$




What is the likelihood?

$$L(\theta|x) = P(x|\theta)$$




<https://seeing-theory.brown.edu/bayesian-inference/index.html>

Bayesian inference: prior and likelihood produce posterior



With increasing sample size, likelihood dominates prior



More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference

More apps to introduce Bayesian inference

- [Wagenmaker's first lesson in Bayesian inference](#)
- [Bayesian Demo](#)

More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference
- Bayesian Demo
- Bayesian inference for a population mean

More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference
- Bayesian Demo
- Bayesian inference for a population mean
- Normal

More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference
- Bayesian Demo
- Bayesian inference for a population mean
- Normal
- Binomial

More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference
- Bayesian Demo
- Bayesian inference for a population mean
- Normal
- Binomial
- Own data

More apps to introduce Bayesian inference

- Wagenmaker's first lesson in Bayesian inference
- Bayesian Demo
- Bayesian inference for a population mean
- Normal
- Binomial
- Own data
- Bayesian t-test

- Integrate information (prior)

- Integrate information (prior)
- Prior regularises unlikely estimates from data

- Integrate information (prior)
- Prior regularises unlikely estimates from data
- Large dataset -> prior effect diminishes


- Integrate information (prior)
- Prior regularises unlikely estimates from data
- Large dataset -> prior effect diminishes
- Uncertainty / Propagate errors

Uncertainty!

Even with same data, different teams reach different conclusions!

How does sibling competition affect nestling growth in blue tits?

Most teams found negative effect, but large variation in effect size. Some teams no or positive effect



73 teams testing the same hypothesis with the same data


PNAS

RESEARCH ARTICLE

SOCIAL SCIENCES

OPEN ACCESS

Observing many researchers using the same data and hypothesis reveals a hidden universe of uncertainty



“This reveals a **universe of uncertainty** that remains hidden when considering a single study in isolation”

“These results call for greater **epistemic humility** and **clarity in reporting** scientific findings”