

Supporting Information for “A nutrient effect on the TEX₈₆ paleotemperature proxy”

Ronnakrit Rattanasriampaipong^{1,2}, Jessica Tierney², Jordan T. Abell³

¹University Corporation for Atmospheric Research, Boulder, CO 80307

²Department of Geosciences, The University of Arizona, Tucson, AZ 85721

³Department of Earth and Environmental Sciences, Lehigh University, Bethlehem, PA 18015

Contents of this file

1. Figures S1 to Sx
2. Tables S1

Additional Supporting Information (Files uploaded separately)

1. Captions for Datasets S1 to S6
2. Captions for Codes S1 to S2

Introduction

This document provides supplementary information for the main text, including (a) supplementary texts for extended methods and materials, (b) a summary table of data sources of the updated global coretop TEX₈₆ dataset, (c) brief descriptions of supple-

Corresponding author: R. Rattanasriampaipong, Department of Geosciences, University of Arizona, Gould-Simpson Building, Room 513A, Tucson, AZ 85721, USA. (rrattan@ucar.edu)

mentary datasets and codes repositied on Zenodo (Rattanasriampaipong et al., 2025) and Github repositories (<https://github.com/PaleoLipidRR/nutrient-effect-on-TEX>).

Text S1: Extended Data and Methods

- 1. New core-top TEX₈₆ data**
- 2. Downcore samples from Tasman Sea**
- 3. GDGT determinations**

Data Set S1

Code S1

References

- Bale, N. J., Villanueva, L., Hopmans, E. C., Schouten, S., & Sinninghe Damsté, J. S. (2013). Different seasonality of pelagic and benthic Thaumarchaeota in the North Sea. *Biogeosciences*, 10(11), 7195–7206. Retrieved from <https://bg.copernicus.org/articles/10/7195/2013/> doi: 10.5194/bg-10-7195-2013
- Castañeda, I. S., Schefuß, E., Pätzold, J., Sinninghe Damsté, J. S., Weldeab, S., & Schouten, S. (2010, mar). Millennial-scale sea surface temperature changes in the eastern Mediterranean (Nile River Delta region) over the last 27,000 years. *Paleoceanography*, 25(1). Retrieved from <https://doi.org/10.1029/2009PA001740> doi: <https://doi.org/10.1029/2009PA001740>
- Ceccopieri, M., Carreira, R. S., Wagener, A. L. R., Hefter, J. H., & Mollenhauer, G. (2018). On the application of alkenone- and GDGT-based temperature proxies in the south-eastern Brazilian continental margin. *Organic Geochemistry*, 126, 43–56. doi: 10.1016/j.orggeochem.2018.10.009

- Chazen, C. R. (2011). *Holocene climate evolution of the eastern tropical Pacific told from high resolution climate records from the Peru margin and equatorial upwelling regions* (Doctoral dissertation, Brown University). doi: <https://doi.org/10.7301/Z0D50K7P>
- Chen, J., Hu, P., Li, X., Yang, Y., Song, J., Li, X., ... Lü, X. (2018). Impact of water depth on the distribution of iGDGTs in the surface sediments from the northern South China Sea: applicability of TEX86 in marginal seas. *Frontiers of Earth Science*, 12(1), 95–107. doi: 10.1007/s11707-016-0620-1
- Chen, W., Mohtadi, M., Schefuß, E., & Mollenhauer, G. (2014). Organic-geochemical proxies of sea surface temperature in surface sediments of the tropical eastern Indian Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 88, 17–29. doi: <https://doi.org/10.1016/j.dsr.2014.03.005>
- Fallet, U., Castañeda, I. S., Henry-Edwards, A., Richter, T. O., Boer, W., Schouten, S., & Brummer, G.-J. (2012, jan). Sedimentation and burial of organic and inorganic temperature proxies in the Mozambique Channel, SW Indian Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 59, 37–53. doi: 10.1016/j.dsr.2011.10.002
- Hagemann, J. R., Lembke-Jene, L., Lamy, F., Vorrath, M.-E., Kaiser, J., Müller, J., ... Tiedemann, R. (2023). *Collection of GDGT-derived temperatures of 22 new sites at the Chilean margin, supplemented by various previously published sites*. [Dataset]. PANGAEA. doi: 10.1594/PANGAEA.964270
- Harning, D. J., Andrews, J. T., Belt, S. T., Cabedo-Sanz, P., Geirsdottir, A., Dildar, N., ... Sepúlveda, J. (2019). Sea Ice Control on Winter Subsurface Temperatures of the North Iceland Shelf During the Little Ice Age: A TEX86 Calibration Case Study. *Paleoceanography and Paleoclimatology*, 34(6), 1006–1021. doi: 10.1029/2018PA003523

- Harning, D. J., Holman, B., Woelders, L., Jennings, A. E., & Sepúlveda, J. (2023). Biomarker characterization of the North Water Polynya, Baffin Bay: implications for local sea ice and temperature proxies. *Biogeosciences*, 20(1), 229–249. Retrieved from <https://doi.org/10.1594/PANGAEA.956212><https://bg.copernicus.org/articles/20/229/2023/> doi: 10.5194/bg-20-229-2023
- Hernández-Sánchez, M. T., Woodward, E. M. S., Taylor, K. W. R., Henderson, G. M., & Pancost, R. D. (2014). Variations in GDGT distributions through the water column in the South East Atlantic Ocean. *Geochimica et Cosmochimica Acta*, 132, 337–348. doi: <https://doi.org/10.1016/j.gca.2014.02.009>
- Ho, S. L., Mollenhauer, G., Fietz, S., Martínez-García, A., Lamy, F., Rueda, G., ... Tiedemann, R. (2014). Appraisal of TEX86 and TEX86L thermometries in subpolar and polar regions. *Geochimica et Cosmochimica Acta*, 131, 213–226. doi: <https://doi.org/10.1016/j.gca.2014.01.001>
- Ho, S. L., Yamamoto, M., Mollenhauer, G., & Minagawa, M. (2011). Core top TEX86 values in the south and equatorial Pacific. *Organic Geochemistry*, 42(1), 94–99. doi: <https://doi.org/10.1016/j.orggeochem.2010.10.012>
- Hu, J., Meyers, P. A., Chen, G., Peng, P., & Yang, Q. (2012). Archaeal and bacterial glycerol dialkyl glycerol tetraethers in sediments from the Eastern Lau Spreading Center, South Pacific Ocean. *Organic Geochemistry*, 43, 162–167. doi: <https://doi.org/10.1016/j.orggeochem.2011.10.012>
- Jaeschke, A., Wengler, M., Hefter, J., Ronge, T. A., Geibert, W., Mollenhauer, G., ... Lamy, F. (2017). A biomarker perspective on dust, productivity, and sea surface temperature in the Pacific sector of the Southern Ocean. *Geochimica et Cosmochimica Acta*, 204, 120–139. doi: 10.1016/j.gca.2017.01.045

- Jia, G., Zhang, J., Chen, J., Peng, P., & Zhang, C. L. (2012). Archaeal tetraether lipids record subsurface water temperature in the South China Sea. *Organic Geochemistry*, 50, 68–77. doi: <https://doi.org/10.1016/j.orggeochem.2012.07.002>
- Kaiser, J., Ruggieri, N., Hefter, J., Siegel, H., Mollenhauer, G., Arz, H. W., & Lamy, F. (2014). Lipid biomarkers in surface sediments from the Gulf of Genoa, Ligurian sea (NW Mediterranean sea) and their potential for the reconstruction of palaeo-environments. *Deep Sea Research Part I: Oceanographic Research Papers*, 89, 68–83. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0967063714000661> doi: <https://doi.org/10.1016/j.dsr.2014.04.009>
- Kaiser, J., Schouten, S., Kilian, R., Arz, H. W., Lamy, F., & Sinninghe Damsté, J. S. (2015). Isoprenoid and branched GDGT-based proxies for surface sediments from marine, fjord and lake environments in Chile. *Organic Geochemistry*, 89-90, 117–127. doi: [10.1016/j.orggeochem.2015.10.007](https://doi.org/10.1016/j.orggeochem.2015.10.007)
- Kim, J.-H., Schouten, S., Hopmans, E. C., Donner, B., & Sinninghe Damsté, J. S. (2008). Global sediment core-top calibration of the TEX86 paleothermometer in the ocean. *Geochimica et Cosmochimica Acta*, 72(4), 1154–1173. doi: <https://doi.org/10.1016/j.gca.2007.12.010>
- Kim, J.-H., Schouten, S., Rodrigo-Gámiz, M., Rampen, S., Marino, G., Huguet, C., ... Sinninghe Damsté, J. S. (2015). Influence of deep-water derived isoprenoid tetraether lipids on the TEX86H paleothermometer in the Mediterranean Sea. *Geochimica et Cosmochimica Acta*, 150, 125–141. doi: [10.1016/j.gca.2014.11.017](https://doi.org/10.1016/j.gca.2014.11.017)
- Kim, J.-H., van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., ... Sinninghe Damsté, J. S. (2010). New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraether lipids: Implications for past

- sea surface temperature reconstructions. *Geochimica et Cosmochimica Acta*, *74*(16), 4639–4654. doi: 10.1016/j.gca.2010.05.027
- Kim, J.-H., Villanueva, L., Zell, C., & Sinninghe Damsté, J. S. (2016). Biological source and provenance of deep-water derived isoprenoid tetraether lipids along the Portuguese continental margin. *Geochimica et Cosmochimica Acta*, *172*, 177–204. doi: 10.1016/j.gca.2015.09.010
- Kusch, S., Rethemeyer, J., Hopmans, E. C., Wacker, L., & Mollenhauer, G. (2016). Factors influencing ^{14}C concentrations of algal and archaeal lipids and their associated sea surface temperature proxies in the Black Sea. *Geochimica et Cosmochimica Acta*, *188*, 35–57. doi: 10.1016/j.gca.2016.05.025
- Lamping, N., Müller, J., Hefter, J., Mollenhauer, G., Haas, C., Shi, X., ... Hillenbrand, C.-D. (2021). Evaluation of lipid biomarkers as proxies for sea ice and ocean temperatures along the Antarctic continental margin. *Climate of the Past*, *17*(5), 2305–2326. doi: 10.5194/cp-17-2305-2021
- Leider, A., Hinrichs, K.-U., Mollenhauer, G., & Versteegh, G. J. M. (2010). Core-top calibration of the lipid-based U37K' and TEX86 temperature proxies on the southern Italian shelf (SW Adriatic Sea, Gulf of Taranto). *Earth and Planetary Science Letters*, *300*(1), 112–124. doi: <https://doi.org/10.1016/j.epsl.2010.09.042>
- Lengger, S. K., Hopmans, E. C., Sinninghe Damsté, J. S., & Schouten, S. (2014). Fossilization and degradation of archaeal intact polar tetraether lipids in deeply buried marine sediments (Peru Margin). *Geobiology*, *12*(3), 212–220. doi: <https://doi.org/10.1111/gbi.12081>
- Liu, X.-L., Zhu, C., Wakeham, S. G., & Hinrichs, K.-U. (2014). In situ production of branched glycerol dialkyl glycerol tetraethers in anoxic marine water columns. *Marine*

- Chemistry*, 166, 1–8. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0304420314001388> doi: <https://doi.org/10.1016/j.marchem.2014.08.008>
- Lo, L., Belt, S. T., Lattaud, J., Friedrich, T., Zeeden, C., Schouten, S., ... Hodell, D. A. (2018). Precession and atmospheric CO₂ modulated variability of sea ice in the central Okhotsk Sea since 130,000 years ago. *Earth and Planetary Science Letters*, 488, 36–45. doi: 10.1016/j.epsl.2018.02.005
- Lü, X., Yang, H., Song, J., Versteegh, G. J. M., Li, X., Yuan, H., ... Xie, S. (2014). Sources and distribution of isoprenoid glycerol dialkyl glycerol tetraethers (GDGTs) in sediments from the east coastal sea of China: Application of GDGT-based paleothermometry to a shallow marginal sea. *Organic Geochemistry*, 75, 24–35. doi: <https://doi.org/10.1016/j.orggeochem.2014.06.007>
- Nieto-Moreno, V., Martínez-Ruiz, F., Willmott, V., García-Orellana, J., Masqué, P., & Sinninghe Damsté, J. S. (2013). Climate conditions in the westernmost Mediterranean over the last two millennia: An integrated biomarker approach. *Organic Geochemistry*, 55, 1–10. doi: <https://doi.org/10.1016/j.orggeochem.2012.11.001>
- Pan, A., Yang, Q., Zhou, H., Ji, F., Wang, H., & Pancost, R. D. (2016). A diagnostic GDGT signature for the impact of hydrothermal activity on surface deposits at the Southwest Indian Ridge. *Organic Geochemistry*, 99, 90–101.
- Park, Y.-H., Yamamoto, M., Nam, S.-I., Irino, T., Polyak, L., Harada, N., ... Saitoh, S.-I. (2014). Distribution, source and transportation of glycerol dialkyl glycerol tetraethers in surface sediments from the western Arctic Ocean and the northern Bering Sea. *Marine Chemistry*, 165, 10–24. doi: <https://doi.org/10.1016/j.marchem.2014.07.001>

- Rattanasriampaipong, R., Tierney, J., & Abell, J. (2025). *Supplementary Data for 'A nutrient effect on the TEX₈₆ paleotemperature proxy'*. [Dataset]. Zenodo. doi: 10.5281/zenodo.14806962
- Rattanasriampaipong, R., Zhang, Y. G., Pearson, A., Hedlund, B. P., & Zhang, S. (2022). Archaeal lipids trace ecology and evolution of marine ammonia-oxidizing archaea. *Proceedings of the National Academy of Sciences*, 119(31), e2123193119. doi: 10.1073/pnas.2123193119
- Richey, J. N., Hollander, D. J., Flower, B. P., & Eglinton, T. I. (2011, mar). Merging late Holocene molecular organic and foraminiferal-based geochemical records of sea surface temperature in the Gulf of Mexico. *Paleoceanography*, 26(1). doi: <https://doi.org/10.1029/2010PA002000>
- Richey, J. N., & Tierney, J. E. (2016). GDGT and alkenone flux in the northern Gulf of Mexico: Implications for the TEX₈₆ and UK₃₇' paleothermometers. *Paleoceanography*, 31(12), 1547–1561. doi: 10.1002/2016pa003032
- Rodrigo-Gámiz, M., Rampen, S. W., de Haas, H., Baas, M., Schouten, S., & Sinninghe Damsté, J. S. (2015). Constraints on the applicability of the organic temperature proxies UK₃₇', TEX₈₆ and LDI in the subpolar region around Iceland. *Biogeosciences*, 12(22), 6573–6590. doi: 10.5194/bg-12-6573-2015
- Schukies, J. (2018). *Concentrations of glycerol dialkyl glycerol tetraethers (GDGTs) in core top sediments from continental slope off Newfoundland, Orphan Knoll Region*. [Dataset]. PANGAEA. doi: 10.1594/PANGAEA.894305
- Seki, O., Bendle, J. A., Harada, N., Kobayashi, M., Sawada, K., Moossen, H., ... Sakamoto, T. (2014). Assessment and calibration of TEX₈₆ paleothermometry in the Sea of Okhotsk and sub-polar North Pacific region: Implications for paleo-

- ceanography. *Progress in Oceanography*, 126, 254–266. doi: <https://doi.org/10.1016/j.pocean.2014.04.013>
- Seki, O., Sakamoto, T., Sakai, S., Schouten, S., Hopmans, E. C., Sinninghe Damsté, J. S., & Pancost, R. D. (2009). Large changes in seasonal sea ice distribution and productivity in the Sea of Okhotsk during the deglaciations. *Geochemistry, Geophysics, Geosystems*, 10(10). doi: <https://doi.org/10.1029/2009GC002613>
- Shevenell, A. E., Ingalls, A. E., Domack, E. W., & Kelly, C. (2011). Holocene Southern Ocean surface temperature variability west of the Antarctic Peninsula. *Nature*, 470(7333), 250–254. doi: [10.1038/nature09751](https://doi.org/10.1038/nature09751)
- Sinninghe Damsté, J. S. (2016). Spatial heterogeneity of sources of branched tetraethers in shelf systems: The geochemistry of tetraethers in the Berau River delta (Kali-mantan, Indonesia). *Geochimica et Cosmochimica Acta*, 186, 13–31. doi: [10.1016/j.gca.2016.04.033](https://doi.org/10.1016/j.gca.2016.04.033)
- Sinninghe Damsté, J. S., Warden, L. A., Berg, C., Jürgens, K., & Moros, M. (2022). Evaluation of the distributions of hydroxylated glycerol dibiphytanyl glycerol tetraethers (GDGTs) in Holocene Baltic Sea sediments for reconstruction of sea surface temperature: the effect of changing salinity. *Climate of the Past*, 18(10), 2271–2288. doi: [10.5194/cp-18-2271-2022](https://doi.org/10.5194/cp-18-2271-2022)
- Smith, M., De Deckker, P., Rogers, J., Brocks, J., Hope, J., Schmidt, S., ... Schouten, S. (2013). Comparison of U37K', TEX86H and LDI temperature proxies for reconstruction of south-east Australian ocean temperatures. *Organic Geochemistry*, 64, 94–104. doi: <https://doi.org/10.1016/j.orggeochem.2013.08.015>
- Tang, W., Ward, B. B., Beman, M., Bristow, L., Clark, D., Fawcett, S., ... Zhang, Y. (2023). Database of nitrification and nitrifiers in the global ocean. *Earth System*

Science Data, 15(11), 5039–5077. doi: 10.5194/essd-15-5039-2023

Tierney, J. E., & Tingley, M. P. (2014). A Bayesian, spatially-varying calibration model for the TEX86 proxy. *Geochimica et Cosmochimica Acta*, 127, 83–106. doi: 10.1016/j.gca.2013.11.026

Tierney, J. E., & Tingley, M. P. (2015). A TEX 86 surface sediment database and extended Bayesian calibration. *Scientific Data*, 2(1), 1–10.

Tierney, J. E., Ummenhofer, C. C., & DeMenocal, P. B. (2015). Past and future rainfall in the Horn of Africa. *Science Advances*, 1(9), e1500682. doi: 10.1126/sciadv.1500682

Trommer, G., Siccha, M., van der Meer, M. T. J., Schouten, S., Sinninghe Damsté, J. S., Schulz, H., ... Kucera, M. (2009). Distribution of Crenarchaeota tetraether membrane lipids in surface sediments from the Red Sea. *Organic Geochemistry*, 40(6), 724–731. doi: 10.1016/j.orggeochem.2009.03.001

Varma, D., Hopmans, E. C., van Kemenade, Z. R., Kusch, S., Berg, S., Bale, N. J., ... Schouten, S. (2024). Evaluating isoprenoidal hydroxylated GDGT-based temperature proxies in surface sediments from the global ocean. *Geochimica et Cosmochimica Acta*, 370, 113–127. doi: 10.1016/j.gca.2023.12.019

Varma, D., Hopmans, E. C., van Kemenda, Z. R., Kusch, S., Berg, S., Bale, N. J., ... Schouten, S. (2024). *OHGDGT global surface sediment data*. [Dataset]. PANGAEA. doi: 10.1594/PANGAEA.964885

Verleye, T. J. (2011). *The late Quaternary palaeoenvironmental changes along the western South-American continental slope: A reconstruction based on dinoflagellate cysts and TEX86* (PhD Thesis, Ghent University). Retrieved from <https://www.vliz.be/en/imis?module=ref&refid=210274>

Wei, B., Jia, G., Hefter, J., Kang, M., Park, E., Wang, S., & Mollenhauer, G. (2020).

- Comparison of the UK₃₇['], LDI, TEX₈₆^H, and RI-OH temperature proxies in sediments from the northern shelf of the South China Sea. *Biogeosciences*, 17(17), 4489–4508. doi: 10.5194/bg-17-4489-2020
- Wei, Y., Wang, J., Liu, J., Dong, L., Li, L., Wang, H., ... Zhang, C. L. (2011). Spatial Variations in Archaeal Lipids of Surface Water and Core-Top Sediments in the South China Sea and Their Implications for Paleoclimate Studies. *Applied and Environmental Microbiology*, 77(21), 7479–7489. doi: 10.1128/aem.00580-11
- Wu, W., Tan, W., Zhou, L., Yang, H., & Xu, Y. (2012, jul). Sea surface temperature variability in southern Okinawa Trough during last 2700 years. *Geophysical Research Letters*, 39(14). doi: <https://doi.org/10.1029/2012GL052749>
- Yang, Y., Gao, C., Dang, X., Ruan, X., Lü, X., Xie, S., ... Yang, H. (2018). Assessing hydroxylated isoprenoid GDGTs as a paleothermometer for the tropical South China Sea. *Organic Geochemistry*, 115, 156–165. doi: 10.1016/j.orggeochem.2017.10.014
- Zell, C., Kim, J.-H., Hollander, D., Lorenzoni, L., Baker, P., Silva, C. G., ... Sinninghe Damsté, J. S. (2014). Sources and distributions of branched and isoprenoid tetraether lipids on the Amazon shelf and fan: Implications for the use of GDGT-based proxies in marine sediments. *Geochimica et Cosmochimica Acta*, 139, 293–312. Retrieved from <http://www.sciencedirect.com/science/article/pii/S0016703714002968> doi: <https://doi.org/10.1016/j.gca.2014.04.038>
- Zhou, H., Hu, J., Spiro, B., Peng, P., & Tang, J. (2014). Glycerol dialkyl glycerol tetraethers in surficial coastal and open marine sediments around China: Indicators of sea surface temperature and effects of their sources. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 395, 114–121. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0031018213005361> doi: <https://doi.org/10.1016/>

j.palaeo.2013.12.006




Figure S1. Ocean surface nitrate concentrations correlate with ammonia oxidation rates. (a) Spatial distribution of gridded ammonia oxidation rates above the thermocline. (b) Relationship between ammonia oxidation rates and nitrate concentrations. Measurements of ammonia oxidation rates and bioavailable nitrogen are co-located. Data were obtained from the global nitrification database (Tang et al., 2023).




Figure S2. The distribution of an extended global core-top TEX₈₆ dataset. Data published in Tierney and Tingley (2015) are plotted in grey. Additional published datasets since 2015 are plotted in blue. New GDGT measurements from this study are plotted in black.




Figure S3. TEX_{86} warming anomalies correlate with bulk sediment $\delta^{15}\text{N}$. Interpolated paleo-records from core sites (left) NIOP905 and (right) SO42-74KL, with a 100-year timestep calculated from the original records shown in **Figures 4B–C**. (a, b) TEX_{86} warming anomalies ($\text{Thermo-T}_{\text{TEX}_{86}} - \text{SST}_{U_{37}^{K'}}$) correlate negatively with bulk sediment $\delta^{15}\text{N}$, with stronger correlations observed during the Younger Dryas and Heinrich Stadial 1 events. (c, d) TEX_{86} residuals respond to varying levels of bulk sediment $\delta^{15}\text{N}$ similarly to TEX_{86} warming anomalies.

Table S1. Summary of coretop GDGT data sources

Remark	Reference	Count
Data from original source	Rattanasriampaipong et al. (2025) (this study)	190
	Ceccopieri et al. (2018)	53
	J. Chen et al. (2018)	28
	Hagemann et al. (2023)	21
	Harning et al. (2019)	12
	Harning et al. (2023)	12
	Kusch et al. (2016)	24
	Liu et al. (2014)	12
	Richey and Tierney (2016)	3
	Rodrigo-Gámiz et al. (2015)*	10
	Schukies (2018)	10
	Tierney and Tingley (2015)	26
	Tierney et al. (2015)	1
	Varma et al. (2024)	224
Data published in Tierney and Tingley (2014, 2015)	B. Wei et al. (2020)	23
	Castañeda et al. (2010)*	1
	Chazen (2011)*	4
	W. Chen et al. (2014)	36
	Fallet et al. (2012)	10
	Hernández-Sánchez (2014)	7
	Ho et al. (2011)	20
	Ho et al. (2014)	161
	Hu et al. (2012)	7
	Jia et al. (2012)	32
	Kaiser et al. (2014)	29
	Kim et al. (2008)	238
	Kim et al. (2010)	158
	Leider et al. (2010)	46
	Lengger et al. (2014)	10
	Lü et al. (2014)	58
	Nieto-Moreno et al. (2013)*	2
	Park et al. (2014)	54
	Richey et al. (2011)*	1
	Seki et al. (2009)*	1
	Seki et al. (2014)	58
	Shevenell et al. (2011)*	8
	Smith et al. (2013)*	14
	Trommer et al. (2009)	20
	Verleye (2011)*	1
	Y. Wei et al. (2011)	11
	Wu et al. (2012)	1
	Zell et al. (2014)	16
	Zhou et al. (2014)	27
Data retrieved from Rattanasriampaipong et al. (2022)	Kim et al. (2015)	146
	Kim et al. (2016)	31
	Pan et al. (2016)	13
Data retrieved from Hagemann et al. (2023)	Jaeschke et al. (2017)	52
	Kaiser et al. (2015)	23
	Lamping et al. (2021)	66
Data retrieved from Varma et al. (2024)	Yang et al. (2018)	23
Reanalyzed samples by Varma et al. (2024)	Bale et al. (2013)	6
	Lo et al. (2018)	12
	Sinninghe Damsté (2016)	39
	Sinninghe Damsté et al. (2022)	13
	Total	2014

* Only reported TEX₈₆ values are available.