

## Assignment 10

### **1. Define the Bayesian interpretation of probability.**

Ans: Bayesian probability is the study of subjective probabilities or belief in an outcome, compared to the frequentist approach where probabilities are based purely on the past occurrence of the event. A Bayesian Network captures the joint probabilities of the events represented by the model.

### **2. Define probability of a union of two events with equation.**

Ans: The general probability addition rule for the union of two events states that  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ , where  $A \cap B$  is the intersection of the two sets.

### **3. What is joint probability? What is its formula?**

Ans: Probabilities are combined using multiplication, therefore the joint probability of independent events is calculated as the probability of event A multiplied by the probability of event B. This can be stated formally as follows:

**Joint Probability:**  $P(A \text{ and } B) = P(A) * P(B)$

### **4. What is chain rule of probability?**

Ans: The chain rule, or general product rule, calculates any component of the joint distribution of a set of random variables using only conditional probabilities. This probability theory is used as a foundation for backpropagation and in creating Bayesian networks.

### **5. What is conditional probability means? What is the formula of it?**

Ans: Conditional probability is defined as the likelihood of an event or outcome occurring, based on the occurrence of a previous event or outcome.

Conditional probability is calculated by multiplying the probability of the preceding event by the updated probability of the succeeding, or conditional, event.

Let's take a real-life example. Probability of selling a TV on a given normal day may be only 30%. But if we consider that given day is Diwali, then there are much more chances of selling a TV. The conditional Probability of selling a TV on a day given that Day is Diwali might be 70%.

### **6. What are continuous random variables?**

Ans: A continuous random variable  $X$  takes all values in a given interval of numbers. ▪ The probability distribution of a continuous random variable is shown by a density curve. ▪ The probability that  $X$  is between an interval of numbers is the area under the density curve between the interval endpoints.

### **7. What are Bernoulli distributions? What is the formula of it?**

Ans: A Bernoulli distribution is a discrete probability distribution for a Bernoulli trial — a random experiment that has only two outcomes (usually called a Success or a Failure). The expected value for a random variable,  $X$ .

For a **Bernoulli distribution** is:  $E[X] = p$ . For example, if  $p = 0.04$ , then  $E[X] = 0.4$

### **8. What is binomial distribution? What is the formula?**

Ans: The binomial is a type of distribution that has two possible outcomes (the prefix “bi” means two, or twice). For example, a coin toss has only two possible outcomes: heads or tails and taking a test could have two possible outcomes: pass or fail.

A Binomial Distribution shows either (S)uccess or (F)ailure.

### **9. What is Poisson distribution? What is the formula?**

Ans: A Poisson distribution is defined as a discrete frequency distribution that gives the probability of the number of independent events that occur in the fixed time.

In statistics, a Poisson distribution is a probability distribution that is used to show how many times an event is likely to occur over a specified period. ... The Poisson distribution is a discrete function, meaning that the variable can only take specific values in a (potentially infinite) list.

### **10. Define covariance ?**

Ans: Covariance is a measure of how much two random variables vary together. It's similar to variance, but where variance tells you how a single variable varies, covariance tells you how two variables vary together.

### **11. Define correlation ?**

Ans: Correlation explains how one or more variables are related to each other. These variables can be input data features which have been used to forecast our target variable. It's a bi-variate analysis measure which describes the association between different variables.

### **12. Define sampling with replacement. Give example.**

Ans: If you sample with replacement, you would choose one person's name, put that person's name back in the hat, and then choose another name. The possibilities for your two-name sample are: John, John. John, Jack.

**13. What is sampling without replacement? Give example.**

Ans: In sampling without replacement, each sample unit of the population has only one chance to be selected in the sample. For example, if one draws a simple random sample such that no unit occurs more than one time in the sample, the sample is drawn without replacement.

**14. What is hypothesis? Give example.**

Ans: A hypothesis (plural hypotheses) is a proposed explanation for a phenomenon. For a hypothesis to be a scientific hypothesis, the scientific method requires that one can test it. ... Even though the words "hypothesis" and "theory" are often used synonymously, a scientific hypothesis is not the same as a scientific theory.

Examples of hypothesis statements: If garlic repels fleas, then a dog that is given garlic every day will not get fleas. Bacterial growth may be affected by moisture levels in the air. If sugar causes cavities, then people who eat a lot of candy may be more prone to cavities.